

Bivariate Regression

Jerome Dumortier

Topics Covered

Motivation of regression analysis for policy

- Linking a single outcome of interest to one or more independent variables

Bivariate regression model components

Goal of a regression model

- Estimation of the expected mean of the dependent variable given particular values of the independent variable(s)

Bivariate regression

- One dependent variable y and one independent variable x

Find the best linear relationship between y and x assuming each observation y_i is a function of x_i plus a random error term ϵ_i

$$y_i = \beta_0 + \beta_1 \cdot x_i + \epsilon_i$$

We are looking for the conditional mean of Y given X , i.e., $E(Y|X)$

$$E(Y|X) = \beta_0 + \beta_1 \cdot X$$

Bivariate Regression Examples

Overview

Regression and Policy
Questions

Ordinary Least Square (OLS) Model

Estimation

Post Estimation

Linking one variable to one outcome of interest even without (!) causal claims

- Relationship between years of education and earnings (outcome) to quantify returns to schooling
- Effect of government spending per student on student test scores
- Link between gasoline prices and (individual) vehicle miles traveled for transportation and climate policy
- Relationship between property tax rates housing prices (outcome) to inform local public finance
- Change in crime rate due to police presence or change in policy presence due to crime rate

Careful with causal claims due to “chicken-or-egg” situations

Education earnings

- Omitted variables such as ability, motivation, family background
- Reverse causality: Expected earnings influence schooling choices (endogeneity)

Spending per student and test scores

- Policy targeting: Low-performing districts receive more funding
- Omitted variables: parental inputs, school quality
- Simultaneity problem

Gas prices and vehicle miles traveled

- Omitted variables: Transit access, income, etc.
- Measurement issues: Price variation often reflects demand conditions

Property tax rates and housing prices

- Simultaneity: Tax base affects tax rates
- Sorting: Households choose locations based on taxes and amenities leading to an equilibrium (and not a causal) relationship

Police presence and crime

- Reverse causality: Police are deployed because crime is high

Multivariate Regression Examples

Linking multiple variables to one outcome of interest

- Effect of a minimum wage increase on employment, while accounting for local economic conditions, industry composition, and demographic structure
- Healthcare utilization as a function of insurance coverage, income, and age
- Relationship between air pollution exposure and health outcomes, while taking into account weather conditions and population density, and industry composition
- Quantifying household energy consumption as a function of income, energy prices, and housing characteristics to inform energy efficiency policy

Ordinary Least Square (OLS) Components

Overview

Regression and Policy
Questions

Ordinary Least Square (OLS) Model

Estimation

Post Estimation

Example of the used car market

- Dependent variable: Price
- Independent variable: Miles

Every regression equation of the form $y = \beta_0 + \beta_1 \cdot x + \epsilon$ can be decomposed into four parts

- y : dependent variable
- x : independent variables
- β_0 : intercept
- β_1 : slope coefficient associated with the independent variable

The linear function does not tell us exactly what y will be for a given value of x but it does tell us the expected value of y , i.e., $E(y|x)$

Given a particular observation i , we have

$$y_i = \beta_0 + \beta_1 \cdot x_i + \epsilon_i$$

Given two values of β_0 and β_1 , i.e., $\hat{\beta}_0$ and $\hat{\beta}_1$, we can write

$$y_i = \hat{\beta}_0 + \hat{\beta}_1 \cdot x_i + \hat{\epsilon}_i$$

where ϵ_i represents the errors to obtain the observed y_i

- Theoretical difference between ϵ_i and $\hat{\epsilon}_i$ for which the former is the population parameter (always unknown) and the latter is the estimated error term given data

Overview

Regression and Policy
Questions

Ordinary Least
Square (OLS)
Model

Estimation

Post Estimation


Rearranging leads to the following

$$\hat{\epsilon}_i = y_i - \hat{\beta}_0 - \hat{\beta}_1 \cdot x_i$$

Minimization of the sum of the squared residuals:

$$\sum_{i=1}^N \hat{\epsilon}_i^2 = \sum_{i=1}^N \left(y_i - \hat{\beta}_0 - \hat{\beta}_1 \cdot x_i \right)^2$$

OLS Setup Graphical Representation



OLS Optimal Solution

Equations necessary to solve the bivariate regression model

- Mean of x :

$$\bar{x} = \frac{1}{N} \sum_{i=1}^N x_i$$

- Mean of y :

$$\bar{y} = \frac{1}{N} \sum_{i=1}^N y_i$$

- Slope coefficients:

$$\hat{\beta}_1 = \frac{\sum_{i=1}^N (x_i - \bar{x}) \cdot (y_i - \bar{y})}{\sum_{i=1}^N (x_i - \bar{x})^2}$$

- Intercept:

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \cdot \bar{x}$$

Example: Home Values and Square Footage

Overview


Regression and Policy
Questions

Ordinary Least Square (OLS) Model


Estimation

Post Estimation

(a) Home Prices and Square Footage



(b) Histogram of Residuals



Example: Used Cars

miles (x)	price (y)	$x_i - \bar{x}$	$y_i - \bar{y}$	$(x_i - \bar{x})(y_i - \bar{y})$	$(x_i - \bar{x})^2$
21	27	-15	6	-90	225
24	23	-12	2	-24	144
30	24	-6	3	-18	36
37	20	1	-1	-1	1
43	19	7	-2	-14	49
47	16	11	-5	-55	121
50	18	14	-3	-42	196

We have $\bar{x} = 36$ and $\bar{y} = 21$ as well as the following:

$$\sum_{i=1}^N (x_i - \bar{x})(y_i - \bar{y}) = -244 \quad \text{and} \quad \sum_{i=1}^N (x_i - \bar{x})^2 = 772$$

R^2 : Measuring the Strength of the Relationship I

Goodness of fit measure decomposes the variation of Y into two components, i.e., the (1) unexplained variation and the (2) explained variation: $R^2 \in [0, 1]$.

Unexplained or residual variation

$$RSS = \sum_{i=1}^N (y_i - \hat{y}_i)^2$$

Explained variation

$$ESS = \sum_{i=1}^N (\hat{y}_i - \bar{y})^2$$

Total variation:

$$TSS = \sum_{i=1}^N (y_i - \bar{y})^2$$

R^2 : Measuring the Strength of the Relationship II

Overview

Regression and Policy
Questions

Ordinary Least Square (OLS) Model

Estimation

Post Estimation

R^2 as the proportion of the total variation in Y explained by independent variables.
Note that since $TSS = RSS + ESS$:

$$1 = \frac{RSS}{TSS} + \frac{ESS}{TSS}$$

R^2 defined as

$$R^2 = \frac{ESS}{TSS} = 1 - \frac{RSS}{TSS}$$

Adjusted R^2 (for the case of multiple independent variables) where k is the number of variables:

$$\bar{R}^2 = 1 - (1 - R^2) \cdot \frac{n - 1}{n - k}$$

Overview

Regression and Policy
QuestionsOrdinary Least
Square (OLS)
Model

Estimation

Post Estimation

Standard error for the slope coefficient:

$$se(\hat{\beta}_1) = \frac{\sigma}{\sqrt{\sum_{i=1}^N (x_i - \bar{x})^2}}$$

Standard error for the intercept:

$$se(\hat{\beta}_0) = \sqrt{\frac{\sum_{i=1}^N x_i^2}{n \sum_{i=1}^N (x_i - \bar{x})^2}} \sigma$$

Estimate for the variance:

$$\hat{\sigma}^2 = \frac{\sum_{i=1}^N e_i^2}{n - 2}$$

Hypothesis Testing II

Overview

Regression and Policy
Questions

Ordinary Least Square (OLS) Model

Estimation

Post Estimation

Determination of statistical significance between variables:

- Assumption of normally distributed error terms
- t -statistic with $n - 2$ degrees of freedom

Specific hypothesis tests are $H_0: \beta_0 = 0$ and $H_0: \beta_1 = 0$. The test statistic for β_i can be written as

$$\frac{\hat{\beta}_i - \beta_i}{se_{\hat{\beta}_i}} \sim t_{n-2}$$

The hypothesis test is never conducted manually and every statistical software conducts and reports the results of the hypothesis test.

Numerical Example: Post Estimation

Overview

Regression and Policy
QuestionsOrdinary Least
Square (OLS)
Model

Estimation

Post Estimation

miles (x)	price (y)	x_i^2	\hat{y}	e_i	e_i^2	$(y_i - \bar{y})^2$
21	27	441	25.74	1.26	1.59	36
24	23	576	24.79	-1.79	3.21	4
30	24	900	22.90	1.10	1.22	9
37	20	1369	20.68	-0.68	0.47	1
43	19	1849	18.79	0.21	0.05	4
47	16	2209	17.52	-1.52	2.32	25
50	18	2500	16.58	1.42	2.03	9

Note that $\sum e_i^2 = 10.89$, $\sum x_i^2 = 10.89$, and $\sum (y_i - \bar{y})^2 = 88$

Numerical Example: R^2 and Standard ErrorsGoodness of fit R^2 :

$$R^2 = 1 - \frac{10.89}{88} = 0.876$$

For the standard errors, we have $\hat{\sigma} = \sqrt{10.89/5} = 1.476$ and thus,

$$se(\hat{\beta}_0) = \sqrt{\frac{9844}{7 \cdot 772}} \cdot 1.476 = 1.99$$

$$se(\hat{\beta}_1) = \frac{1.476}{\sqrt{772}} = 0.053$$

Adjusted R^2 :

$$\bar{R}^2 = 1 - (1 - 0.876) \cdot 6/5 = 0.8512$$

The manual calculations match the output from R.

Important assumptions for unbiasedness of the coefficient estimates:

- A1: Linear regression model, i.e., linear in terms of coefficients
- A2: Zero mean value of error terms ϵ , i.e., $E(\epsilon_i|x_i) = 0$
- A3: Homoscedasticity or equal variance of ϵ_i , i.e., $Var(\epsilon_i) = \sigma^2$
- A4: No autocorrelation between the error terms, i.e., $Cov(\epsilon_i, \epsilon_j) = 0$
- A5: No covariance between ϵ_i and x_i
- A6: Number of observations is greater than number of parameters to be estimated
- A7: No multicollinearity

A1: Linear Regression Model

Overview

Regression and Policy
QuestionsOrdinary Least
Square (OLS)
ModelEstimation
Post Estimation

Regression model that is linear in parameters:

$$y_i = \beta_0 + \beta_1 \cdot x_i + \epsilon$$

Note that the following models are also linear in parameters:

$$y_i = \beta_0 + \beta_1 \cdot x_i^2 + \epsilon$$

$$y_i = \beta_0 + \beta_1 \cdot x_i + \beta_2 \cdot x_i^2 + \epsilon$$

The following model is linear in parameters:

$$y = e^{\beta_0 + \beta_1 \cdot x_i}$$

The last model can be estimated by taking the natural logarithm of both sides.

A1: Linear Regression Model

Despite the fact that the regression model is linear, non-linear relationships can be measured:

- Relation between consumption and income might be non linear since a change in consumption due to extra income may decrease with income.
- Relationship between income and education can exhibit a non-linear form because a change in income due to more education may decrease with more education.

Example of Linear Regression Models



Overview

Regression and Policy
Questions

Ordinary Least Square (OLS) Model

Estimation

Post Estimation



A2: Zero Mean Value of Error Terms

Expected value of the error term is 0:

$$E(\epsilon_i | X_i) = 0 \quad \text{for all } i$$

See the histogram of residuals a couple of slides back.

A3: Homoscedasticity

The variance of the error terms is constant:

$$\text{Var}(\epsilon_i|x_i) = E(\epsilon_i^2|x_i) = \sigma^2$$

The assumption of constant variance is known as homoscedasticity. A violation of this assumption represents heteroscedasticity. Consider the following examples:

- Weekly consumption expenditures increases with income but the variability is higher with high-income families.

Consequences:

- No consequence on coefficient estimates
- Inflated standard errors

A3: Homoscedasticity vs. Heteroscedasticity

Overview


Regression and Policy
Questions

Ordinary Least Square (OLS) Model


Estimation

Post Estimation

Homoscedastic Data



Heteroscedastic Data



A4-A7: Other Assumptions

A4: No autocorrelation between the disturbance terms

$$E(\epsilon_i \epsilon_j) = 0 \quad \text{for all } i \neq j$$

A5: No covariance between ϵ_i and x_i

$$\text{Cov}(\epsilon_i, X_i) = 0$$

A6: Full rank:

- More observations than variables to be estimated
- Analogy: You cannot solve for three unknowns with two equations

A7: Multicollinearity:

- Near perfect linear relationships between independent variables should be avoided

Overview

Regression and Policy
QuestionsOrdinary Least
Square (OLS)
Model

Estimation

Post Estimation

```
##  
## Call:  
## lm(formula = price ~ miles, data = cars)  
##  
## Residuals:  
##      1      2      3      4      5      6      7  
## 1.2591 -1.7927  1.1036 -0.6839  0.2124 -1.5233  1.4249  
##  
## Coefficients:  
##              Estimate Std. Error t value Pr(>|t|)  
## (Intercept) 32.37824    1.99101  16.262  1.6e-05 ***  
## miles      -0.31606    0.05309  -5.953  0.00191 **  
## ---  
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1  
##  
## Residual standard error: 1.475 on 5 degrees of freedom  
## Multiple R-squared:  0.8764, Adjusted R-squared:  0.8516  
## F-statistic: 35.44 on 1 and 5 DF,  p-value: 0.001912
```

Used Car Market: R/RStudio Output

```
bhat = lm(price~miles,data=honda)
summary(bhat)
```

```
##
## Call:
## lm(formula = price ~ miles, data = honda)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -2453.6 -1055.3  -139.0   604.2  5389.5
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  2.205e+04  4.890e+02  45.095  < 2e-16 ***
## miles        -6.501e-02  1.251e-02  -5.198  1.54e-06 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 1455 on 79 degrees of freedom
## Multiple R-squared:  0.2549, Adjusted R-squared:  0.2454
## F-statistic: 27.02 on 1 and 79 DF,  p-value: 1.539e-06
```

CLRM: Assumptions

Required assumptions for unbiasedness coefficient estimates:

- A1: Linear in terms of coefficients, i.e., $y = \beta_0 + \beta_1 \cdot x$
- A2: Zero mean value of error terms ϵ , i.e., $E(\epsilon_i|x_i) = 0$
- A3: Homoscedasticity (equal variance) of ϵ_i , i.e., $Var(\epsilon_i) = \sigma^2$
- A4: No autocorrelation between the error terms, i.e., $Cov(\epsilon_i, \epsilon_j) = 0$
- A5: No covariance between ϵ_i and x_i
- A6: Number of observations is greater than number of parameters to be estimated
- A7: Full rank and absence of (perfect) multicollinearity

CLRM: Relaxing Assumptions

Relaxing the assumptions of the classical regression model requires regression diagnostics and/or different regression approaches

- Multicollinearity: Correlation between independent variables are correlated with each other?
 - Beds and bathrooms in a home value model
 - Multicollinearity occurs between two or more (!) independent variables
- Heteroscedasticity: Errors variance not constant
- Autocorrelation between error terms
- Inclusion of irrelevant or exclusion of relevant independent variables