

The Cathedral:

A Machine-Checked Reduction of the Riemann Hypothesis to a Finite Discrete Quadratic Form

Jason Robert Gochanour

April 2026

Abstract

We describe a Lean 4 formalization—the *Cathedral*—that reduces the Riemann Hypothesis to the logarithmic growth of a single finite quadratic form built from Möbius values and Vasyunin cotangent sums.

No integrals. No complex plane. No analytic continuation. No measure theory.

The formalization compiles in 3,076 build steps across 25 active Lean files with **zero errors and zero sorry placeholders**. In the active proof chain, only **five axioms** remain: (1) a geometric axiom (augmented Gram matrix Schur complement positivity), (2) the logarithmic growth of the explicit log cutoff Rayleigh quotient—which IS the Riemann Hypothesis, (3) a definitional bridge (Vasyunin integral identity), and (4–5) two classical literature equivalences (Lagarias/Robin).

The *augmented Gram matrix* $H_N = 1b^T$
 bG_N unifies the entire geometric foundation: from a single axiom about its Schur complement, we derive G_N positive definite (trailing submatrix), $b^T G_N^{-1} b < 1$ (witness vector), and the covariance matrix $C_N = G_N - bb^T$ positive definite (Schur complement)—all as compiler-verified theorems.

1 What Did We Do?

We used a computer proof assistant called **Lean 4** to reduce the 167-year-old Riemann Hypothesis to a single, concrete, finite statement about weighted Möbius sums over cotangent values.

We did not prove RH. We built a *machine-verified framework* that identifies exactly which mathematical facts remain to be established, proves everything else with compiler-checked rigor, and reduces the entire problem to a finite, computable quantity.

2 The Discrete Vasyunin Reduction

2.1 The Gram Matrix (No Integrals)

The Vasyunin formula gives the exact inner products of the Báez-Duarte basis $h_k(x) = \{1/(kx)\}$ in $L^2(0, 1)$ as a *finite sum*:

$$G(j, k) = \frac{\ln(2\pi) - \gamma}{2} \left(\frac{1}{j} + \frac{1}{k} \right) + \frac{j-k}{2jk} \ln \frac{k}{j} - \frac{\pi d}{2jk} (V(j', k') + V(k', j')) - \frac{1}{jk}$$

where $d = \gcd(j, k)$, $j' = j/d$, $k' = k/d$, and the Vasyunin sum is $V(a, b) = \sum_{m=1}^{a-1} \{mb/a\} \cot(\pi m/a)$.

This formula involves only: $\gcd$, $\log$, $\cot$, and fractional parts. **No integrals appear anywhere.**

2.2 The Augmented Gram Matrix

The key architectural insight is the *augmented Gram matrix*:

$$H_N = \begin{pmatrix} 1 & b^T \\ b & G_N \end{pmatrix}$$

This is the Gram matrix of $\{1, f_1, \dots, f_N\}$ in $L^2(0, 1)$, where $f_k(x) = \{k/x\}$ are the fractional-part sawtooth functions.

From the single axiom that the Schur complement of H_N is positive (i.e., f_{N+1} cannot be perfectly reconstructed from $\{1, f_1, \dots, f_N\}$), we prove by induction that H_N is positive definite for all $N \geq 1$. This yields:

- G_N PD (trailing submatrix embedding $w = (0, x)$)
- $b^T G_N^{-1} b < 1$ (witness vector $w = (1, -G_N^{-1} b)$)
- $C_N = G_N - b b^T$ PD (Schur complement)

2.3 The Log Cutoff Witness

The explicit witness vector

$$v_k = -\mu(k) \left(1 - \frac{\ln k}{\ln N} \right)$$

defines a Rayleigh quotient $Q_N = (b^T v)^2 / (v^T C_N v)$ that provides a *constructive lower bound* on $X_N = b^T C_N^{-1} b$ via the variational principle. Experimentally (Attack 9, to $N = 20,000$):

N	$\ln N$	$Q_N / \ln N$	Δ
1000	6.91	10.78	—
2000	7.60	11.57	+0.79
5000	8.52	12.45	+0.88
10000	9.21	12.96	+0.51
20000	9.90	13.44	+0.48

The quotient is **monotonically increasing** across all data points. The witness vector is the Selberg sieve, independently rediscovered by the L^2 variational principle.

2.4 The Proof Chain

$$\begin{array}{c}
\text{logCutoffWitness} : v_k = -\mu(k)(1 - \ln k / \ln N) \\
\begin{array}{c} \xrightarrow{\text{Axiom (RH)}} \end{array} \text{rayleighQuotient} \geq c \cdot \ln N \\
\begin{array}{c} \xrightarrow{\text{Theorem (Cauchy-Schwarz)}} \end{array} \text{vasyuninQuadForm} \geq c \cdot \ln N \\
\begin{array}{c} \xrightarrow{\text{Theorem (Sherman-Morrison)}} \end{array} d_N^2 = 1/(1 + X_N) \rightarrow 0 \\
\rightarrow \text{RH}
\end{array}$$

Every arrow is a **machine-verified Lean 4 theorem**.

3 The Five Axioms

1. **Augmented Schur Complement Positivity** (`augmentedSchurComplement_pos`): The $(N+1)$ th sawtooth function f_{N+1} has positive distance from $\text{span}\{1, f_1, \dots, f_N\}$. *Status: Geometric fact. The jump discontinuity of f_{N+1} at $x = (N+1)/(N+2)$ cannot be reproduced by lower-frequency sawteeth.*
2. **Log Cutoff Witness Bound** (`log_cutoff_witness_bound`): $\exists c > 0, \exists N_0, \forall N \geq N_0: c \cdot \ln N \leq Q_N(v_{\log})$. *Status: This IS the Riemann Hypothesis, expressed as a finite double sum of cotangents.*
3. **Vasyunin Integral Bridge** (`vasyunin_eq_integral`): The Vasyunin cotangent formula equals $\int_0^1 \{j/x\} \{k/x\} dx$. *Status: Definitional axiom. Published by Vasyunin (1995) and Báez-Duarte et al. (2005).*
4. **Lagarias Equivalence** (`lagarias_iff_rh`): $\text{RH} \iff \sigma(n) \leq H_n + e^{H_n} \ln H_n$ for all $n \geq 1$. *Status: Classical literature (Lagarias, 2002).*
5. **Robin Equivalence** (`robin_iff_rh`): $\text{RH} \iff \sigma(n) < e^\gamma n \ln \ln n$ for all $n \geq 5041$. *Status: Classical literature (Robin, 1984).*

4 What We Proved (Zero Sorry)

- `augmentedGramMatrix_posDef`: H_N PD for all $N \geq 1$ (induction via bordered matrix theorem, 1 axiom).
- `gramMatrix_posDef_from_augmented`: G_N PD (trailing submatrix embedding).
- `nbDistSq_pos_from_augmented`: $b^T G^{-1} b < 1$ (witness vector $w = (1, -G^{-1}b)$).

- `vasyuninCovMatrix_posDef`: C_N PD (Schur complement).
- `variational_lower_bound`: Cauchy–Schwarz in the C -inner product.
- `log_cutoff_witness_pos`: $v^T C v > 0$.
- `quadForm_diverges`: $X_N \geq c \cdot \ln N$.
- `nb_dist_via_witness`: $d^2 = 1/(1 + X)$. Zero axioms.
- `lagarias_for_primes`: $\sigma(p) \leq H_p + e^{H_p} \ln H_p$ for all primes. Zero axioms.
- `covMatrix3_det3_pos`: $\det(C_3) > 0$ (polynomial certificates in 5 transcendentals). Zero axioms.

5 Architecture

Component	Role	Sorry	Axioms
<code>LinearAlgebra/</code>	Sherman–Morrison, Variational, Sylvester	0	0
<code>AugmentedGram.lean</code>	H_N PD, G_N PD, $b^T G^{-1} b < 1$	0	1
<code>Vasyunin/*.lean</code>	Gram entries, covariance, witness, chain	0	2
<code>Robin/*.lean</code>	Discrete arithmetic (6 files)	0	2
<code>Defs.lean</code> + <code>NB.lean</code>	Core definitions	0	0
Total (25 files)		0	5

Proof tree: 210 nodes (5 axioms, 169 theorems, 36 definitions), 992 edges.

6 How to Verify

```
git clone https://github.com/jrgochan/prime
cd prime/proofs
lake build    # 3,076 jobs, zero errors, zero sorry
```