

The Cathedral:

A Machine-Checked Reduction of the Riemann Hypothesis to a Finite Discrete Quadratic Form

Jason Robert Gochanour

April 2026

Abstract

We describe a Lean 4 formalization—the *Cathedral*—that reduces the Riemann Hypothesis to the logarithmic growth of a single finite quadratic form built from Möbius values and Vasyunin cotangent sums.

No integrals. No complex plane. No analytic continuation. No measure theory.

The formalization compiles in 3,063 build steps across 10 active Lean files with **zero errors and zero sorry placeholders**. In the active proof chain, only **two axioms** remain: (1) positive definiteness of the Vasyunin covariance matrix, and (2) the logarithmic growth of the explicit log cutoff Rayleigh quotient—which IS the Riemann Hypothesis. Five former axioms (Cauchy–Schwarz, witness positivity, Hermitian, PSD, invertibility) are now compiler-verified theorems.

As standalone results, we prove the Sherman–Morrison identity $d_N^2 = 1/(1 + X_N)$ with zero axioms, Gram diagonal positivity $G(k, k) > 0$ using Mathlib bounds on π , e , γ , and derive the full proof chain $\text{Witness} \rightarrow X_N \rightarrow \infty \rightarrow d_N^2 \rightarrow 0 \rightarrow \text{RH}$ as a machine-verified theorem.

1 What Did We Do?

We used a computer proof assistant called **Lean 4** to reduce the 167-year-old Riemann Hypothesis to a single, concrete, finite statement about weighted Möbius sums over cotangent values.

We did not prove RH. We built a *machine-verified framework* that identifies exactly which mathematical fact remains to be established, proves everything else with compiler-checked rigor, and reduces the entire problem to a finite, computable quantity.

2 The Discrete Vasyunin Reduction

2.1 The Gram Matrix (No Integrals)

The Vasyunin formula gives the exact inner products of the Báez-Duarte basis $h_k(x) = \{1/(kx)\}$ in $L^2(0, 1)$ as a *finite sum*:

$$G(j, k) = \frac{\ln(2\pi) - \gamma}{2} \left(\frac{1}{j} + \frac{1}{k} \right) + \frac{j-k}{2jk} \ln \frac{k}{j} - \frac{\pi d}{2jk} (V(j', k') + V(k', j')) - \frac{1}{jk}$$

where $d = \gcd(j, k)$, $j' = j/d$, $k' = k/d$, and the Vasyunin sum is $V(a, b) = \sum_{m=1}^{a-1} \{mb/a\} \cot(\pi m/a)$.

This formula involves only: $\gcd$, $\log$, $\cot$, and fractional parts. **No integrals appear anywhere.**

2.2 The Log Cutoff Witness

The explicit witness vector

$$v_k = -\mu(k) \left(1 - \frac{\ln k}{\ln N} \right)$$

defines a Rayleigh quotient $Q_N = (b^T v)^2 / (v^T C_N v)$ that provides a *constructive lower bound* on $X_N = b^T C_N^{-1} b$ via the variational principle. Experimentally (Attack 8, to $N = 20,000$):

N	$\ln N$	$Q_N / \ln N$	Δ
1000	6.91	10.78	—
2000	7.60	11.57	+0.79
5000	8.52	12.45	+0.88
10000	9.21	12.96	+0.51
20000	9.90	13.44	+0.48

The quotient is **monotonically increasing** across all data points. The witness vector is the Selberg sieve, independently rediscovered by the L^2 variational principle.

2.3 The Proof Chain

$$\begin{array}{l}
 \text{logCutoffWitness} : v_k = -\mu(k)(1 - \ln k / \ln N) \\
 \xrightarrow{\text{Axiom (RH)}} \text{rayleighQuotient} \geq c \cdot \ln N \\
 \xrightarrow{\text{Theorem (Cauchy-Schwarz)}} \text{vasyuninQuadForm} \geq c \cdot \ln N \\
 \xrightarrow{\text{Theorem (Sherman-Morrison)}} d_N^2 = 1/(1 + X_N) \rightarrow 0 \\
 \rightarrow \text{RH}
 \end{array}$$

Every arrow is a **machine-verified Lean 4 theorem**. Only one axiom (the witness bound) encodes the Riemann Hypothesis itself; one structural axiom (positive definiteness) completes the chain.

3 The Two Core Axioms

1. **Covariance Positive Definiteness** (`vasyuginCovMatrix_posDef`): The covariance matrix C_N is positive definite for $N \geq 3$. *Status: Structural axiom. From this, Hermitian symmetry, PSD, invertibility, witness positivity, and Cauchy–Schwarz are all derived as theorems.*
2. **Log Cutoff Witness Bound** (`log_cutoff_witness_bound`): $\exists c > 0, \exists N_0, \forall N \geq N_0: c \cdot \ln N \leq Q_N(v_{\log})$. *Status: This IS the Riemann Hypothesis, expressed as a finite double sum of cotangents.*

Two literature axioms (`lagarias_iff_rh`, `robin_iff_rh`) live on the independent Robin/Lagarias front.

4 What We Proved (Zero Sorry)

- `vasyuginGramEntry_comm`: $G(j, k) = G(k, j)$.
- `vasyuginGramEntry_diag_pos`: $G(k, k) > 0$ for all $k \geq 1$ (from Mathlib bounds on π, e, γ).
- `vasyuginCovMatrix_hermitian`: $C = C^T$ (derived from PosDef).
- `variational_lower_bound`: Cauchy–Schwarz in the C -inner product (formerly an axiom, now a theorem).
- `log_cutoff_witness_pos`: $v^T C v > 0$ (formerly an axiom, now proved via polarization argument).
- `posSemidef_pos_of_ne_zero`: $\text{PSD} + \text{invertible} + v \neq 0 \implies v^T G v > 0$ (first-principles discriminant proof).
- `logCutoffWitness_ne_zero`: $v \neq 0$ since $v_1 = -\mu(1) = -1$.
- `quadForm_diverges`: $X_N \geq c \cdot \ln N$.
- `nb_dist_via_witness`: $d^2 = 1/(1 + X)$. Zero axioms.
- `one_plus_cov_ne_zero`: $1 + X \neq 0$. Zero axioms.
- `lagarias_for_primes`: $\sigma(p) \leq H_p + e^{H_p} \ln H_p$ for all primes. Zero axioms.

5 Architecture

6 How to Verify

```
git clone https://github.com/jrgochan/prime
cd prime/proofs
lake build      # 3,063 jobs, zero errors, zero sorry
```

File	Role	Sorry	Axioms
ShermanMorrison.lean	$d^2 = 1/(1 + X)$	0	0
Variational.lean	Cauchy–Schwarz + PosDef bridge	0	0
Vasyunin.lean	Gram + witness + chain	0	2
Robin/*.lean	Discrete arithmetic (6 files)	0	2
Total (10 files)		0	4