



University of Antwerp  
Faculty of Social Sciences

# Objective rating method: Entropy

Speech intelligibility estimation

Jose Rivera

April 16, 2022

# What are we going to talk about?

## 1 Preliminars

- Research question
- Research hypothesis production

## 2 Estimand and Process model

## 3 Synthetic data generation

## 4 References

# 1. Preliminars

Research question

# Research question

On two fronts:

1. Can comparative judgement (CJ) methods be used to assess speech intelligibility (SI)?,

To investigate this we need:

- an objective measure of SI

2. where CJ stands versus absolute holistic judgement (HJ) methods?,

In terms of:

- validity
- reliability
- statistical efficiency
- time efficiency

# Objective measure of SI

the **most objective** measure of SI (we know of) comes from a **transcription task**:

1. transcribing children's utterances (made by multiple judges),
2. align transcriptions at the utterance level,
3. calculate an entropy measure ( $H$ ), defined as

$$H = H(\mathbf{p}) = \frac{-\sum_{i=1}^n p_i \cdot \log_2(p_i)}{\log_2(N)}$$

4. characteristics of  $H$  [1, 2]
  - bounded in  $[0, 1]$  space,
  - utterances with more agreement are more intelligible, and therefore  $H \rightarrow 0$ ,
  - utterances with low agreement are less intelligible, and therefore  $H \rightarrow 1$ .

# 1. Preliminars

Research hypothesis production

# A typical scientific lab<sup>1</sup>

What is needed?

1. Quality of theory
2. Quality of data
3. Reliable procedures and code
4. Quality of data analysis
5. Documentation
6. Reporting

What we will deal with:

1. Quality of theory
2. Quality of data
3. Reliable procedures and code
4. Quality of data analysis
5. Documentation
6. Reporting

---

<sup>1</sup>McElreath [5], lecture 20 and McElreath [6], chapter 17

# Research hypothesis production<sup>2</sup>

## Well known challenges

- Insufficient data
- Wrong population
- Measurement error
- Selection bias
- Confounding

## Known challenges in our research;

- Insufficient data (possibly)
- Wrong population
- Measurement error
- Selection bias
- Confounding


---

<sup>2</sup>Hernán [4], lesson 4



# Research hypothesis schematics<sup>3</sup>

- Estimand and process model
- Synthetic data generation
- Statistical model design and testing
- Apply statistical model to data



<sup>3</sup>McElreath [6], lecture 20, Pearl [9]. Follow Fogarty et al. [3] on item (c).


# Why do we need to follow this?

Because the improvement of:

- A clear definition of the estimand and process model (assumptions).
- An improved the reliability of your procedures.
- As a documentation procedure.

leads to:


- A sound analysis, and sound results (even when we cannot answer our question).
- An improved planning to get data.



## 2. Estimand and Process model

# The theory behind our research

- $H_{ik}$  = (observed) entropy replicates
- $H_i$  = (latent) child's entropy
- $SI_i$  = (latent) child's SI score  
(inversely related to  $H_i^T$ )
- $A_i$  = child's "hearing" age
- $E_i$  = child's etiology of disease
- $HS_i$  = child's hearing status
- $PTA_i$  = child's pure tone average
- variables **assumed independent**,  
beyond the described relationships,




General causal diagram


$$\begin{aligned} P(\mathbf{U}) &= P(U_{ik}, U_i, U_A, U_E, U_{HS}, U_P) \\ &= P(U_{ik})P(U_i)P(U_A)P(U_E)P(U_{HS})P(U_P) \end{aligned}$$

# Interested in two effects

1. **total effects** model inherits:
  - children's characteristics that lead to the fitting of specific apparatus,
  - the (convenience of) sample selection  
(fixed with post-stratification)
2. to do the last, we stratify for all variables that explain variability, ergo, use a **direct effects** model
3. We have two levels: replicates (k), child's level (i), denoted by squares
4.  $U_{ik}$  = replicates measurement error  
 $U_i$  = within child SI variability



(b) total effects



(a) direct effects

# Causal and probabilistic model

$$H_{ik}^O \sim \text{BetapProp}(H_i^T, df_{ik})$$

$$H_i^T = \text{inv\_logit}(-SI_i)$$

$$SI_i \sim \text{Normal}(\mu_{SI}, \sigma_{U_i})$$

$$\begin{aligned} \mu_{SI} = & a_i + \alpha + \alpha_{HS[i]} + \alpha_{E[i]} \\ & + \beta_{A,HS[i]}(A_i - \bar{A}) + \beta_P PTA_i \end{aligned}$$

$$HS_i \sim \text{data}$$

$$A_i \sim \text{data}$$

$$E_i \sim \text{data}$$

$$PTA_i \sim \text{data}$$

$$U \sim \text{unobservable}$$

(a) general probabilistic model

$$H_{ik}^O \leftarrow f(H_i^T, U_{ik})$$

$$H_i^T \leftarrow f(SI_i)$$

$$SI_i \leftarrow f(HS_i, A_i, E_i, PTA_i, U_i)$$

$$HS_i \leftarrow f(U_{HS})$$

$$A_i \leftarrow f(U_A)$$

$$E_i \leftarrow f(U_E)$$

$$PTA_i \leftarrow f(U_P)$$

$$U \sim P(\mathbf{U})$$

(a) general structural model

### 3. Synthetic data generation


# Intervention

- Purpose: to keep a control on all the factors responsible for the outcome's variation (understand the system).
- It is modeled by modifying the structural model (and causal diagram).
- remember:  $\mathbf{V} = \{Z, X, Y\}$ ,  $\mathbf{U} = \{U_z, U_x, U_y\}$ , and  $\mathbf{F} = \{f_z, f_x, f_y\}$ .
- Intervention on  $X$  can be written in do-calculus<sup>a</sup> as:  $P(\mathbf{V} \mid \text{do}(X = x))$ .

<sup>a</sup>we are not delving into this (the usual suspects [7, 8, 10, 11])

$$M = \begin{cases} Z \leftarrow f_z(U_z) \\ X \leftarrow f_x(U_x) \\ Y \leftarrow f_y(X, Z, U_y) \\ U \sim P(\mathbf{U}) \end{cases}$$

(a) structural model



(b) causal diagram



## 4. References

## 4. References

- [1] Boonen, N., Kloots, H., Nurzia, P. and Gillis, S. [2021]. Spontaneous speech intelligibility: early cochlear implanted children versus their normally hearing peers at seven years of age, *Journal of Child Language* pp. 1–26.  
doi: <https://doi.org/10.1017/S0305000921000714>.
- [2] Faes, J., De Maeyer, S. and Gillis, S. [2021]. Speech intelligibility of children with an auditory brainstem implant: a triple-case study, pp. 1–50. (submitted).
- [3] Fogarty, L., Madeleine, A., Holding, T., Powell, A. and Kandler, A. [2022]. Ten simple rules for principled simulation modelling, *PLOS Computational Biology* 18(3): 1–8.  
doi: <https://doi.org/10.1371/journal.pcbi.1009917>.
- [4] Hernán, M. [2020]. Causal diagrams: Draw your assumptions before your conclusions.  
url: <https://www.edx.org/course/causal-diagrams-draw-your-assumptions-before-your>.
- [5] McElreath, R. [2020]. *Statistical Rethinking: A Bayesian Course with Examples in R and STAN*, Chapman and Hall/CRC.
- [6] McElreath, R. [2022]. *Statistical rethinking, 2022 course*.  
url: [https://github.com/rmcelreath/stat\\_rethinking\\_2022](https://github.com/rmcelreath/stat_rethinking_2022).

- [7] Pearl, J. [1988]. Probabilistic reasoning in intelligent systems: Networks of plausible inference, *The Journal of Philosophy* 88(8): 434–437.  
doi: <https://doi.org/10.2307/2026705>.  
url: <https://www.jstor.org/stable/2026705>.
- [8] Pearl, J. [2009]. *Causality: Models, Reasoning and Inference*, Cambridge University Press.
- [9] Pearl, J. [2019]. The seven tools of causal inference, with reflections on machine learning, *Communications of the ACM* 62(3): 54–60.  
doi: <https://doi.org/10.1177/0962280215586010>.
- [10] Pearl, J., Glymour, M. and Jewell, N. [2016]. *Causal Inference in Statistics: A Primer*, John Wiley Sons, Inc.
- [11] Pearl, J. and Mackenzie, D. [2018]. *The Book of Why: The New Science of Cause and Effect*, 1st edn, Basic Books, Inc.