

# Let's talk about Thurstone & Co.: An information-theoretical model for comparative judgments, and its statistical translation

Jose Manuel Rivera Espejo<sup>a,\*</sup>, Tine van Daal<sup>a</sup>, Sven De Maeyer<sup>a</sup>, Steven Gillis<sup>b</sup>

<sup>a</sup>*University of Antwerp, Training and education sciences,*

<sup>b</sup>*University of Antwerp, Linguistics,*

---

## Abstract

This study revisits Thurstone's law of comparative judgment (CJ) by addressing two central issues in the CJ literature. First, it critiques the widespread reliance on Thurstone's Case V assumptions and, by extension, the Bradley-Terry-Luce (BTL) model in the analysis of CJ data. Specifically, this study argues that while the assumptions of equal discriminial dispersions and zero correlation between stimuli simplify the trait measurement model, they may fail to capture the complexity of some traits or account for heterogeneous stimuli, potentially leading to unreliable and inaccurate trait estimates. Second, the study highlights the apparent disconnect between trait measurement and hypothesis testing procedures in CJ applications. It contends that while separating these procedures simplifies the analysis of CJ data, this practice can also undermine the resulting statistical inferences.

To address these issues, this study aims to extend Thurstone's general form using a systematic and integrated approach based on causal and Bayesian inference methods. This extension integrates Thurstone's core theoretical principles alongside key CJ assessment design features. It then translates these elements into a probabilistic statistical model for analyzing dichotomous CJ data, overcoming the rigid assumptions of Case V and the BTL model.

Finally, the study emphasizes the relevance of this extension for contemporary empirical CJ research, particularly stressing the need for bespoke CJ models tailored to the experiments and data assumptions. It also lays the foundation for broader applications, encouraging researchers across the social sciences to adopt more robust and interpretable methodologies.

*Keywords:* causal inference, directed acyclic graphs, structural causal models, bayesian statistical methods, thurstonian model, comparative judgement, probability, statistical modeling

---

## 1. Introduction

In *comparative judgment* (CJ) studies, judges assess a specific trait or attribute across different stimuli by performing pairwise comparisons (Thurstone, 1927b,a). Each comparison produces a dichotomous outcome, indicating which stimulus is perceived to have a higher trait level. For example, judges can compare pairs of written texts (the stimuli) to determine the relative writing quality each text exhibit (the trait) (Pollitt, 2012b; van Daal et al., 2016; Lesterhuis, 2018; Coertjens et al., 2017; Goossens and De Maeyer, 2018; Bouwer et al., 2023).

Numerous studies have documented the effectiveness of CJ in assessing traits and competencies over the past decade. These studies have highlighted three key strengths of the method: its reliability, its validity, and practical applicability. Research on reliability suggests that CJ requires a relatively modest number of pairwise comparisons (Verhavert et al., 2019; Cromptvoets et al., 2022) to generate trait scores that are as precise and consistent as those generated by other assessment methods (Coertjens et al., 2017; Goossens and De Maeyer, 2018; Bouwer et al., 2023). In addition, the evidence suggests that the reliability and time efficiency of CJ are comparable, if not superior, to those of other assessment methods when employing adaptive comparison algorithms (Pollitt, 2012b; Verhavert et al., 2022; Mikhailiuk et al., 2021). Meanwhile, research on validity indicates the capacity of CJ scores to represent accurately the traits under measurement (Whitehouse, 2012; van Daal et al., 2016; Lesterhuis, 2018; Bartholomew et al., 2018; Bouwer et al., 2023). Lastly, research on its practical applicability highlights CJ’s versatility across educational and non-educational contexts (Kimbell, 2012; Jones and Inglis, 2015; Bartholomew et al., 2018; Jones et al., 2019; Marshall et al., 2020; Bartholomew and Williams, 2020; Boonen et al., 2020).

Nevertheless, despite the increasing number of CJ studies, research in this domain remains unsystematic and fragmented, leaving several critical issues unresolved. This study identifies and discusses two prominent issues in the CJ literature that can undermine the reliability and validity of CJ’s trait estimates. First, it critiques the widespread reliance on Thurstone’s Case V assumptions (Thurstone, 1927a) and, by extension, the Bradley-Terry-Luce (BTL) model (Bradley and Terry, 1952; Luce, 1959) in the analysis of CJ data. Specifically, this study argues that while the assumptions of equal discriminial dispersions and zero correlation between stimuli simplify the trait measurement model, they may fail to capture the complexity of some traits or account for heterogeneous stimuli,

---

\*Corresponding author

Email addresses: JoseManuel.RiveraEspejo@uantwerpen.be (Jose Manuel Rivera Espejo),  
tine.vandaal@uantwerpen.be (Tine van Daal), sven.demaeyer@uantwerpen.be (Sven De Maeyer),  
steven.gillis@uantwerpen.be (Steven Gillis)

potentially leading to unreliable and inaccurate trait estimates. Second, the study highlights the disconnect between trait measurement and hypothesis testing procedures in CJ applications. In particular, it contends that while separating these procedures simplifies the analysis of CJ data, this practice can also undermine the resulting statistical inferences (McElreath, 2020; Kline, 2023; Hoyle, 2023).

To address these issues, this study aims to extend Thurstone’s general form through a systematic and integrated approach that combines causal and Bayesian inference methods. This extension integrates Thurstone’s core theoretical principles alongside key CJ assessment design features, such as the selection of judges, stimuli, and comparisons. In addition to potentially enhancing measurement reliability and validity, and improving statistical accuracy in hypothesis testing, this approach offers two key advantages. First, it clarifies the interactions among all actors and processes involved in CJ assessments. Second, it shifts the current comparative data analysis paradigm from passively accepting Case V and the BTL model assumptions to actively testing whether those assumptions fit the data under analysis.


The remainder of this study is organized into seven sections. Section 2 provides an overview of Thurstone’s theory. Section 3 examines the central issues identified in the CJ literature. Section 4 extends Thurstone’s general form to address these issues. The extension integrates core theoretical principles alongside key CJ assessment design features, such as the selection of judges, stimuli, and comparisons. Section 5 translates these theoretical and design elements into a probabilistic statistical model to analyze dichotomous pairwise comparison data. Section 6 reviews the findings, outline directions for future research, discusses the study’s limitations and details the challenges that applied researchers may encounter. Finally, Section 7 provides the concluding remarks.

## 2. Thurstone’s theory

In its *general form*, Thurstone’s theory addresses pairwise comparisons of a single judge who evaluates multiple stimuli (Thurstone, 1927b,a). The theory posits that two key factors determine the dichotomous outcome of these comparisons: the discriminial process of each stimulus and their discriminial difference. The *discriminial process* captures the psychological impact each stimulus exerts on the judge or, more simply, his perception of the stimulus trait. The theory assumes that the discriminial process for any given stimulus forms a Normal distribution along the trait continuum (Thurstone, 1927a). The mode (mean) of this distribution, known as the *modal discriminial process*, indicates the stimulus position on this continuum, while its dispersion, referred to as the *discriminial*

*dispersion*, reflects variability in the perceived trait of the stimulus.

These ideas become clearer with an example. Figure 1a illustrates the hypothetical discriminial processes for two written texts along a *quality* trait continuum. The figure shows that the modal discriminial process for Text B lies further along the continuum than that of Text A ( $T_B > T_A$ ), suggesting that Text B exhibits higher quality. The figure also shows that Text B has a broader distribution than Text A, indicating a greater variability in perception due to its larger discriminial dispersion ( $\sigma_B > \sigma_A$ ).



(a) Hypothetical discriminial processes for two written texts (b) Hypothetical discriminial difference of two written texts

Figure 1: Hypothetical discriminial processes and discriminant difference along a quality trait continuum for two written texts.

However, since the individual discriminial processes of the stimuli are not directly observable (Thurstone, 1927b), the theory introduces the *law of comparative judgment*. This law posits that in pairwise comparisons, a judge perceives the stimulus with a discriminial process positioned further along the trait continuum as possessing more of the trait (Bramley, 2008). This suggests that pairwise comparison outcomes depend on the relative distance between stimuli, not their absolute positions on the continuum. Indeed, the theory assumes that the difference between the underlying discriminial processes of the stimuli, referred to as the *discriminal difference*, determines the observed dichotomous outcome. Furthermore, the theory assumes that because the individual discriminial processes form a Normal distribution on the continuum, the discriminial difference will also conform to a Normal distribution (Andrich, 1978). In this distribution, the mode (mean) represents the average relative separation between the stimuli ( $T_B - T_A$ ), and its dispersion  $\sigma_{BA}$  indicates the variability of that separation.

Figure 1b illustrates the distribution of the discriminial difference for the two hypothetical texts. The figure indicates that the judge perceives Text B as having significantly higher quality than

Text A. Two key observations support this conclusion: the positive difference between their modal discriminial processes ( $T_B - T_A > 0$ ) and the probability area where the discriminial difference distinctly favors Text B over Text A, represented by the shaded gray area denoted as  $P(B > A)$ . As a result, the dichotomous outcome of this comparison is more likely to favor Text B over A.

### 3. Two Prominent Issues in Traditional CJ Practice

Thurstone noted from the outset that his general formulation, described in Section 2, led to a *trait scaling problem*. Specifically, the model required estimating more “unknown” parameters than the number of available pairwise comparisons (Thurstone, 1927a). For instance, in a CJ assessment with five texts, the general form would require estimating 20 parameters: five modal discriminial processes, five discriminial dispersions, and 10 correlations—one per comparison. However, a single judge could only provide  $\binom{5}{2} = 10$  unique comparisons, an “insufficient” data set to estimate the required parameters.

To address this issue and facilitate the practical implementation of the theory, Thurstone developed five cases derived from this general form, each progressively incorporating additional simplifying assumptions (Thurstone, 1927a). In Case I, Thurstone postulated that pairs of stimuli would maintain a constant correlation across all comparisons. In Case II, he allowed multiple judges to undertake comparisons instead of confining evaluations to a single judge. In Case III, he posited that there was no correlation between stimuli. In Case IV, he assumed that the stimuli exhibited similar dispersions. Finally, in Case V, he replaced this assumption with the condition that stimuli had equal discriminial dispersions. Table 1 summarizes the assumptions of the general form and the five cases. For a detailed discussion of these cases and their growing simplification, refer to Thurstone (1927a) and Bramley (2008).

Table 1: Thurstones cases and their assumptions

Assumption	General form	Thurstone’s					BTL model
		Case I	Case II	Case III	Case IV	Case V	
Discriminal process (distribution)	Normal	Normal	Normal	Normal	Normal	Normal	Logistic
Discriminal dispersion (between stimuli)	Different	Different	Different	Different	Similar	Equal	Equal
Correlation (between stimuli)	One per pair	Constant	Constant	Zero	Zero	Zero	Zero
How many judges compare?	Single	Single	Multiple	Multiple	Multiple	Multiple	Multiple

As the table suggest, Thurstone developed Case V with an emphasis on statistical simplicity, but

this simplicity comes at the expense of accurate and precise trait measurement and practical guidance for inference. Specifically, Thurstone cautioned that Case V use “should not be made without (an) experimental test” (Thurstone, 1927a, p. 270), as it imposes the most extensive set of simplifying assumptions (Bramley, 2008; Kelly et al., 2022) (see Table 1). Moreover, because Thurstone’s primary goal was to produce a “rather coarse scaling” of traits and “allocate the compared stimuli on this continuum” (Thurstone, 1927a, p. 269), his theory did not support formal statistical inference. Nevertheless, despite Case V’s limitations, CJ research has predominantly relied on it to measure various traits, raising significant concerns about the reliability and validity of such measurements in contexts where its assumptions may not hold (Kelly et al., 2022; Andrich, 1978). Furthermore, although the CJ tradition has attempted to address the gap in statistical inference by relying on the point estimates of traits—or their transformations—the statistical literature cautions against using these estimates as the sole basis for inference, as such practices introduce bias and reduce the precision of hypothesis tests (McElreath, 2020; Kline, 2023; Hoyle, 2023). The next sections discuss both issues in greater depth.


### *3.1. The Case V and the statistical analysis of CJ data*

Case V is the most widely used model in the CJ literature. This preference largely stems from the widespread adoption of the BTL model, which provides a simplified statistical representation of the case. The BTL model incorporates most of Case V’s assumptions, with one notable exception. While Case V assumes a Normal distribution for the stimuli’s discriminial processes, the BTL model uses the more mathematically tractable Logistic distribution (Andrich, 1978; Bramley, 2008) (see Table 1). However, this substitution has minimal impact on trait estimation or model interpretation because the scale of the discriminial process (i.e., the latent trait) is arbitrary up to a non-monotonic transformation (van der Linden, 2017a; McElreath, 2021). In other words, as long as the substitution (transformation) preserves the rank order of the data, the choice of distribution for the discriminial processes is inconsequential. The BTL model satisfies this condition because the Normal and Logistic distributions exhibit analogous statistical properties, differing only by a scaling factor of approximately 1.7 (van der Linden, 2017a).

However, Thurstone acknowledged that some assumptions of Case V are problematic when researchers assess complex traits or heterogeneous stimuli (Thurstone, 1927b). Thus, given that contemporary CJ applications often involve such traits and stimuli, two key assumptions of Case V, and by extension, the BTL model, may not always hold in theory or practice. These assumptions are the equal dispersion and zero correlation between stimuli.

### 3.1.1. The assumption of equal dispersions between stimuli

According to the theory, discrepancies in the discriminial dispersions of stimuli shape the distribution of the discriminial difference, directly influencing the outcome of pairwise comparisons. A thought experiment can help illustrate this idea. Suppose researchers observe the discriminial processes for two texts, A and B, assuming that the dispersion for Text A remains constant and that the two texts are uncorrelated ( $\rho = 0$ ). Figure 2a demonstrates that an increase in the uncertainty associated with the perception of Text B relative to Text A ( $\sigma_B - \sigma_A$ ), broadens the distribution of their discriminial difference. This broadening affects the probability area where the discriminial difference distinctly favors Text B over Text A, expressed as  $P(B > A)$ , ultimately influencing the comparison outcome. Additionally, the figure reveals that when the discriminial dispersions of the texts are equal, as in the BTL model ( $\sigma_B - \sigma_A = 0$ ), the discriminial difference distribution is more narrow than when the dispersions differ. As a result, the discriminial difference is more likely to favor Text B over Text A, as it is represented by the shaded gray area.



(a) Discriminal Difference distribution under varying discrepancies in stimuli dispersions (b) Discriminal Difference distribution under varying levels of correlation between stimuli

Figure 2: The effect of dispersion discrepancies and stimuli correlation on the distribution of the discriminial difference.

In experimental practice, however, the thought experiment occurs in reverse. Researchers first observe the comparison outcome and then use the BTL model to infer the discriminial difference between stimuli and their respective discriminial processes (Thurstone, 1927b). Consequently, the effectiveness of the outcome to reflect *true* differences between stimuli largely depends on the validity of the model's assumptions (Kohler et al., 2019), in this case, the assumption of equal dispersions. When the assumption accurately captures the complexity of the data, the BTL model estimates a discriminial difference distribution that accurately represents the *true* discriminial difference between the texts. This scenario is illustrated in Figure 2a, when the model's discriminial difference distribution aligns with the *true* discriminial difference distribution, represented by the thick continuous line

corresponding to  $\sigma_B - \sigma_A = 0$ . The accuracy of this discriminial difference then ensures reliable estimates for the texts’ discriminial processes.

Notably, while assuming equal dispersions simplifies the trait measurement model, evidence from the CJ literature suggests that this assumption may not hold for heterogeneous stimuli, such as handwritten texts or English compositions (Thurstone, 1927b; Andrich, 1978; Bramley, 2008; Kelly et al., 2022). The presence of the so-called misfit texts signals this limitation. *Misfit texts* are those that elicit more judgment discrepancies than others (Pollitt, 2004, 2012b,a; Goossens and De Maeyer, 2018), and these discrepancies may arise from larger discriminial dispersions caused by the stimulus’ heterogeneity or because the texts are genuine outliers—i.e., texts with distinctive characteristics that deviate markedly from the rest of the sample in which they occur (Grubbs, 1969). In either case, the BTL model’s assumptions prevent it from adequately accounting for or addressing these anomalies, leaving exclusion of such “problematic” texts as the primary remedy (Pollitt, 2012a,b).

Significant statistical and measurement issues can arise when the assumption of equal dispersions between stimuli does not hold. Specifically, the BTL model may overestimate the trait’s reliability, that is, the degree to which the outcome accurately reflects the *true* discriminial differences between stimuli. This overestimation, in turn, results in spurious conclusions about these differences (McElreath, 2020; Wu et al., 2022) and, by extension, about the underlying discriminial processes of stimuli. Figure 2a also illustrates this scenario when the model’s discriminial difference distribution aligns with the thick continuous line for  $\sigma_B - \sigma_A = 0$ , while the *true* discriminial difference follows any discontinuous line where  $\sigma_B - \sigma_A \neq 0$ . Furthermore, if researchers acknowledge that *misfit statistics* may represent texts with different dispersions or outlying observations, the common CJ practice of excluding stimuli based on these statistics may unintentionally discard valuable information (Miller, 2023), and introduce bias into the trait estimates (Zimmerman, 1994; McElreath, 2020). The direction and magnitude of these biases remain unpredictable, as they depend on which stimuli researchers exclude from the analysis.

### 3.1.2. The assumption of zero correlation between stimuli

The correlation between two stimuli  $\rho$  measures how much the judges’ perception of a specific trait in one stimulus depends on their perception of the same trait in other stimulus. Similar to the discriminial dispersions, this correlation shapes the distribution of the discriminial difference, directly impacting the outcomes of pairwise comparisons. Assuming that the discriminial dispersions for a couple of texts remain constant, Figure 2b shows that as the correlation between the two



texts increases, the distribution of their discriminial difference becomes narrower. This narrowing, in turn, affects the probability that the discriminial difference distinctly favors Text B over Text A—denoted as  $P(B > A)$ —and thus directly influences the comparison outcome. Furthermore, the figure shows that when two texts are independent or uncorrelated, as assumed in the BTL model ( $\rho = 0$ ), the distribution of their discriminial difference is less narrow than in scenarios where the texts are positively correlated. As a result, it becomes less likely for the comparison to favor Text B over Text A, as indicated by the larger shaded area.

Despite these notable differences in the distribution of the discriminial difference under various correlational assumptions, in practice, assessment designs often adopt the assumption of no correlation between stimuli based on Thurstone’s early theoretical justification (Thurstone, 1927a). He argued that stimuli could be treated as uncorrelated because judges’ biases—arising from two opposing and equally weighted effects occurring during the pairwise comparisons—would cancel each other out. This idea was later formalized by Andrich (1978), who provided a mathematical demonstration of this cancellation using the BTL model under the assumption of discriminial processes with additive biases. However, evidence from the CJ literature indicates that the assumption of zero correlation does not hold in practice in at least two cases: (1) when intricate aspects of multidimensional, complex traits or heterogeneous stimuli influence judges’ perceptions, (2) when additional hierarchical structures are relevant to the stimuli.

Regarding the first case, research on text quality assessments suggests that when judges evaluate complex, multidimensional traits or heterogeneous stimuli, they often rely on a variety of intricate stimulus characteristics to inform their judgments (van Daal et al., 2016; Lesterhuis et al., 2018; Chambers and Cunningham, 2022). Regardless of their relevance, these characteristics may not be equally weighted or consistently opposed across comparisons. As a result, they may exert a disproportionate influence on judges’ perceptions, generating biases that persist rather than cancel out. For example, this might occur when a judge assessing the argumentative quality of a text is disproportionately influenced by the clarity of the handwriting, thereby favoring neatly written texts even if their arguments are weaker. Moreover, because the discriminial process of stimuli becomes an observable outcome only through the judges’ perceptions, these biases could introduce dependencies between the stimuli (van der Linden, 2017b). While direct evidence for this exact scenario is limited, existing studies document the presence of judge’s biases in CJ contexts (Pollitt and Elliott, 2003; van Daal et al., 2016; Bartholomew et al., 2020), reinforcing the argument that the factors influencing pairwise comparisons do not always cancel each other out.

In the second case, the shared context or inherent connections introduced by additional hierarchical structures may create dependencies between stimuli—a statistical phenomenon known as clustering (Everitt and Skrondal, 2010). For instance, when the same individual produces multiple texts, those texts often share several features such as writing style or overall quality. Although some CJ studies acknowledge the presence of such hierarchical structures and account for them (e.g., Boonen et al., 2020), the treatment of this additional source of dependence in other research has often been insufficient. For instance, when CJ data include multiple samples of stimuli from the same individuals, researchers frequently rely on (averaged) point estimates of the BTL scores to conduct subsequent analyses and tests at the individual level (Bramley and Vitello, 2019; Bouwer et al., 2023; van Daal et al., 2017; Jones et al., 2019; Gijzen et al., 2021).

Thus, erroneously assuming zero correlation between stimuli can also lead to significant statistical and measurement issues. In particular, neglecting judges’ biases or relevant hierarchical structures can create dimensional mismatches in the model, leading to the over- or underestimation of trait reliability (Ackerman, 1989; Hoyle, 2023) and even introduce statistical biases (Wu et al., 2022). These inaccuracies can result in spurious conclusions about the discriminial differences (McElreath, 2020) and, by extension, the underlying discriminial processes of the stimuli. One such spurious conclusion could be the incorrect classification of stimuli (or judges) as *misfits*. Figure 2b illustrates how assuming zero correlation can undermine trait reliability: the discriminial difference distribution of the BTL scores follows the thick continuous line ( $\rho = 0$ ), while the *true* discriminial difference may correspond to any discontinuous line where  $\rho \neq 0$ .

Finally, similar to misfit stimuli, removing *misfit judges* risks discarding valuable information and introducing bias into trait estimates (Miller, 2023). The direction and magnitude of these biases remain unpredictable, as they depend on which judges researchers exclude from the analysis (Zimmerman, 1994; O’Hagan, 2018; McElreath, 2020). *Misfit* judges are those whose assessments deviate markedly from the shared consensus (Pollitt, 2012a,b; van Daal et al., 2016; Goossens and De Maeyer, 2018; Wu et al., 2022), often appearing as outliers under the BTL model (Wu et al., 2022).

### 3.2. The disconnect between trait measurement and hypothesis testing

Researchers in CJ studies typically use the BTL model to measure traits and position the compared stimuli along a latent continuum (Thurstone, 1927a). The CJ literature shows that research frequently relies on point estimates of these traits—typically the BTL scores or its transformations—to conduct

statistical inference or hypothesis testing. For example, researchers have used these scores to identify ‘misfit’ judges and stimuli (Pollitt, 2012b; van Daal et al., 2016; Goossens and De Maeyer, 2018), detect biases in judges’ ratings (Pollitt and Elliott, 2003; Pollitt, 2012b), calculate correlations with other assessment methods (Goossens and De Maeyer, 2018; Bouwer et al., 2023), or test hypotheses related to the underlying trait of interest (Casalicchio et al., 2015; Bramley and Vitello, 2019; Boonen et al., 2020; Bouwer et al., 2023; van Daal et al., 2017; Jones et al., 2019; Gijzen et al., 2021).

Nevertheless, while separating the trait measurement and hypothesis testing procedures simplifies the analysis of CJ data, the statistical literature cautions against relying solely on the point estimates of BTL scores to conduct statistical inference or hypothesis tests, as this practice can undermine the resulting statistical inferences. A key consideration is that BTL scores are parameter estimates that inherently carry uncertainty (measurement error). Ignoring this uncertainty can bias the analysis and reduce the precision of hypothesis tests. The direction and magnitude of such biases are often unpredictable. Results may be attenuated, exaggerated, or remain unaffected depending on the degree of uncertainty in the scores and the actual effects being tested (McElreath, 2020; Kline, 2023; Hoyle, 2023). Furthermore, the reduced precision in hypothesis tests diminishes their statistical power, increasing the likelihood of committing type-I or type-II errors (McElreath, 2020).

In aggregate, the heavy reliance on Thurstone’s Case V assumptions in the statistical analysis of comparative data can compromise the reliability of trait estimates. This overreliance may also undermine their validity (Perron and Gillespie, 2015), particularly when coupled with the disconnect between the trait measurement and hypothesis testing procedures. The structural approach to causal inference can address these issues by offering a systematic and integrated framework to extend Thurstone’s general form. This approach can also strengthen measurement reliability and validity while enhancing the statistical accuracy of hypothesis tests.

#### 4. Extending Thurstone’s general form

The *structural approach* to causal inference provides a formal framework for identifying causes and estimating their effects using data. The approach uses structural causal models (SCMs) and directed acyclic graphs (DAGs) (Pearl, 2009; Pearl et al., 2016; Gross et al., 2018; Neal, 2020) to formally and graphically represent the assumed causal structure of a system, such as the one found in CJ assessments. Essentially, SCMs and DAGs function as *conceptual models* on which identification analysis rests. *Identification analysis* determines whether an estimator can accurately compute an estimand based solely on its (causal) assumptions, regardless of random variability (Schuessler

and Selb, 2023). Here, *estimands* represent the specific quantities researchers aim to determine (i.e., a parameter) (Everitt and Skrondal, 2010). *Estimators* denote the methods or functions that transform data into an estimate (e.g., a statistical model), while *estimates* are the numerical values approximating the estimand (Neal, 2020; Everitt and Skrondal, 2010).

A motivating example that will appear throughout this study clarifies these concepts. In this example, researchers aim to answer the question: “To what extent do different teaching methods influence students’ ability to produce high-quality written texts?” To investigate this, a researcher designs a CJ assessment by randomly assigning students (individuals) to two groups, each receiving a different teaching method. Judges then compare pairs of students’ written texts (stimuli) to produce a dichotomous outcome reflecting the relative quality of each text (trait). Based on this setup, researchers can reformulate the research question as the estimand: “*On average*, is there a difference in the ability to produce high-quality written texts between the two groups of students?”.

Following standard CJ practices, researchers would typically use estimates from the BTL model—or its transformations—to approximate this estimand. However, as discussed in Section 3, Thurstone’s Case V and the BTL model exhibit several statistical and measurement limitations. These limitations hinder the model’s ability to identify various estimands relevant to CJ inquiries, including the one described in the example.

Fortunately, SCMs and DAGs support identification analysis through two key advantages<sup>1</sup>. First, regardless of complexity, they can represent various causal structures using only five fundamental building blocks (Neal, 2020; McElreath, 2024). This feature allows researchers to decompose complex structures into manageable components, facilitating their analysis. Second, they depict causal relationships in a non-parametric way. This flexibility enables feasible identification strategies without requiring specification of the types of variables, the functional forms relating them, or the parameters of those functional forms (Pearl et al., 2016).

Thus, this section addresses the issues identified in Section 3 by extending Thurstone’s general form using the structural approach to causal inference. Specifically, it combines Thurstone’s core

---

<sup>1</sup>In depth explanation of these topics is beyond the scope of this study, thus, readers seeking a more profound understanding can refer to introductory papers such as Pearl (2010), Rohrer (2018), Pearl (2019), and Cinelli et al. (2020), and introductory books like Pearl and Mackenzie (2018), Neal (2020), and McElreath (2020) are useful. For more advanced study, seminal papers such as Neyman (1923), Rubin (1974), Spirtes et al. (1991), and Sekhon (2009), along with books such as Pearl (2009), Morgan and Winship (2014), and Hernán and Robins (2025), are recommended.

theoretical principles outlined in Section 2 with key CJ assessment design features, such as the selection of judges, stimuli, and comparisons. Section 4.1 introduces the *conceptual-population model*, which incorporates these theoretical principles and assumes an idealized setting where researchers observe a *conceptual population* of comparative judgment data—that is, data representing all repeated judgments made by every available judge for each pair of stimuli produced by each pair of individuals in the population. Conversely, Section 4.2 presents the *sample-comparison model*, which integrates the assessment design features and reflects a more realistic setting where researchers access only a sample of judges, individuals, stimuli, and comparisons from the conceptual population.

#### 4.1. The conceptual-population model

In the conceptual-population model, the idealized scenario of a *conceptual population* of comparative data enables the integration of Thurstone’s theoretical principles and provides a foundation for proposing innovations aimed at addressing some of the issues discussed in Section 3.

##### 4.1.1. Integrating the first theoretical principles

Before incorporating the first theoretical principles of Thurstone’s theory, it is essential to further define SCMs. SCMs are formal mathematical models characterized by a set of *endogenous* variables  $V$ , a set of *exogenous* variables  $E$ , and a set of functions  $F$  (Pearl, 2009; Pearl et al., 2016; Cinelli et al., 2020). Endogenous variables are those whose causal mechanisms a researcher chooses to model (Neal, 2020). In contrast, exogenous variables represent *errors* or *disturbances* arising from omitted factors that the investigator chooses not to model explicitly (Pearl, 2009). Lastly, the functions, referred to as *structural equations*, express the endogenous variables as non-parametric functions of other endogenous and exogenous variables. These functions use the symbol ‘:=’ to denote the asymmetrical causal dependence between variables and the symbol ‘ $\perp$ ’ to represent *d-separation*, a concept akin to statistical (conditional) independence.

SCM 3a presents the first theoretical principles embedded in the conceptual-population model, which evaluates the impact of different teaching methods on students’ writing ability. This SCM outlines the relationship between the conceptual-population outcome ( $O_{iahbjk}^{cp}$ ) and several related variables. The subscripts  $i$  and  $h$  identify the students who authored the texts (i.e., the individuals). The indices  $a$  and  $b$  represent the texts under comparison (i.e., the stimuli). The index  $j$  indicates the judge conducting the comparison, while the index  $k$  accounts for assessment conditions where a judge compares the same pair of stimuli multiple times, i.e., a *repeated measures designs* (Lawson, 2015, p. 366-376). Thus, the indexing system supports comparisons between different texts written

by the same student ( $i = h$ ;  $a \neq b$ ) and between texts written by distinct students ( $i \neq h$ ; where  $a = b$  is permitted), each compared once or repeatedly by all judges ( $j = 1, \dots, n_J$ ;  $k = 1, \dots, n_K$ ; where  $n_J > 1$  and  $n_K \geq 1$ ). However, it excludes cases where a judge compares a student’s text to itself, whether once or multiple times ( $i = h$ ;  $a = b$ ;  $j = 1, \dots, n_J$ ;  $k = 1, \dots, n_K$ ; where  $n_J > 1$  and  $n_K \geq 1$ ), as such comparison lacks practical relevance within the CJ framework. Here,  $n_J$  indicates the total number of judges, and  $n_K$  denotes the number of repeated judgments each judge performs.




Figure 3: Conceptual-population model, scalar form.

In line with Thurstone’s theory, SCM 3a depicts the texts’ discriminative processes ( $T_{ia}, T_{hb}$ ) and their discriminative difference ( $D_{iahbjk}$ ) (see Section 2). Additionally, the SCM incorporates a key CJ design feature: the judges’ biases ( $B_{kj}$ ). This extension builds on the arguments presented in Section 3.1.2, contending that the discriminative difference becomes an observable outcome only through judges’ perceptions. Given that such perceptions may be imperfect—and that each judge may carry some degree of bias (see Pollitt and Elliott, 2003; van Daal et al., 2016)—it is reasonable that judges’ perceptions (bias) should be treated as an integral component of the CJ system from the outset, as this leads to a more accurate representation of the data-generating process underlying the pairwise comparisons. This model defines the preliminary set of endogenous variables,  $V = \{O_{iahbjk}, D_{iahbjk}, T_{ia}, T_{hb}, B_{kj}\}$ , and the preliminary set of structural equations,  $F = f_O, f_D$ , which capture the non-parametric dependencies among these variables.

Notably, every SCM has an associated DAG (Pearl et al., 2016; Cinelli et al., 2020). A DAG is a *graph* consisting of nodes connected by edges, where nodes represent random variables. The term *directed* indicates that edges or arrows extend from one node to another, indicating the direction of causal influence. The absence of an edge implies no direct relationship between the nodes. The term *acyclic* means that the causal influences do not form loops, ensuring the influences do not cycle back on themselves (McElreath, 2020). DAGs conventionally depict observed variables as solid black circles and unobserved (latent) variables as open circles (Morgan and Winship, 2014). Although

DAGs conventionally omit exogenous variables for simplicity, the DAGs presented in this section includes exogenous variables to improve clarity and reveal potential issues related to conditioning and confounding (Cinelli et al., 2020).

Figure 3b displays the DAG corresponding to SCM 3a, illustrating the expected causal relationships outlined in Thurstone’s theory. The graph shows that the discriminial processes of the texts ( $T_{ia}, T_{hb}$ ) influence their discriminial difference ( $D_{iahbjk}$ ), which in turn determines the outcome ( $O_{iahbjk}^{cp}$ ). It also highlights the influence of judges’ biases ( $B_{kj}$ ) on the discriminial difference. Additionally, the DAG differentiates between observed endogenous variables, such as the outcome (solid black circle), and latent endogenous variables, including the texts’ discriminial processes, their discriminial difference, and the judges’ biases (open circles).

#### 4.1.2. The conceptual-population data structure

Although specifying a data structure is not mandatory when using SCMs and DAGs, defining one improves clarity and facilitates the description of the system. Thus, to re-express the scalar form of the CJ system shown in Figure 3 into an equivalent vectorized form, we first define the vectors  $I$  and  $J$ , along with the matrices  $IA$  and  $JK$ , as in Equation (1). Here, each element of  $I$  represents a unique individual  $i$  or  $h$ , where  $n_I$  denotes the total number of individuals. Similarly, each element of  $J$  corresponds to a unique judge  $j$ , with  $n_J$  indicating the total number of judges. Moreover, each row of  $IA$  represents a unique pairing of individuals  $i, h$  with stimuli  $a, b$ . As a result, the matrix  $IA$  contains  $n_I \cdot n_A$  rows and 2 columns, where  $n_A$  specifies the number of stimuli available per individual. Likewise, each row of  $JK$  associates a judge  $j$  with a (repeated) judgment index  $k$ . Consequently, the matrix  $JK$  has  $n_J \cdot n_K$  rows and 2 columns, where  $n_K$  indicates the number of repeated judgments each judge makes.

Additionally, we construct the matrix  $R$  to map each row of the  $IA$  matrix with a corresponding row from the  $JK$  matrix. This matrix has  $n$  rows and 6 columns, where  $n = \binom{n_I \cdot n_A}{2} \cdot n_J \cdot n_K$ . Here, the term  $\binom{n_I \cdot n_A}{2}$  represents the binomial coefficient, which quantifies the total number of unique comparisons possible between every pair of stimuli generated by each pair of individuals in the population. Thus, we define the matrix as in Equation (1).

$$\begin{aligned}
I = \begin{bmatrix} 1 \\ \vdots \\ i \\ \vdots \\ h \\ \vdots \\ n_I \end{bmatrix}; J = \begin{bmatrix} 1 \\ \vdots \\ j \\ \vdots \\ n_J \end{bmatrix}; IA = \begin{bmatrix} 1 & 1 \\ \vdots & \vdots \\ 1 & n_A \\ \vdots & \vdots \\ i & a \\ \vdots & \vdots \\ h & b \\ \vdots & \vdots \\ n_I & 1 \\ \vdots & \vdots \\ n_I & n_A \end{bmatrix}; JK = \begin{bmatrix} 1 & 1 \\ \vdots & \vdots \\ 1 & n_K \\ \vdots & \vdots \\ j & k \\ \vdots & \vdots \\ n_J & 1 \\ \vdots & \vdots \\ n_J & n_K \end{bmatrix}; R = \begin{bmatrix} 1 & 1 & 1 & 2 & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & 1 & 2 & 1 & n_K \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ i & a & h & b & j & k \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ n_I & n_A - 1 & n_I & n_A & n_J & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ n_I & n_A - 1 & n_I & n_A & n_J & n_K \end{bmatrix}
\end{aligned} \tag{1}$$

It is easier to visualize the structure of the previously defined vectors and matrices by considering an example. Assuming  $n_I = 5$ ,  $n_A = 2$ ,  $n_J = 3$ , and  $n_K = 3$ , the vectors and matrices described in Equation (1) take the form as in Equation (2).

$$\begin{aligned}
I = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \end{bmatrix}; J = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}; IA = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 2 & 1 \\ 2 & 2 \\ 3 & 1 \\ 3 & 2 \\ 4 & 1 \\ 4 & 2 \\ 5 & 1 \\ 5 & 2 \end{bmatrix}; JK = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \\ 2 & 1 \\ 2 & 2 \\ 2 & 3 \\ 3 & 1 \\ 3 & 2 \\ 3 & 3 \end{bmatrix}; R = \begin{bmatrix} 1 & 1 & 1 & 2 & 1 & 1 \\ 1 & 1 & 1 & 2 & 1 & 2 \\ 1 & 1 & 1 & 2 & 1 & 3 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 1 & 5 & 2 & 1 & 1 \\ 1 & 1 & 5 & 2 & 1 & 2 \\ 1 & 1 & 5 & 2 & 1 & 3 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 4 & 2 & 5 & 2 & 3 & 1 \\ 4 & 2 & 5 & 2 & 3 & 2 \\ 4 & 2 & 5 & 2 & 3 & 3 \\ 5 & 1 & 5 & 2 & 3 & 1 \\ 5 & 1 & 5 & 2 & 3 & 2 \\ 5 & 1 & 5 & 2 & 3 & 3 \end{bmatrix}
\end{aligned} \tag{2}$$

Now, using Equation (1), we can re-express SCM 3a and DAG 3b in an equivalent vectorized form, as shown in Figure 4. In this depiction, the outcome  $O_R^{cp}$ , the texts' discriminial difference  $D_R$ , their



discriminal processes  $T_{IA}$ , and the judges' biases  $B_{JK}$  are represented as vectors rather than scalar values. These vectors capture all the observations from the conceptual population. Specifically,  $O_R^{cp}$  and  $D_R$  are observed and latent vectors of length  $n$ , respectively. Moreover,  $T_{IA}$  and  $B_{JK}$  are latent vectors of lengths  $n_I \cdot n_A$  and  $n_J \cdot n_K$ , respectively.




Figure 4: Conceptual-population model, initial vectorized form.

#### 4.1.3. Integrating hierarchical structural components

Building on the principles of Structural Equation Modeling (SEM) (Hoyle, 2023) and Item Response Theory (IRT) (Fox, 2010; van der Linden, 2017a), the conceptual-population model integrates two *hierarchical structural components* to examine how different *relevant*<sup>2</sup> variables—whether observed or latent—affect the primary latent variable of interest (Everitt and Skrondal, 2010). This hierarchical design enables researchers to formulate and test hypotheses that account for both the nested structure of stimuli and the uncertainties inherent in trait estimation (see Section 3.1.2 and Section 3.2 for a discussion of these considerations).

The top branch of DAG 5b illustrates the first component, where *relevant*<sup>3</sup> student-related variables  $X_I$ , such as teaching method, and students' idiosyncratic errors  $e_I$  causally influence the latent variable representing students' writing-quality trait  $T_I$ . The error term  $e_I$  captures variations in students' traits unexplained by  $X_I$ . Here,  $X_I$  is an observed matrix with  $n_I$  rows and  $q_I$  independent columns (variables), and both  $e_I$  and  $T_I$  are latent vectors of length  $n_I$ . Additionally,

<sup>2</sup>*Relevant variables* are those that satisfy the *backdoor criterion* (Neal, 2020, pp 37), that is, they belong to a *sufficient adjustment set* (Pearl, 2009; Pearl et al., 2016; Morgan and Winship, 2014). A *sufficient set* (potentially empty) blocks all non-causal paths between a predictor and an outcome without opening new ones (Pearl, 2009).

Refer also to footnote 1.

<sup>3</sup>refer to footnote 2.

this branch shows how  $T_I$ , along with *relevant*<sup>4</sup> text-related variables  $X_{IA}$  (e.g., text length), and texts' idiosyncratic errors  $e_{IA}$  causally influence the texts' written-quality trait  $T_{IA}$ , the first primary latent variable of interest. The error term  $e_{IA}$  captures variations in the texts' traits that remain unexplained by  $T_I$  or  $X_{IA}$ . Here,  $X_{IA}$  is an observed matrix with dimensions  $n_I \cdot n_A$  rows and  $q_{IA}$  independent columns (variables), while  $e_{IA}$  and  $T_{IA}$  are latent matrices with  $n_I$  rows and  $n_A$  columns.

Similarly, the bottom branch of DAG 5b depicts the second component, where *relevant*<sup>5</sup> judge-related variables  $Z_J$ , such as judgment expertise, and judges' idiosyncratic errors  $e_J$  causally influence the latent variable representing judges' bias  $B_J$ . The error  $e_J$  captures variations in judges' bias unexplained by  $Z_J$ . Here,  $Z_J$  is an observed matrix with  $n_J$  rows and  $q_J$  independent columns (variables), and both  $e_J$  and  $B_J$  are latent vectors of length  $n_J$ . Furthermore, the branch shows how  $B_J$ , along with *relevant*<sup>6</sup> judgment-related variables  $Z_{JK}$  (e.g., the number of judgments a judge makes), and judgments' idiosyncratic errors  $e_{JK}$  causally influence the judges' biases associated with each text  $B_{JK}$ , the second primary latent variable of interest. The error  $e_{JK}$  captures variations in judgments unexplained by  $B_J$  or  $Z_{JK}$ . Here,  $Z_{JK}$  is an observed matrix with dimension  $n_J \cdot n_K$  rows and  $q_{JK}$  independent columns (variables), while  $e_{JK}$  and  $B_{JK}$  are latent matrices with  $n_J$  rows and  $n_K$  columns.

Notably, all variables and functions shown in SCM 5a and DAG 5b are part of the set of endogenous variables  $V$ , structural equations  $F$ , and exogenous variables  $E$  for the conceptual-population model. Additionally, the figures demonstrate that all exogenous variables are independent of one another, as indicated by the relationships  $e_{IA} \perp \{e_I, e_{JK}, e_J\}$ ,  $e_I \perp \{e_{JK}, e_J\}$  and  $e_{JK} \perp e_J$  and the absence of connecting arrows.

Overall, the conceptual-population model extends Thurstone's general form by introducing key innovations to address the limitations discussed in Section 3.1.2 and Section 3.2. These enhancements include accounting for judges' biases and integrating hierarchical structural components. Nevertheless, despite its promise of enhancing measurement accuracy and precision, the model still depends on the unrealistic assumption that researchers have access to data from the *conceptual population*. Since researchers rarely meet this assumption in practice, they must consider a more realistic scenario.

---

<sup>4</sup>refer to footnote 2.

<sup>5</sup>refer to footnote 2.

<sup>6</sup>refer to footnote 2.

$$O_R^{cp} := f_O(D_R)$$

$$D_R := f_D(T_{IA}, B_{JK})$$

$$T_{IA} := f_T(T_I, X_{IA}, e_{IA})$$

$$T_I := f_T(X_I, e_I)$$

$$B_{JK} := f_B(B_J, Z_{JK}, e_{JK})$$


$$B_J := f_B(Z_J, e_J)$$

$$e_I \perp \{e_J, e_{IA}, e_{JK}\}$$

$$e_J \perp \{e_{IA}, e_{JK}\}$$

$$e_{IA} \perp e_{JK}$$

(a) SCM



(b) DAG

Figure 5: Conceptual-population model, final vectorized form.

#### 4.2. The sample-comparison model

The sample-comparison model presents a more realistic scenario than the conceptual-population model. First, it explicitly assumes researchers work with a data sample consisting of a limited

number of repeated judgments ( $n_K^s$ ) from a sample of judges ( $n_J^s$ ) and a specific number of texts ( $n_A^s$ ) from a sample of students ( $n_I^s$ ), all drawn from the conceptual population (Section 4.2.1). Second, the model assumes that judges do not perform *all repeated judgments* within the data sample (Section 4.2.2). Instead, they conduct a sufficient number of stimuli comparisons,  $n_C$ , to ensure an accurate estimation of the proportion  $P(B > A)$ , as proposed by [Thurstone \(1927a\)](#).

#### 4.2.1. The sample mechanism

To incorporate the sampling mechanism and facilitate the interpretation of the sample-comparison model, we first define the *data sampling process* using the binary vector variables  $S_I$ ,  $S_J$ ,  $S_{IA}$ , and  $S_{JK}$  as follows:

$$S_I = \begin{bmatrix} i_{(1)} \\ \vdots \\ i_{(i)} \\ \vdots \\ i_{(h)} \\ \vdots \\ i_{(nI)} \end{bmatrix} ; S_J = \begin{bmatrix} j_{(1)} \\ \vdots \\ j_{(j)} \\ \vdots \\ j_{(nJ)} \end{bmatrix} ; S_{IA} = \begin{bmatrix} ia_{(1,1)} \\ \vdots \\ ia_{(1,n_A)} \\ \vdots \\ ia_{(i,a)} \\ \vdots \\ ia_{(h,b)} \\ \vdots \\ ia_{(nI,1)} \\ \vdots \\ ia_{(nI,n_A)} \end{bmatrix} ; S_{JK} = \begin{bmatrix} jk_{(1,1)} \\ \vdots \\ jk_{(1,n_K)} \\ \vdots \\ jk_{(j,k)} \\ \vdots \\ jk_{(nJ,1)} \\ \vdots \\ jk_{(nJ,n_K)} \end{bmatrix} \quad (3)$$

Where each element of  $S_I$  is a binary value indicating the presence or absence of corresponding elements in the vector  $I$ , as in Equation (4). We apply the same logic to  $S_J$  using vector  $J$  (not shown). Thus, the vectors  $S_I$  and  $S_J$  contains  $n_I$  and  $n_J$  elements, respectively.

$$i_{(i)} = \begin{cases} 1 & \text{if data element } i \text{ from } I \text{ is sampled} \\ 0 & \text{if data element } i \text{ from } I \text{ is missing} \end{cases} \quad (4)$$

Similarly, each element of  $S_{IA}$  is a binary value indicating the presence or absence of data rows in the matrices  $IA$ , as defined in Equation (5). We apply the same logic to  $S_{JK}$  using the matrix  $JK$  (not shown). Thus, the vectors  $S_{IA}$  and  $S_{JK}$  contains  $n_I \cdot n_A$  and  $n_J \cdot n_K$  elements, respectively.

$$ia_{(i,a)} = \begin{cases} 1 & \text{if data elements } i, a \text{ from } IA \text{ are sampled} \\ 0 & \text{if data elements } i, a \text{ from } IA \text{ are missing} \end{cases} \quad (5)$$

We can illustrate the structure of these vectors more clearly with an example. Suppose researchers exclude the second student, the second text from each student, and the third judge from the setup shown in Equation (2). Given  $n_I = 5$ ,  $n_A = 2$ ,  $n_J = 3$ , and  $n_K = 3$ , the resulting vectors would have the following structure:

$$S_I = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 1 \\ 1 \end{bmatrix} ; S_J = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} ; S_{IA} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 1 \\ 0 \\ 1 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} ; S_{JK} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \quad (6)$$

Notably, Equation (6) shows that missing observations in the vectors  $S_I$  and  $S_J$ —which represent unsampled students and judges—directly determine which observations are missing in  $S_{IA}$  and  $S_{JK}$ . In other words, researchers can only observe texts and judgments from students and judges initially included in the sample. The equation also shows that the sum of observed elements in  $S_I$  equals the number of sampled students ( $n_I^s$ ) and that a similar sum in vector  $S_J$  equals the sampled judges ( $n_J^s$ ). Conversely, the sum of observed elements in  $S_{IA}$  represents the total sampled texts across all sampled students ( $n_I^s \cdot n_A^s$ ), while a similar sum in vector  $S_{JK}$  represents the total sampled repeated judgments across all sampled judges ( $n_J^s \cdot n_K^s$ ). Notice that in this example, because the design systematically excludes every third repeated judgment, researchers can also express  $S_{JK}$  using  $n_K = n_K^s = 2$ .

Finally, we define the *sample mechanism*  $S$  in Equation (8), which maps each element of  $S_{IA}$  to every element of  $S_{JK}$ . Each element  $s_{(i,a,h,b,j,k)}$  is a binary value indicating the presence or absence of data rows in the matrix  $R$  resulting from the sample mechanism, as in Equation (7). Thus, the vector contains  $n$  elements, matching the number of rows in  $R$ , and the sum of its elements represents the total data sample:  $n^s = \binom{n_I^s \cdot n_A^s}{2} \cdot n_J^s \cdot n_K^s$ . Here, the term  $\binom{n_I^s \cdot n_A^s}{2}$  represents the binomial coefficient, which quantifies the total number of unique comparisons possible between every pair of sampled stimuli generated by each pair of sampled individuals.

$$s_{(i,a,h,b,j,k)} = \begin{cases} 1 & \text{if data elements } i, a, h, b, j, k \text{ from } R \text{ are sampled} \\ 0 & \text{if data elements } h, i, a, b, j, k \text{ from } R \text{ are missing} \end{cases} \quad (7)$$

$$S = \begin{bmatrix} s_{(1,1,1,2,1,1)} \\ \vdots \\ s_{(1,1,1,2,1,n_K)} \\ \vdots \\ s_{(i,a,h,b,j,k)} \\ \vdots \\ s_{(n_I,n_A-1,n_I,n_A,n_J,1)} \\ \vdots \\ s_{(n_I,n_A-1,n_I,n_A,n_J,1)} \end{bmatrix} \quad (8)$$

With the definition of  $S$ , we incorporate the sample mechanism into the conceptual-population model. Following the convention of [McElreath \(2020\)](#) and [Deffner et al. \(2022\)](#), DAG 6b represents the conceptual-population outcome  $O_R^{cp}$  as unobserved, emphasizing that researchers cannot directly access this outcome due to the sampling mechanism. The DAG also depicts the *sample design* vector  $S$  as a causal factor influencing the sample-comparison outcome  $O_R^{sc}$ . A square encloses  $S$ , indicating that it is a conditioned variable. In this context, *conditioning* means that researchers restrict their focus to the elements of  $O_R^{cp}$  that satisfy  $s_{(i,a,h,b,j,k)} = 1$  ([Neal, 2020](#); [McElreath, 2020](#)). In essence,  $S$  is a vector that selects *all repeated judgments made by a subset of judges for a subset of stimuli produced by the sampled individuals*.

Notably, the DAG shows that  $S$  is independent of all other variables in the model. This implies that DAG 6b applies exclusively to Simple Random Sampling (SRSg) designs. In these designs, each repeated judgment, judge, stimulus, and individual has the same probability of being included in the sample as any other observation within their respective groups ([Lawson, 2015](#)).

However, due to concerns about the practical feasibility of the comparison task ([Boonen et al., 2020](#)), CJ assessments rarely implement an exhaustive pairings of sampled judges, stimuli, and individuals. Thus, a realistic scenario must account for the fact that judges typically compare only a subset of stimuli authored by a sample of individuals.

$$O_R := f_C(O_R^{sc}, C)$$

$$O_R^{sc} := f_S(O_R^{cp}, S)$$

$$O_R^{cp} := f_O(D_R)$$

$$D_R := f_D(T_{IA}, B_{JK})$$

$$T_{IA} := f_T(T_I, X_{IA}, e_{IA})$$

$$T_I := f_T(X_I, e_I)$$

$$B_{JK} := f_B(B_J, Z_{JK}, e_{JK})$$


$$B_J := f_B(Z_J, e_J)$$

$$e_I \perp \{e_J, e_{IA}, e_{JK}\}$$

$$e_J \perp \{e_{IA}, e_{JK}\}$$

$$e_{IA} \perp e_{JK}$$

(a) SCM



(b) DAG

Figure 6: Sample-comparison model, final vectorized form

#### 4.2.2. The comparison mechanism

As in the previous section, we begin defining the *comparison mechanism* using the binary vector variable  $C$  to facilitate the interpretation of the sample-comparison model. Equation (9) shows that  $C$  contains  $n$  elements corresponding to the number of rows in the  $R$  matrix, with each element  $c_{(i,a,h,b,j,k)}$  being a binary value indicating the presence or absence of data rows in  $R$ , a definition similar to that of  $s_{(i,a,h,b,j,k)}$  in Equation (7).

$$C = \begin{bmatrix} c_{(1,1,1,2,1,1)} \\ \vdots \\ c_{(1,1,1,2,1,n_K)} \\ \vdots \\ c_{(i,a,h,b,j,k)} \\ \vdots \\ c_{(n_I,n_A-1,n_I,n_A,n_J,1)} \\ \vdots \\ c_{(n_I,n_A-1,n_I,n_A,n_J,1)} \end{bmatrix} \quad (9)$$

The DAG 6b also incorporates the *comparison mechanism*  $C$  into the conceptual-population model. It shows the sample-comparison outcome  $O_R^{sc}$  as unobserved, emphasizing that researchers cannot directly access this variable because of the comparison mechanism. The DAG further shows  $C$  as a conditioned variable (enclosed in a square) that causally influences the observed outcome  $O_R$ . This structure implies that  $C$  determines *which repeated judgments judges make for the stimuli produced by the individuals*. In essence,  $C$  reflects the assumption that judges *do not* perform all possible repeated judgments but instead complete a sufficient number,  $n_C$ , to enable the accurate estimation of the proportion  $P(B > A)$  for each stimulus pair (Thurstone, 1927a, p. 267).

Notably, DAG 6b also shows that  $C$  is independent of all other variables in the model. This independence implies that the conceptual model represented by the DAG applies exclusively to Random Allocation Comparative Designs (Bramley, 2015), or Incomplete Block Designs (Lawson, 2015), where every repeated judgment has an equal probability of being included in the sample.

Finally, since it is standard to assume that the distribution of the conceptual-population outcome  $O_R^{cp}$  also holds for  $O_R^{sc}$  and  $O_R$ , we can reformulate the sample-comparison model in Figure 6 into the equivalent form shown in Figure 7. This reformulation produces a model that applies directly to a sample of comparative data. In this version, the unobserved outcomes  $O_R^{cp}$  and  $O_R^{sc}$  are omitted, and



$O_R$  inherits the structural equation  $f_O$  that originally defined  $O_R^{cp}$ . Moreover, the definition of  $O_R$  now reflects its direct dependence on the discriminial difference  $D_R$  and the sample and comparison mechanisms,  $S$  and  $C$ .

$$O_R := f_O(D_R, S, C)$$

$$D_R := f_D(T_{IA}, B_{JK})$$

$$T_{IA} := f_T(T_I, X_{IA}, e_{IA})$$

$$T_I := f_T(X_I, e_I)$$

$$B_{JK} := f_B(B_J, Z_{JK}, e_{JK})$$


$$B_J := f_B(Z_J, e_J)$$

$$e_I \perp \{e_J, e_{IA}, e_{JK}\}$$

$$e_J \perp \{e_{IA}, e_{JK}\}$$

$$e_{IA} \perp e_{JK}$$

(a) SCM



(b) DAG

Figure 7: Comparative judgment model

In summary, the SCM 7a and DAG 7b extend Thurstone’s general form to address several limitations of the BTL model. These extensions account for judge biases (see Section 4.1.1), reflect the hierarchical structure of stimuli and incorporate measurement error in trait estimation and hypothesis testing (see Section 4.1.3), and even clarify the role of the sample and comparison mechanisms in CJ assessments (see Section 4.2). However, they do not resolve concerns about the assumption of equal dispersions among stimuli discussed in Section 3.1.1. Since this concern relates to the statistical assumption underlying the distribution of the discriminial process, we develop a formal statistical model to address it in the next section.

## 5. From SCM to statistical model

Using the structural causal model (SCM) 7a, we can derive a statistical model that addresses violations of the equal dispersion assumption (see Section 3.1.1). This derivation is possible because a fully specified SCM encodes functional and probabilistic information, which we can replace with suitable functions and probabilistic assumptions (Pearl et al., 2016). Specifically, SCM 7a allows us to express the joint distribution of our complex CJ system as a product of simpler conditional probability distributions (CPDs)<sup>7</sup>, as shown in Equation (10). For clarity, we treat expressions such as  $Y := f_Y(X)$ ,  $P(Y | X)$ , and  $Y \sim f(Y | X)$  as equivalent, where  $P(Y | X)$  and  $f(Y | X)$  represent the CPD of  $Y$  given  $X$ .

$$\begin{aligned}
&P(O_R, S, C, D_R, T_{IA}, X_{IA}, e_{IA}, T_I, X_I, e_I, B_{JK}, Z_{JK}, e_{JK}, B_J, Z_J, e_J) \\
&= P(O_R | D_R, S, C) \cdot P(S) \cdot P(C) \cdot P(D_R | T_{IA}, B_{JK}) \\
&\quad \cdot P(T_{IA} | T_I, X_{IA}, e_{IA}) \cdot P(T_I | X_I, e_I) \\
&\quad \cdot P(B_{JK} | B_J, Z_{JK}, e_{JK}) \cdot P(B_J | Z_J, e_J) \\
&\quad \cdot P(X_{IA}) \cdot P(X_I) \cdot P(Z_{JK}) \cdot P(Z_J) \\
&\quad \cdot P(e_{IA}) \cdot P(e_I) \cdot P(e_{JK}) \cdot P(e_J)
\end{aligned} \tag{10}$$

Each CPD in Equation (10) rests on specific assumptions, which we outline in the statistical model presented in Figure 8c. The model starts by assuming that  $O_R$  follows a Bernoulli distribution<sup>8</sup>,

<sup>7</sup>This re-expression is possible because the *chain rule* of probability and the *Bayesian Network Factorization (BNF)* property. For further details, see Pearl et al. (2016) and Neal (2020).

<sup>8</sup>The binomial distribution—including its special case, the Bernoulli distribution—represent a maximum entropy distribution for binary events (McElreath, 2020, p. 34). This means that the Bernoulli distribution is the most consistent alternative when only two un-ordered outcomes are possible and their expected frequencies are assumed to be constant (McElreath, 2020, p. 310). For a detailed discussion of the binomial as a maximum entropy distribution, see McElreath (2020, sec. 10.1.2).

reflecting the binary nature of CJ outcomes. Furthermore, following the conventions of Generalized Linear Models (GLMs) (McCullagh and Nelder, 1983; Lee and Nelder, 1996; Agresti, 2015), the distribution links  $O_R$  to the latent discriminial difference vector  $D_R$  using an inverse-logit function:  $\text{inv\_logit}(x) = 1/(1 + \exp(-x))$ .

$O_R := f_O(D_R, S, C)$	$P(O_R   D_R, S, C)$	$O_R \stackrel{iid}{\sim} \text{Bernoulli}[\text{inv\_logit}(D_R)]$
$D_R := f_D(T_{IA}, B_{JK})$	$P(D_R   T_{IA}, B_{JK})$	$D_R = (T_{IA}[i, a] - T_{IA}[h, b]) + B_{JK}[j, k]$
$T_{IA} := f_T(T_I, X_{IA}, e_{IA})$	$P(T_{IA}   T_I, X_{IA}, e_{IA})$	$T_{IA} = T_I + \beta_{XA}X_{IA} + e_{IA}$
$T_I := f_T(X_I, e_I)$	$P(T_I   X_I, e_I)$	$T_I = \beta_{XI}X_I + e_I$
$B_{JK} := f_B(B_J, Z_{JK}, e_{JK})$	$P(B_{JK}   B_J, Z_{JK}, e_{JK})$	$B_{JK} = B_J + \beta_{ZK}Z_{JK} + e_{JK}$
$B_J := f_B(Z_J, e_J)$	$P(B_J   Z_J, e_J)$	$B_J = \beta_{ZJ}Z_J + e_J$
$e_I \perp \{e_J, e_{IA}, e_{JK}\}$	$P(e_I)P(e_{IA})P(e_J)P(e_{JK})$	$e \sim \text{Multi-Normal}(\mu, \Sigma)$
$e_J \perp \{e_{IA}, e_{JK}\}$		$\Sigma = VQV$
$e_{IA} \perp e_{JK}$		
(a) SCM	(b) Probabilistic model	(c) Statistical model

Figure 8: Comparative judgment model, SCM, probabilistic and statistical model assuming different discriminial dispersions for the student’s traits

While the joint distribution in Equation (10) includes the probability distributions of the sampling and comparison mechanisms,  $P(S)$  and  $P(C)$ , as well as those of the predictor variables— $P(X_{IA})$ ,  $P(X_I)$ ,  $P(Z_{JK})$ , and  $P(Z_J)$ —all of these probabilities are omitted from the statistical model 8c. This omission is justified because, while these distributions contribute to the overall joint distribution of the data, the variables  $S$ ,  $C$ ,  $X_{IA}$ ,  $X_I$ ,  $Z_{JK}$ , and  $Z_J$  are observed and independent of any other variable in the model. As observed variables, they do not require distributional assumptions in the same way the idiosyncratic errors do. Their independence follows from the underlying random selection procedures that govern the variables<sup>9</sup>.

<sup>9</sup>Randomization ensures that data—and, by extension, an estimator—satisfies several key identification properties, such as common support, no interference, and consistency. The most critical property, however, is the elimination of confounding. *Confounding* occurs when an external variable, such as  $X_I$ , simultaneously influences both the outcome (e.g.,  $O_R$ ) and a variable of interest (e.g.,  $S$ ), resulting in spurious associations between the latter two (Everitt and Skrondal, 2010). Randomization ensure the absence of confounding by effectively decoupling the association between

Next  $D_R$  is defined as the difference between the discriminial processes  $T_{IA}[i, a]$  and  $T_{IA}[h, b]$ , representing the underlying written-quality trait of the compared texts, plus the corresponding repeated judge bias  $B_{JK}[j, k]$ . Note that if it is assumed that  $B_{JK}[j, k]$  reflects the difference in stimulus-specific biases, i.e.,  $B_{JK}[j, k] = B_{JK}[i, a, j, k] - B_{JK}[h, b, j, k]$ , the discriminial difference can be re-written as:

$$\begin{aligned} D_R &= (T_{IA}[i, a] - T_{IA}[h, b]) + B_{JK}[j, k] \\ &= (T_{IA}[i, a] + B_{JK}[i, a, j, k]) - (T_{IA}[h, b] + B_{JK}[h, b, j, k]) \\ &= T_{IA}^*[i, a] - T_{IA}^*[h, b] \end{aligned} \tag{11}$$

This formulation reveals that the discriminial difference captures a *pure interaction effect*, in which neither the texts' discriminial processes nor the judges' biases alone determine the outcome, but their interaction does (Attia et al., 2022). Put simply, this mathematical description captures the idea that the stimuli' discriminial processes become an observable outcome only through the lens of judges' perceptions (i.e., their biases). For clarity, the square brackets in  $D_R$  indicate the relevant indices for each trait vector; they do not imply any subsetting of the data.

Now the functional forms for  $T_{IA}$ ,  $T_I$ ,  $B_{JK}$ , and  $B_J$  are specified.  $T_{IA}$  is modeled as a linear combination of the students' underlying writing-quality traits  $T_I$ , the effects of relevant text-related variables on quality assessment  $\beta_{XA}X_{IA}$  (such as the influence of text length), and the text-specific idiosyncratic errors  $e_{IA}$ . Similarly,  $T_I$  is expressed as a linear combination of relevant student-related variables affecting the quality assessment  $\beta_{XI}X_I$ , and student-specific idiosyncratic errors  $e_I$ . For the judge-specific terms,  $B_{JK}$  is modeled as a linear combination of the judge's individual bias  $B_J$ , the influence of relevant judgment-related variables on quality assessment  $\beta_{ZK}Z_{JK}$  (e.g., how the number of judgments affect the evaluation), and judgment-specific idiosyncratic errors  $e_{JK}$ . Finally,  $B_J$  is defined as a linear combination of relevant judge-level variables influencing the quality assessment  $\beta_{ZJ}Z_J$  (such as judgment expertise) and judge-specific idiosyncratic errors  $e_J$ .

Next, the probabilistic assumptions for the idiosyncratic errors  $e_I$ ,  $e_{IA}$ ,  $e_J$ , and  $e_{JK}$  are specified. Unlike other variables in the model, these error terms exhibit indeterminacies in their *location*, *orientation*, and *scale* due to the lack of an inherent scale in the associated latent variables  $T_I$ ,  $T_{IA}$ ,  $B_J$ , and  $B_{JK}$ . Thus, to identify the latent variable model these indeterminacies must be resolved (Depaoli, 2021; de Ayala, 2009). Drawing on principles from SEM (Hoyle, 2023), the vector of

---

the variable of interest and any other variable, except for the outcome itself. For a more detailed discussion on the benefits of randomization, see Pearl (2009), Morgan and Winship (2014), Neal (2020), and Hernán and Robins (2025).

idiosyncratic errors  $e = [e_I, e_{IA}, e_J, e_{JK}]^T$  are assumed to follow a Multivariate Normal distribution with mean vector  $\mu$  and a covariance matrix  $\Sigma = VQV$ , with  $V$  denoting a diagonal matrix of standard deviations and  $Q$  a correlation matrix. To address the *location* indeterminacy, the errors' mean vector is set to zero:

$$\mu = [0, 0, 0, 0]^T \quad (12)$$

Following SCM 8a, the *orientation* indeterminacy is solved by assuming that the errors are uncorrelated. This assumption leads to the definition of the error's correlation matrix,  $Q$ , as the identity matrix:

$$Q = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad (13)$$

To resolve the *scale* indeterminacy, the diagonal matrix  $V$  is defined as follows:

$$V = \begin{bmatrix} s_{XI} & 0 & 0 & 0 \\ 0 & p_{IA} & 0 & 0 \\ 0 & 0 & s_{ZJ} & 0 \\ 0 & 0 & 0 & p_{JK} \end{bmatrix} \quad (14)$$

Here,  $s_{XI}$  represents the standard deviation for the individuals,  $p_{IA}$  for the stimuli,  $s_{ZJ}$  for the judges, and  $p_{JK}$  for the judgments. It is assumed  $s_{XI}$  varies depending on the teaching method group to which each student belongs. Using the example from Section 4, where the teaching method  $X_I = \{1, 2\}$ , the model sets the constraint according to Equation (15). This constraint anchors the scale of the individuals' latent trait while relaxing the assumption of equal dispersion for the stimuli, thereby addressing the concerns raised in Section 3.1.1.

$$\sum_{g=1}^2 s_{XI}[g]/2 = 1 \quad (15)$$

Because the error vector  $e$  follows an uncorrelated Multivariate Normal distribution, the marginal distribution of  $e_{IA}$  is a univariate Normal distribution with mean zero and standard deviation  $p_{IA}$ . Thus,  $p_{IA}$  is set as a proportion of 1 to establish the scale of the stimuli' latent trait relative to the scale of the individuals' trait. Note that as a result,  $T_{IA}$  is also normally distributed. This configuration effectively reinstates Thurstone's original assumption of Normal discriminial processes for the stimuli (see Table 1).

Similarly, it is assumed that  $s_{ZJ}$  varies depending on the groups to which each judge belongs. For instance, if  $Z_J = \{1, 2, 3\}$  represents three groups of judges with varying expertise, the model sets the constraint according to Equation (16). This constraint anchors the scale of the judges' latent trait and relaxes the assumption of equal dispersion for the judgments.

$$\sum_{g=1}^2 s_{ZJ}[g]/3 = 1 \quad (16)$$

Conversely,  $p_{JK}$  is defined as a proportion of 1 to establish the scale of the judgments' latent trait relative to the scale of the judges' trait.

Finally, we use *Bayesian inference methods* to convert the statistical model 8c into a practical statistical tool for analyzing paired comparison data. Bayesian inference offers three main advantages in this context. First, it handles complex and overparameterized models, where the number of parameters exceeds the number of observations (Baker, 1998; Kim and Cohen, 1999). This feature is essential for our implementation, as the proposed model is indeed overparameterized. Second, it incorporates prior information to constrain parameter estimates within plausible bounds, thereby mitigating estimation issues like non-convergence or improper solutions that often affect frequentist methods (Martin and McDonald, 1975; Seaman III et al., 2011). Prior distributions are used to define the error distribution and set the scale of latent variables (Depaoli, 2014). Third, Bayesian inference supports robust inferences from small samples, where the asymptotic properties underlying frequentist methods are less reliable (Baldwin and Fellingham, 2013; Lambert et al., 2006; Depaoli, 2014). This feature is particularly relevant in CJ assessments, as researchers often collect large volumes of paired comparisons but work with relatively small samples of judges, stimuli, and individuals to test hypotheses.

The **Declarations** section of this document provides a link to the model code, along with an alternative specification that assumes equal discriminial dispersions. We tested both versions of the model with success using **Stan** (Stan Development Team., 2021, version 2.26.1).

## 6. Discussion

Thurstone introduced the Law of Comparative Judgment to measure psychological traits of stimuli through pairwise comparisons (Thurstone, 1927b,a). In its general form, the theory models single-judge comparisons across multiple, potentially correlated stimuli. Each comparison produces a dichotomous outcome indicating which stimulus the judge perceives as having a higher trait level.

However, Thurstone identified one key challenge in this general formulation: the measurement model required estimating more “unknown” parameters than the number of available pairwise comparisons (Thurstone, 1927a). To address this issue and facilitate the theory’s practical applicability, he formulated five cases, each progressively incorporating several simplifying assumptions.

Among these, Case V remains the most widely used model in empirical CJ research, mainly due to the widespread adoption of the BTL model. The BTL model mirrors the core assumptions of Case V—namely, equal discriminial dispersions and zero correlation among stimuli’ discriminial processes—but replaces the processes’ normal distribution with the more mathematically tractable logistic distribution (Andrich, 1978; Bramley, 2008). Although this substitution has minimal impact on trait estimation or model interpretation (van der Linden, 2017a; McElreath, 2021), the simplifying assumptions of the BTL model—and by extension, of Case V—may fail to capture the complexity of some traits or account for heterogeneous stimuli (Thurstone, 1927b; Andrich, 1978; Bramley, 2008; Kelly et al., 2022), potentially leading to unreliable and inaccurate trait estimates (Ackerman, 1989; Zimmerman, 1994; McElreath, 2020; Hoyle, 2023).

Moreover, because Thurstone’s original goal was to produce a “coarse scaling” of traits and allocate stimuli along this continuum (Thurstone, 1927b, p. 269), his theory offered no guidance on how to use trait estimates for statistical inference. The CJ tradition has attempted to address this gap by separating trait estimation from hypothesis testing, relying on point estimates, such as BTL scores or their transformations, for inference. While this approach simplifies analysis, it can also introduce bias and compromise the reliability of the resulting inferences (McElreath, 2020; Kline, 2023; Hoyle, 2023).

To address the limitations of Thurstone’s Case V and the BTL model, this study extends Thurstone’s general form using a systematic, integrated approach that combines causal and Bayesian inference methods. The approach begins with the development of a conceptual model, represented as a Structural Causal Model (SCM) and a Directed Acyclic Graph (DAG) (Pearl, 2009; Pearl et al., 2016; Gross et al., 2018; Neal, 2020). This model integrates Thurstone’s core theoretical principles, such as the discriminial processes of stimuli, alongside key CJ assessment design features, including judges’ bias, sampling procedures, and comparison mechanisms, to disentangle the causal processes underlying the CJ system.

The approach then translates the SCM into a bespoke statistical model that allows researchers to analyze CJ data when violations to the assumptions of equal dispersion and zero correlation

occur, and when statistical inference is necessary. In particular, this model accounts for judge biases, captures the hierarchical structure of stimuli, incorporates measurement error into the hypothesis testing process, and accommodates heterogeneity in discriminial dispersions. By addressing all these issues, these methodological innovations have the potential to enhance the reliability and validity of trait measurement in CJ (Perron and Gillespie, 2015), while also improving the accuracy of statistical inferences.

Beyond these potential benefits, the approach offers two additional advantages. First, it clarifies the roles and interactions of all actors and processes involved in CJ assessments. Second, it shifts the analytic paradigm from passively accepting the assumptions of Case V and the BTL model to actively testing their fit with observed data. Together, these advantages establish a principled framework for evaluating best practices in designing CJ assessments, one that better aligns with the demands of contemporary CJ contexts (Kelly et al., 2022), offering new insights into existing research and opening promising avenues for future inquiry.

#### *6.1. Future research directions using our approach*

Among the many potential directions for future research, three avenues deserve particular attention due to their direct impact on the reliability and validity of CJ trait estimates, as well as on the accuracy of statistical inferences. The following sections outline these avenues and explain how our approach facilitates their investigation.

##### *6.1.1. The impact of sampling and comparison mechanisms*

Although sampling and comparison mechanisms are central to modern CJ assessments, it is striking that most CJ literature has examined them within a limited scope. Researchers have primarily investigated the effects of adaptive comparative judgment (ACJ) designs on trait reliability (Pollitt, 2012a,b; Bramley, 2015; Verhavert et al., 2022; Mikhailiuk et al., 2021; Gray et al., 2024) or proposed practical guidelines for the number of comparisons judges should make (Verhavert et al., 2019; Cromptvoets et al., 2022). While these studies offer valuable insights, they also overlook the broader role that these mechanisms play within the CJ system. As this oversight likely stems from a more fundamental lack of conceptual clarity about how these mechanisms function within the system, this study integrates these mechanisms into the conceptual model of CJ.

The explicit integration of the sampling and comparison mechanisms offers a new perspective on how these mechanisms shape the CJ process. Specifically, it clarifies their role as sources of missing data in CJ's data-generating process, that is, as mechanisms that determine which observations are



missing from the final data sample. This new perspective encourages the application of Little and Rubin’s principled missing data framework (2020), allowing a more rigorous evaluation of existing claims about these missing data mechanisms, their influence on CJ outcomes, and their implications for designing and evaluating more complex assessments setups.

This study circumvents the need to apply this missing data framework by deliberately structuring the sampling and comparison mechanisms to be independent of any observed or unobserved variables, including the outcome. In other words, these mechanisms are designed to produce data that are *missing completely at random* (MCAR) (Little and Rubin, 2020). This design offers one key advantage: it generates simple random samples that satisfy the condition of *ignorability*, allowing researchers to legitimately *ignore* missing data during analysis without introducing bias (Everitt and Skrondal, 2010; Kohler et al., 2019; Neal, 2020).

However, many modern CJ applications rely on more complex assessment designs, in which the sampling and comparison mechanisms introduce more intricate forms of missingness such as *missing at random* (MAR) or *missing not at random* (MNAR) (Little and Rubin, 2020). A prominent example is the previously discussed ACJ design, where prior judgment outcomes inform the selection of stimulus pairs for subsequent comparisons (Pollitt, 2012a,b; Bramley, 2015). This pair selection procedure suggests that ACJ’s comparison mechanism is outcome-dependent, potentially classifying the method as a generator of MNAR data. If this classification holds, the mixed findings on ACJ’s capabilities become more comprehensible: some studies find that the method improves trait reliability (Pollitt and Elliott, 2003; Pollitt, 2012a,b), while others contend that it artificially inflates these gains (Bramley, 2015; Bramley and Vitello, 2019; Cromptvoets et al., 2020, 2022).

Regardless of the underlying missingness mechanisms, any CJ assessment design would benefit from explicitly defining its assumptions—a practice supported by our approach. This clarity enables researchers to evaluate how the sampling and comparison mechanisms affect trait estimation and statistical inference within each design. Such assessments are particularly relevant given the common misconception in the CJ literature that Thurstone’s model can naturally handle even non-random missing data without compromising the reliability or validity of trait estimates (Bramley, 2008).

#### 6.1.2. The effects of judges’ bias on the reliability of traits

Despite the growing notion that various stimulus-related factors influence judges’ perceptions (van Daal et al., 2016; Lesterhuis et al., 2018; Chambers and Cunningham, 2022) and that these influences may not always cancel each other out, few studies in the CJ literature provide empirical

evidence for judges’ biases (Pollitt and Elliott, 2003; van Daal et al., 2016; Bartholomew et al., 2020). This gap likely persists not due to a lack of interest or research but because researchers often rely on ad-hoc detection methods, such as ‘misfit’ statistics, that may not be well-suited for the task (Kelly et al., 2022). To address this limitation, this study treats judges’ biases as an integral component of the CJ system from the outset. This approach offers one key advantage: it provides a more accurate representation of the data-generating process behind pairwise comparisons, one that acknowledges that the discriminative processes of stimuli become an observable outcome only through judges’ perceptions, which may exhibit bias.

The explicit integration of judges’ bias into CJ’s conceptual model then paves the way for investigating several relevant research questions. One key question is whether researchers can validly analyze CJ data under the assumption of “sample-free” trait calibration, specifically under the hypothesis that judges exhibit no systematic bias. This question is particularly relevant because many researchers still regard “sample-freeness” as an inherent property of the BTL model (Bramley, 2008; Andrich, 1978) despite growing evidence of persistent biases. Another critical question is whether training or expertise can help judges avoid focusing on irrelevant stimulus features, such as handwriting, over more central criteria like argumentative quality in writing assessments (Kelly et al., 2022). Exploring these questions may also provide insights into what it truly means to be an “expert” within the CJ context (Kelly et al., 2022).

Moreover, since judges rely on these stimulus-related factors when evaluating complex, multidimensional traits (van Daal et al., 2016; Lesterhuis et al., 2018; Chambers and Cunningham, 2022), and these factors account for variation in judgment accuracy (Gill and Bramley, 2013; van Daal et al., 2017; van Daal, 2020; Gijzen et al., 2021), it is reasonable to expect that assessments also vary according to judge-specific attributes such as gender, age, culture, income, education, training, or expertise (Kelly et al., 2022). Prior studies support this view (Bartholomew et al., 2020; McMahon and Jones, 2015). Thus, building on the discussion in Section 6.1.1, researchers could further explore how judges’ selection influences the formation of a “shared consensus” and whether these attributes introduce systematic biases or distortions in the observed trait distribution (Deffner et al., 2022). Furthermore, if such attributes indeed undermine the assumption of “sample-freeness,” it becomes essential to explore strategies for mitigating their influence and to determine how many judges (and how many judgments per judge) are needed to produce reliable trait estimates under these conditions. In addition, it is worth considering whether *repeated measures designs*, in which judges evaluate the same stimulus pairs multiple times (Lawson, 2015), can improve judgment consistency

and accuracy. As anticipated, the approach presented in this study provides the necessary structure to investigate rigorously these questions.

### 6.1.3. The identification of ‘misfitting’ judges and stimuli

Although the CJ literature clearly defines *misfit* judges and stimuli, CJ researchers have rarely examined how these observations relate to Thurstonian theory. In particular, they have not identified which elements of Thurstone’s theory might account for the occurrence of misfits. This disconnect likely stems from the fact that CJ researchers derive misfit statistics from residual analysis and outlier detection methods rather than from Thurstonian principles. Specifically, *misfit judges* are typically defined as those whose assessments diverge significantly from the “shared consensus” (Pollitt, 2012a,b; van Daal et al., 2016; Goossens and De Maeyer, 2018; Wu et al., 2022), while *misfit stimuli* are those that elicit more judgment discrepancies than others (Pollitt, 2004, 2012a,b; Goossens and De Maeyer, 2018). Both definitions closely mirror the statistical concept of outliers, that is, observations that deviate markedly from the rest of the sample in which they occur (Grubbs, 1969). But this resemblance extends beyond the definitions themselves, as CJ researchers often identify misfits using conventional outlier detection procedures, such as transforming BTL model residuals into diagnostic statistics and comparing them against predefined thresholds (Pollitt, 2012a,b; Wu et al., 2022).

Nevertheless, it is not the classification of misfits as outliers that raises concerns for trait measurement and inference. Rather, the concern lies in the prevailing CJ practice of classify them through ad-hoc detection procedures and then excluding them from analysis (Pollitt, 2012a,b), often without providing empirical evidence for the various hypotheses proposed to explain their occurrence. In this regard, it is essential to recognize that outliers can only be defined relative to a specific model (McElreath, 2020). As such, detection procedures based on models like the BTL, which relies on strong assumptions, should be approached with caution, as they may not be well-suited for the task (Kelly et al., 2022). In particular, because the BTL model may not always accurately reflect the underlying data-generating process of the CJ system, misfit classification based on this model risks misidentifying observations. Furthermore, the exclusion of these observations carries additional risks. The statistical literature cautions that removing outliers can discard valuable information (Miller, 2023) and introduce bias into trait estimates. The direction and magnitude of this bias are often unpredictable, as they depend on which observations researchers exclude from the analysis (Zimmerman, 1994; O’Hagan, 2018; McElreath, 2020). Finally, precisely because of its rigid assumptions, the BTL model lacks the flexibility to adequately test many of the hypotheses proposed to explain the presence of misfits.

In contrast, the approach presented in this study provides a rigorous framework that enables the examination of several relevant hypotheses. For instance, researchers could investigate whether misfit judges are those who exhibit (an outlying degree of) systematic bias or greater variability in their judgments compared to their peers. Similarly, researchers might explore whether misfit stimuli exhibit more variable discriminative processes relative to other stimuli or if they are genuinely outlying cases. Moreover, since outliers are not necessarily “bad data” (McElreath, 2020), our approach also offers a principled alternative that retains misfits in the analysis without compromising trait estimation or inference. This alternative involves adapting the proposed model into robust measurement models (McElreath, 2020), a broad class of procedures designed to reduce the sensitivity of parameter estimates to mild or moderate departures from model assumptions (Everitt and Skrondal, 2010). This alternative also relates to a broader discussion in social science, which is: that when researchers face low predictive capacity from their models, they should not only search for “new” variables or alternative procedures but also consider employing more sophisticated measurement models (Wainer et al., 1978).

## 6.2. Study limitations and practical challenges for applied CJ researchers

The process of deriving conclusions from observed data always requires assumptions, whether the data are observational or experimental (Kohler et al., 2019; Deffner et al., 2022). The proposed approach is not an exception to this fundamental principle. As with all approaches grounded on causal inference, it relies on expert knowledge and assumptions about the variables’ causal structure that are often untestable at the outset (Hernán and Robins, 2025). As such, the approach does not seek to yield automatic answers when applied to a given CJ dataset. Instead, it aims to encourage researchers to formulate precise questions and to make their assumptions explicit, fostering a generalizable understanding of the CJ system under study (Rohrer et al., 2022; Deffner et al., 2022; Sterner et al., 2024). This clarity is critical because a model’s ability to produce accurate estimates and valid inferences depends heavily on how well the data and inferential goals align with its underlying assumptions (Kohler et al., 2019). Although this alignment remains empirically untested for the proposed models, the theory-driven nature of our approach provides a solid foundation for future empirical evaluation of its causal assumptions (Deffner et al., 2022), which are well supported by both theory and existing evidence.

Moreover, this theoretical commitment to causal inference also introduces several practical challenges that applied CJ researchers must navigate. These fall into two main categories: first, acquiring the foundational knowledge necessary to apply the approach effectively; and second, dedicating greater

attention to both conceptual and statistical modeling.

#### 6.2.1. Required foundational knowledge

To apply the approach effectively, CJ researchers require foundational knowledge in two areas. First, they need a solid understanding of causal inference principles. Second, they must learn to translate the functional and probabilistic content of conceptual models into bespoke statistical models. One example that illustrates the importance of a sound knowledge of causal inference is the recurrent assumption that predictor variables are “relevant” to the research context—interpreting this relevance as their inclusion in a *sufficient adjustment set* (Pearl, 2009; Pearl et al., 2016; Morgan and Winship, 2014). However, we do not fully explore what this assumption entails or its implications for model specification, estimation, and inference. As a result, CJ researchers must deeply engage with these and other complex ideas, including how SCMs encode functional and probabilistic information. To assist this effort, this study includes key references throughout the study to guide applied CJ researchers toward a deeper understanding of these concepts <sup>10</sup>.

Developing the skills to translate conceptual models into bespoke statistical models also presents challenges. Although Bayesian inference methods offer a more accessible path for many applied CJ researchers to develop these skills—by reducing the need for specialized knowledge in areas like optimization theory, which frequentist methods often require—they still demand a working knowledge of other technical concepts such as probabilistic programming languages (PPLs), probability distributions, and convergence. These requirements can be non-trivial to master. To support CJ researchers in developing these skills, this study provides a link to the statistical model code and alternative model specifications in the **Declarations** section of this document <sup>11</sup>.

#### 6.2.2. Attention to conceptual and statistical modeling

Even after acquiring the necessary foundational knowledge, CJ researchers still face two additional challenges when applying our approach to their specific research context. First, they must verify whether the conceptual model provides a *faithful* <sup>12</sup> representation of the CJ system under study. Second, they need to assess whether the statistical translation of the model can accurately estimate the intended estimands (i.e., parameters) from empirical data. To ensure a faithful representation of

---

<sup>10</sup>refer to footnote 1, 2, and the detailed online document referenced in the Declarations section

<sup>11</sup>Seminal texts on Bayesian inference methods, such as Gelman et al. (2014) and McElreath (2020), offer valuable support for developing a deeper understanding of these models.

<sup>12</sup>Avoid confusing this term with the *faithfulness assumption* (Neal, 2020; Hernán and Robins, 2025) described in the causal inference literature. For further details, see footnote 1.

the CJ system, we encourage researchers to treat our conceptual and statistical models as starting points rather than universal solutions for all CJ designs or datasets. While these models may prove adequate in some contexts, researchers cannot assume this apriori. Instead, they need to adapt these models to their specific contexts, paying close attention to assessment design features and causal assumptions. As discussed in Section 6.2, our approach facilitates this process by offering a transparent framework for articulating new assumptions and guiding the design of CJ assessments.

Conversely, to evaluate the estimation capabilities of the statistical model, researchers must tackle the challenge of performing identification analysis. As outlined in Section 4, *identification analysis* determines whether a statistical model can accurately compute a given estimand (e.g., a parameter) based solely on its (causal) assumptions, independent of random variability (Schuessler and Selb, 2023). Identification is crucial because it is a necessary condition for consistency. *Consistency* is the property of an estimator (e.g., a statistical model) whose estimates converge to the “true” value of an estimand as data size approaches infinity (Everitt and Skrondal, 2010). Without identification, consistency is impossible—even with infinite, error-free data—and thus, meaningful inference from finite samples cannot be achieved (Schuessler and Selb, 2023). Although performing identification analysis through formal derivations may seem like a natural next step, the complexity of the CJ system renders such an approach impractical at the outset. Instead, simulation-based methods, such as power analysis, provide a more practical and flexible alternative, enabling researchers to examine the consistency of estimates without relying on cumbersome mathematical proofs. Nonetheless, the development of formal identification proofs remains a significant goal for future CJ research. Notably, our approach supports both strategies by providing the probabilistic foundation for formal derivations and the necessary statistical structure for simulation-based methods.

## 7. Conclusion

This paper underscores the need to extend Thurstone’s theory to meet the demands of contemporary empirical CJ research. It specifically advocates for the development of bespoke CJ models aligned with the underlying data-generating processes of specific CJ assessment designs, as these models have the potential to strengthen the robustness of trait estimates and the clarity of inferences. The work also outlines a clear trajectory for advancing CJ research in both theoretical and applied domains. Finally, the study also lays the groundwork for broader applications, encouraging researchers across the social sciences to adopt more robust and interpretable methodologies.

## Declarations

**Funding:** The Research Fund (BOF) of the University of Antwerp funded this project.

**Financial interests:** The authors declare no relevant financial interests.

**Non-financial interests:** The authors declare no relevant non-financial interests.

**Ethics approval:** The University of Antwerp Research Ethics Committee confirmed that this study does not require ethical approval.

**Consent to participate:** Not applicable

**Consent for publication:** All authors have read and approved the final version of the manuscript for publication.

**Data availability:** This study did not use any data.

**Materials and code availability:** The CODE LINK section at the top of the digital document located at: [https://jriveraespejo.github.io/paper2\\_manuscript/](https://jriveraespejo.github.io/paper2_manuscript/) provides access to all materials and code.

**AI-assisted technologies in the writing process:** The authors used various AI-based language tools to refine phrasing, optimize wording, and enhance clarity and coherence throughout the manuscript. They take full responsibility for the final content of the publication.

**CRedit authorship contribution statement:** *Conceptualization:* J.M.R.E, T.vD., S.DM., and S.G.; *Methodology:* J.M.R.E, T.vD., and S.DM.; *Software:* J.M.R.E.; *Validation:* J.M.R.E.; *Formal Analysis:* J.M.R.E.; *Investigation:* J.M.R.E; *Resources:* T.vD. and S.DM.; *Data curation:* J.M.R.E.; *Writing - original draft:* J.M.R.E.; *Writing - review and editing:* T.vD., S.DM., and S.G.; *Visualization:* J.M.R.E.; *Supervision:* S.G. and S.DM.; *Project administration:* S.G. and S.DM.; *Funding acquisition:* S.G. and S.DM.

## 8. Appendix

### 8.1. Statistical and Causal inference

This section introduces fundamental statistical and causal inference concepts necessary for understanding the core theoretical principles described in this document. It does not, however, offer a comprehensive overview of causal inference methods. Readers seeking more in-depth understanding may wish to explore introductory papers such as [Pearl \(2010\)](#), [Rohrer \(2018\)](#), [Pearl \(2019\)](#), and [Cinelli et al. \(2020\)](#). They may also find it helpful to consult introductory books like [Pearl and Mackenzie \(2018\)](#), [Neal \(2020\)](#), and [McElreath \(2020\)](#). For more advanced study, readers may refer to seminal intermediate papers such as [Neyman \(1923\)](#), [Rubin \(1974\)](#), [Spirtes et al. \(1991\)](#), and [Sekhon \(2009\)](#), as well as books such as [Pearl \(2009\)](#), [Morgan and Winship \(2014\)](#), and [Hernán and Robins \(2025\)](#).

#### 8.1.1. Empirical research and randomized experiments

Empirical research uses evidence from observation and experimentation to address real-world challenges. In this context, researchers typically formulate their research questions as *estimands* or *targets of inference*, i.e., the specific quantities they seek to determine ([Everitt and Skrondal, 2010](#)). For instance, researchers might be interested in answering the following question: “To what extent do different teaching methods ( $T$ ) influence students’ ability to produce high-quality written texts ( $Y$ )?” To investigate this, researchers could randomly assign students to two groups, each exposed to a different teaching method ( $T_i = \{1, 2\}$ ). Then, they would perform pairwise comparisons, generating a dichotomous outcome ( $Y_i = \{0, 1\}$ ) showing which student exhibits more of the ability. In this scenario, the research question can be rephrased as the estimand, “*On average*, is there a difference in the ability to produce high-quality written texts between the two groups of students?” and this estimand can be mathematically represented by the random associational quantity in Equation 17, where  $E[\cdot]$  denotes the expected value.

$$E[Y_i | T_i = 1] - E[Y_i | T_i = 2] \tag{17}$$

Researchers then proceed to identify the estimands. *Identification* determines whether an estimator can accurately compute the estimand based solely on its assumptions, regardless of random variability ([Schuessler and Selb, 2023](#), p. 4). An *estimator* refers to a method or function that transforms data into an estimate ([Neal, 2020](#)). *Estimates* are numerical values that approximate the estimand derived through the process of *estimation*, which integrates data with an estimator ([Everitt and](#)



Skron dal, 2010). The Identification-Estimation flowchart (McElreath, 2020; Neal, 2020), shown in Figure 9, visually represents the transition from estimands to estimates.




Figure 9: Identification-Estimation flowchart. Extracted and slightly modified from Neal (2020, p. 32)

Identification is a necessary condition to ensure *consistent* estimators. An estimator achieves *consistency* when it converges to the “true” value of an estimand as the data size approaches infinity (Everitt and Skron dal, 2010). Without identification, researchers cannot achieve consistency, even with “infinite” and error-free data. As a result, deriving meaningful insights about an estimand from finite data becomes impossible (Schuessler and Selb, 2023, p. 5). Therefore, to ensure accurate and reliable estimates, researchers prioritize estimators with desirable identification properties. For instance, the Z-test is a widely used estimator for comparing group proportions, yielding accurate estimates when its underlying assumptions are satisfied (Kanji, 2006). Furthermore, researchers can interpret estimates from the Z-test as causal, provided the data is collected through a randomized experiment.

Randomized experiments are widely recognized as the gold standard in evidence-based science (Hariton and Locascio, 2018; Hansson, 2014). This recognition stems from their ability to enable researchers interpret associational estimates as causal. They achieve this by ensuring data, and by extension an estimator, satisfies several key identification properties, such as common support, no interference, and consistency (Morgan and Winship, 2014; Neal, 2020). The most critical property, however, is the elimination of confounding. *Confounding* occurs when an external variable  $X$  simultaneously influences the outcome  $Y$  and the variable of interest  $T$ , resulting in spurious associations (Everitt and Skron dal, 2010). Randomization addresses this issue by decoupling the association between the intervention allocation  $T$  from any other variable  $X$  (Morgan and Winship, 2014; Neal, 2020).

Nevertheless, researchers often face constraints that limit their ability to conduct randomized

experiments. These constraints include ethical concerns, such as the assignment of individuals to potentially harmful interventions, and practical limitations, such as the infeasibility of, for example, assigning individuals to genetic modifications or physical impairments (Neal, 2020). In these cases, causal inference offers a valuable alternative for generating causal estimates and understanding the mechanisms underlying specific data. In addition, the framework can provide significant theoretical insights that can enhance the design of experimental and observational studies (McElreath, 2020).

### 8.1.2. Identification under causal inference

Unlike classical statistical modeling, which focuses primarily on summarizing data and inferring associations, the *causal inference* framework is designed to identify causes and estimate their effects using data (Shaughnessy et al., 2010; Neal, 2020). The framework uses rigorous mathematical techniques to address the *fundamental problem of causality* (Pearl, 2009; Pearl et al., 2016; Morgan and Winship, 2014). This problem revolves around the question, “What would have happened ‘in the world’ under different circumstances?” This question introduces the concept of counterfactuals, which are instrumental in defining and identifying causal effects.

*Counterfactuals* are hypothetical scenarios that are *contrary to fact*, where alternative outcomes resulting from a given cause are neither observed nor observable (Neal, 2020; Counterfactual, 2024). The structural approach to causal inference (Pearl, 2009; Pearl et al., 2016) provides a formal framework for defining counterfactuals. For instance, in the scenario described in Section 8.1.1, the approach begins by defining the *individual causal effect* (ICE) as the difference between each student’s potential outcomes, as in Equation 18.

$$\tau_i = Y_i \mid do(T_i = 1) - Y_i \mid do(T_i = 2) \quad (18)$$

where  $do(T_i = t)$  represents the intervention operator,  $Y_i \mid do(T_i = 1)$  represents the potential outcome under intervention  $T_i = 1$ , and  $Y_i \mid do(T_i = 2)$  represents the potential outcome under intervention  $T_i = 2$ . Here, an *intervention* involves assigning a constant value to the treatment variable for each student’s potential outcomes. Note that if a student is assigned to intervention  $T_i = 1$ , the potential outcome under  $T_i = 2$  becomes a counterfactual, as it is no longer observed nor observable. To address this challenge, the structural approach extends the ICE to the *average causal effect* (ACE, Equation 19), representing the average difference between the students’ observed potential outcomes and their counterfactual counterparts.

$$\begin{aligned}
\tau &= E[\tau_i] \\
&= E[Y_i \mid do(T_i = 1)] - E[Y_i \mid do(T_i = 2)]
\end{aligned} \tag{19}$$

Even though counterfactuals are unobservable, researchers can still identify the ACE from associational estimates by leveraging the structural approach. The approach identifies the ACE by statistically conditioning data on a *sufficient adjustment set* of variables  $X$  (Pearl, 2009; Pearl et al., 2016; Morgan and Winship, 2014). This *sufficient* set (potentially empty) must block all non-causal paths between  $T$  to  $Y$  without opening new ones. When such a set exists, then  $T$  and  $Y$  are *d-separated* by  $X$  ( $T \perp Y \mid X$ ) (Pearl, 2009), and  $X$  satisfies the *backdoor criterion* (Neal, 2020, pp 37). Here, *conditioning* describes the process of restricting the focus to the subset of the population defined by the conditioning variable (Neal, 2020, p. 32) (see Equation 20).

Conditioning on a sufficient adjustment set enables researchers to estimate the ACE, even when the data comes from an observational study. This process is feasible because such conditioning ensures that the ACE estimator satisfies several critical properties, including confounding elimination (Morgan and Winship, 2014). Naturally, the validity of claims about the causal effects of  $T$  on  $Y$  now hinges on the assumption that  $X$  serves as a sufficient adjustment set. However, as Kohler et al. (2019, p. 150) noted, drawing conclusions about the real world from observed data inevitably requires assumptions. This requirement holds true for both observational and experimental data.

For instance, if researchers cannot conduct the randomized experiments described in Section 8.1.1 and must instead rely on observational data, they can still identify the ACE as long as an observed variable  $X$ , such as the socio-economic status of the school, satisfies the backdoor criterion. Under these circumstances, researchers first identify the *conditional average causal effect* (CACE, Equation 20)

$$CACE_t = E[Y_i \mid T_i = t, X] \tag{20}$$

From the CACE, researchers can identify the ACE from associational quantities as in Equation 21. This identification process is commonly known as the *backdoor adjustment*. Here,  $E_X[\cdot]$  represents the marginal expected value over  $X$  (Morgan and Winship, 2014).

$$\begin{aligned}
\tau &= E[Y_i \mid do(T_i = 1)] - E[Y_i \mid do(T_i = 2)] \\
&= E_X[CACE_1 - CACE_2] \\
&= E_X[E[Y_i \mid T_i = 1, X] - E[Y_i \mid T_i = 2, X]]
\end{aligned} \tag{21}$$

Notably, the approach extends the ACE identification for a continuous variable  $T$  as in Equation 22, ensuring broad applicability across different causal scenarios (Neal, 2020, p. 45)

$$\begin{aligned}
\tau &= E[Y_i \mid do(T_i = t)] \\
&= dE_X[E[Y_i \mid T_i = t, X]] / dt
\end{aligned} \tag{22}$$

### 8.1.3. Diving into the specifics

The structural approach to causal inference uses SCMs and DAGs to formally and graphically represent the presumed causal structure underlying the ACE (Pearl, 2009; Pearl et al., 2016; Gross et al., 2018; Neal, 2020). Essentially, these tools serve as *conceptual (theoretical) models* on which identification analysis rests (Schuessler and Selb, 2023, p. 4). Thus, using these tools, researchers can determine which statistical models can identify (ACE, CACE, or other), assuming the depicted causal structure is correct (McElreath, 2020), thus enabling valid causal inference. Figure 9 shows the role of theoretical models in the inference process.

SCMs and DAGs support identification analysis through two key advantages. First, regardless of complexity, they can represent various causal structures using only five fundamental building blocks (Neal, 2020; McElreath, 2020). This feature allows researchers to decompose complex structures into manageable components, facilitating their analysis (McElreath, 2020). Second, they depict causal relationships in a non-parametric and fully interactive way. This flexibility enables feasible ACE identification strategies without defining the variables' data types, the functional form between them, or their parameters (Pearl et al., 2016, p. 35).

Thus, Section 8.1.3.1 and Section 8.1.3.2 elaborate on the first advantage, while Section 8.1.3.2 and Section 8.1.3.3 do so for the second. Finally, Section 8.1.3.4 explains how researchers use SCMs and DAGs alongside Bayesian inference methods in the estimation process.

#### 8.1.3.1. The five fundamental block for SCMs and DAGs.

Figures 10, 11, 12, 13, and 14 display the five fundamental building blocks for SCMs and DAGs. The left panels of the figures show the formal mathematical models, represented by the SCMs,

defined in terms of a set of *endogenous* variables  $V = \{X_1, X_2, X_3\}$ , a set of *exogenous* variables  $E = \{e_{X_1}, e_{X_2}, e_{X_3}\}$ , and a set of functions  $F = \{f_{X_1}, f_{X_2}, f_{X_3}\}$  (Pearl, 2009; Cinelli et al., 2020). Endogenous variables are those whose causal mechanisms a researcher chooses to model (Neal, 2020). In contrast, exogenous variables represent *errors* or *disturbances* arising from omitted factors that the investigator chooses not to model explicitly (Pearl, 2009, p. 27,68). Lastly, the functions, referred to as *structural equations*, express the endogenous variables as non-parametric functions of other variables. These functions use the symbol ‘ $:=$ ’ to denote the asymmetrical causal dependence of the variables and the symbol ‘ $\perp$ ’ to represent *d-separation*, a concept akin to (conditional) independence.

Notably, every SCM has an associated DAG (Pearl et al., 2016; Cinelli et al., 2020). The right panels of the figures display these DAGs. A DAG is a graph consisting of nodes connected by edges, where the nodes represent random variables. The term *directed* means that the edges extend from one node to another, with arrows indicating the direction of causal influence. The term *acyclic* implies that the causal influences do not form loops, ensuring the influences do not cycle back on themselves (McElreath, 2020). DAGs represent observed variables as solid black circles, while they use open circles for unobserved (latent) variables (Morgan and Winship, 2014). Although the *standard representation* of DAGs typically omits exogenous variables for simplicity, the *magnified representation* depicted in the figures offers one key advantage: including exogenous variables can help researchers highlight potential issues related to conditioning and confounding (Cinelli et al., 2020).




Figure 10: Two unconnected nodes

$$X_1 := f_{X_1}(e_{X_1})$$

$$X_3 := f_{X_3}(X_1, e_{X_3})$$

$$e_{X_1} \perp e_{X_3}$$

(a) SCM



(b) DAG

Figure 11: Two connected nodes or descendant

$$X_1 := f_{X_1}(e_{X_1})$$

$$X_2 := f_{X_2}(X_1, e_{X_2})$$


$$X_3 := f_{X_3}(X_2, e_{X_3})$$

$$e_{X_1} \perp e_{X_2}$$

$$e_{X_1} \perp e_{X_3}$$

$$e_{X_2} \perp e_{X_3}$$

(a) SCM



(b) DAG

Figure 12: Chain or mediator

$$X_1 := f_{X_1}(X_2, e_{X_1})$$

$$X_2 := f_{X_2}(e_{X_2})$$


$$X_3 := f_{X_3}(X_2, e_{X_3})$$

$$e_{X_1} \perp e_{X_2}$$

$$e_{X_1} \perp e_{X_3}$$

$$e_{X_2} \perp e_{X_3}$$

(a) SCM



(b) DAG

Figure 13: Fork or confounder

$$X_1 := f_{X1}(e_{X1})$$

$$X_2 := f_{X2}(X_1, X_3, e_{X2})$$


$$X_3 := f_{X3}(e_{X3})$$

$$e_{X1} \perp e_{X2}$$

$$e_{X1} \perp e_{X3}$$

$$e_{X2} \perp e_{X3}$$

(a) SCM



(b) DAG

Figure 14: Collider or immorality

A careful examination of these building blocks highlights the theoretical assumptions underlying their observed variables. SCM 10a and DAG 10b depict two unconnected nodes, representing a scenario where variables  $X_1$  and  $X_3$  are independent or not causally related. SCM 11a and DAG 11b illustrate two connected nodes, representing a scenario where a *parent* node  $X_1$  exerts a causal influence on a *child* node  $X_3$ . In this setup,  $X_3$  is considered a *descendant* of  $X_1$ . Additionally,  $X_1$  and  $X_3$  are described as *adjacent* because there is a *direct path* connecting them. SCM 12a and DAG 12b depict a *chain*, where  $X_1$  influences  $X_2$ , and  $X_2$  influences  $X_3$ . In this configuration,  $X_1$  is a parent node of  $X_2$ , which is a parent node of  $X_3$ . This structure creates a *directed path* between  $X_1$  and  $X_3$ . Consequently,  $X_1$  is an *ancestor* of  $X_3$ , and  $X_2$  fully *mediates* the relationship between the two. SCM 13a and DAG 13b illustrate a *fork*, where variables  $X_1$  and  $X_3$  are both influenced by  $X_2$ . Here,  $X_2$  is a parent node that *confounds* the relationship between  $X_1$  and  $X_3$ . Finally, SCM 14a and DAG 14b show a *collider*, where variables  $X_1$  and  $X_3$  are concurrent causes of  $X_2$ . In this configuration,  $X_1$  and  $X_3$  are not causally related to each other but both influence  $X_2$  (an *immorality*). Notably, all building blocks assume the errors are independent of each other and from all other variables in the graph, as evidenced by the pairwise relations  $e_{X1} \perp e_{X2}$ ,  $e_{X1} \perp e_{X3}$ , and  $e_{X2} \perp e_{X3}$ .

Researchers can then use these building blocks to represent the scenario outlined in Section 8.1.2. SCM 15a and DAG 15b depict the plausible causal structure for this example. In this context, the variable  $X$  (socio-economic status of the school) is thought to be a confounder in the relationship between the teaching method  $T$  and the outcome  $Y$ . The figures display multiple descendant relationships such as  $X \rightarrow T$ ,  $X \rightarrow Y$ , and  $T \rightarrow Y$ . They also highlight unconnected node pairs,

evident from the relationships  $e_T \perp e_X$ ,  $e_T \perp e_Y$ , and  $e_X \perp e_Y$ . Additionally, the figures show one fork,  $X \rightarrow \{T, Y\}$ , and two colliders:  $\{X, e_T\} \rightarrow T$  and  $\{X, T, e_Y\} \rightarrow Y$ .




Figure 15: Plausible causal structure the scenario outlined in Section 8.1.2.

#### 8.1.3.2. The probabilistic implications of these blocks.

Beyond their graphical capabilities, researchers can translate SCMs and DAGs into statistical models because these conceptual models encode functional and probabilistic information, which they can then replace with suitable functions and probabilistic assumptions (Pearl et al., 2016). Notably, this encoding relies on two fundamental assumptions: the causal Markov and the faithfulness assumption. The *causal Markov assumption* links the graph structure to the joint probability distribution of variables by asserting that any variable in a graph is independent of all its non-descendants conditional on its parents (Neal, 2020, p. 20; Hernán and Robins, 2025, p. 80). The *faithfulness assumption*, in turn, states that the observed statistical (in)dependencies in the data reflect the underlying causal structure among variables—except in rare degenerate cases (Neal, 2020, p. 100; Hernán and Robins, 2025, p. 81).

A notable implication of the assumptions underlying the probabilistic encoding is that any conceptual model described by an SCM and DAG can represent the joint distribution of variables more efficiently (Pearl et al., 2016, p. 29). This expression takes the form of a product of conditional probability distributions (CPDs) of the type  $P(\text{child} \mid \text{parents})$ . This property is formally known as the *Bayesian Network factorization* (BNF, Equation 23) (Pearl et al., 2016, p. 29; Neal, 2020,



p. 21). In this expression,  $pa(X_i)$  denotes the set of variables that are the parents of  $X_i$ .

$$\begin{aligned} P(X_1, X_2, \dots, X_P) &= X_1 \cdot \prod_{p=2}^P P(X_i \mid X_{i-1}, \dots, X_1) \quad (\text{by chain rule}) \\ &= X_1 \cdot \prod_{p=2}^P P(X_i \mid pa(X_i)) \quad (\text{by BNF}) \end{aligned} \tag{23}$$

This encoding enables researchers with conceptual (theoretical) knowledge in the form of an SCM and DAG to predict patterns of (in)dependencies in the data. As highlighted by [Pearl et al. \(2016, p. 35\)](#), these predictions depend solely on the structure of these conceptual models without requiring the quantitative details of the equations or the distributions of the errors. Moreover, once researchers observe empirical data, the patterns of (in)dependencies in the data can provide significant insights into the validity of the proposed conceptual model.

The five fundamental building blocks described in Section 8.1.3.1 clearly illustrate which (in)dependencies can SMCs and DAGs predict. For instance, applying the BNF to the causal structure shown in the SCM 10a and DAG 10b enables researchers to express the joint probability distribution of the observed variables as  $P(X_1, X_3) = P(X_1)P(X_3)$ , supporting the theoretical assumption that the observed variables  $X_1$  and  $X_3$  are unconditionally independent ( $X_1 \perp X_3$ ) ([Neal, 2020, p. 24](#)). Conversely, when  $X_3$  is unconditionally dependent on  $X_1$  ( $X_1 \not\perp X_3$ ), as depicted in the SCM 11a and DAG 11b, the BNF express their joint probability distribution as  $P(X_1, X_3) = P(X_3 \mid X_1)P(X_1)$ . Notably, these descriptions demonstrate the clear correspondence between the structural equations illustrated in Section 8.1.3.1 and the CPDs.

Beyond the insights gained from two-node structures, researchers can uncover more nuanced patterns of (in)dependencies from chains, forks, and colliders. These (in)dependencies apply to any data set generated by a causal model with those structures, regardless of the specific functions attached to the SCM ([Pearl et al., 2016, p. 36](#)). For instance, applying the BNF to the chain structure depicted in the SCM 12a and DAG 12b allow researchers to represent the joint distribution for the observed variables as  $P(X_1, X_2, X_3) = P(X_1)P(X_2 \mid X_1)P(X_3 \mid X_2)$ . This expression implies that  $X_1$  and  $X_3$  are unconditionally dependent ( $X_1 \not\perp X_3$ ), but conditionally independent when controlling for  $X_2$  ( $X_1 \perp X_3 \mid X_2$ ). Moreover, in the fork structure shown in the SCM 13a and DAG 13b, researchers can express the joint distribution of the observed variables as  $P(X_1, X_2, X_3) = P(X_1 \mid X_2)P(X_2)P(X_3 \mid X_2)$ . Similar to the chain structure, this expression allows researchers to further infer that  $X_1$  and  $X_3$  are unconditionally dependent ( $X_1 \not\perp X_3$ ), but conditionally independent when controlling for  $X_2$  ( $X_1 \perp X_3 \mid X_2$ ). Finally, researchers analyzing

the collider structure illustrated in the SCM 14a and DAG 14b can express the joint distribution of the observed variables as  $P(X_1, X_2, X_3) = P(X_1)P(X_2 | X_1, X_3)P(X_3)$ . This representation allows researchers to infer that  $X_1$  and  $X_3$  are unconditionally independent ( $X_1 \perp X_3$ ), but conditionally dependent when controlling for  $X_2$  ( $X_1 \not\perp X_3 | X_2$ ). The authors Pearl et al. (2016, p. 37, 40, 41) and Neal (2020, p. 25–26) provide the mathematical proofs for these conclusions.

Using these additional probabilistic insights, researchers can revisit the scenario in Section 8.1.2. In this context, applying the BNF to the SCM 16a structure, enables the representation of the joint probability distribution of the observed variables as  $P(Y, T, X) = P(Y | T, X)P(T | X)P(X)$ . From this expression, researchers can infer that the outcome  $Y$  is unconditionally dependent on the teaching method  $T$  ( $Y \not\perp T$ ). This dependency arises from two key structures: a direct causal path from the teaching method  $T$  to the outcome  $Y$ , represented by the two-connected-nodes structure  $T \rightarrow Y$  (black path in DAG 16b), and a confounding non-causal path from the teaching method  $T$  to the outcome  $Y$  through the socio-economic status of the school  $X$ , represented by the fork structure  $T \leftarrow X \rightarrow Y$  (gray path in DAG 16b).

$$X := x$$

$$T := f_T(x, e_T)$$


$$Y := f_Y(T, x, e_Y)$$

$$e_T \perp e_X$$

$$e_T \perp e_Y$$

$$e_X \perp e_Y$$

(a) SCM



(b) Conditioned DAG

Figure 16: Plausible causal structure the scenario outlined in Section 8.1.2.

### 8.1.3.3. From probability to causality.

The structural approach to causal inference translates probabilistic insights into actionable strategies seeking to identify the ACE from associational quantities. The approach achieves this by relying on the *modularity assumption*, which posits that intervening on a node alters only the causal mechanism of that node, leaving others unchanged (Neal, 2020, p. 34).

The modularity assumption underpins the concepts of manipulated graphs and Truncated Factorization, which are essential for representing interventions  $P(Y_i | do(T_i = t))$  within SCMs and

DAGs. *Manipulated graphs* simulate physical interventions by removing specific edges from a DAG, while preserving the remaining structure unchanged (Neal, 2020, p. 34). In parallel, *Truncated Factorization* (TF) achieves a similar simulation by removing specific functions from the conceptual model and replacing them with constants, while keeping the rest of the structure unchanged (Pearl, 2010). The probabilistic implications of this factorization are formalized in Equation 24, where  $S$  represents the subset of variables  $X_p$  directly influenced by the intervention, while an example illustrating these concepts follows below.

$$P(X_1, X_2, \dots, X_P | do(S)) = \begin{cases} \prod P(X_p | pa(X_p)) & \text{if } p \notin S \\ 1 & \text{otherwise} \end{cases} \quad (24)$$

Using the TF, researchers can define the *backdoor adjustment* to identify the ACE. This adjustment states that if a variable  $X_p \in S$  serves as a *sufficient adjustment set* for the effect of  $X_a$  on  $X_b$ , then the ACE can be identified using Equation 25. The sufficient adjustment set (potentially empty) must block all non-causal paths between  $X_a$  and  $X_b$  without introducing new paths. If such a set exists, then  $X_a$  and  $X_b$  are *d-separated* by  $X_p$  ( $X_a \perp X_b | X_p$ ) (Pearl, 2009), and  $X_p$  satisfies the *backdoor criterion* (Neal, 2020, p. 37).

$$P(X_a | do(X_b = x)) = \sum_{X_p} P(X_a | X_b = x, X_p) P(X_p) \quad (25)$$

Ultimately, the backdoor adjustment enables researchers to express the ACE as:

$$\begin{aligned} \tau &= E[X_a | do(X_b = 1)] - E[X_a | do(X_b = 2)] \\ &= E_{X_p} [E[X_a | do(X_b = 1), X_p] - E[X_a | do(X_b = 2), X_p]] \\ &= \sum_{X_p} X_a \cdot P(X_a | X_b = 1, X_p) \cdot P(X_p) - \sum_{X_p} X_a \cdot P(X_a | X_b = 2, X_p) \cdot P(X_p) \end{aligned} \quad (26)$$

With these new insights, researchers revisiting the scenario in Section 8.1.3.2 can infer that the socio-economic status of the school,  $X$ , satisfies the backdoor criterion, assuming the causal structure depicted by the SCM 16a and DAG 16b is correct. This means that  $X$  serves as a sufficient adjustment set, as it effectively blocks all confounding non-causal paths introduced by the fork structure. Nevertheless, since  $Y$  remains dependent on  $T$  even after conditioning ( $Y \not\perp T | X$ ), this dependency can only be attributed to the direct causal effect  $T \rightarrow Y$ . Notably, for the purpose of

identification, the conditioned DAG 16b is equivalent to the manipulated DAG 17b, because  $X$  satisfies the backdoor criterion.




Figure 17: Plausible causal structure the scenario outlined in Section 8.1.3.2.

Researchers can then apply the *backdoor adjustment* to identify the ACE of  $T$  on  $Y$ . They achieve this by first identifying the CACE of  $T$  on  $Y$  by conditioning on  $X$ , and then marginalizing this effect over  $X$  to obtain the ACE. This process is expressed in Equation 27 (see Section 8.1.2).

$$\begin{aligned}
\tau &= E[Y_i \mid do(T_i = 1)] - E[Y_i \mid do(T_i = 2)] \\
&= E_X [E[Y_i \mid T_i = 1, X] - E[Y_i \mid T_i = 2, X]] \\
&= \sum_X Y_i \cdot P(Y_i \mid T_i = 1, X) \cdot P(X) - \sum_X Y_i \cdot P(Y_i \mid T_i = 2, X) \cdot P(X)
\end{aligned} \tag{27}$$

#### 8.1.3.4. The estimation process.

Ultimately, researchers can use Bayesian inference methods to estimate the ACE. The approach begins by defining two probability distributions: the likelihood of the data,  $P(X_1, X_2, \dots, X_P \mid \theta)$ , and the prior distribution,  $P(\theta)$  (Everitt and Skrongdal, 2010), where  $X_P$  represents a random variable, and  $\theta$  represents a one-dimensional parameter space for simplicity. After observing empirical data, researchers can update the priors to posterior distributions using Bayes' rule in Equation 28:

$$P(\theta \mid X_1, X_2, \dots, X_P) = \frac{P(X_1, X_2, \dots, X_P \mid \theta) \cdot P(\theta)}{P(X_1, X_2, \dots, X_P)} \tag{28}$$

Given that the denominator on the right-hand side of Equation 28 serves as a normalizing constant independent of the parameter  $\theta$ , researchers can simplify the posterior updating process into three

steps. First, they integrate new empirical data through the likelihood. Second, they update the parameters' priors to a posterior distribution according to Equation 29. Ultimately, they normalize these results to obtain a valid probability distribution.

$$P(\theta \mid X_1, X_2, \dots, X_P) \propto P(X_1, X_2, \dots, X_P \mid \theta) \cdot P(\theta) \quad (29)$$

Temporarily setting aside the definition of prior distributions  $P(\theta)$ , note that the posterior updating process depends heavily on the assumptions underlying the likelihood of the data. However, as the number of random variables,  $P$ , increases, this joint distribution quickly becomes intractable (Neal, 2020). This intractability is evident from Equation 30, where the likelihood distribution is expressed by multiple chained CPDs.

$$P(X_1, X_2, \dots, X_P \mid \theta) = P(X_1 \mid \theta) \prod_{p=2}^P P(X_p \mid X_{p-1}, \dots, X_1, \theta) \quad (30)$$

Nevertheless, researchers can manage the complexity of the likelihood by assuming specific local (in)dependencies among variables. SCMs and DAGs provide a formal framework to represent these assumptions, as detailed in Section 8.1.3.2. These assumptions improve model tractability and simplify the estimation process by enabling the derivation of the BNF of the likelihood (Equation 31). With this simplified structure, any probabilistic programming language can model the system and compute the parameter's posterior distribution using Equation 28.

$$P(X_1, X_2, \dots, X_P \mid \theta) = P(X_1 \mid \theta) \prod_{p=2}^P P(X_p \mid pa(X_p), \theta) \quad (31)$$

## References

- Ackerman, T., 1989. Unidimensional irt calibration of compensatory and noncompensatory multidimensional items. *Applied Psychological Measurement* 13, 113–127. doi:[10.1177/014662168901300201](https://doi.org/10.1177/014662168901300201).
- Agresti, A., 2015. Foundations of linear and generalized linear models. Wiley series in probability and statistics, John Wiley & Sons.
- Andrich, D., 1978. Relationships between the thurstone and rasch approaches to item scaling. *Applied Psychological Measurement* 2, 451–462. doi:[10.1177/014662167800200319](https://doi.org/10.1177/014662167800200319).
- Attia, J., Holliday, E., Oldmeadow, C., 2022. A proposal for capturing interaction and effect modification using dags. *International Journal of Epidemiology* 51, 1047–1053. doi:[10.1093/ije/dyac126](https://doi.org/10.1093/ije/dyac126).
- Baker, F., 1998. An investigation of the item parameter recovery characteristics of a gibbs sampling procedure. *Applied Psychological Measurement* 22, 153–169. doi:[10.1177/01466216980222005](https://doi.org/10.1177/01466216980222005).
- Baldwin, S., Fellingham, G., 2013. Bayesian methods for the analysis of small sample multilevel data with a complex variance structure. *Journal of Psychological Methods* 18, 151–164. doi:[10.1037/a0030642](https://doi.org/10.1037/a0030642).
- Bartholomew, S., Nadelson, L., Goodridge, W., Reeve, E., 2018. Adaptive comparative judgment as a tool for assessing open-ended design problems and model eliciting activities. *Educational Assessment* 23, 85–101. doi:[10.1080/10627197.2018.1444986](https://doi.org/10.1080/10627197.2018.1444986).
- Bartholomew, S., Williams, P., 2020. Stem skill assessment: An application of adaptive comparative judgment, in: Anderson, J., Li, Y. (Eds.), *Integrated Approaches to STEM Education. Advances in STEM Education*. Springer, pp. 331–349. doi:[10.1007/978-3-030-52229-2\\_18](https://doi.org/10.1007/978-3-030-52229-2_18).
- Bartholomew, S., Yoshikawa, E., Hartell, E., Strimel, G., 2020. Identifying design values across countries through adaptive comparative judgment. *International Journal of Technology and Design Education* doi:[10.1007/s10798-019-09506-8](https://doi.org/10.1007/s10798-019-09506-8).
- Boonen, N., Kloots, H., Gillis, S., 2020. Rating the overall speech quality of hearing-impaired children by means of comparative judgements. *Journal of Communication Disorders* 83, 1675–1687. doi:[10.1016/j.jcomdis.2019.105969](https://doi.org/10.1016/j.jcomdis.2019.105969).
- Bouwer, R., Lesterhuis, M., De Smedt, F., Van Keer, H., De Maeyer, S., 2023. Comparative approaches to the assessment of writing: Reliability and validity of benchmark rating and comparative judgement. *Journal of Writing Research* 15, 497–518. doi:[10.17239/jowr-2024.15.03.03](https://doi.org/10.17239/jowr-2024.15.03.03).
- Bradley, R., Terry, M., 1952. Rank analysis of incomplete block designs: I. the method of paired comparisons. *Biometrika* 39, 324–345. doi:[10.2307/2334029](https://doi.org/10.2307/2334029).
- Bramley, T., 2008. Paired comparison methods, in: Newton, P., Baird, J., Goldsteing, H., Patrick, H., Tymms, P. (Eds.), *Techniques for monitoring the comparability of examination standards*. GOV.UK., pp. 246–300. URL: <https://assets.publishing.service.gov.uk/media/5a80d75940f0b62305b8d734/2007-comparability-exam-standards-i-chapter7.pdf>.
- Bramley, T., 2015. Investigating the reliability of adaptive comparative judgment. URL: <http://www.cambridgeassessment.org.uk/Images/232694-investigating-the-reliability-of-adaptive-comparative-judgment.pdf>. cambridge Assessment Research Report.
- Bramley, T., Vitello, S., 2019. The effect of adaptivity on the reliability coefficient in adaptive comparative judgement. *Assessment in Education: Principles, Policy & Practice* 26, 43–58. doi:[10.1080/0969594X.2017.1418734](https://doi.org/10.1080/0969594X.2017.1418734).
- Casalicchio, G., Tutz, G., Schauberger, G., 2015. Subject-specific bradley–terry–luce models with implicit variable selection. *Statistical Modelling* 15, 526–547. doi:[10.1177/1471082X15571817](https://doi.org/10.1177/1471082X15571817).
- Chambers, L., Cunningham, E., 2022. Exploring the validity of comparative judgement: Do judges attend to

- construct-irrelevant features? *Frontiers in Education* doi:[10.3389/feduc.2022.802392](https://doi.org/10.3389/feduc.2022.802392).
- Cinelli, C., Forney, A., Pearl, J., 2020. A crash course in good and bad controls. SSRN URL: <https://ssrn.com/abstract=3689437>, doi:[10.2139/ssrn.3689437](https://doi.org/10.2139/ssrn.3689437).
- Coertjens, L., Lesterhuis, M., Verhavert, S., Van Gasse, R., De Maeyer, S., 2017. Teksten beoordelen met criterialijsten of via paarsgewijze vergelijking: een afweging van betrouwbaarheid en tijdsinvestering. *Pedagogische Studien* 94, 283–303. URL: <https://repository.uantwerpen.be/docman/irua/e71ea9/147930.pdf>.
- Counterfactual, 2024. Merriam-webster.com dictionary. URL: <https://www.merriam-webster.com/dictionary/hacker>. retrieved July 23, 2024.
- Crompvoets, E., Béguin, A., Sijtsma, K., 2020. Adaptive pairwise comparison for educational measurement. *Journal of Educational and Behavioral Statistics* 45, 316–338. doi:[10.3102/1076998619890589](https://doi.org/10.3102/1076998619890589).
- Crompvoets, E., Béguin, A., Sijtsma, K., 2022. On the bias and stability of the results of comparative judgment. *Frontiers in Education* 6. doi:[10.3389/feduc.2021.788202](https://doi.org/10.3389/feduc.2021.788202).
- de Ayala, R., 2009. *The Theory and Practice of Item Response Theory*. Methodology in the Social Sciences, The Guilford Press.
- Deffner, D., Rohrer, J., McElreath, R., 2022. A causal framework for cross-cultural generalizability. *Advances in Methods and Practices in Psychological Science* 5. doi:[10.1177/25152459221106366](https://doi.org/10.1177/25152459221106366).
- Depaoli, S., 2014. The impact of inaccurate “informative” priors for growth parameters in bayesian growth mixture modeling. *Journal of Structural Equation Modeling* 21, 239–252. doi:[10.1080/10705511.2014.882686](https://doi.org/10.1080/10705511.2014.882686).
- Depaoli, S., 2021. *Bayesian Structural Equation Modeling*. Methodology in the social sciences, The Guilford Press.
- Everitt, B., Skrondal, A., 2010. *The Cambridge Dictionary of Statistics*. Cambridge University Press.
- Fox, J., 2010. *Bayesian Item Response Modeling, Theory and Applications*. Statistics for Social and Behavioral Sciences, Springer.
- Gelman, A., Carlin, J., Stern, H., Dunson, D., Vehtari, A., Rubin, D., 2014. *Bayesian Data Analysis*. Texts in Statistical Science. 3rd ed., Chapman and Hall/CRC.
- Gijzen, M., van Daal, T., Lesterhuis, M., Gijbels, D., De Maeyer, S., 2021. The complexity of comparative judgments in assessing argumentative writing: An eye tracking study. *Frontiers in Education* 5. doi:[10.3389/feduc.2020.582800](https://doi.org/10.3389/feduc.2020.582800).
- Gill, T., Bramley, T., 2013. How accurate are examiners’ holistic judgements of script quality? *Assessment in Education: Principles, Policy and Practice* 20, 308–324. doi:[10.1080/0969594X.2013.779229](https://doi.org/10.1080/0969594X.2013.779229).
- Goossens, M., De Maeyer, S., 2018. How to obtain efficient high reliabilities in assessing texts: Rubrics vs comparative judgement, in: Ras, E., Guerrero Roldán, A. (Eds.), *Technology Enhanced Assessment*, Springer International Publishing. pp. 13–25. doi:[10.1007/978-3-319-97807-9\\_2](https://doi.org/10.1007/978-3-319-97807-9_2).
- Gray, A., Rahat, A., Crick, T., Lindsay, S., 2024. A bayesian active learning approach to comparative judgement within education assessment. *Computers and Education: Artificial Intelligence* 6, 100–245. URL: <https://www.sciencedirect.com/science/article/pii/S2666920X24000481>, doi:[10.1016/j.caeai.2024.100245](https://doi.org/10.1016/j.caeai.2024.100245).
- Gross, J., Yellen, J., Anderson, M., 2018. *Graph Theory and Its Applications*. Textbooks in Mathematics, Chapman and Hall/CRC. doi:<https://doi.org/10.1201/9780429425134>. 3rd edition.
- Grubbs, F., 1969. Procedures for detecting outlying observations in samples. *Technometrics* 11, 1–21. URL: <https://www.tandfonline.com/doi/abs/10.1080/00401706.1969.10490657>, doi:[10.1080/00401706.1969.10490657](https://doi.org/10.1080/00401706.1969.10490657).
- Hansson, S., 2014. Why and for what are clinical trials the gold standard? *Scandinavian Journal of Public Health* 42, 41–48. doi:[10.1177/1403494813516712](https://doi.org/10.1177/1403494813516712). PMID: 24553853.
- Hariton, E., Locascio, J., 2018. Randomised controlled trials – the gold standard for effectiveness research. *BJOG*:

- An International Journal of Obstetrics & Gynaecology 125, 1716–1716. URL: <https://obgyn.onlinelibrary.wiley.com/doi/abs/10.1111/1471-0528.15199>, doi:10.1111/1471-0528.15199.
- Hernán, M., Robins, J., 2025. Causal Inference: What If. 1 ed., Chapman and Hall/CRC. URL: [https://miguelhernan.org/s/hernanrobins\\_WhatIf\\_27may25.pdf](https://miguelhernan.org/s/hernanrobins_WhatIf_27may25.pdf). last accessed 30 may 2025.
- Hoyle, R.e., 2023. Handbook of Structural Equation Modeling. Guilford Press.
- Jones, I., Bisson, M., Gilmore, C., Inglis, M., 2019. Measuring conceptual understanding in randomised controlled trials: Can comparative judgement help? British Educational Research Journal 45, 662–680. doi:10.1002/berj.3519.
- Jones, I., Inglis, M., 2015. The problem of assessing problem solving: can comparative judgement help? Educational Studies in Mathematics 89, 337–355. doi:10.1007/s10649-015-9607-1.
- Kanji, G., 2006. 100 Statistical Tests. Introduction to statistics, SAGE Publications.
- Kelly, K., Richardson, M., Isaacs, T., 2022. Critiquing the rationales for using comparative judgement: a call for clarity. Assessment in Education: Principles, Policy & Practice 29, 674–688. doi:10.1080/0969594X.2022.2147901.
- Kim, S., Cohen, A., 1999. Accuracy of parameter estimation in gibbs sampling under the two-parameter logistic model. URL: <https://eric.ed.gov/?id=ED430012>. annual Meeting of the American Educational Research Association.
- Kimbell, R., 2012. Evolving project e-scape for national assessment. International Journal of Technology and Design Education 22, 135–155. doi:10.1007/s10798-011-9190-4.
- Kline, R., 2023. Principles and Practice of Structural Equation Modeling. Methodology in the Social Sciences, Guilford Press.
- Kohler, U., Kreuter, F., Stuart, E., 2019. Nonprobability sampling and causal analysis. Annual Review of Statistics and Its Application 6, 149–172. URL: <https://www.annualreviews.org/content/journals/10.1146/annurev-statistics-030718-104951>, doi:https://doi.org/10.1146/annurev-statistics-030718-104951.
- Lambert, P., Sutton, A., Burton, P., Abrams, K., Jones, D., 2006. How vague is vague? a simulation study of the impact of the use of vague prior distributions in mcmc using winbugs. Journal of Statistics in Medicine 24, 2401–2428. doi:10.1002/sim.2112.
- Lawson, J., 2015. Design and Analysis of Experiments with R. Chapman and Hall/CRC.
- Lee, Y., Nelder, J.A., 1996. Hierarchical generalized linear models. Journal of the Royal Statistical Society: Series B (Methodological) 58, 619–656. URL: <https://rss.onlinelibrary.wiley.com/doi/abs/10.1111/j.2517-6161.1996.tb02105.x>, doi:10.1111/j.2517-6161.1996.tb02105.x.
- Lesterhuis, M., 2018. The validity of comparative judgement for assessing text quality: An assessor's perspective. Ph.D. thesis. University of Antwerp. URL: <https://hdl.handle.net/10067/1548280151162165141>.
- Lesterhuis, M., van Daal, T., Van Gasse, R., Coertjens, L., Donche, V., De Maeyer, S., 2018. When teachers compare argumentative texts: Decisions informed by multiple complex aspects of text quality. L1-Educational Studies in Language and Literature 18, 1–22. doi:10.17239/L1ESLL-2018.18.01.02.
- Little, R., Rubin, D., 2020. Statistical analysis with missing data. Wiley Series in Probability and Statistics, John Wiley & Sons. doi:10.1002/9781119482260. third Edition.
- Luce, R., 1959. On the possible psychophysical laws. The Psychological Review 66, 482–499. doi:10.1037/h0043178.
- Marshall, N., Shaw, K., Hunter, J., Jones, I., 2020. Assessment by comparative judgement: An application to secondary statistics and english in new zealand. New Zealand Journal of Educational Studies 55, 49–71. doi:10.1007/s40841-020-00163-3.
- Martin, J., McDonald, R., 1975. Bayesian estimation in unrestricted factor analysis: A treatment for heywood cases. Psychometrika , 505–517doi:10.1007/BF02291552.



- McCullagh, P., Nelder, J., 1983. Generalized Linear Models. Monographs on Statistics and Applied Probability, Routledge. doi:[10.1201/9780203753736](https://doi.org/10.1201/9780203753736).
- McElreath, R., 2020. Statistical Rethinking: A Bayesian Course with Examples in R and STAN. Chapman and Hall/CRC. doi:<https://doi.org/10.1201/9780429029608>.
- McElreath, R., 2021. Science before statistics: Causal inference. <https://www.youtube.com/watch?v=KNPYUVmY3NM>. Last accessed 30 April 2024.
- McElreath, R., 2024. Statistical rethinking, 2024 course. URL: [https://github.com/rmcelreath/stat\\_rethinking\\_2024](https://github.com/rmcelreath/stat_rethinking_2024). last accessed 15 March 2025.
- McMahon, S., Jones, I., 2015. A comparative judgement approach to teacher assessment. Assessment in Education: Principles, Policy & Practice 22, 368–389. doi:[10.1080/0969594X.2014.978839](https://doi.org/10.1080/0969594X.2014.978839).
- Mikhailiuk, A., Wilmot, C., Perez-Ortiz, M., Yue, D., Mantiuk, R., 2021. Active sampling for pairwise comparisons via approximate message passing and information gain maximization, in: 2020 25th International Conference on Pattern Recognition (ICPR), pp. 2559–2566. doi:[10.1109/ICPR48806.2021.9412676](https://doi.org/10.1109/ICPR48806.2021.9412676).
- Miller, J., 2023. Outlier exclusion procedures for reaction time analysis: The cures are generally worse than the disease. Journal of Experimental Psychology: General 152, 3189–3217. doi:[10.1037/xge0001450](https://doi.org/10.1037/xge0001450).
- Morgan, S., Winship, C., 2014. Counterfactuals and Causal Inference: Methods and Principles for Social Research. Analytical Methods for Social Research. 2 ed., Cambridge University Press.
- Neal, B., 2020. Introduction to causal inference from a machine learning perspective. URL: [https://www.bradynear.com/Introduction\\_to\\_Causal\\_Inference-Dec17\\_2020-Neal.pdf](https://www.bradynear.com/Introduction_to_Causal_Inference-Dec17_2020-Neal.pdf). last accessed 30 April 2024.
- Neyman, J., 1923. On the application of probability theory to agricultural experiments. essay on principles. section 9. Statistical Science 5, 465–472. URL: <http://www.jstor.org/stable/2245382>. translated by Dabrowska, D. and Speed, T. (1990).
- O’Hagan, A., 2018. On outlier rejection phenomena in bayes inference. Journal of the Royal Statistical Society: Series B (Methodological) 41, 358–367. URL: [https://academic.oup.com/jrsssb/article-pdf/41/3/358/49097051/jrsssb\\_41\\_3\\_358.pdf](https://academic.oup.com/jrsssb/article-pdf/41/3/358/49097051/jrsssb_41_3_358.pdf), doi:[10.1111/j.2517-6161.1979.tb01090.x](https://doi.org/10.1111/j.2517-6161.1979.tb01090.x).
- Pearl, J., 2009. Causality: Models, Reasoning and Inference. Cambridge University Press.
- Pearl, J., 2010. An introduction to causal inference. The international journal of biostatistics 6, 855–859. URL: <https://www.degruyter.com/document/doi/10.2202/1557-4679.1203/html>, doi:[10.2202/1557-4679.1203](https://doi.org/10.2202/1557-4679.1203).
- Pearl, J., 2019. The seven tools of causal inference, with reflections on machine learning. Communications of the ACM 62, 54–60. doi:[10.1177/0962280215586010](https://doi.org/10.1177/0962280215586010).
- Pearl, J., Glymour, M., Jewell, N., 2016. Causal Inference in Statistics: A Primer. John Wiley & Sons, Inc.
- Pearl, J., Mackenzie, D., 2018. The Book of Why: The New Science of Cause and Effect. 1st ed., Basic Books, Inc.
- Perron, B., Gillespie, D., 2015. Reliability and Measurement Error, in: Key Concepts in Measurement. Oxford University Press. Pocket guides to social work research methods. chapter 4. doi:[10.1093/acprof:oso/9780199855483.003.0004](https://doi.org/10.1093/acprof:oso/9780199855483.003.0004).
- Pollitt, A., 2004. Let’s stop marking exams, in: Proceedings of the IAEA Conference, University of Cambridge Local Examinations Syndicate, Philadelphia. URL: <https://www.cambridgeassessment.org.uk/images/109719-let-s-stop-marking-exams.pdf>.
- Pollitt, A., 2012a. Comparative judgement for assessment. International Journal of Technology and Design Education 22, 157–170. doi:[10.1007/s10798-011-9189-x](https://doi.org/10.1007/s10798-011-9189-x).
- Pollitt, A., 2012b. The method of adaptive comparative judgement. Assessment in Education: Principles, Policy and

- Practice 19, 281—300. doi:[10.1080/0969594X.2012.665354](https://doi.org/10.1080/0969594X.2012.665354).
- Pollitt, A., Elliott, G., 2003. Finding a proper role for human judgement in the examination system. URL: <https://www.cambridgeassessment.org.uk/Images/109707-monitoring-and-investigating-comparability-a-proper-role-for-human-judgement.pdf>. research & Evaluation Division.
- Rohrer, J., 2018. Thinking clearly about correlations and causation: Graphical causal models for observational data. *Advances in Methods and Practices in Psychological Science* 1, 27–42. doi:[10.1177/2515245917745629](https://doi.org/10.1177/2515245917745629).
- Rohrer, J., Schmukle, S., McElreath, R., 2022. The only thing that can stop bad causal inference is good causal inference. *Behavioral and Brain Sciences* 45, e91. doi:[10.1017/S0140525X21000789](https://doi.org/10.1017/S0140525X21000789).
- Rubin, D., 1974. Estimating causal effects of treatments in randomized and nonrandomized studies. *Journal of Educational Psychology* 66, 688–701. doi:[10.1037/h0037350](https://doi.org/10.1037/h0037350).
- Schuessler, J., Selb, P., 2023. Graphical causal models for survey inference. *Sociological Methods and Research* 0. doi:[10.1177/00491241231176851](https://doi.org/10.1177/00491241231176851).
- Seaman III, J., Seaman Jr., J., Stamey, J., 2011. Hidden dangers of specifying noninformative priors. *The American Statistician* 66, 77–84. doi:[10.1080/00031305.2012.695938](https://doi.org/10.1080/00031305.2012.695938).
- Sekhon, J., 2009. The neyman-rubin model of causal inference and estimation via matching methods, in: Box-Steffensmeier, J., Brady, H., Collier, D. (Eds.), *The Oxford Handbook of Political Methodology*. Oxford University Press, pp. 271–299. doi:[10.1093/oxfordhb/9780199286546.003.0011](https://doi.org/10.1093/oxfordhb/9780199286546.003.0011).
- Shaughnessy, J., Zechmeister, E., Zechmeister, J., 2010. *Research Methods in Psychology*. McGraw-Hill. URL: [https://web.archive.org/web/20141015135541/http://www.mhhe.com/socscience/psychology/shaugh/ch01\\_concepts.html](https://web.archive.org/web/20141015135541/http://www.mhhe.com/socscience/psychology/shaugh/ch01_concepts.html). retrieved July 23, 2024.
- Spiroes, P., Glymour, C., Scheines, R., 1991. From probability to causality. *Philosophical Studies* 64, 1–36. URL: <https://www.jstor.org/stable/4320244>.
- Stan Development Team., 2021. *Stan Modeling Language Users Guide and Reference Manual*, version 2.26. Vienna, Austria. URL: <https://mc-stan.org>.
- Sterner, P., Pargent, F., Deffner, D., Goretzko, D., 2024. A causal framework for the comparability of latent variables. *Structural Equation Modeling: A Multidisciplinary Journal* 31, 747–758. doi:[10.1080/10705511.2024.2339396](https://doi.org/10.1080/10705511.2024.2339396).
- Thurstone, L., 1927a. A law of comparative judgment. *Psychological Review* 34, 482–499. doi:[10.1037/h0070288](https://doi.org/10.1037/h0070288).
- Thurstone, L., 1927b. Psychophysical analysis. *American Journal of Psychology* , 368–89URL: [https://brocku.ca/MeadProject/Thurstone/Thurstone\\_1927g.html](https://brocku.ca/MeadProject/Thurstone/Thurstone_1927g.html). last accessed 20 december 2024.
- van Daal, T., 2020. Making a choice is not easy?!: Unravelling the task difficulty of comparative judgement to assess student work. Ph.D. thesis. University of Antwerp.
- van Daal, T., Lesterhuis, M., Coertjens, L., Donche, V., De Maeyer, S., 2016. Validity of comparative judgement to assess academic writing: examining implications of its holistic character and building on a shared consensus. *Assessment in Education: Principles, Policy & Practice* 26, 59–74. doi:[10.1080/0969594X.2016.1253542](https://doi.org/10.1080/0969594X.2016.1253542).
- van Daal, T., Lesterhuis, M., Coertjens, L., van de Kamp, M., Donche, V., De Maeyer, S., 2017. The complexity of assessing student work using comparative judgment: The moderating role of decision accuracy. *Frontiers in Education* 2. doi:[10.3389/feduc.2017.00044](https://doi.org/10.3389/feduc.2017.00044).
- van der Linden, W. (Ed.), 2017a. *Handbook of Item Response Theory: Models*. volume 1 of *Statistics in the Social and Behavioral Sciences Series*. CRC Press.
- van der Linden, W. (Ed.), 2017b. *Handbook of Item Response Theory: Statistical Tools*. volume 2 of *Statistics in the Social and Behavioral Sciences Series*. CRC Press.

- Verhavert, S., Bouwer, R., Donche, V., De Maeyer, S., 2019. A meta-analysis on the reliability of comparative judgement. *Assessment in Education: Principles, Policy and Practice* 26, 541–562. doi:[10.1080/0969594X.2019.1602027](https://doi.org/10.1080/0969594X.2019.1602027).
- Verhavert, S., Furlong, A., Bouwer, R., 2022. The accuracy and efficiency of a reference-based adaptive selection algorithm for comparative judgment. *Frontiers in Education* 6. doi:[10.3389/feduc.2021.785919](https://doi.org/10.3389/feduc.2021.785919).
- Wainer, H., TimbersFairbank, D., Hough, R., 1978. Predicting the impact of simple and compound life change events. *Applied Psychological Measurement* 2, 313–322. doi:[10.1177/014662167800200301](https://doi.org/10.1177/014662167800200301).
- Whitehouse, C., 2012. Testing the validity of judgements about geography essays using the adaptive comparative judgement method. URL: [https://filestore.aqa.org.uk/content/research/CERP\\_RP\\_CW\\_24102012\\_0.pdf?download=1](https://filestore.aqa.org.uk/content/research/CERP_RP_CW_24102012_0.pdf?download=1). aQA Education.
- Wu, W., Niezink, N., Junker, B., 2022. A diagnostic framework for the bradley–terry model. *Journal of the Royal Statistical Society Series A: Statistics in Society* 185, S461–S484. URL: [https://academic.oup.com/jrssa/article-pdf/185/Supplement\\_2/S461/49421054/jrssa\\_185\\_supplement\\_2\\_s461.pdf](https://academic.oup.com/jrssa/article-pdf/185/Supplement_2/S461/49421054/jrssa_185_supplement_2_s461.pdf), doi:[10.1111/rssa.12959](https://doi.org/10.1111/rssa.12959).
- Zimmerman, D., 1994. A note on the influence of outliers on parametric and nonparametric tests. *The Journal of General Psychology* 121, 391–401. doi:[10.1080/00221309.1994.9921213](https://doi.org/10.1080/00221309.1994.9921213).