

Variational Inference with Continuously-Indexed Normalizing Flows

ICML 2020 INN⁺ Workshop

Anthony Caterini, Rob Cornish, Dino Sejdinovic, Arnaud Doucet

University of Oxford

July 18, 2020

Variational Inference

Given a joint target $p_{X,Z}(x, z)$ over data $x \in \mathcal{X}$ and latent variables $z \in \mathcal{Z}$, **variational inference (VI)** seeks to approximate the intractable posterior $p_{Z|X}$

Variational Inference

Given a joint target $p_{X,Z}(x, z)$ over data $x \in \mathcal{X}$ and latent variables $z \in \mathcal{Z}$, **variational inference (VI)** seeks to approximate the intractable posterior $p_{Z|X}$

VI methods introduce an **approximate posterior** q_Z , and are trained to maximize the **evidence lower bound (ELBO)** objective

Variational Inference

Given a joint target $p_{X,Z}(x, z)$ over data $x \in \mathcal{X}$ and latent variables $z \in \mathcal{Z}$, **variational inference (VI)** seeks to approximate the intractable posterior $p_{Z|X}$

VI methods introduce an **approximate posterior** q_Z , and are trained to maximize the **evidence lower bound (ELBO)** objective

Can also **amortize** inference across observations

- e.g. Encoder in a VAE [Kingma and Welling, 2014]

Explicit VI with Normalizing Flows

Explicit VI involves specifying q_Z with tractable sampling and density evaluation

Explicit VI with Normalizing Flows

Explicit VI involves specifying q_Z with tractable sampling and density evaluation

Normalizing flows (NFs) [Rezende and Mohamed, 2015] define q_Z explicitly according to the generative process

$$W \sim q_W, \quad Z := g(W)$$

- g is a diffeomorphism, q_W is a simple base density

Explicit VI with Normalizing Flows

Explicit VI involves specifying q_Z with tractable sampling and density evaluation

Normalizing flows (NFs) [Rezende and Mohamed, 2015] define q_Z explicitly according to the generative process

$$W \sim q_W, \quad Z := g(W)$$

- g is a diffeomorphism, q_W is a simple base density

Although powerful, NFs can suffer from expressiveness issues, even becoming **pathological** when the topologies of q_W and $p_{Z|X}$ don't match [Cornish et al., 2020]

Implicit VI with Continuously-Indexed Normalizing Flows

Main contribution: We can gain expressiveness in VI by instead using **continuously indexed flows (CIFs)** [Cornish et al., 2020], with generative process

$$W \sim q_W, \quad U \mid W \sim q_{U|W}(\cdot \mid W), \quad Z := G(W; U),$$

where $q_{U|W}$ is trainable, and **each** $G(\cdot; u)$ is a normalizing flow

Implicit VI with Continuously-Indexed Normalizing Flows

Main contribution: We can gain expressiveness in VI by instead using **continuously indexed flows (CIFs)** [Cornish et al., 2020], with generative process

$$W \sim q_W, \quad U \mid W \sim q_{U \mid W}(\cdot \mid W), \quad Z := G(W; U),$$

where $q_{U \mid W}$ is trainable, and **each** $G(\cdot; u)$ is a normalizing flow

Any existing normalizing flow g can be used to construct G , e.g.

$$G(w; u) := e^{s(u)} \odot g(w) + t(u)$$

Implicit VI with Continuously-Indexed Normalizing Flows

Main contribution: We can gain expressiveness in VI by instead using **continuously indexed flows (CIFs)** [Cornish et al., 2020], with generative process

$$W \sim q_W, \quad U \mid W \sim q_{U|W}(\cdot \mid W), \quad Z := G(W; U),$$

where $q_{U|W}$ is trainable, and **each** $G(\cdot; u)$ is a normalizing flow

Any existing normalizing flow g can be used to construct G , e.g.

$$G(w; u) := e^{s(u)} \odot g(w) + t(u)$$

Now, $q_Z(z) := \int q_{Z,U}(z, u) \mathrm{d}u$ is defined **implicitly** over **auxiliary variables** $u \in \mathcal{U}$, rendering the original ELBO intractable

Auxiliary Variational Inference

Since $q_{Z,U}$ is tractable for CIFs, we can train using the framework of **auxiliary variational inference** [Agakov and Barber, 2004]

Auxiliary Variational Inference

Since $q_{Z,U}$ is tractable for CIFs, we can train using the framework of **auxiliary variational inference** [Agakov and Barber, 2004]

New objective is the **auxiliary ELBO**, having **auxiliary inference distribution** $r_{U|Z}$

Auxiliary Variational Inference

Since $q_{Z,U}$ is tractable for CIFs, we can train using the framework of **auxiliary variational inference** [Agakov and Barber, 2004]

New objective is the **auxiliary ELBO**, having **auxiliary inference distribution** $r_{U|Z}$

CIFs **already** provide an auxiliary inference distribution

- For multi-layer CIFs, this has the correct **factorization structure**

Results - Mixture of Gaussians

We compare explicit VI with a **neural spline flow (NSF)** [Durkan et al., 2019] to a **CIF-NSF** on a mixture of Gaussians inference problem with varied initial noise σ_0

Results - Mixture of Gaussians

We compare explicit VI with a **neural spline flow (NSF)** [Durkan et al., 2019] to a **CIF-NSF** on a mixture of Gaussians inference problem with varied initial noise σ_0




Figure 1: CIFs on top, NSF on bottom.

Results - Image Datasets

We further compare the performance of generative models which use either **NSFs**, **CIF-NSFs**, or baseline **VAEs** as an **amortized** inference method

Results - Image Datasets

We further compare the performance of generative models which use either NSF_s, CIF-NSF_s, or baseline VAE_s as an amortized inference method

Table 1: Test-set marginal log-likelihood averaged over three runs.

	VAE	NSF	CIF-NSF
MNIST	$-87.37_{\pm 0.15}$	$-82.95_{\pm 0.11}$	$-82.22_{\pm 0.13}$
FASHION-MNIST	$-217.82_{\pm 0.07}$	$-215.45_{\pm 0.08}$	$-214.50_{\pm 0.11}$

Thank you!



Figure 2: Joint work with Rob Cornish, Dino Sejdinovic, and Arnaud Doucet

References

- Anthony Caterini, Rob Cornish, Dino Sejdinovic, and Arnaud Doucet. Variational inference with continuously-indexed normalizing flows. In *ICML Workshop on Invertible Neural Networks and Normalizing Flows*, 2020.
- Rob Cornish, Anthony L Caterini, George Deligiannidis, and Arnaud Doucet. Relaxing bijectivity constraints with continuously-indexed normalising flows. In *International Conference on Machine Learning*, 2020.
- Diederik P. Kingma and Max Welling. Auto-encoding variational bayes. In *2nd International Conference on Learning Representations, ICLR*, 2014.
- Danilo Rezende and Shakir Mohamed. Variational inference with normalizing flows. In *International Conference on Machine Learning*, pages 1530–1538, 2015.
- Felix V Agakov and David Barber. An auxiliary variational method. In *International Conference on Neural Information Processing*, pages 561–566. Springer, 2004.
- Conor Durkan, Artur Bekasov, Iain Murray, and George Papamakarios. Neural spline flows. In *Advances in Neural Information Processing Systems*, pages 7509–7520, 2019.