## Nature network mapping

**The Challenge:** to identify the most important areas of mainland Cornwall for biodiversity and the provision of ecosystem services that are currently excluded from any statutory protection.

**What does the map show?** The nature network map identifies the most valuable 20% of mainland Cornwall not currently subject to statutory protection. The area is sub-divided into three exclusive bands representing the most valued 5%, 5-10% and 10-20% of unprotected areas.

**Intended uses:** The map is intended to inform a range of strategic planning application that could include:

- the expansion of the existing network of protected areas;
- the strategic targeting of resources to protect biodiversity and ecosystem services:
- identification of those areas which could result in a strategic loss of biodiversity or service provision as a result of any future change in land use or development.

## Notes on the use and interpretation of the nature network map:

- The spatial unit of prioritization is 100 square metre cells although some finer resolution features are accounted for (such as ancient and venerable trees).
- Rankings are assigned uniquely on the basis of <u>current</u> land cover and other indicators of ecosystem services and biodiversity or factors like population or industrial activity that may affect the value of ecosystem services.
- Cell rankings are dependent upon maintaining the landcover and existing services of more highly ranked cells. This dependence is most evident by considering how the value of a habitat 'corridor' is only meaningful if the two habitats that it links are maintained.
- Existing protected areas (irrespective of their condition) are assumed to be preserved and maintained. Their presence informs the prioritization of non-protected areas.
- The exclusion of an area from the top 20% of cells dose *not* imply that the area lacks any strategic biodiversity value or service provision.
- Similarly, a high ranking does not necessarily imply that the biodiversity or services provided cannot be augmented by habitat modification of changes to management.

## Method of spatial prioritization

The mapping is produced using the Zonation model of spatial prioritization (Lehtomäki *et al.* 2016), which has been widely used for the identification of new protected areas or conservation zones, and the expansion of existing networks. The approach requires data and expert judgement concerning, for example, the relative importance of different land features and habitats.

Inevitably mapping is constrained by the data and information that is available. The method does not pretend to capture every possible factor that might inform the relative importance of areas. The method focuses on key factors that (i) we believe should inform spatial prioritization and (ii) for which there is sufficiently reliable information available.

It is important to recognize that the data informing this prioritization is not comprehensive. Nevertheless, use of the model and its outputs provides an improvement on *ad hoc* decisions that do not exploit the available evidence and data concerning the state of the natural environment in Cornwall. Basing decisions on the Zonation prioritization should not be considered sufficient, but can contribute to more informed decision-making.

The approach allows easy updating or replacement of source data layers, adjustments to the perceived importance of different services, and can be applied to different spatial areas and scales.

# Feature weighting

The relative importance of different habitats and features informing the prioritization is determined by assigning weights. Positive weights describe the relative value or benefits of different services and features. Negative weights indicate a constraint or cost of realising benefits. Weightings vary from -3 to +5. Full weightings are provided in Table \*\*.

The dominant land cover/habitat of an area is determinant of both of the biodiversity potential of an area and its capacity to deliver ecosystem services.

**Landcover weightings:** were attributed according to a judgment of their capacity to support biodiversity. In addition, the dominant landcover is often determinant of the capacity of an area to deliver many ecosystem services (accounted for separately).

The following general weightings were used for the exclusive landcover/habitat categories:

- BAP priority habitats: +2
- Other semi-natural habitats: +1
- Urban, arable and improved grassland: 0

In effect therefore approximately 77% of mainland Cornwall is attributed a zero weighting in terms of its dominant landcover type.

**Land features:** features providing additional habitats such as hedgerows or open waterways were given a 0 to +2 weighting dependent upon length and/or quality of the feature.

**Habitat designations:** land designations relating to biodiversity were used as strong indicators of the biodiversity value of areas, whether these were statutory

protections (e.g. SSSIs) or other designations such as County wildlife sites or ancient woodland. Weightings varied from +1 to +5.

**Ecosystem services:** were weighted between +1 and +5 according to a judgement of the relative importance of the service and confidence in its estimation. Service layers were calculated on the basis of existing demands such as current sources of drinking water, location of aquaculture or buildings at risk of flooding.


**Constraints and opportunity costs:** opportunity costs have not been included as doing so would result in a prioritization that maximises the difference between costs and benefits, rather than one seeking to maximise ecological or biodiversity benefits which is the primary intension of the nature network map. Nevertheless, a negative weighting has been applied to certain land features likely to reduce the potential biodiversity or other benefits of a cell including:

- man-made infrastructure, such as main roads and railways;
- the percentage of an area that is covered with buildings;
- certain recreational uses such as golf courses;
- china clay extraction areas.

#### Cell removal iteration

The spatial prioritization works by iteratively removing the least 'valuable', lowest ranked cells (see figure \*) on the basis of:

- Weighted fraction of all remaining features that values cells containing a
  high overall proportion of individual features (*ie* it will most value cells
  containing a high proportion of a habitat type or service).
- An 'added benefit function' that sums the normalised distribution across all features and so tends to remove cells with low feature diversity before cells with higher feature diversity, such as cells providing *both* valuable habitat *and* ecosystem services.
- Aggregation of protected areas a 'boundary length penalty' seeks to reduce edge to area ratio of *all* remaining cells (irrespective of the type of features within those cells). In effect this will aggregate and connect together protected areas rather than specific habitat types.
- Existing protected areas are designated as the last areas to be removed, ensuring that they inform the ranking of all other areas.



**Figures \*** show the iterative 'removal' in grey of the least valued or lowest ranked (i) 20% (predominantly urban and infrastructure areas), (ii) 60% and (iii) 80% of cells for the Fowey valley region, the remaining cells shown in (iv) corresponding to the highest 20% of ranked cells (red) and areas under statutory protection (yellow).

#### **Issues and Considerations**

**Vulnerability and zonation:** the most highly ranked cells are assigned to areas supporting biodiversity and delivering ecosystem services. It is important to note that these do *not* equate to the most vulnerable habitats. For example, it could be argued that many areas of Bodmin moor are less vulnerable than habitats around urban centres as the pressures for development will be greatest for these latter areas. The lack of opportunity costs in the prioritization method means that pressures associated with alternative land uses are not accounted for.

Valuing urban and agricultural land - currently all cells with a urban, suburban, arable and improved grassland land cover type are given a zero weighting for dominant land cover. Negative weightings are applied to several land features associated with urban areas, including the % of each cell covered by buildings (but excludes man-made surface and excludes some recent developments), allocation areas, china clay pits, rail and main roads. Equally the presence of positively weighted features, such as hedgerows and rivers, as well as the ecosystem services they can provide, will lead to some of these areas being more highly valued.

In effect these land cover types are unlikely to be heavily represented in the proposed nature network but are not excluded *a priori*. In certain areas, such as

areas that deliver significant flood mitigation services, such land cover may be valued due to the presence of such features even though alternative land cover and uses would deliver even greater service value.

The biodiversity potential of urban and agricultural land cover is very variable. Urban and suburban areas often have the capacity to support higher levels of biodiversity than many arable and improved grassland landscapes. Equally however, methods of land management of rural environments have significant implications for their biodiversity value. Biodiversity indicators for urban and agricultural landscapes could include participation in stewardship schemes (in the case of agriculture) or measures of canopy cover in the case of urban environments.

How these areas are best represented within the spatial model needs to reflect the policy applications of the nature network mapping. Currently, for example, the zero weighting applied to most agricultural and urban areas does not imply they lack any biodiversity value, but that such value is unlikely to change if these areas do not receive any additional protection, or that their value depends on policies or management options that are unaffected (or not targeted) by the nature network mapping.

#### References

Lehtomäki J, Moilanen A, Toivonen T Leathwick J 2016. Running a Zonation planning project. Unigrafia, Helsinki. ISBN: 978-951-51-1923-0 (PDF).

Moilanen A, Pouzols FM, Meller L, Veach V, Arponen A, Leppanen J, Kujala H 2014 Zonation - Spatial conservation planning methods and software. Version 4. User Manual. University of Helsinki, Helsinki.

Moilanen, A. 2013. Planning impact avoidance and biodiversity offsetting using software for spatial conservation prioritization. Wildlife Research, 40: 153-162.

Joona Lehtomaki J, Tomppo E, Kuokkanen P, Hanski I, Moilanen A 2009 Applying spatial conservation prioritization software and high-resolution GIS data to a national-scale study in forest conservation. Forest Ecology and Management 258, 2439–2449

#### Datasets and resources used

The type of data used for the spatial prioritization included

- Mutually exclusive presence / absence data, such as the main habitat or land cover types.
- Non-exclusive presence / absence data that might be combined with other habitat types, such as the presence of open rivers or main roads
- Non-exclusive scaled quantitative data that provided an indication of the relative importance of different features (for example hedgerow lengths) or service provision (quantified estimates of ecosystem services).

## Land cover and habitat type

Landcover types were derived from a compilation of several landcover data sources. Each cell was attributed to a single landcover type derived from these classifications (ie exclusive categorization).

		LANDCOVER & HABITATS TYPE DATA	
Data source	Data owner	Source link	Usage Notes
Land cover map 2015 v1.2 (2017)	СЕН	https://eip.ceh.ac.uk/lcm/lcmdata	
Corine LC2012	EU	http://land.copernicus.eu/pan-european/corine- land-cover	Only used for validating other types and comparison
Priority habitat inventory	NE	https://naturalengland- defra.opendata.arcgis.com/datasets/priority- habitat-inventory-england-south	Compilation of BAP priority habitats
National forest inventory	FC	https://data.gov.uk/dataset/national-forest-inventory-woodland-england-2015	Largely based on satellite imagery up to 3 yrs old. Woodland categorization only.
Plymouth Coastal Observatory habitat data.	PCO	http://southwest.coastalmonitoring.org/data- types/habitat-mapping/	Coastal landcover data
Open mosaic on previously developed land	NE & Buglife	https://data.gov.uk/dataset/8509c11a-de20-42e8-9ce4-b47e0ba47481/open-mosaic-habitat-draft	DRAFT data
Woodland pasture and parkland	NE	https://data.gov.uk/dataset/61e6f611-af12-471e-8732-76b4d66dbb1f/wood-pasture-and-parkland-bap-priority-habitat-inventory-for-england	Recent addition to priority habitats

#### Species presence data

Point data of species observations or recordings have <u>not</u> been used in the analysis. It was considered that the observational records showed high spatial bias resulting from the very uneven recording effort across the county. In the

future it may be possible to use such point information to inform models of a species distribution that in turn could be used for spatial prioritization.

#### **Habitat features**

In addition to land cover data describing cell habitat types, additional data on habitat features contained within cells was also used. These included linear and point features, such as open river, hedgerows, and the presence of ancient and venerable trees. Such data was non-exclusive (a cell could contain several features of the same or different types) and could be recorded as the presence/absence of a feature (such as a river bank) or scaled according to a measure of the abundance of a feature, such as the total length of hedgerows within a cell or number of recorded ancient/venerable trees.

		HABITAT & LANDCOVER FEATURES	
Data source	Data	Source link and reference	Usage
	owner		Notes
OS Open	OS	https://www.ordnancesurvey.co.uk/opendatadownl	
Rivers		oad/products.html	
Open	OS	https://www.ordnancesurvey.co.uk/opendatadownl	golf
Greenspace		oad/products.html	courses &
			playing
			fields
Hedges	ERCCIS	ERCCIS Hedges and Field Boundaries Project hedges	
		layer: erccis@cornwallwildlifetrust.org.uk	
Ancient Tree	Woodland	https://ati.woodlandtrust.org.uk	
Inventory	Trust		
Priority	NE	https://naturalengland-	No main
habitats		defra.opendata.arcgis.com/datasets/priority-habitat-	habitat
inventory		<u>inventory-england-south</u>	category

#### Habitat and biodiversity designations

A range of statutory and non-statutory designations, primarily derived from the biodiversity richness of an area, were used as further indicators of biodiversity. These designations are generally non-exclusive, and the same areas can be classified under several different designations. For example, many SSSI areas were also designated as SAC or SPA areas. Undue bias towards multiple-designated areas was prevented by either suitable weighting scores or by creating synthesis layers brining together several different designations. Statutory designated areas were defined as all areas designated as SSSI, SAC, SPA, national or local nature reserves.

		HABITAT & BIODIVERSITY DESIGNATIONS	
Data source	Data	Source link and reference	Usage Notes
	owner		
Ancient	NE	https://data.gov.uk/dataset/9461f463-c363-4309-	
woodlands		ae77-fdcd7e9df7d3/ancient-woodlands-england	
County	ERCCIS	https://erccis.org.uk/CountySites	
wildlife sites			
AONB (2018)	NE	https://naturalengland-	
		defra.opendata.arcgis.com/datasets/areas-of-	
		outstanding-natural-beauty-england	
SSSI (2018)	NE	https://naturalengland-	
		defra.opendata.arcgis.com/datasets/sites-of-special-	

	1		
		<u>scientific-interest-england</u>	
SPA (2018)	NE	https://naturalengland-	
		defra.opendata.arcgis.com/datasets/special-	
		protection-areas-england	
SAC (2018)	NE	https://naturalengland-	
, ,		defra.opendata.arcgis.com/datasets/special-areas-	
		of-conservation-england	
National	NE	https://naturalengland-	
Nature		defra.opendata.arcgis.com/datasets/national-nature-	
Reserves		reserves-england	
(2018)			
Local Nature	NE	https://naturalengland-	
Reserves	NE		
		defra.opendata.arcgis.com/datasets/local-nature-	
(2018)		<u>reserves-england</u>	
Important	Plantlife	http://www.plantlifeipa.org/home	
Plant Areas			
RSPB	RSPB	https://www.arcgis.com/home/item.html?id=60767	
reserves		15cb76d4c388fa38b87db7d9d24	

# **Ecosystem service benefits**

A comprehensive list of data sources used in the calculation of ecosystem services is detailed under their individual methodologies.

No ecosystem services dependent upon public access to the land (*eg* recreation, tourism, health benefits) or

	ECOSYSTEM SERVICE
Service	Description
Flood mitigation	Mitigation of flood surface water and river/sea flood risk on downstream potential vulnerable populated areas
Aquaculture water quality	Mitigation of water pollution upstream of major aquaculture sites
Drinking water quality	Mitigation of water pollution in drinking water sensitive areas
Bathing water quality	Mitigation of water pollution upstream of beaches
Soil loss mitigation	Mitigation of soil loss risk derived from ESDAC soil erosion by water estimates: <a href="https://esdac.jrc.ec.europa.eu/content/soil-erosion-water-rusle2015">https://esdac.jrc.ec.europa.eu/content/soil-erosion-water-rusle2015</a> )
Soil carbon	LINK SOURCE
Pollination	Pollinator friendly habitats close to agricultural arable areas.
Air quality mitigation	Mitigation of landcover in air quality monitoring zones

#### Facilitating factors & other non-biodiversity benefits

Data sources were used to identify key factors that could restrict alternative uses (reduce opportunity costs) or bring additional non-biodiversity benefits, such as sites of geological or heritage value. These layers might be considered facilitating factors that would help realise biodiversity benefits and/or reduce opportunity costs as they restrict alternative land uses. Additional factors that could be considered include:

- Additional indicators of sympathetic land ownership or management.
- Unsuitability for alternative land uses including topography such as steep slopes, aspect, elevation and wind exposure.

		FACILITATING FACTORS & DESIGNATIONS	
Data source	Data	Source link and reference	Usage Notes
	owner		
County	ERCCIS	https://erccis.org.uk/DataHoldings	Not used
geology sites			
World	EH	https://data.gov.uk/dataset/3ac5c299-6805-476b-	Not used
Heritage Site		af9b-90aadec5e7b4/world-heritage-sites-gis-data	
Heritage	NE	https://naturalengland-	Not used
coast		defra.opendata.arcgis.com/datasets/heritage-coasts-	
		<u>england</u>	
Scheduled	EH	https://historicengland.org.uk/listing/the-list/data-	Not used
monuments		downloads/	
Battlefields	EH	https://historicengland.org.uk/listing/the-list/data-	Not used
		downloads/	
Doorstep	NE	https://data.gov.uk/dataset/6a80e5a7-017e-49ba-	
greens		a981-5cd0c727086f/doorstep-greens-polygons	

# Additional habitat condition and quality indicators

Very little spatial data is available on the current condition of habitats in Cornwall. Where available, catchment ecological quality data was used as indicators of aquatic habitat condition and SSSI condition data was used as indicator of habitat condition within those sites. Areas participating in Higher level or Organic countryside stewardship schemes were also considered indicative of good condition of arable and grassland areas.

		HABITAT CONDITION INDICATORS	
Data source	Data	Source link and reference	Usage Notes
	owner		
SSSI (2018)	NE	https://naturalengland-	SSSI habitat
		defra.opendata.arcgis.com/datasets/sites-of-special-	condition
		scientific-interest-england	
WFD	EA	https://environment.data.gov.uk/catchment-	Aquatic
Catchment		<u>planning/ui/reference</u>	habitat
data			condition.
Countryside	NE	https://naturalengland-	Higher and
stewardship		defra.opendata.arcgis.com/datasets/countryside-	Organic level
participation		stewardship-scheme-2016-management-areas-	only
		<u>england</u>	

### **Constraints and Costs**

Factors that could restrict the potential biodiversity value or impede the protection of sites can be attributed a negative weighting. These can include existing infrastructure and/or buildings, future planning commitments, certain kinds of recreational use (golf courses, playing fields) or suitability for alternative land use such as the agricultural land grade.

		HABITAT CONDITION INDICATORS	
Data source	Data	Source link and reference	Usage Notes
	owner		
Agricultural	NE	http://naturalengland-	Not used
land		defra.opendata.arcgis.com/datasets/agricultural-	
classification		land-classification-alc-grades-post-1988-survey-	
		polygons-england	

OS Open Map Local	OS	https://www.ordnancesurvey.co.uk/opendatadownload/products.html	Rail network and % built-
Vector		oad/products.html	up area
OS Open Roads Vector	OS	https://www.ordnancesurvey.co.uk/opendatadownload/products.html	Road network and type
China clay pits	OSM	https://www.openstreetmap.org/#map=6/54.910/-3.432	Complemente d by aerial photography and remote sensing data
Allocation areas	CC		

# Feature layers and weightings used by Zonation

Dominant landcover type	Type of data	Weighting
Coniferous woodland	Presence / Absence	1
Deciduous woodland	Presence / Absence	2
Scrub and other woodland	Presence / Absence	1
Traditional orchard	Presence / Absence	2
Woodland pasture	Presence / Absence	2
Seminatural grassland	Presence / Absence	1
Acid grassland	Presence / Absence	1
Lowland dry acid grassland	Presence / Absence	2
Calcareous grassland	Presence / Absence	2
Neutral grassland	Presence / Absence	1
Meadows upland and lowland	Presence / Absence	2
Grazing marsh pasture	Presence / Absence	2
Maritime grassland	Presence / Absence	1
Fen Marsh	Presence / Absence	1
Reedbeds	Presence / Absence	2
Fens upland and lowland	Presence / Absence	2
Purple moor grass	Presence / Absence	2
Dwarf Shrub Heath Bog	Presence / Absence	1
Blanket bog	Presence / Absence	2
Bracken	Presence / Absence	1
Heathland upland and lowland	Presence / Absence	2
Inland rock	Presence / Absence	1
Calaminarian grassland	Presence / Absence	2
Quarry	Presence / Absence	1
Open mosaic	Presence / Absence	1
Supralittoral rock	Presence / Absence	1
Maritime cliff and slope	Presence / Absence	2
Littoral rock priority	Presence / Absence	2
Littoral rock	Presence / Absence	1
Supralittoral sediment	Presence / Absence	1
Coastal sand dunes	Presence / Absence	2
Coastal vegetated shingle	Presence / Absence	2
Littoral sediment	Presence / Absence	1
Saltmarsh	Presence / Absence	2
Intertidal mudflats	Presence / Absence	2
Sheltered muddy gravels	Presence / Absence	2
Blue mussel beds	Presence / Absence	2
Saltwater	Presence / Absence	1
Saline lagoons	Presence / Absence	2
Freshwater	Presence / Absence	1

River or stream	Presence / Absence	2
Standing freshwater	Presence / Absence	1
Arable	Presence / Absence	0
Improved grassland	Presence / Absence	0
Builtup	Presence / Absence	0
Suburban	Presence / Absence	0
Urban	Presence / Absence	0
Land features / linear habitats	Type of data	Weighting
Hedges	Total length	2
Open rivers	Presence / Absence	1
A roads	Presence / Absence	-3
B roads	Presence / Absence	0
Dual carriageway roads	Presence / Absence	0
Minor roads	Presence / Absence	0
Railways	Presence / Absence	-3
Building cover percentage	Scaled 0:100	-5
Doorstep greens	Presence / Absence	0
Parks and Gardens	Presence / Absence	0
Golf courses	Presence / Absence	-1
Playing field	Presence / Absence	-1
China claypit area	Presence / Absence	-5
Biodiversity designations	Type of data	Weighting
Ancient woodlands	Presence / Absence	5
High level stewardship	Presence / Absence	0
Organic stewardship	Presence / Absence	0
Plantlife designated area	Presence / Absence	5
RSPB reserve	Presence / Absence	5
County wildlife site	Presence / Absence	3
Non-dominant priority habitats	Presence / Absence	1
Any statutory protected areas	Presence / Absence	5
(SSSI, SPA, SAC, national or local		
nature reserve)		_
AONB	Presence / Absence	0
World heritage area	Presence / Absence	0
Heritage coast	Presence / Absence	0
Allocation zones	Presence / Absence	0
Ecosystem Services	Type of data	Weighting
Water quality service for	Scaled 0:100	3
aquaculture	Cooled 0.100	2
Water quality service for bathing waters	Scaled 0:100	3
Water quality service for drinking	Scaled 0:100	4
water	364/64 0.100	7
Water quality service for other	Scaled 0:100	0
atc. quanty service for other	· <del></del>	-

|--|

Heritage coast

Flood mitigation service	Scaled 0:100	5
Pollination service	Scaled 0:100	3
Soil loss mitigation service	Scaled 0:100	4
Air pollution mitigation service	Scaled 0:100	3
Topography	Type of data	Weighting
Elevation	Metres elevation	0
North facing aspect	Presence / Absence	0
South facing aspect	Presence / Absence	0
Medium slope	Presence / Absence	0
Stoon clans	Presence / Absence	0
Steep slope	Presence / Absence	· ·
Soil	Type of data	Weighting
· · · · · · · · · · · · · · · · · · ·		
Soil	Type of data	Weighting
Soil High peat content soil	Type of data Presence / Absence	Weighting 2
Soil  High peat content soil  Agricultural grade 1 land	Type of data  Presence / Absence  Presence / Absence	Weighting 2 0
Soil  High peat content soil Agricultural grade 1 land Agricultural grade 2 land	Type of data  Presence / Absence  Presence / Absence  Presence / Absence	Weighting  2  0 0
Soil  High peat content soil Agricultural grade 1 land Agricultural grade 2 land Agricultural grade 3 land	Presence / Absence	Weighting  2  0  0  0
Soil  High peat content soil Agricultural grade 1 land Agricultural grade 2 land Agricultural grade 3 land Agricultural grade 4 land	Type of data  Presence / Absence	Weighting  2  0  0  0  0
Soil  High peat content soil Agricultural grade 1 land Agricultural grade 2 land Agricultural grade 3 land Agricultural grade 4 land Agricultural grade 5 land	Type of data  Presence / Absence  Presence / Absence	Weighting  2  0  0  0  0  0

Presence / Absence 0