

Integrative Medicine Research 9 (2020) 100593**Anti-inflammatory active constituents of *Cicadiae Periostracum***

Seung Mok Ryu, Hye-sun Lim, Gunhyuk Park,
Jun-ho Song, Byeong Cheol Moon, Jun Lee

*Herbal Medicine Resources Research Center, Korea
Institute of Oriental Medicine, Naju, Korea*

Cicadiae Periostracum, the cast-off shell of *Cryptotympana* sp., is used in Korean traditional medicine for its spasmolytic, antiferbrile, and an antiphlogistic effects. Two new N-acetyldopamine dimers (1 and 2) were isolated from a 70% EtOH extract of *Cicadiae Periostracum* together with nine known compounds (3-11). The structures of the isolated compounds were identified by NMR and MS experiments. All isolated compounds were evaluated for their potential to inhibit LPS-induced nitric oxide production in BV-2 cells.

<https://doi.org/10.1016/j.imr.2020.100593>

Integrative Medicine Research 9 (2020) 100594**To investigate the meridian through electroacupuncture for postoperative pain relief in patients with arthroscopic repair for triangular fibrocartilage complex (TFCC) tears**

Chiu-Ming Chang, Yung-Cheng Chiu, Yu-Chen Lee

China Medical University Hospital, Taichung, Taiwan

Background: The purpose of the study is to investigate the meridian. **Methods:** After the patients accepted arthroscopic repair for triangular fibrocartilage complex (TFCC) tears, the investigator use electroacupuncture for postoperative pain relief. It is expected that 20 patients will be randomly assigned to the following groups: electroacupuncture group (EAG), EA at points; non-point group (NPG), EA at non-points. Acupoint selection: *true acupoints: needles were inserted to Kunlun (BL60) and Sanyinjiao (SP6) contralateral to the operated leg and deqi sensation elicited at acupoints *false point: the needles were inserted to Zusanli (ST36) and Shangjuxu (ST37) contralateral to the operated leg and deqi sensation elicited at acupoints Data collection: the investigator recorded the data including the time required for the first injection of analgesics and the number of doses of injection painkiller used in 24 hours after surgery (count).

Results: The group of electroacupuncture group (EAG) significant increases in the time required for the first injection of analgesics and decrease in the number of doses of injection painkiller than the group of non-point group (NPG).

Conclusion: We think electroacupuncture group (EAG) is effective because the meridian is right.

<https://doi.org/10.1016/j.imr.2020.100594>

Integrative Medicine Research 9 (2020) 100595**Effectiveness of sacral acupuncture for sleep depth: a randomized controlled cross-over trial**

Hisashi Honjo, Masato Uemura, Hikari Takahashi,
Eizen Ryu, Yoko Shiozaki, Masahito Miyake,
Masato Nagata, Fumihiko Fukuda,
Kazuhiko Taniguchi

Meiji School of Oriental Medicine, Suita, Japan

Background: We have reported the effects of sacral acupuncture on primary nocturnal enuresis and nocturia due to overactive bladder (OAB). Indeed, sacral acupuncture could represent the effectiveness for nocturnal frequency or enuresis by improvement of bladder storage functions, but the therapeutic mechanisms for the sleep functions remain unknown. In this study we investigated the effectiveness of sacral acupuncture for the sleep depth by a single-blinded, randomized controlled cross-over trials.

Methods: This is a randomized, single-blinded, cross-over study designed to examine the effects of acupuncture for the sleep functions. The randomization was performed using a randomization database programmed by a data manager A total of 10 participants (6 males, 4 females, mean 27.4 years) without severe insomnia. Sacral acupuncture was undergone with disposable stainless steel needles (0.30 × 60 mm) inserted into the bilateral BL33 points, and controlled acupuncture was done with disposable stainless steel needles (0.18 × 40 mm) inserted into the bilateral PC6 and HT7 points. The acupuncture performed total 4 weeks, every 2 weeks each. Global sleep disturbance was assessed using the Pittsburgh Sleep Quality Index (PSQI), and sleep depth was measured using smart watch. Major assessments were at baseline and post-acupuncture, alternately.

Results: No side effects were recognized throughout the acupuncture. Total PSQI score decreased significantly from 5.5 to 3.8 ($p=0.03$) 2 weeks after sacral acupuncture, however, the score did not change from 5.5 to 4.0 ($p=0.10$) 2 weeks after controlled acupuncture. On the sleep depth, the proportion of deep sleep increased significantly from 53.8% to 60.5% ($p=0.04$) 2 weeks after sacral acupuncture, however, the proportion did not change 57.7% to 56.4% ($p=0.68$) 2 weeks after controlled acupuncture.

Conclusion: Sacral acupuncture could represent increase deep sleep, may be excellent in the maintenance of sleep. These results suggest the potential for new therapeutic contributions to sleep disturbances and insomnia.

<https://doi.org/10.1016/j.imr.2020.100595>

Integrative Medicine Research 9 (2020) 100596**Development of a SCAR marker-based diagnostic method for the detection of four species of mantis**

Min Jee Kim, Wook Jin Kim, Pureum Noh,
Inkyu Park, Byeong Cheol Moon

*Herbal Medicine Resources Research Center, Korea
Institute of Oriental Medicine, Naju, Korea*

Recently, the adulteration of medicinal materials has become a primary concern of both consumers and industry for reasons of safety and efficacy. Therefore, it is essential to verify that the materials are sourced from an authentic species and necessary to monitor the commercial medicinal product for consumers. The egg case of the mantis (called *Mantidis ootheca* at pharmacopeia) is one example of a medicinal material derived from an insect, and it