

ACUPUNCTURE & ELECTRO-THERAPEUTICS RES., INT. J., Vol. 37, pp.125-138 , 2012
Copyright ©2012 Cognizant Communication Corp. Printed in the USA.
0360-1293/12 \$60.00 + .00
DOI: <http://dx.doi.org/10.3727/036012912X13831831256249>

Effects of Combining Electroacupuncture with General Anesthesia Induced by Sevoflurane in Patients Undergoing Supratentorial Craniotomy and Improvements in Their Clinical Recovery Profile & Blood Enkephalin

Caixia Yang, Ph.D.

Staff Scientist, Dept. of Laboratory Diagnosis Centre, Beijing Tiantan Hospital of Capital Medical University, Beijing, China.

Lixin An, MD.

Professor of Clinical Anesthesiology, Dept. of Neurosurgical Anesthesiology, Beijing Tiantan Hospital of Capital Medical University, Beijing, China.

Ruquan Han, MD.

Professor & Director of Clinical Anesthesiology, Dept. of Neurosurgical Anesthesiology, Beijing Tiantan Hospital of Capita lMedical University, Beijing, China.

Song Lin, MD.

Professor & Director, Dept. of Neurosurgery, Beijing Tiantan Hospital of Capital Medical University, Beijing, China.

Xixiong Kang, Ph.D.

Professor & Director, Dept. of Laboratory Diagnosis Centre, Beijing Tiantan Hospital of CapitalMedicalUniversity, Beijing, China.

Baoguo Wang, MD.

Professor of Clinical Anesthesiology, Dept. of Neurosurgical Anesthesiology, Beijing Tiantan Hospital of Capital Medical University, Beijing, China.

(Correspondence: Xixiong Kang, Ph.D. Tel. 86-010-67096881, FAX.86-010-67096881. Baoguo Wang, MD. Tel. 86-010-62856766, FAX.86-010-62856766.)

(Received: Feb 10, 2012; Accepted with Revisions: June 20, 2012)

ABSTRACT:

Drug-induced anesthesia combined with electroacupuncture (EA) in patients has been put into practice in recent years in China. In this study, we showed the effectiveness of EA on the speed of post-operative recovery of patients undergoing supratentorial craniotomy and the potential clinical mechanism of EA. Dual channel electrical stimulator made by HANS Beijing connected the following acupoints respectively: LI4 (Hegu), SJ5 (Waiguan), ST36 (Zusanli), BL63 (Jinmen), LR3 (Taichong), and GB40 (Qiuxu). Disperse-dense and

ACUPUNCTURE & ELECTRO-THERAPEUTICS RES., INT. J., Vol. 37, pp.125-138 , 2012
Copyright ©2012 Cognizant Communication Corp. Printed in the USA.
0360-1293/12 \$60.00 + .00
DOI: <http://dx.doi.org/10.3727/036012912X13831831256249>

Effects of Combining Electroacupuncture with General Anesthesia Induced by Sevoflurane in Patients Undergoing Supratentorial Craniotomy and Improvements in Their Clinical Recovery Profile & Blood Enkephalin

Caixia Yang, Ph.D.

Staff Scientist, Dept. of Laboratory Diagnosis Centre, Beijing Tiantan Hospital of Capital Medical University, Beijing, China.

Lixin An, MD.

Professor of Clinical Anesthesiology, Dept. of Neurosurgical Anesthesiology, Beijing Tiantan Hospital of Capital Medical University, Beijing, China.

Ruquan Han, MD.

Professor & Director of Clinical Anesthesiology, Dept. of Neurosurgical Anesthesiology, Beijing Tiantan Hospital of Capita lMedical University, Beijing, China.

Song Lin, MD.

Professor & Director, Dept. of Neurosurgery, Beijing Tiantan Hospital of Capital Medical University, Beijing, China.

Xixiong Kang, Ph.D.

Professor & Director, Dept. of Laboratory Diagnosis Centre, Beijing Tiantan Hospital of CapitalMedicalUniversity, Beijing, China.

Baoguo Wang, MD.

Professor of Clinical Anesthesiology, Dept. of Neurosurgical Anesthesiology, Beijing Tiantan Hospital of Capital Medical University, Beijing, China.

(Correspondence: Xixiong Kang, Ph.D. Tel. 86-010-67096881, FAX.86-010-67096881. Baoguo Wang, MD. Tel. 86-010-62856766, FAX.86-010-62856766.)

(Received: Feb 10, 2012; Accepted with Revisions: June 20, 2012)

ABSTRACT:

Drug-induced anesthesia combined with electroacupuncture (EA) in patients has been put into practice in recent years in China. In this study, we showed the effectiveness of EA on the speed of post-operative recovery of patients undergoing supratentorial craniotomy and the potential clinical mechanism of EA. Dual channel electrical stimulator made by HANS Beijing connected the following acupoints respectively: LI4 (Hegu), SJ5 (Waiguan), ST36 (Zusanli), BL63 (Jinmen), LR3 (Taichong), and GB40 (Qiuxu). Disperse-dense and

symmetric biphasic pulse waves were selected, frequency of waves (pulse rates) were 2Hz/100Hz, altered/3sec; pulse duration was 0.6ms/0.2ms, 2Hz: 0.6ms, 100Hz: 0.2ms; symmetric biphasic pulse wave. We found that the EA-group required 9.62% less sevoflurane than the sham EA-group (P<0.05). During recovery from anesthesia, the autonomous respiration recovery time, tracheo-tube removal time, eye-opening time, voluntary motor recovery time, orientation force recovery time, and the operating-room departure time of the EA-group were all significantly shortened 35.86%, 27.07%, 38.38%, 30.11%, 34.95%, 28.80% than the corresponding sham EA-group, respectively (P<0.05). The serum enkephalin values were elevated in the EA group versus the sham EA-group.

Keywords: Glioma; Craniotomy; Electroacupuncture; Analgesia; Sevoflurane; Anesthesia; Recovery time after surgery; Enkephalin.

INTRODUCTION

Gliomas are the most serious type of brain tumors in adults, including astrocytic tumor、oligodendroglioma tumors、mixed gliomas and ependymal tumors. They originate in neuroectoderm, grow in an uninhibited fashion, diffusely infiltrate the brain, make surgical resectioning difficult or even impossible. Among them, glioblastoma multiforme (GBM) is the most common form of malignant brain cancer in adults. Patients with GBM have a uniformly poor prognosis, with a median survival of one year[1]. Surgical resection is the most effective therapeutic method to date. Despite the availability of newer analgesic drugs and techniques, concerns remain on the side effects of both opioid and nonopioid analgesic techniques, such as nausea and vomiting, hypotension, urinary retention and ileus, pruritis, constipation and even respiratory depression. For patients undergoing craniotomy, these side effects may lead to devastating results owing to the increased risk of intracranial hemorrhage and neurological dysfunction [2]. The incidence of side effects is related in part to the total dose of opioid analgesic medication [3]. These side effects can be reduced by using minimal flow rates of the opioid dosage but cannot be completely eliminated due to the need of analgesic medication [4]. Acupuncture is a complementary non-pharmacological therapy with few side effects of analgesic medication. Its importance for health care in Asian culture can be traced back almost 3000 yrs, and it has been practiced to treat a variety of pathological symptoms and conditions [5]. Acupuncture in conjunction with electrical stimulation techniques (e.g., electroacupuncture, transcutaneous electrical nerve stimulation [TENS]) are based on traditional Chinese analgesic methods. EA is an inexpensive, safe, and easy stimulation technique that has been used for the management of pain following a wide variety of surgical procedures [6,7]. The present study was randomized, sham controlled, and blinded design to objectively test the hypothesis that EA would: (1) decrease the amount of sevoflurane needed for anesthesia; (2) facilitate regulation of the physiological and pathological changes of the human body during surgery; and (3) improve the overall recovery profiles for the post-operative patient. Further, we have assayed serum levels of enkephalin to assess physiological changes in response to EA.

symmetric biphasic pulse waves were selected, frequency of waves (pulse rates) were 2Hz/100Hz, altered/3sec; pulse duration was 0.6ms/0.2ms, 2Hz: 0.6ms, 100Hz: 0.2ms; symmetric biphasic pulse wave. We found that the EA-group required 9.62% less sevoflurane than the sham EA-group (P<0.05). During recovery from anesthesia, the autonomous respiration recovery time, tracheo-tube removal time, eye-opening time, voluntary motor recovery time, orientation force recovery time, and the operating-room departure time of the EA-group were all significantly shortened 35.86%, 27.07%, 38.38%, 30.11%, 34.95%, 28.80% than the corresponding sham EA-group, respectively (P<0.05). The serum enkephalin values were elevated in the EA group versus the sham EA-group.

Keywords: Glioma; Craniotomy; Electroacupuncture; Analgesia; Sevoflurane; Anesthesia; Recovery time after surgery; Enkephalin.

INTRODUCTION

Gliomas are the most serious type of brain tumors in adults, including astrocytic tumor、oligodendroglioma tumors、mixed gliomas and ependymal tumors. They originate in neuroectoderm, grow in an uninhibited fashion, diffusely infiltrate the brain, make surgical resectioning difficult or even impossible. Among them, glioblastoma multiforme (GBM) is the most common form of malignant brain cancer in adults. Patients with GBM have a uniformly poor prognosis, with a median survival of one year[1]. Surgical resection is the most effective therapeutic method to date. Despite the availability of newer analgesic drugs and techniques, concerns remain on the side effects of both opioid and nonopioid analgesic techniques, such as nausea and vomiting, hypotension, urinary retention and ileus, pruritis, constipation and even respiratory depression. For patients undergoing craniotomy, these side effects may lead to devastating results owing to the increased risk of intracranial hemorrhage and neurological dysfunction [2]. The incidence of side effects is related in part to the total dose of opioid analgesic medication [3]. These side effects can be reduced by using minimal flow rates of the opioid dosage but cannot be completely eliminated due to the need of analgesic medication [4]. Acupuncture is a complementary non-pharmacological therapy with few side effects of analgesic medication. Its importance for health care in Asian culture can be traced back almost 3000 yrs, and it has been practiced to treat a variety of pathological symptoms and conditions [5]. Acupuncture in conjunction with electrical stimulation techniques (e.g., electroacupuncture, transcutaneous electrical nerve stimulation [TENS]) are based on traditional Chinese analgesic methods. EA is an inexpensive, safe, and easy stimulation technique that has been used for the management of pain following a wide variety of surgical procedures [6,7]. The present study was randomized, sham controlled, and blinded design to objectively test the hypothesis that EA would: (1) decrease the amount of sevoflurane needed for anesthesia; (2) facilitate regulation of the physiological and pathological changes of the human body during surgery; and (3) improve the overall recovery profiles for the post-operative patient. Further, we have assayed serum levels of enkephalin to assess physiological changes in response to EA.

MATERIALS AND METHODS

Eighty patients having an American Society of Anesthesiologists (ASA) classification of physical status I - II and with supratentorial tumors were recruited for this study. Written informed consents were obtained from all patients. The study protocol was approved by the Institutional Review Board at Beijing Tiantan Hospital. Equal numbers of patients were randomly assigned to the sham EA-group and the EA- group. The acupoints included for the study were LI4 (Hegu), SJ5 (Waiguan), ST36 (Zusanli), BL63 (Jinmen), LR3 (Taichong), GB40 (Qixu) (Fig. 1).





Fig. 1A: The groups and acupoints of the study.



MATERIALS AND METHODS

Eighty patients having an American Society of Anesthesiologists (ASA) classification of physical status I - II and with supratentorial tumors were recruited for this study. Written informed consents were obtained from all patients. The study protocol was approved by the Institutional Review Board at Beijing Tiantan Hospital. Equal numbers of patients were randomly assigned to the sham EA-group and the EA- group. The acupoints included for the study were LI4 (Hegu), SJ5 (Waiguan), ST36 (Zusanli), BL63 (Jinmen), LR3 (Taichong), GB40 (Qixu) (Fig. 1).




Fig. 1A: The groups and acupoints of the study.




Fig. 1B: The locations of LI4 (Hegu) and SJ5 (Waiguan): Hegu (LI4), on the dorsum of the hand, between the first and the second metacarpal bones, on the radial side of the midpoint of the second metacarpal bone. Waiguan (SJ5), on the dorsum of the forearm, 2 cun proximal to the dorsal wrist crease, between the radius and the ulna.




Fig. 1C: The locations of ST36 (Zusanli), BL63 (Jinmen), LR3 (Taichong), and GB40 (Qixu): Zusanli(ST36): On the anterior aspect of lower leg, one finger width (thumb finger) lateral to the anterior border of tibia. Jinmen (BL63): On the lateral of the foot, anterior to the lateral malleolar, inferior to the cuboid bone. Taichong (LR3): On the dorsum of the foot, at the depression of the posterior side of the first interosseous metatarsal space. Qixu (GB40): anterior to the lateral malleolar,in the depression lateral to the vaginae tendinum musculi extensoris digitorum pedis longi.

Excluded from the study were: (1) pregnant and/or lactating women; (2) patients with clinically significant cardiovascular problems or pulmonary diseases; (3) long term smokers; (4) obese patients (BMI > 30); (5) emergency patients; and (6) patients with bleeding > 2,500 mL during surgery and/or patients having a surgical duration > 8 h.

Anesthesia was induced with 0.3 µg/kg sufentanil i.v.and 2 mg/kg propofol i.v. Tracheal intubation was facilitated with 0.1 mg/kg vecuronium. Ventilation was controlled to maintain end-tidal PaCO₂ at 28-33 mm Hg after intubation. General anesthesia was initially maintained with sevoflurane 2% in oxygen 1L/Min. Vecuronium top-up doses were administered as guided by TOF monitor (TOF watch SX monitor). Before incision,0.5% ropivacaine was used for scalp infiltration, 0.15 µg/kg sufentanil was administered before fixing the head brace and after

Fig. 1B: The locations of LI4 (Hegu) and SJ5 (Waiguan): Hegu (LI4), on the dorsum of the hand, between the first and the second metacarpal bones, on the radial side of the midpoint of the second metacarpal bone. Waiguan (SJ5), on the dorsum of the forearm, 2 cun proximal to the dorsal wrist crease, between the radius and the ulna.




Fig. 1C: The locations of ST36 (Zusanli), BL63 (Jinmen), LR3 (Taichong), and GB40 (Qixu): Zusanli(ST36): On the anterior aspect of lower leg, one finger width (thumb finger) lateral to the anterior border of tibia. Jinmen (BL63): On the lateral of the foot, anterior to the lateral malleolar, inferior to the cuboid bone. Taichong (LR3): On the dorsum of the foot, at the depression of the posterior side of the first interosseous metatarsal space. Qixu (GB40): anterior to the lateral malleolar,in the depression lateral to the vaginae tendinum musculi extensoris digitorum pedis longi.

Excluded from the study were: (1) pregnant and/or lactating women; (2) patients with clinically significant cardiovascular problems or pulmonary diseases; (3) long term smokers; (4) obese patients (BMI > 30); (5) emergency patients; and (6) patients with bleeding > 2,500 mL during surgery and/or patients having a surgical duration > 8 h.

Anesthesia was induced with 0.3 µg/kg sufentanil i.v.and 2 mg/kg propofol i.v. Tracheal intubation was facilitated with 0.1 mg/kg vecuronium. Ventilation was controlled to maintain end-tidal PaCO₂ at 28-33 mm Hg after intubation. General anesthesia was initially maintained with sevoflurane 2% in oxygen 1L/Min. Vecuronium top-up doses were administered as guided by TOF monitor (TOF watch SX monitor). Before incision,0.5% ropivacaine was used for scalp infiltration, 0.15 µg/kg sufentanil was administered before fixing the head brace and after

suturing the duramater. The dose of sufentanil was fixed for all patients. At the end of surgery, sevoflurane was stopped.

A trained acupuncturist determined the locations for acupoints: LI4 (Hegu), SJ 5(Waiguan), ST36 (Zusanli), BL63 (Jinmen), LR3 (Taichong), GB40 (Qixu) ,based on the Traditional Chinese Medicine meridian theory. The needles were inserted at the specific points (17–33 mm depth, depending upon the build of patient) in the electroacupuncture group. A De Qi sensation was ensured by attaching small crocodile clips to the end of the needles to connect them to an electroacupuncture device (HANS dual-channel unit; LH202H, Huawei, Beijing) [8] (Fig. 2).




Fig. 2A: HANS dual-channel unit, LH202H, rechargeable batteries electrical stimulator made in Beijing, China.



Fig. 2B: Patient's acupoints: LI4 (Hegu) and SJ5 (Waiguan) connected to HANS electrical stimulator in EA-group. Small crocodile clips attached to the end of the needles to connect them to an electroacupuncture device.

suturing the duramater. The dose of sufentanil was fixed for all patients. At the end of surgery, sevoflurane was stopped.

A trained acupuncturist determined the locations for acupoints: LI4 (Hegu), SJ 5(Waiguan), ST36 (Zusanli), BL63 (Jinmen), LR3 (Taichong), GB40 (Qixu) ,based on the Traditional Chinese Medicine meridian theory. The needles were inserted at the specific points (17–33 mm depth, depending upon the build of patient) in the electroacupuncture group. A De Qi sensation was ensured by attaching small crocodile clips to the end of the needles to connect them to an electroacupuncture device (HANS dual-channel unit; LH202H, Huawei, Beijing) [8] (Fig. 2).




Fig. 2A: HANS dual-channel unit, LH202H, rechargeable batteries electrical stimulator made in Beijing, China.



Fig. 2B: Patient's acupoints: LI4 (Hegu) and SJ5 (Waiguan) connected to HANS electrical stimulator in EA-group. Small crocodile clips attached to the end of the needles to connect them to an electroacupuncture device.



Fig. 2C: Patient’s acupoints:ST36 (Zusanli), BL63(Jinmen), LR3(Taichong), GB40 (Qixu) connected to HANS electrical stimulator in EA-group. Small crocodile clips attached to the end of the needles to connect them to an electroacupuncture device.

The electroacupuncture device was a biphasic pulse generator and was used with a maximum tolerable intensity of current and frequency: disperse-dense waves, frequency of waves(pulse rates)were 2 Hz/ 100 Hz[9], altered / 3second, pulse duration was 0.6ms/0.2ms; 2 Hz : 0.6ms; 100 Hz: 0.2ms; symmetric biphasic pulse wave (Fig. 3).




Fig.3: Diagram of the waveform of electroacupuncture.

The acupuncture was applied and activated before the induction of anesthesia in each EA patient, and was maintained until completion of the operation (completion of skin closure). In controls (the sham EA- group), the electric wire was placed at the surface of the chosen acupoint sites, however, no needles were inserted or current applied.

The end-tidal sevoflurane concentration and the speed of recovery were recorded, including the autonomous respiration recovery time (ARRT), tracheo-tube removal time (TTRT), eye opening time (EOT), voluntary motion recovery time (VMRT), orientation force recovery time (OFRT), and operating room departure time (ORDT). The start point for timing the above events was the cessation of sevoflurane administration for each patient in the 2 groups.



Fig. 2C: Patient’s acupoints:ST36 (Zusanli), BL63(Jinmen), LR3(Taichong), GB40 (Qixu) connected to HANS electrical stimulator in EA-group. Small crocodile clips attached to the end of the needles to connect them to an electroacupuncture device.

The electroacupuncture device was a biphasic pulse generator and was used with a maximum tolerable intensity of current and frequency: disperse-dense waves, frequency of waves(pulse rates)were 2 Hz/ 100 Hz[9], altered / 3second, pulse duration was 0.6ms/0.2ms; 2 Hz : 0.6ms; 100 Hz: 0.2ms; symmetric biphasic pulse wave (Fig. 3).




Fig.3: Diagram of the waveform of electroacupuncture.

The acupuncture was applied and activated before the induction of anesthesia in each EA patient, and was maintained until completion of the operation (completion of skin closure). In controls (the sham EA- group), the electric wire was placed at the surface of the chosen acupoint sites, however, no needles were inserted or current applied.

The end-tidal sevoflurane concentration and the speed of recovery were recorded, including the autonomous respiration recovery time (ARRT), tracheo-tube removal time (TTRT), eye opening time (EOT), voluntary motion recovery time (VMRT), orientation force recovery time (OFRT), and operating room departure time (ORDT). The start point for timing the above events was the cessation of sevoflurane administration for each patient in the 2 groups.

A five milliliter blood sample was drawn from each patient using the femoral vein under aseptic conditions, during general anesthesia at 0, 30, 45, 60, 120 and 240 min. Blood samples were immediately transferred to vacutainers and centrifuged to facilitate serum collection. Serum was transferred to storage vials and kept at -80°C until analysis. Quantification of serum enkephalin was performed using ELISA, i.e. (Human Enkephalin Elisa Kit. Catalog No:E01E0312). Valid samples were 13 and 17 in control and EA-group respectively.

The data were entered into EXCEL data sheets and analyzed using the SPSS 13.0 statistical package. Data from the control and EA groups (sevoflurane requirement and the recover profiles) were compared using the Student *t* test. The time-dependent intragroup data for serum enkephalin were evaluated by repeated-measures and analysis of variance. *P* < 0.05 was considered statistically significant. Data were expressed as the mean ± SD.

RESULTS

In this study no patient had previously received an EA treatment. All patients were enrolled based on the inclusion criteria (above) and were randomly assigned to the two groups. No complications related to EA procedures were observed during the operation. The two treatment groups were comparable with respect to both demographic and clinical variables. There were no significant intergroup differences in patients' age, weight, height, or anesthesia time (Table 1).

General conditions	Sham EA-group (n=40)	EA- group (n=40)
Age (years)	39±11	43±11
Weigh (kg)	66±10	69±15
Male/female (cases)	21/ 19	17/ 23
Anesthesia time (min)	211±56	233±56
Meningioma	17	11
Glioma	11	23
Metastatic tumor	2	6
Intraoperative sufentanil (µg)	38±4	37±1
Data presented as mean±SD or as a number (n) of patients.		

Table 1: Patients' characteristics and perioperative use of analgesics.

The consumption of sevoflurane during operation was shown in Table 2.

Stages of surgery	Sevoflurane end-tidal concentration in Sham EA-group (%); n=40	Sevoflurane end-tidal concentration in EA-group (%); n=40	Sham EA-group vs EA-group sevoflurane reduction (%)	P
Skin incising	1.60 ± 0.37	1.50 ± 0.26	6.25	0.263
Skull drilling	1.72 ± 0.43	1.58 ± 0.32	8.13	0.175
After skull drilling	1.74 ± 0.39	1.57 ± 0.31	9.19	0.032*

A five milliliter blood sample was drawn from each patient using the femoral vein under aseptic conditions, during general anesthesia at 0, 30, 45, 60, 120 and 240 min. Blood samples were immediately transferred to vacutainers and centrifuged to facilitate serum collection. Serum was transferred to storage vials and kept at -80°C until analysis. Quantification of serum enkephalin was performed using ELISA, i.e. (Human Enkephalin Elisa Kit. Catalog No:E01E0312). Valid samples were 13 and 17 in control and EA-group respectively.

The data were entered into EXCEL data sheets and analyzed using the SPSS 13.0 statistical package. Data from the control and EA groups (sevoflurane requirement and the recover profiles) were compared using the Student *t* test. The time-dependent intragroup data for serum enkephalin were evaluated by repeated-measures and analysis of variance. *P* < 0.05 was considered statistically significant. Data were expressed as the mean ± SD.

RESULTS

In this study no patient had previously received an EA treatment. All patients were enrolled based on the inclusion criteria (above) and were randomly assigned to the two groups. No complications related to EA procedures were observed during the operation. The two treatment groups were comparable with respect to both demographic and clinical variables. There were no significant intergroup differences in patients' age, weight, height, or anesthesia time (Table 1).

General conditions	Sham EA-group (n=40)	EA- group (n=40)
Age (years)	39±11	43±11
Weigh (kg)	66±10	69±15
Male/female (cases)	21/ 19	17/ 23
Anesthesia time (min)	211±56	233±56
Meningioma	17	11
Glioma	11	23
Metastatic tumor	2	6
Intraoperative sufentanil (µg)	38±4	37±1
Data presented as mean±SD or as a number (n) of patients.		

Table 1: Patients' characteristics and perioperative use of analgesics.

The consumption of sevoflurane during operation was shown in Table 2.

Stages of surgery	Sevoflurane end-tidal concentration in Sham EA-group (%); n=40	Sevoflurane end-tidal concentration in EA-group (%); n=40	Sham EA-group vs EA-group sevoflurane reduction (%)	P
Skin incising	1.60 ± 0.37	1.50 ± 0.26	6.25	0.263
Skull drilling	1.72 ± 0.43	1.58 ± 0.32	8.13	0.175
After skull drilling	1.74 ± 0.39	1.57 ± 0.31	9.19	0.032*

After skull opening	1.75 ± 0.39	1.58 ± 0.30	9.19	0.035*
Dura incising	1.79 ± 0.43	1.58 ± 0.28	11.73	0.014*
Intracranial surgery 10 min	1.78 ± 0.45	1.56 ± 0.26	12.35	0.010*
Intracranial surgery 20 min	1.76 ± 0.44	1.54 ± 0.26	12.50	0.010*
Intracranial surgery 30 min	1.80 ± 0.44	1.59 ± 0.28	11.66	0.022*
Dura suturing	1.83 ± 0.39	1.70 ± 0.28	7.10	0.110
Skin suturing	1.73 ± 0.37	1.59 ± 0.23	8.09	0.620
*P<0.05				

Table 2: The consumption of sevoflurane during operation. 9.62% decreased in sevoflurane requirement were produced in EA-group compared to sham EA-group. ARRT, TTRT, EOT, VMRT, OFRT and ORDT after stopping administration of sevoflurane in EA group were all less than in control group.

The recovery profiles were shown in Table3.

Recovery of anesthesia	Sham EA-group	n	EA-group	n	Sham EA-group vs EA-group sevoflurane reduction (%)	P
RRT	12.69	32	8.14	29	35.86	0.013*
TTRT	21.32	31	15.55	29	27.07	0.005*
EOT	28.52	31	17.57	28	38.38	0.000*
VMRT	29.47	30	20.59	27	30.11	0.002*
OFRT	34.17	30	22.22	27	34.95	0.000*
ORDT	39.32	31	28.00	28	28.80	0.000*
*p < 0.05						

Table 3: The recovery profiles (min): ARRT: autonomous respiration recovery time; TTRT: tracheo-tube removal time; EOT: eye-opening time; VMRT: voluntary motor recovery time; OFRT: orientation force recovery time; and ORDT: operating room departure time.

Blood enkephalin, a pain relieving neurotransmitter, was consistently higher in the control than in the EA-group during the operative period according to ELISA measurements (P=0.011; Fig. 4).

After skull opening	1.75 ± 0.39	1.58 ± 0.30	9.19	0.035*
Dura incising	1.79 ± 0.43	1.58 ± 0.28	11.73	0.014*
Intracranial surgery 10 min	1.78 ± 0.45	1.56 ± 0.26	12.35	0.010*
Intracranial surgery 20 min	1.76 ± 0.44	1.54 ± 0.26	12.50	0.010*
Intracranial surgery 30 min	1.80 ± 0.44	1.59 ± 0.28	11.66	0.022*
Dura suturing	1.83 ± 0.39	1.70 ± 0.28	7.10	0.110
Skin suturing	1.73 ± 0.37	1.59 ± 0.23	8.09	0.620
*P<0.05				

Table 2: The consumption of sevoflurane during operation. 9.62% decreased in sevoflurane requirement were produced in EA-group compared to sham EA-group. ARRT, TTRT, EOT, VMRT, OFRT and ORDT after stopping administration of sevoflurane in EA group were all less than in control group.

The recovery profiles were shown in Table3.

Recovery of anesthesia	Sham EA-group	n	EA-group	n	Sham EA-group vs EA-group sevoflurane reduction (%)	P
RRT	12.69	32	8.14	29	35.86	0.013*
TTRT	21.32	31	15.55	29	27.07	0.005*
EOT	28.52	31	17.57	28	38.38	0.000*
VMRT	29.47	30	20.59	27	30.11	0.002*
OFRT	34.17	30	22.22	27	34.95	0.000*
ORDT	39.32	31	28.00	28	28.80	0.000*
*p < 0.05						

Table 3: The recovery profiles (min): ARRT: autonomous respiration recovery time; TTRT: tracheo-tube removal time; EOT: eye-opening time; VMRT: voluntary motor recovery time; OFRT: orientation force recovery time; and ORDT: operating room departure time.

Blood enkephalin, a pain relieving neurotransmitter, was consistently higher in the control than in the EA-group during the operative period according to ELISA measurements (P=0.011; Fig. 4).




Fig. 4: Detected enkephalin in EA-group and sham EA-group. Enkephalin was consistently higher in sham EA-group than in EA-group during the operative period according to ELISA measurements ($P=0.011$).

DISCUSSION

Omura [10] reports that the best pulse rates for average human adult is same as pulse rate of the patient which is synchronized with local blood circulation between 1 and 1.5 pulses / sec. Any pulse rate over 2 pulses/sec in human adult creates various degrees of local circulatory disturbance and reduces pain relief time duration depending on pulse repetition rate and other stimulating parameters. For prolonged (post-stimulation) effects, typically equating to 1.2-3pps [11-13]. Richard P. Francis [14] believes that the characteristics of acupuncture-like TENS include pulse rate between 1-5pps; a pulse duration between 100 μ s and 200 μ s. During our study, according to the release of analgesic substances and synergistic effect, we selected 2Hz/100 Hz frequency to work. Han JS [9] indicated that enkephalin and endorphin could be released with 2Hz electric stimulation, and the release of dynorphin need 100Hz electric stimulation. 2Hz EA produced a 7 fold increase in met-enkephalin [15]. Radioimmunoassay has shown that rat brain levels of beta -endorphin are lower than those of the enkephalins [16]. Enkephalin within vagal nuclei of the medulla might play a role in visceral reflexes affected by opiates, such as the cough and vomiting reflexes. Within the periaqueductal gray, enkephalin may mediate electrical stimulation-induced analgesia [17-19]. Enkephalin in the infundibulum and periventricular nuclei of the hypothalamus may reflect loci where opiates influence endocrine functions. Within certain limbic and cortical areas, especially the central nucleus of the amygdala, enkephalin might be




Fig. 4: Detected enkephalin in EA-group and sham EA-group. Enkephalin was consistently higher in sham EA-group than in EA-group during the operative period according to ELISA measurements ($P=0.011$).

DISCUSSION

Omura [10] reports that the best pulse rates for average human adult is same as pulse rate of the patient which is synchronized with local blood circulation between 1 and 1.5 pulses / sec. Any pulse rate over 2 pulses/sec in human adult creates various degrees of local circulatory disturbance and reduces pain relief time duration depending on pulse repetition rate and other stimulating parameters. For prolonged (post-stimulation) effects, typically equating to 1.2-3pps [11-13]. Richard P. Francis [14] believes that the characteristics of acupuncture-like TENS include pulse rate between 1-5pps; a pulse duration between 100 μ s and 200 μ s. During our study, according to the release of analgesic substances and synergistic effect, we selected 2Hz/100 Hz frequency to work. Han JS [9] indicated that enkephalin and endorphin could be released with 2Hz electric stimulation, and the release of dynorphin need 100Hz electric stimulation. 2Hz EA produced a 7 fold increase in met-enkephalin [15]. Radioimmunoassay has shown that rat brain levels of beta -endorphin are lower than those of the enkephalins [16]. Enkephalin within vagal nuclei of the medulla might play a role in visceral reflexes affected by opiates, such as the cough and vomiting reflexes. Within the periaqueductal gray, enkephalin may mediate electrical stimulation-induced analgesia [17-19]. Enkephalin in the infundibulum and periventricular nuclei of the hypothalamus may reflect loci where opiates influence endocrine functions. Within certain limbic and cortical areas, especially the central nucleus of the amygdala, enkephalin might be

associated with the euphoric effects of opiates[16]. Beta -endorphin is a potent analgetic, being 18 to 33 times a more potent than morphineon a molar basis. Chronic, localized infusion of beta-endorphin into the rat brain can induce morphine-like physical dependence [20]. Many brain nuclei composing a complicated network are involved in processing acupuncture analgesia, including the nucleus raphe magnus (NRM), periaqueductal grey (PAG), locus coeruleus, arcuate nucleus (Arc), preoptic area, nucleus submedius, habenular nucleus, accumbens nucleus, caudate nucleus, septal area, amygdale, etc[21]. The result of our study confirmed the release of serum enkephalin. More research will continue to study the effects of 100Hz electric stimulation on circulatory disturbance.

This study suggests that EA at the acupoints (Hegu, Waiguan, Zusanli, Jinmen, Taichong, and Qiu xu) combined with sevoflurane could improve recovery profiles of patients undergoing supratentorial craniotomy. Patients from EA group required 9.62% less sevoflurane than the control patients during surgery ($p<0.05$). It indicates that EA had effect of anesthesia. Acupuncture analgesia is the basis of acupuncture anesthesia [22-26]. Previous studies have shown the effects of EA on painful, such as: neck pain, shoulder pain, elbow pain, low back pain, knee pain etc [27-30]. EA may be a complementary method for anesthesia that could minimize the side effects from anesthetics. However, EA cannot replace the role of anesthetics. Our results show that patients' recovery profiles of EA, including autonomous respiration recovery time (ARRT), tracheotube removing time (TTRT), eye opening time (EOT), voluntary motion recovery time (VMRT), orientation force recovery time (OFRT), and operating room departure time (ORDT) after stopping administration of sevoflurane, are all less than control group. These effects may be related to the decreased sevoflurane requirement to maintain anesthesia. The most important function about EA may be the regulation for body.

Acupuncture is an intricate therapeutic system. Its mechanisms include traditional Chinese theories along with western anatomy and physiology. Traditional Chinese acupuncture presumes that there are two opposing and complementary forces that coexist in nature: Yin and Yang. These two forces interact to regulate the flow of “vital energy” known as Qi. It is defined that good health is a balance between yin and yang, in this situation, Qi flows freely through a system of channels (meridians), which bring Qi from the internal organs to the skin's surface. Along these meridians there are acupuncture points that can be stimulated to correct the imbalance and restore the body to normal health. In disease this flow is disrupted causing an imbalance between yin and yang. Needles inserted into specific points along meridians are said to regulate imbalances in the flow of Qi and treat diseases. According to the experiences of curing headaches using acupuncture, the traditional Chinese theories propose that: distant acupoints treat proximal disease. We have chosen the distant acupoints: Hegu, Waiguan, Jinmen, Taichong, Zusanli and Qiu xu to insert needles. Acupoints usage is effective for treatment, the traditional Chinese explanation has little “scientific” basis. Western scientific explanations of EA are based on the presumption that acupuncture induces signals in afferent nerves that modulate spinal signal transmission and pain perception in brain. Neural mechanisms that have been proposed for short-term analgesic effects of acupuncture include spinal gating of descending inhibitory pathways. Spinal gating pain is mediated by two types of peripheral fibers: myelinated A-fibers responsible for sharp, well-localized pain (‘fast’pain) and smaller unmyelinated C-fibers responsible for prolonged and ill-defined pain (‘slow’pain) [31]. EA stimulates the large A, β -fibers and activates the gating mechanism. This may inhibit central pain transmission

associated with the euphoric effects of opiates[16]. Beta -endorphin is a potent analgetic, being 18 to 33 times a more potent than morphineon a molar basis. Chronic, localized infusion of beta-endorphin into the rat brain can induce morphine-like physical dependence [20]. Many brain nuclei composing a complicated network are involved in processing acupuncture analgesia, including the nucleus raphe magnus (NRM), periaqueductal grey (PAG), locus coeruleus, arcuate nucleus (Arc), preoptic area, nucleus submedius, habenular nucleus, accumbens nucleus, caudate nucleus, septal area, amygdale, etc[21]. The result of our study confirmed the release of serum enkephalin. More research will continue to study the effects of 100Hz electric stimulation on circulatory disturbance.

This study suggests that EA at the acupoints (Hegu, Waiguan, Zusanli, Jinmen, Taichong, and Qiu xu) combined with sevoflurane could improve recovery profiles of patients undergoing supratentorial craniotomy. Patients from EA group required 9.62% less sevoflurane than the control patients during surgery ($p<0.05$). It indicates that EA had effect of anesthesia. Acupuncture analgesia is the basis of acupuncture anesthesia [22-26]. Previous studies have shown the effects of EA on painful, such as: neck pain, shoulder pain, elbow pain, low back pain, knee pain etc [27-30]. EA may be a complementary method for anesthesia that could minimize the side effects from anesthetics. However, EA cannot replace the role of anesthetics. Our results show that patients' recovery profiles of EA, including autonomous respiration recovery time (ARRT), tracheotube removing time (TTRT), eye opening time (EOT), voluntary motion recovery time (VMRT), orientation force recovery time (OFRT), and operating room departure time (ORDT) after stopping administration of sevoflurane, are all less than control group. These effects may be related to the decreased sevoflurane requirement to maintain anesthesia. The most important function about EA may be the regulation for body.

Acupuncture is an intricate therapeutic system. Its mechanisms include traditional Chinese theories along with western anatomy and physiology. Traditional Chinese acupuncture presumes that there are two opposing and complementary forces that coexist in nature: Yin and Yang. These two forces interact to regulate the flow of “vital energy” known as Qi. It is defined that good health is a balance between yin and yang, in this situation, Qi flows freely through a system of channels (meridians), which bring Qi from the internal organs to the skin's surface. Along these meridians there are acupuncture points that can be stimulated to correct the imbalance and restore the body to normal health. In disease this flow is disrupted causing an imbalance between yin and yang. Needles inserted into specific points along meridians are said to regulate imbalances in the flow of Qi and treat diseases. According to the experiences of curing headaches using acupuncture, the traditional Chinese theories propose that: distant acupoints treat proximal disease. We have chosen the distant acupoints: Hegu, Waiguan, Jinmen, Taichong, Zusanli and Qiu xu to insert needles. Acupoints usage is effective for treatment, the traditional Chinese explanation has little “scientific” basis. Western scientific explanations of EA are based on the presumption that acupuncture induces signals in afferent nerves that modulate spinal signal transmission and pain perception in brain. Neural mechanisms that have been proposed for short-term analgesic effects of acupuncture include spinal gating of descending inhibitory pathways. Spinal gating pain is mediated by two types of peripheral fibers: myelinated A-fibers responsible for sharp, well-localized pain (‘fast’pain) and smaller unmyelinated C-fibers responsible for prolonged and ill-defined pain (‘slow’pain) [31]. EA stimulates the large A, β -fibers and activates the gating mechanism. This may inhibit central pain transmission

through enkephalin dependent mechanisms [32,33]. Peripheral noxious stimulus, such as pinpricks, results in impulses ascending in the spinothalamic tracts to the thalamus. At the midbrain, collateral pathways project to the periaqueductal grey area (PAG). Electrical stimulation of the PAG leads to analgesia that is reversed by naloxone [34]. Neurons from the PAG project to the nucleus raphe magnus of the medulla and from there serotonergic fibers descend to the stalked cells of the dorsal horn where release of enkephalin inhibits SG (spiral ganglion) cells [32,35], thereby blocking the centripetal transmission of pain from C-fibers. Pomeranz proposed that acupuncture stimulation activates A δ -and C afferent fibers in muscle, causing signals to be transmitted to the spinal cord and results in release of dynorphin and enkephalins. Pomeranz's theory was confirmed by a large series of experiments in his research laboratory and other investigators [36-41]. The regulation of enkephalin peptide synthesis was mainly at the mRNA level [42] and EA was shown to enhance preproenkephalin mRNA expression in rostral ventrolateral medulla of rats [43]. It suggests that there would be more peptides to produce following EA [44]. In our study, enkephalin did not increase in EA-group compared with sham EA-group. The reasons maybe include the flowing items:(1) the blood samples detected enkephalin could not consistent with the quantity of EA-group (n=40) and sham EA-group(n=40) repectively; this maybe the limitation of our study; (2) the exact mechanisms of anesthetics remain unclear to date; maybe the change of enkephalin in EA-group is the result of the combination EA and anesthetics. Our results indicate that dynamic change of enkephalin during operations in EA is different from control. The tendency of increased enkephalin in EA suggests that enkephalin might play a role for general anesthesia in pain reduction.

CONCLUSION

EA anesthesia could not replace anesthetics completely. However, EA has become the most popular complementary and alternative medicine modality and is highly praised by physicians and anesthetists. Due to the limited number of blood samples available to us, future research maybe needed to assess enkephalin and mechanisms of EA treatment.

ACKNOWLEDGEMENTS

This work was supported by funding from the 973 traditional Chinese medicine theory of special (2007 CB 512503) and Dr. Nathaniel Mills, Dept. of Biology, Texas Woman's University.

REFERENCES

1. Ohgaki, H., & Kleihues, P. Epidemiology and etiology of gliomas. Acta Neuropathologica Vol.109, pp. 93–108, 2005.
2. Wang, X.Q., Yu, J.L., Du, Z.Y., Xu, R., Jiang, C.C., & Gao, X. Electroacupoint Stimulation for Postoperative Nausea and Vomiting in Patients Undergoing Supratentorial Craniotomy. Journal of Neurosurgical Anesthesiology Vol. 22, pp. 128-131, 2010.
3. White, P.F. Mishaps with patient-controlled analgesia. Anesthesiology Vol. 65, pp. 81-83,

through enkephalin dependent mechanisms [32,33]. Peripheral noxious stimulus, such as pinpricks, results in impulses ascending in the spinothalamic tracts to the thalamus. At the midbrain, collateral pathways project to the periaqueductal grey area (PAG). Electrical stimulation of the PAG leads to analgesia that is reversed by naloxone [34]. Neurons from the PAG project to the nucleus raphe magnus of the medulla and from there serotonergic fibers descend to the stalked cells of the dorsal horn where release of enkephalin inhibits SG (spiral ganglion) cells [32,35], thereby blocking the centripetal transmission of pain from C-fibers. Pomeranz proposed that acupuncture stimulation activates A δ -and C afferent fibers in muscle, causing signals to be transmitted to the spinal cord and results in release of dynorphin and enkephalins. Pomeranz's theory was confirmed by a large series of experiments in his research laboratory and other investigators [36-41]. The regulation of enkephalin peptide synthesis was mainly at the mRNA level [42] and EA was shown to enhance preproenkephalin mRNA expression in rostral ventrolateral medulla of rats [43]. It suggests that there would be more peptides to produce following EA [44]. In our study, enkephalin did not increase in EA-group compared with sham EA-group. The reasons maybe include the flowing items:(1) the blood samples detected enkephalin could not consistent with the quantity of EA-group (n=40) and sham EA-group(n=40) repectively; this maybe the limitation of our study; (2) the exact mechanisms of anesthetics remain unclear to date; maybe the change of enkephalin in EA-group is the result of the combination EA and anesthetics. Our results indicate that dynamic change of enkephalin during operations in EA is different from control. The tendency of increased enkephalin in EA suggests that enkephalin might play a role for general anesthesia in pain reduction.

CONCLUSION

EA anesthesia could not replace anesthetics completely. However, EA has become the most popular complementary and alternative medicine modality and is highly praised by physicians and anesthetists. Due to the limited number of blood samples available to us, future research maybe needed to assess enkephalin and mechanisms of EA treatment.

ACKNOWLEDGEMENTS

This work was supported by funding from the 973 traditional Chinese medicine theory of special (2007 CB 512503) and Dr. Nathaniel Mills, Dept. of Biology, Texas Woman's University.

REFERENCES

1. Ohgaki, H., & Kleihues, P. Epidemiology and etiology of gliomas. Acta Neuropathologica Vol.109, pp. 93–108, 2005.
2. Wang, X.Q., Yu, J.L., Du, Z.Y., Xu, R., Jiang, C.C., & Gao, X. Electroacupoint Stimulation for Postoperative Nausea and Vomiting in Patients Undergoing Supratentorial Craniotomy. Journal of Neurosurgical Anesthesiology Vol. 22, pp. 128-131, 2010.
3. White, P.F. Mishaps with patient-controlled analgesia. Anesthesiology Vol. 65, pp. 81-83,

- 1987.
4. Parker, R.K, Holtmann B., & White P.F. Patient-controlled analgesia: does a concurrent opioid infusion improve pain management after surgery? The Journal of the American Medical Association Vol. 266, pp. 1947-1952, 1991.
 5. Wang, S.M., Kain, Z.N., & White, P. Acupuncture Analgesia: I. The Scientific Basis. Anesthesia & Analgesia Vol. 106, pp. 602- 610, 2008.
 6. Chen, L., Tang, J., White, P.F., Sloninsky, A., Wender, R.H., & Naruse, R. The Effect of Location of Transcutaneous Electrical Nerve Stimulation on Postoperative Opioid Analgesic Requirement: Acupoint Versus Nonacupoint Stimulation. Anesthesia & Analgesia Vol. 87, pp. 1129-1134, 1998.
 7. Wang, B., Tang, J., White, P.F., Naruse, R., Sloninsky, A., Kariger, R., Gold, J., & Wender, R.H. Effect of the Intensity of Transcutaneous Acupoint Electrical Stimulation on the Postoperative Analgesic Requirement. Anesthesia & Analgesia Vol. 85, pp. 406- 413, 1997.
 8. Ahsin, S., Saleem, S., Bhatti, A.M., Iles, R.K., & Aslam, M. Clinical and endocrinological changes after electro-acupuncture treatment in patients with osteoarthritis of the knee. Pain Vol. 147, pp. 60- 66, 2009.
 9. Han, J.S., Chen, X.H., Sun, S.L., Xu, X.J., Yuan, Y., Yan, S.C., Hao, J.X., & Terenius, L. Effect of low-and high-frequency TENS on Met-human lumbar CSF. Pain Vol. 47, pp. 295-298, 1991.
 10. Omura, Y., Chen, Y., Lermant, O., Jones, M., Duvvi, H. & Shimotsuura, Y. Effects of transcutaneous electrical stimulation (1 pulse/sec) through custom-made disposable surface electrodes covering Omura's ST36 area of both legs on normal cell telomeres, oncogen C-fosAb2, integrin alpha5beta1, chlamydia trachomatis, etc. in breast cancer & alzheimer patients. Acupuncture & Electro-Electrotherapeutics Research, International Journal of Integrated Medicine Vol 35, pp. 147-85, 2010.
 11. Omura, Y. Electro-acupuncture: its electro-physiological basis and criteria for effectiveness and safety - part 1. Acupuncture & Electro-Electrotherapeutics Research, International Journal of Integrated Medicine Vol 1, pp. 157-181, 1975.
 12. Omura, Y. Patho-physiology of acupuncture treatment: effects of acupuncture on cardiovascular and nervous systems. Acupuncture & Electro-Electrotherapeutics Research, International Journal of Integrated Medicine Vol 1, pp. 51-140, 1975.
 13. Omura, Y. Basic electrical parameters for safe and effective electro-therapeutics [electro-acupuncture, TES, TENMS (or TEMS), TENS and electro-magnetic field stimulation with or without drug field] for pain, neuromuscular skeletal problems, and circulatory disturbances. Acupuncture & Electro-Electrotherapeutics Research, International Journal of Integrated Medicine Vol 12, pp. 201-25, 1987.
 14. Francis, R.P., & Johnson, M.I. The characteristics of acupuncture-like transcutaneous electrical nerve stimulation (acupuncture-like TENS): a literature review. Acupuncture & Electro-Electrotherapeutics Research, International Journal of Integrated Medicine Vol. 36, pp. 231-58, 2011.
 15. Fei, H., Xie, G.X., & Han, J.S. Low and high frequency electroacupuncture stimulation release met-enkephalin and dynorphin A in rat spinalcord. Chinese Science Bulletin Vol.32,pp.1496-1501, 1987.
 16. Simantov, R., Kuhar, M.J., Uhl, G.R., & Snyder, S.H. Opioid peptide enkephalin: Immunohistochemical mapping in ratcentral nervous system. Proceedings of the National Academy of Sciences Vol. 74, No. 5, pp. 2167-2171, 1977.

- 1987.
4. Parker, R.K, Holtmann B., & White P.F. Patient-controlled analgesia: does a concurrent opioid infusion improve pain management after surgery? The Journal of the American Medical Association Vol. 266, pp. 1947-1952, 1991.
 5. Wang, S.M., Kain, Z.N., & White, P. Acupuncture Analgesia: I. The Scientific Basis. Anesthesia & Analgesia Vol. 106, pp. 602- 610, 2008.
 6. Chen, L., Tang, J., White, P.F., Sloninsky, A., Wender, R.H., & Naruse, R. The Effect of Location of Transcutaneous Electrical Nerve Stimulation on Postoperative Opioid Analgesic Requirement: Acupoint Versus Nonacupoint Stimulation. Anesthesia & Analgesia Vol. 87, pp. 1129-1134, 1998.
 7. Wang, B., Tang, J., White, P.F., Naruse, R., Sloninsky, A., Kariger, R., Gold, J., & Wender, R.H. Effect of the Intensity of Transcutaneous Acupoint Electrical Stimulation on the Postoperative Analgesic Requirement. Anesthesia & Analgesia Vol. 85, pp. 406- 413, 1997.
 8. Ahsin, S., Saleem, S., Bhatti, A.M., Iles, R.K., & Aslam, M. Clinical and endocrinological changes after electro-acupuncture treatment in patients with osteoarthritis of the knee. Pain Vol. 147, pp. 60- 66, 2009.
 9. Han, J.S., Chen, X.H., Sun, S.L., Xu, X.J., Yuan, Y., Yan, S.C., Hao, J.X., & Terenius, L. Effect of low-and high-frequency TENS on Met-human lumbar CSF. Pain Vol. 47, pp. 295-298, 1991.
 10. Omura, Y., Chen, Y., Lermant, O., Jones, M., Duvvi, H. & Shimotsuura, Y. Effects of transcutaneous electrical stimulation (1 pulse/sec) through custom-made disposable surface electrodes covering Omura's ST36 area of both legs on normal cell telomeres, oncogen C-fosAb2, integrin alpha5beta1, chlamydia trachomatis, etc. in breast cancer & alzheimer patients. Acupuncture & Electro-Electrotherapeutics Research, International Journal of Integrated Medicine Vol 35, pp. 147-85, 2010.
 11. Omura, Y. Electro-acupuncture: its electro-physiological basis and criteria for effectiveness and safety - part 1. Acupuncture & Electro-Electrotherapeutics Research, International Journal of Integrated Medicine Vol 1, pp. 157-181, 1975.
 12. Omura, Y. Patho-physiology of acupuncture treatment: effects of acupuncture on cardiovascular and nervous systems. Acupuncture & Electro-Electrotherapeutics Research, International Journal of Integrated Medicine Vol 1, pp. 51-140, 1975.
 13. Omura, Y. Basic electrical parameters for safe and effective electro-therapeutics [electro-acupuncture, TES, TENMS (or TEMS), TENS and electro-magnetic field stimulation with or without drug field] for pain, neuromuscular skeletal problems, and circulatory disturbances. Acupuncture & Electro-Electrotherapeutics Research, International Journal of Integrated Medicine Vol 12, pp. 201-25, 1987.
 14. Francis, R.P., & Johnson, M.I. The characteristics of acupuncture-like transcutaneous electrical nerve stimulation (acupuncture-like TENS): a literature review. Acupuncture & Electro-Electrotherapeutics Research, International Journal of Integrated Medicine Vol. 36, pp. 231-58, 2011.
 15. Fei, H., Xie, G.X., & Han, J.S. Low and high frequency electroacupuncture stimulation release met-enkephalin and dynorphin A in rat spinalcord. Chinese Science Bulletin Vol.32,pp.1496-1501, 1987.
 16. Simantov, R., Kuhar, M.J., Uhl, G.R., & Snyder, S.H. Opioid peptide enkephalin: Immunohistochemical mapping in ratcentral nervous system. Proceedings of the National Academy of Sciences Vol. 74, No. 5, pp. 2167-2171, 1977.

17. Pert, A., & Yaksh, T. Sites of morphine induced analgesia in the primate brain: relation to pain pathways. Brain Research Vol. 80, pp.135-140, 1974.
18. Jacquet, Y.F., & Lajtha, A. Morphine action at central nervous system sites in rat: analgesia or hyperalgesia depending on site and dose. Science Vol.182, pp. 490-492, 1973.
19. Akil, H., Mayer, D. J., & Liebeskind, J. Antagonism of stimulation-produced analgesia by naloxone, a narcotic antagonist. Science Vol.191, pp.961-962, 1976.
20. Loh, H.H., Tseng, L.F., Weit, E., & Lit, C.H. Endorphin is a potent analgesic agent. Proceedings of the National Academy of Sciences Vol. 73, pp. 2895-2898, 1976.
21. Zhao, Z.Q. Neural mechanism underlying acupuncture analgesia. Progress in Neurobiology Vol. 85, pp.355-375, 2008.
22. Zhang, W.T., Jin, Z., Huang, J., Zhang, L., Zeng, Y.W., Luo, F., Chen, A.C. & Han, J.S. Modulation of cold pain in human rain by electric acupoint stimulation: evidence from fMRI. Neuroreport Vol.14, pp. 1591- 1596, 2003.
23. Hamza, M.A., White, P.F., Ahmend, H.E., & Ghoname, E.A. Effect of the frequency of transcutaneous electrical nerve stimulation on the postoperative opioid analgesic requirement and recovery profile. Anesthesiology Vol. 91, pp.1232- 1238, 1999.
24. Ghoname, E.S., Craing, W.F., White, P.F., Ahmed, H.E., Hamza, M.A., Gajraj, N.M., Vakharia, A.S., & Noe, C.E. The effect of stimulus frequency on the analgesic response to percutaneous electrical nerve stimulation in patients with chronic low back pain. Anesthesia & Analgesia Vol. 88, pp. 841- 846, 1999.
25. Lin, J.G., Lo, M.W., Wen, Y.R., Hsieh, C.L., Tsai, S.K., & Sun, W.Z. The effect of high and low frequency electroacupuncture in pain after lower abdominal surgery. Pain Vol. 99, pp. 509- 514, 2002.
26. Wang, Y., Zhang, Y., Wang, W., Cao, Y., & Han, J.S. Effects of synchronous or asynchronous electroacupuncture stimulation with low versus high frequency on spinal opioid release and tail flick nociception. Experimental Neurology Vol. 192, pp. 156-162, 2005.
27. Trinh, K.V., Graham, N., Gross, A.R., Goldsmith, C.H., Wang, E., Cameron, I.D., Kay, T., & Cervical Overview Group. Acupuncture for neck disorders. Cochrane Database of Systematic Reviews Vol. 3, CD004870, 2006.
28. Green, S., Buchbinder, R., & Hetrick, S. Acupuncture for shoulder pain. Cochrane Database of Systematic Reviews Vol.2, CD005319, 2005.
29. Trinh, K.V., Phillips, S.D., Ho, E., & Damsma, K. Acupuncture for the alleviation of lateral epicondyle pain: a systematic review. Rheumatology (Oxford) Vol. 43, pp. 1085-1090, 2004.
31. Pyne, D., & Shenker, N.G. Demystifying acupuncture. Rheumatology (Oxford) Vol. 47, pp.1132- 1136, 2008.
32. Gobel, S., Falls, W.M., Bennett, G.J. Abdelmoumene, M., Hayashi, H., & Humphrey, E. An E.M. Analysis of the synaptic connections of horseradish peroxidase filled stalked cells and islet cells in the substantia gelatinosa of the adult cat spinal cord. Journal of Comparative Neurology Vol.194, pp.781-807, 1980.
33. Sugiura, Y., Lee, C.L., & Perl, E.R. Central projection of identified, unmyelinated (C) afferent fibers innervating mammalian skin. Science Vol. 234, pp. 358-361, 1986.
34. Hosobuchi, Y., Adams, J.E., & Linchitz, R. Pain relief by electrical stimulation of the centra grey matter in humans and its reversal by naloxone. Science Vol. 197, pp. 18-186, 1977.
35. Guowei, L., Rongzhao, L., Jingqiang, X., Yuanshen, W., & Guorui, H. Role of peripheral afferent nerve fiber in acupuncture analgesia elicited by needling point zusanli. Acta

17. Pert, A., & Yaksh, T. Sites of morphine induced analgesia in the primate brain: relation to pain pathways. Brain Research Vol. 80, pp.135-140, 1974.
18. Jacquet, Y.F., & Lajtha, A. Morphine action at central nervous system sites in rat: analgesia or hyperalgesia depending on site and dose. Science Vol.182, pp. 490-492, 1973.
19. Akil, H., Mayer, D. J., & Liebeskind, J. Antagonism of stimulation-produced analgesia by naloxone, a narcotic antagonist. Science Vol.191, pp.961-962, 1976.
20. Loh, H.H., Tseng, L.F., Weit, E., & Lit, C.H. Endorphin is a potent analgesic agent. Proceedings of the National Academy of Sciences Vol. 73, pp. 2895-2898, 1976.
21. Zhao, Z.Q. Neural mechanism underlying acupuncture analgesia. Progress in Neurobiology Vol. 85, pp.355-375, 2008.
22. Zhang, W.T., Jin, Z., Huang, J., Zhang, L., Zeng, Y.W., Luo, F., Chen, A.C. & Han, J.S. Modulation of cold pain in human rain by electric acupoint stimulation: evidence from fMRI. Neuroreport Vol.14, pp. 1591- 1596, 2003.
23. Hamza, M.A., White, P.F., Ahmend, H.E., & Ghoname, E.A. Effect of the frequency of transcutaneous electrical nerve stimulation on the postoperative opioid analgesic requirement and recovery profile. Anesthesiology Vol. 91, pp.1232- 1238, 1999.
24. Ghoname, E.S., Craing, W.F., White, P.F., Ahmed, H.E., Hamza, M.A., Gajraj, N.M., Vakharia, A.S., & Noe, C.E. The effect of stimulus frequency on the analgesic response to percutaneous electrical nerve stimulation in patients with chronic low back pain. Anesthesia & Analgesia Vol. 88, pp. 841- 846, 1999.
25. Lin, J.G., Lo, M.W., Wen, Y.R., Hsieh, C.L., Tsai, S.K., & Sun, W.Z. The effect of high and low frequency electroacupuncture in pain after lower abdominal surgery. Pain Vol. 99, pp. 509- 514, 2002.
26. Wang, Y., Zhang, Y., Wang, W., Cao, Y., & Han, J.S. Effects of synchronous or asynchronous electroacupuncture stimulation with low versus high frequency on spinal opioid release and tail flick nociception. Experimental Neurology Vol. 192, pp. 156-162, 2005.
27. Trinh, K.V., Graham, N., Gross, A.R., Goldsmith, C.H., Wang, E., Cameron, I.D., Kay, T., & Cervical Overview Group. Acupuncture for neck disorders. Cochrane Database of Systematic Reviews Vol. 3, CD004870, 2006.
28. Green, S., Buchbinder, R., & Hetrick, S. Acupuncture for shoulder pain. Cochrane Database of Systematic Reviews Vol.2, CD005319, 2005.
29. Trinh, K.V., Phillips, S.D., Ho, E., & Damsma, K. Acupuncture for the alleviation of lateral epicondyle pain: a systematic review. Rheumatology (Oxford) Vol. 43, pp. 1085-1090, 2004.
31. Pyne, D., & Shenker, N.G. Demystifying acupuncture. Rheumatology (Oxford) Vol. 47, pp.1132- 1136, 2008.
32. Gobel, S., Falls, W.M., Bennett, G.J. Abdelmoumene, M., Hayashi, H., & Humphrey, E. An E.M. Analysis of the synaptic connections of horseradish peroxidase filled stalked cells and islet cells in the substantia gelatinosa of the adult cat spinal cord. Journal of Comparative Neurology Vol.194, pp.781-807, 1980.
33. Sugiura, Y., Lee, C.L., & Perl, E.R. Central projection of identified, unmyelinated (C) afferent fibers innervating mammalian skin. Science Vol. 234, pp. 358-361, 1986.
34. Hosobuchi, Y., Adams, J.E., & Linchitz, R. Pain relief by electrical stimulation of the centra grey matter in humans and its reversal by naloxone. Science Vol. 197, pp. 18-186, 1977.
35. Guowei, L., Rongzhao, L., Jingqiang, X., Yuanshen, W., & Guorui, H. Role of peripheral afferent nerve fiber in acupuncture analgesia elicited by needling point zusanli. Acta

- Pharmacologica Sinica Vol. 22, pp. 680, 1979.
36. Pomeranz, B., & Chiu, D. Naloxone blockade of acupuncture analgesia: endorphin implicated. Life Sciences Vol.19, pp. 1757-1762, 1976.
 37. Lim, T., Loh, T., Kranz, H., & Scott, D. Acupuncture-effect on normal subjects. Medical Journal of Australia Vol.1, pp. 440-442, 1977.
 38. Cheng, R.S., & Pomeranz, B. Monoaminergic mechanism of electroacupuncture analgesia. Brain Research Vol. 215, pp.77-92, 1981.
 39. Cheng, R.S., & Pomeranz, B. Electroacupuncture analgesia is mediate by sterosppecific opiate receptors and is reversed by antagonists of type I receptors. Life Sciences Vol. 26, pp. 631-638, 1980.
 40. Han, J.S., & Terenius, L. Neurochemical basis of acupuncture analgesia. Annual Review of Pharmacology and Toxicology Vol.22, pp. 193-220, 1982.
 41. Horrigan, B. Acupuncture and the Raison D'etre for alterative medicine. Alternative Therapies in Health and Medicine Vol.2, pp. 85- 91, 1996.
 42. Borsook, D., & Hyman, E.S. Proenkephalin gene regulation in the neuroendocrine hypothalamus: model of gene regulation in the CNS. American Journal of Physiology Vol. 269, pp. E393-E408, 1995.
 43. Li, P., Tjen-A-Looi, S., & Longhurst, J.C. Rostral ventrolateral medullary opioid receptor subtypes in the inhibitory effect of electroacupuncture on reflex autonomic response in cats. Autonomic Neuroscience: Basic & Clinical Vol.89, pp. 38- 47, 2001.
 44. Li, M., Tjen-A-Looi, S.C., & Longhurst, J.C. Electroacupuncture enhances preproenkephalin mRNA expression in rostral ventrolateral medulla of rats. Neuroscience Letters Vol. 477, pp. 61- 65, 2010.

- Pharmacologica Sinica Vol. 22, pp. 680, 1979.
36. Pomeranz, B., & Chiu, D. Naloxone blockade of acupuncture analgesia: endorphin implicated. Life Sciences Vol.19, pp. 1757-1762, 1976.
 37. Lim, T., Loh, T., Kranz, H., & Scott, D. Acupuncture-effect on normal subjects. Medical Journal of Australia Vol.1, pp. 440-442, 1977.
 38. Cheng, R.S., & Pomeranz, B. Monoaminergic mechanism of electroacupuncture analgesia. Brain Research Vol. 215, pp.77-92, 1981.
 39. Cheng, R.S., & Pomeranz, B. Electroacupuncture analgesia is mediate by sterosppecific opiate receptors and is reversed by antagonists of type I receptors. Life Sciences Vol. 26, pp. 631-638, 1980.
 40. Han, J.S., & Terenius, L. Neurochemical basis of acupuncture analgesia. Annual Review of Pharmacology and Toxicology Vol.22, pp. 193-220, 1982.
 41. Horrigan, B. Acupuncture and the Raison D'etre for alterative medicine. Alternative Therapies in Health and Medicine Vol.2, pp. 85- 91, 1996.
 42. Borsook, D., & Hyman, E.S. Proenkephalin gene regulation in the neuroendocrine hypothalamus: model of gene regulation in the CNS. American Journal of Physiology Vol. 269, pp. E393-E408, 1995.
 43. Li, P., Tjen-A-Looi, S., & Longhurst, J.C. Rostral ventrolateral medullary opioid receptor subtypes in the inhibitory effect of electroacupuncture on reflex autonomic response in cats. Autonomic Neuroscience: Basic & Clinical Vol.89, pp. 38- 47, 2001.
 44. Li, M., Tjen-A-Looi, S.C., & Longhurst, J.C. Electroacupuncture enhances preproenkephalin mRNA expression in rostral ventrolateral medulla of rats. Neuroscience Letters Vol. 477, pp. 61- 65, 2010.