

Let C_1, \dots, C_K denote sets containing the indices of the observations in each cluster. These sets satisfy two properties:

1. $C_1 \cup C_2 \cup \dots \cup C_K = \{1, \dots, n\}$. In other words, each observation belongs to at least one of the K clusters.
2. $C_k \cap C_{k'} = \emptyset$. In other words, the clusters are non-overlapping: no observation belongs to more than one cluster.

For instance, if the i th observation is in the k th cluster, then $i \in C_k$.

- The idea behind K -means clustering is that a *good* clustering is one for which the *within-cluster variation* is as small as possible.
- The within-cluster variation for cluster C_k is a measure $\text{WCV}(C_k)$ of the amount by which the observations within a cluster differ from each other.
- Hence we want to solve the problem

$$\underset{C_1, \dots, C_K}{\text{minimize}} \left\{ \sum_{k=1}^K \text{WCV}(C_k) \right\}$$

- This formula says that we want to partition the observations into K clusters such that the total within-cluster variation, summed over all K clusters, is as small as possible.

- Typically we use Euclidean distance

$$\text{WCV}(C_k) = \frac{1}{|C_k|} \sum_{i,i' \in C_k} \sum_{j=1}^p (x_{ij} - x_{i'j})^2$$

where $|C_k|$ denotes the number of observations in the k th cluster.

- Combining the minimization task expression with the previous equation gives the optimization problem that defines K -means clustering:

$$\underset{C_1, \dots, C_K}{\text{minimize}} \left\{ \sum_{k=1}^K \frac{1}{|C_k|} \sum_{i,i' \in C_k} \sum_{j=1}^p (x_{ij} - x_{i'j})^2 \right\}$$