

Statistics 630 – Final Exam
Wednesday, 2 December 2020

Printed Name: _____ **Email:** _____

INSTRUCTIONS FOR THE STUDENT:

1. You have 2 hours to complete the exam (after taking a moment to read these instructions). Please indicate your start time: _____, and your end time: _____
2. There are 11 pages including this cover sheet and the formula sheets.
3. Questions 1–7 are multiple choice and worth 5 points each.
4. Questions 8–10 are multiple selection and worth 5 points each.
5. Questions 11–15 require solutions to be worked out and are 10 points each. Please write out your answers in *the spaces provided*, explaining your steps. You may refer to theorems by name/description rather than by its number in the book.
6. If you *cannot* print out the exam, please write your answers on blank sheet of paper – in order.
7. You may use the *attached formula sheets*. No other resources are allowed. Do not use the textbook, the class notes, homework or formula sheets that were posted online.
8. *Do not use a calculator.* You may leave answers in forms that can easily be put into a calculator such as $\frac{12}{19}$, $\binom{40}{5}$, e^{-3} , $\Phi(1.5)$, etc.
9. Do not discuss or provide any information to anyone concerning any of the questions on this exam until your solutions are returned or I post my solutions.

I attest that I spent no more than 2 hours to complete the exam. I used only the materials allowed above. I did not receive assistance from or provide assistance to anyone either before or while taking this exam.

Student's Signature _____

Questions 1–7 are multiple choice: circle the single correct answer. No partial credit!

1. (5 points) $\text{Var}(X) = 4$, $\text{Var}(Y) = 1$ and $\text{Cov}(X, Y) = -2$. What is $\text{Var}(X - Y)$?
 - (a) 1.
 - (b) 9.
 - (c) 3.
 - (d) 0.
 - (e) 5.

2. (5 points) Sally Scientist obtained a 98% confidence interval for μ from a random sample of $\text{normal}(\mu, 37.0)$ data (σ^2 known). The sample size was $n = 25$ and the interval she computed was $9.54 \pm 2.83 = (6.71, 12.37)$. The confidence level 98% is the probability, before sampling,
 - (a) that \bar{X} will be between 6.71 and 12.37.
 - (b) that μ is between 6.71 and 12.37.
 - (c) that \bar{X} will be within 2.83 of 9.54.
 - (d) that \bar{X} will be within 2.83 of μ .
 - (e) that a Type I error is not made.

3. (5 points) T is a statistic with cdf $F(t) = \theta t + (1 - \theta)t^2$, $0 \leq t \leq 1$, $0 \leq \theta \leq 1$. Suppose $H_0 : \theta = 0$ is rejected when $T \leq \sqrt{\alpha}$, which is a size α test. Then the power function for the test is
 - (a) $\theta T + (1 - \theta)T^2$.
 - (b) $\theta\sqrt{\alpha} + (1 - \theta)\alpha$.
 - (c) $\Phi\left(\frac{\sqrt{\alpha} - .5}{\sqrt{1/12}}\right)$.
 - (d) $\theta\alpha + (1 - \theta)\alpha^2$.
 - (e) $\Phi\left(\frac{\sqrt{\alpha} - \theta}{\sqrt{1/12}}\right)$.

4. (5 points) An estimator $\hat{\theta}$ is asymptotically normal($\theta, \frac{1}{n\theta}$). Consequently, a level γ Wald confidence interval for θ is
 - (a) $\hat{\theta} \pm z_{(1+\gamma)/2} \frac{1}{\sqrt{n\hat{\theta}}}$.
 - (b) $\hat{\theta} \pm \sqrt{\frac{\chi_\gamma^2}{n\theta}}$.
 - (c) $\hat{\theta} \pm z_{(1+\gamma)/2} \frac{\hat{\theta}}{\sqrt{n}}$.
 - (d) $\hat{\theta} \pm z_{(1+\gamma)/2} \frac{1}{\sqrt{n\theta}}$.
 - (e) $\hat{\theta} \pm \sqrt{\frac{z_{(1+\gamma)/2}^2}{n\hat{\theta}}}$.

5. (5 points) A and B are independent events with probabilities α and β , respectively. The conditional probability that both occur, given at least one occurs, (that is, $P(A \cap B \mid A \cup B)$) is
- (a) $\frac{\alpha\beta}{\alpha+\beta}$.
 - (b) $\frac{\alpha+\beta}{\alpha\beta}$.
 - (c) $\frac{\alpha\beta}{(1-\alpha)(1-\beta)}$.
 - (d) $\frac{\alpha+\beta}{2-\alpha-\beta}$.
 - (e) $\frac{\alpha\beta}{\alpha+\beta-\alpha\beta}$.
6. (5 points) In the spirit of diversity, independent random variables $X \sim \text{normal}(1, 2)$, $Y \sim \text{Poisson}(3)$ and $Z \sim \text{gamma}(4, 5)$ get together for a (socially distanced) party. Let $T = X + Y + Z$ be their sum. The variance of T is
- (a) $2 \times 3 \times 5$.
 - (b) $2 + 3 + 5$.
 - (c) $2 + 3 + 20$.
 - (d) $2 + 3 + \frac{4}{5}$.
 - (e) $2 + 3 + \frac{4}{25}$.
7. (5 points) Consider the basic two-sided test for a mean (with variance known) having rejection criterion $\sqrt{n} \left| \frac{\bar{X} - \mu_0}{\sigma} \right| > z_{1-\alpha/2}$. The power of the test increases
- (a) when α decreases or n increases.
 - (b) only when both α decreases and n increases.
 - (c) when α increases or n increases.
 - (d) only when n increases.
 - (e) only when α decreases.

Questions 8–10 can have 0, 1 or more than 1 answer: circle all that are correct.

8. (5 points) X_1, \dots, X_n are iid normal(2,1) random variables. Which of the following hold?
- (a) $(X_1 - 2)^2 \sim \text{chi-square}(1)$.
 - (b) $\frac{(X_1 - 2)^2}{((X_1 - 2)^2 + \dots + (X_n - 2)^2)/n} \sim F(1, n)$.
 - (c) $\frac{(X_1 - 2)^2}{(X_2 - 2)^2 + \dots + (X_n - 2)^2} \sim F(1, n - 1)$.
 - (d) $\sum_{i=1}^n (X_i - \bar{X})^2 \sim \text{chi-square}(n - 1)$.
 - (e) $\sum_{i=1}^n X_i^2 \sim \text{chi-square}(n)$.

9. (5 points) Q_1, \dots, Q_n is a random sample from a distribution with mean $\sqrt{\beta}$ and variance 2β . Which of the following are correct?
- (a) \bar{Q} is unbiased for $\sqrt{\beta}$.
 - (b) $\frac{1}{n} \sum_{i=1}^n Q_i^2$ is unbiased for 2β .
 - (c) \bar{Q}^2 is unbiased for β .
 - (d) $\frac{\bar{Q}^2}{1+2/n}$ is unbiased for β .
 - (e) $\frac{1}{2(n-1)} \sum_{i=1}^n (Q_i - \bar{Q})^2$ is unbiased for β .
10. (5 points) A Poisson(λ) random sample N_1, \dots, N_n has MLE $\hat{\lambda} = \bar{N}$, score $S(\lambda) = n(\frac{\bar{N}}{\lambda} - 1)$ and Fisher information $I(\lambda) = \frac{n}{\lambda}$. Which of the following hold, as $n \rightarrow \infty$?
- (a) $\frac{\bar{N} - \lambda}{\sqrt{n/\lambda}} \xrightarrow{D} \text{normal}(0, 1)$.
 - (b) $\frac{(\bar{N} - \lambda)^2}{\lambda/n} \xrightarrow{D} \text{chi-square}(1)$.
 - (c) $\frac{S^2(\lambda)}{I(\lambda)} \xrightarrow{D} \text{chi-square}(n)$.
 - (d) $\bar{N} \xrightarrow{D} \text{normal}(\lambda, \lambda/n)$.
 - (e) $\frac{S(\lambda)}{\sqrt{I(\lambda)}} \xrightarrow{D} \text{normal}(0, 1)$.

Provide solutions to Questions 11–15, to the point of a calculable expression.

11. X has pdf $f_X(x) = \frac{e^{-x}}{c(1+x)}$, $x > 0$ (where $c = \int_0^\infty \frac{e^{-x}}{1+x} dx$), and the conditional pdf for Y , given $X = x$, is $f_{Y|X}(y|x) = (1+x)e^{-y-xy}$, $y > 0$. Find the conditional pdf for X , given $Y = y$.

12. (10 points) R_1, R_2, \dots are iid random variables from the distribution with pdf $f(r) = 2\theta r e^{-\theta r^2}$, $r > 0, \theta > 0$. Determine the MLE for θ and its asymptotic variance.

13. (10 points) Assume as in the previous problem but suppose θ has prior distribution gamma(2,1) with pdf $\theta e^{-\theta}$. Identify the posterior distribution for θ (by name and parameter values).

14. (10 points) Consider an exponential(λ) random sample X_1, \dots, X_n . Determine a moment estimator for $e^{-\lambda} = e^{-1/\mu}$, based on \bar{X} , and use the delta method to derive an asymptotic variance for it.

15. (10 points) Let Y_1, Y_2, \dots, Y_n be a random sample from the geometric(θ) distribution with pmf $\theta(1 - \theta)^y$, $y = 0, 1, 2, \dots$, $0 < \theta < 1$. Determine the size α (asymptotic) generalized likelihood ratio test for $H_0 : \theta = \frac{1}{2}$ versus $H_a : \theta \neq \frac{1}{2}$.

Formulas for Final Exam

Bayes' rule $P(B_j | A) = \frac{P(A|B_j)P(B_j)}{\sum_{k=1}^n P(A|B_k)P(B_k)}$ if B_1, \dots, B_n are disjoint and $\bigcup_{k=1}^n B_k = S$.

quantile function $Q_X(p)$ satisfies $F_X(x) \leq p \leq F(Q_X(p))$ for all $x < Q_X(p)$. $F(Q_X(p)) = p$ if X is a continuous rv.

distribution of a function of X $F_Y(y) = P(h(X) \leq y)$ for $Y = h(X)$.

If X is a discrete rv or $h(x)$ takes only countably many values then Y has pmf $p_Y(y) = P(h(X) = y)$.

If X is a continuous rv and $h(x)$ is a continuous function then Y has pdf $f_Y(y) = \frac{dx}{dy} P(h(X) \leq y)$.

binomial theorem $\sum_{k=0}^n \binom{n}{k} a^k b^{n-k} = (a+b)^n$.

geometric sum $\sum_{k=n}^{\infty} a^k = \frac{a^n}{1-a}$ if $-1 < a < 1$.

exponential expansion $\sum_{k=0}^{\infty} \frac{a^k}{k!} = e^a$.

gamma integral $\int_0^{\infty} x^a e^{-x} dx = \Gamma(a+1) = a!$ for $a > -1$.

Bernoulli pmf $p(x) = (1-\theta)^{1-x} \theta^x I_{\{0,1\}}(x)$ for $0 < \theta < 1$, same as binomial(1, θ).

beta(a, b) pdf $f(x) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} x^{a-1} (1-x)^{b-1} I_{(0,1)}(x)$ for $a > 0, b > 0$; $E(X) = \frac{a}{a+b}$ $\text{Var}(X) = \frac{ab}{(a+b)^2(a+b+1)}$.

binomial(n, θ) pmf $p(x) = \binom{n}{x} \theta^x (1-\theta)^{n-x} I_{\{0,1,\dots,n\}}(x)$ for $0 < \theta < 1$. $E(X) = n\theta$, $\text{Var}(X) = n\theta(1-\theta)$, $m(s) = (1-\theta + \theta e^s)^n$.

chi-square(n) same as gamma($\frac{n}{2}, \frac{1}{2}$), the distribution of $X = Z_1^2 + \dots + Z_n^2$ for iid standard normal Z_1, \dots, Z_n . $E(X) = n$, $\text{Var}(X) = 2n$.

In particular, if $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} \text{normal}(\mu, \sigma^2)$ then $\frac{(n-1)S^2}{\sigma^2} \sim \text{chi-square}(n-1)$.

discrete uniform(N) pmf $p(x) = \frac{1}{N} I_{\{1,2,\dots,N\}}(x)$. $E(X) = \frac{N+1}{2}$, $\text{Var}(X) = \frac{N^2-1}{12}$.

exponential(λ) pdf $f(x) = \lambda e^{-\lambda x} I_{(0,\infty)}(x)$ for $\lambda > 0$, same as gamma(1, λ). $E(X) = \frac{1}{\lambda}$, $\text{Var}(X) = \frac{1}{\lambda^2}$.

F(m, n) the distribution of $W = \frac{X/m}{Y/n}$ where $X \sim \text{chi-square}(m)$, $Y \sim \text{chi-square}(n)$, independent. $E(W) = \frac{n}{n-2}$ if $n > 2$.

gamma(α, λ) pdf $f(x) = \frac{\lambda^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\lambda x} I_{(0,\infty)}(x)$ for $\lambda > 0, \alpha > 0$; $E(X) = \frac{\alpha}{\lambda}$, $\text{Var}(X) = \frac{\alpha}{\lambda^2}$, $m(s) = \left(\frac{\lambda}{\lambda-s}\right)^\alpha$ if $s < \lambda$.

geometric(θ) pmf $p(x) = \theta(1-\theta)^x I_{\{0,1,2,\dots\}}(x)$ for $0 < \theta < 1$, same as negative binomial(1, θ). $E(X) = \frac{1-\theta}{\theta}$, $\text{Var}(X) = \frac{1-\theta}{\theta^2}$.

hypergeometric(N, M, n) **pmf** $p(x) = \frac{\binom{M}{x}\binom{N-M}{n-x}}{\binom{N}{n}} I_{\{0,1,\dots,n\}}(x)$ for $M < N$. $E(X) = np$ where $p = \frac{M}{N}$, $\text{Var}(X) = \frac{N-n}{N-1} np(1-p)$.

negative binomial(r, θ) **pmf** $p(x) = \binom{r+x-1}{r-1} \theta^r (1-\theta)^x I_{\{0,1,2,\dots\}}(x)$ for $0 < \theta < 1$. $E(X) = \frac{r(1-\theta)}{\theta}$, $\text{Var}(X) = \frac{r(1-\theta)}{\theta^2}$, $m(s) = \left(\frac{\theta}{1-(1-\theta)e^s}\right)^r$ if $s < -\log(1-\theta)$.

normal(μ, σ^2) **pdf** $f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2/(2\sigma^2)} I_{(-\infty,\infty)}(x)$ for $\sigma^2 > 0$; $E(X) = \mu$, $\text{Var}(X) = \sigma^2$, $m(s) = e^{\mu s + \sigma^2 s^2/2}$.

Poisson(λ) **pmf** $p(x) = \frac{\lambda^x}{x!} e^{-\lambda} I_{\{0,1,2,\dots\}}(x)$ for $\lambda > 0$. $E(X) = \lambda$, $\text{Var}(X) = \lambda$, $m(s) = e^{\lambda(e^s-1)}$.

t(n) the distribution of $T = \frac{Z}{\sqrt{Y/n}}$ where $Z \sim \text{normal}(0, 1)$, $Y \sim \text{chi-square}(n)$, independent.

$E(T) = 0$, $\text{Var}(T) = \frac{n}{n-2}$ if $n > 2$. In particular, if $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} \text{normal}(\mu, \sigma^2)$ then $\frac{\bar{X}-\mu}{S/\sqrt{n}} \sim t(n-1)$.

uniform(a, b) **pdf** $f(x) = \frac{1}{b-a} I_{(a,b)}(x)$ for $a < b$. $E(X) = \frac{a+b}{2}$, $\text{Var}(X) = \frac{(b-a)^2}{12}$.

Weibull(α, β) **pdf** $f(x) = \frac{\alpha}{\beta} (x/\beta)^{\alpha-1} e^{-(x/\beta)^\alpha} I_{(0,\infty)}(x)$ for $\alpha > 0$, $\beta > 0$. $E(X^k) = \beta^k \Gamma(1 + \frac{k}{\alpha})$.

marginal pmf/pdf $p_X(x) = \sum_y p(x, y)$; $f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy$.

conditional pmf/pdf $p_{X|Y}(x|y) = \frac{p(x,y)}{p_Y(y)}$; $f_{X|Y}(x|y) = \frac{f(x,y)}{f_Y(y)}$.

independent random variables $p(x, y) = p_X(x)p_Y(y)$ if (X, Y) is discrete;
 $f(x, y) = f_X(x)f_Y(y)$ if (X, Y) is continuous.

discrete convolution $p_{X+Y}(z) = \sum_x p_X(x)p_Y(z-x)$ for independent X, Y .

continuous convolution $f_{X+Y}(z) = \int_{-\infty}^{\infty} f_X(x)f_Y(z-x) dx$ for independent X, Y .

covariance and correlation $\text{Cov}(X, Y) = E((X-\mu_X)(Y-\mu_Y)) = E(XY) - \mu_X\mu_Y$; $\text{Corr}(X, Y) = \frac{\text{Cov}(X, Y)}{\sigma_X\sigma_Y}$. For independent X and Y , $\text{Cov}(X, Y) = \text{Corr}(X, Y) = 0$.

expectation of a sum $E(a_1X_1 + \dots + a_nX_n) = a_1E(X_1) + \dots + a_nE(X_n)$.

expectation of a product If X_1, \dots, X_n are independent, $E\left(\prod_{i=1}^n h_i(X_i)\right) = \prod_{i=1}^n E(h_i(X_i))$.

variance of a sum $\text{Var}(aX + bY) = a^2 \text{Var}(X) + 2ab \text{Cov}(X, Y) + b^2 \text{Var}(Y)$.

variance of a sum of independent rvs $\text{Var}(a_1X_1 + \dots + a_nX_n) = a_1^2 \text{Var}(X_1) + \dots + a_n^2 \text{Var}(X_n)$.

moments k -th moment is $\mu_k = E(X^k)$, $k = 1, 2, \dots$

moment generating function $m_X(s) = E(e^{sX})$; $E(X^k) = \left. \frac{d^k m_X(s)}{ds^k} \right|_{s=0}$.

mgf of a sum If X and Y are independent, $m_{aX+bY}(s) = E(e^{(aX+bY)s}) = m_X(as)m_Y(bs)$.

conditional expectation $E(h(Y)|X=x) = \sum_y h(y)p_{Y|X}(y|x)$ or
 $E(h(Y) | X=x) = \int_{-\infty}^{\infty} h(y)f_{Y|X}(y|x) dy$.

iterated expectation $E(h(Y)) = E(E(h(Y) | X))$, $E(g(X)h(Y)) = E(g(X)E(h(Y) | X))$.

conditional variance $\text{Var}(Y | X) = E(Y^2 | X) - (E(Y | X))^2$.

variance partition formula $\text{Var}(Y) = E(\text{Var}(Y | X)) + \text{Var}(E(Y | X))$.

Markov's inequality $P(|X| \geq x) \leq \frac{E(|X|)}{x}$ for $x > 0$.

Chebyshev's inequality $P(|X - \mu_X| \geq x) \leq \frac{\text{Var}(X)}{x^2}$ for $x > 0$.

sample mean, variance, k -th moment $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$; $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$;
 $m_k = \frac{1}{n} \sum_{i=1}^n X_i^k$.

unbiased sample variance $S^2 = \frac{n}{n-1} \hat{\sigma}^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$.

law of large numbers For iid X_1, X_2, \dots with mean μ , $\bar{X}_n \rightarrow \mu$ as $n \rightarrow \infty$.

central limit theorem For iid X_1, X_2, \dots with mean μ and variance σ^2 ,
 $P\left(\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \leq z\right) = P\left(\frac{X_1 + \dots + X_n - n\mu}{\sqrt{n}\sigma} \leq z\right) \rightarrow \Phi(z)$ (normal(0,1) cdf), as $n \rightarrow \infty$.

bias and standard error $\text{Bias}(\hat{\theta}) = E(\hat{\theta}) - \theta$; $\text{SE}(\hat{\theta}) = \sqrt{\text{Var}(\hat{\theta})}$.

mean squared error $\text{MSE}(\hat{\theta}) = E((\hat{\theta} - \theta)^2) = \text{Var}(\hat{\theta}) + (\text{Bias}(\hat{\theta}))^2$.

consistency $\hat{\theta}$ is consistent if $\text{MSE}(\hat{\theta}) \rightarrow 0$ as $n \rightarrow \infty$.

method of moments for iid sample match the k -th population moment $E(X^k)$ with the k -th sample moment m_k , and solve for the desired parameter estimates.

likelihood function $L(\theta | X_1, \dots, X_n) = \prod_{i=1}^n f_\theta(X_i)$ for iid sample $\underline{X} = (X_1, \dots, X_n)$.

maximum likelihood for iid sample maximize the likelihood function $L(\theta | X_1, \dots, X_n) = \prod_{i=1}^n f_\theta(X_i)$ or the log-likelihood $\ell(\theta | X_1, \dots, X_n) = \log L(\theta | X_1, \dots, X_n) = \sum_{i=1}^n \log f_\theta(X_i)$.

If $\log L(\theta)$ is differentiable and concave at θ , the MLE is a solution to $S(\theta) = \frac{d}{d\theta} \log L(\theta) = 0$.
(For a multidimensional parameter θ this is a system of equations.)

score function $S(\theta | X_1, \dots, X_n) = \frac{d}{d\theta} \ell(\theta)$.

Fisher information $I_n(\theta) = \text{Var}\left(\frac{d}{d\theta} \ell(\theta)\right) = -E\left(\frac{d^2}{d\theta^2} \ell(\theta)\right)$, if ℓ has two derivatives.

For an iid sample, $I_n(\theta) = nI_1(\theta)$ and $I_1(\theta) = \text{Var}\left(\frac{d}{d\theta} \log f_\theta(X_1)\right) = -E\left(\frac{d^2}{d\theta^2} \log f_\theta(X_1)\right)$.

sufficient statistic $T = T(\underline{X})$ is sufficient if $L(\theta | \underline{X}) = h(\underline{X})g(T(\underline{X}), \theta)$ for some functions $h(\underline{x})$ and $g(t, \theta)$.

exponential family The pdf/pmf has the form $f_X(x|\theta) = d(\theta)h(x)e^{c(\theta)t(x)}$ for all x, θ . In this case, with an iid random sample, $T(\underline{X}) = \sum_{i=1}^n t(X_i)$ is a sufficient statistic and $I_n(\theta) = n(c'(\theta))^2 \text{Var}(t(X_1))$.

asymptotics for MLE Assuming Fisher information exists and $\hat{\theta}$ is the MLE, $\hat{\theta} \rightarrow \theta$ in probability and $\sqrt{I_n(\hat{\theta})}(\hat{\theta} - \theta) \rightarrow \text{normal}(0, 1)$ in distribution as $n \rightarrow \infty$.

asymptotic normality $\hat{\theta}$ is asymptotic normal(θ, V_n) if $\frac{\hat{\theta} - \theta}{\sqrt{V_n}} \rightarrow \text{normal}(0, 1)$ in distribution as $n \rightarrow \infty$. V_n may depend on θ or other parameters. If \hat{V}_n is an estimator such that $\hat{V}_n/V_n \rightarrow 1$ then $\frac{\hat{\theta} - \theta}{\sqrt{\hat{V}_n}} \rightarrow \text{normal}(0, 1)$ in distribution.

delta method If $g(\theta)$ is continuously differentiable and estimator $\hat{\theta}$ is asymptotic normal(θ, V_n), then $g(\hat{\theta})$ is asymptotic normal($g(\theta), (g'(\theta))^2 V_n$).

level γ confidence interval $(L(\underline{X}), U(\underline{X}))$ such that $P_\theta(L(\underline{X}) \leq \theta \leq U(\underline{X})) = \gamma$.

confidence interval from pivot If $h(\underline{X}, \theta)$ has a distribution that does not depend on θ , a level γ confidence interval is defined by $\{\theta : h(\underline{X}, \theta) \in A\}$ where $P_\theta(h(\underline{X}, \theta) \in A) = \gamma$.

Wald confidence interval If $\hat{\theta}$ is asymptotic normal(θ, V_n) and \hat{V}_n is an estimator for V_n , an approximate level γ confidence interval for θ has endpoints $\hat{\theta} \pm z_{(1+\gamma)/2} \sqrt{\hat{V}_n}$, where $z_{(1+\gamma)/2}$ is the $(1 + \gamma)/2$ quantile of the normal(0,1) distribution.

score confidence interval For MLE $\hat{\theta}$, an approximate level γ confidence interval defined by $\{\theta : -z_{(1+\gamma)/2} \leq (\frac{d}{d\theta} \ell(\theta)) / \sqrt{I_n(\theta)} \leq z_{(1+\gamma)/2}\}$, where $z_{(1+\gamma)/2}$ is the $(1 + \gamma)/2$ quantile of the normal(0,1) distribution.

A related method is the interval given by $\{\theta : -z_{(1+\gamma)/2} \leq \sqrt{I_n(\theta)}(\hat{\theta} - \theta) \leq z_{(1+\gamma)/2}\}$.

Type I and II errors, level and power A Type I error is rejecting H_0 when it is true. The level of a test is $\alpha = \max_{\theta \in H_0} P_\theta(H_0 \text{ is rejected})$ computed with values of θ such that H_0 true.

A Type II error is not rejecting H_0 when H_a is true. The power of a test is $\beta = \beta(\theta) = P_\theta(H_0 \text{ is rejected})$ computed with parameter value θ (satisfying H_a).

P-value The smallest level α for which H_0 will still be rejected – it is a statistic (function of the data).

Neyman-Pearson likelihood ratio test For simple hypotheses $H_0 : \theta = \theta_0$ vs. $H_a : \theta = \theta_1$,

reject H_0 if $LR = \frac{L(\theta_1)}{L(\theta_0)} \geq c_\alpha$ where $P(LR \geq c_\alpha) = \alpha$ when H_0 is true.

If, for each c , $LR \geq c \iff T \geq k$ (or $R \geq c \iff T \leq k$) for some statistic T and some value k then it suffices to find k_α such that $P(T \geq k_\alpha) = \alpha$ (resp., $P(T \leq k_\alpha) = \alpha$) when H_0 is true.

generalized likelihood ratio test For hypotheses H_0 and H_a about parameter θ and MLE $\hat{\theta}$,

reject H_0 if $LR = \frac{L(\hat{\theta})}{\max_{\theta \in H_0} L(\theta)} \geq c_\alpha$ where $\max_{\theta \in H_0} P(LR \geq c_\alpha) = \alpha$.

If $H_0 : \theta = \theta_0$ and $H_a : \theta \neq \theta_0$ then $LR = \frac{L(\hat{\theta})}{L(\theta_0)}$.

uniformly most powerful test A test is UMP if it has maximum possible power for every parameter value θ that satisfies H_a .

In particular, if the test is the same as the Neyman-Pearson test for each θ satisfying H_a then it is UMP.

Wald test If $\hat{\theta}$ is asymptotic normal(θ, V_n) and \hat{V}_n is an estimator for V_n , reject $H_0 : \theta = \theta_0$ when $\frac{|\hat{\theta} - \theta_0|}{\sqrt{\hat{V}_n}} \geq z_{1-\alpha/2}$, where $z_{1-\alpha/2}$ is the $(1 - \alpha/2)$ quantile of the normal(0,1) dist. Equivalently,

reject H_0 if $\frac{(\hat{\theta} - \theta_0)^2}{\hat{V}_n} \geq \chi_{1,1-\alpha}^2$.

Important case: $\hat{\theta}$ is the MLE and $\hat{V}_n = 1/I_n(\hat{\theta})$.

(asymptotic) score test For MLE $\hat{\theta}$, reject $H_0 : \theta = \theta_0$ when $\frac{|\frac{d}{d\theta} \ell(\theta_0)|}{\sqrt{I_n(\theta_0)}} \geq z_{1-\alpha/2}$, where $z_{1-\alpha/2}$ is

the $(1 - \alpha/2)$ quantile of the normal(0,1) dist. Equivalently, reject H_0 if $\frac{(\frac{d}{d\theta} \ell(\theta_0))^2}{I_n(\theta_0)} \geq \chi_{1,1-\alpha}^2$.

A related test is to reject $H_0 : \theta = \theta_0$ when $\sqrt{I_n(\theta_0)} |\hat{\theta} - \theta_0| \geq z_{1-\alpha/2}$, and $\hat{\theta}$ is the MLE.

asymptotic likelihood ratio test Using the generalized LR statistic, reject $H_0 : \theta = \theta_0$ when $2 \log(LR) \geq z_{1-\alpha/2}^2 = \chi_{1,1-\alpha}^2$.

test equivalent to interval Define a test from an interval (or an interval from a test) by: reject $H_0 : \theta = \theta_0$ at level $\alpha \iff \theta_0$ is not in the $1 - \alpha$ confidence interval.

prior and posterior distributions If the prior density (or pmf) for θ is $f_{\Theta}(\theta)$ then the posterior density (or pmf) is $f_{\Theta}(\theta|\underline{X}) = c(\underline{X})f_{\underline{X}}(\underline{X}|\theta)f_{\Theta}(\theta)$, with $c(\underline{X})$ chosen so that $f_{\Theta}(\theta|\underline{X})$ is a proper pdf (pmf) in θ .

Bayes estimator Either the mean or the mode of the posterior distribution.

Bayes γ credible interval An interval $(L(\underline{X}), U(\underline{X}))$ such that, under the posterior distribution, $P(L(\underline{X}) \leq \theta \leq U(\underline{X}) | \underline{X}) = \gamma$.

The interval is HPD (highest posterior density) if it equals the set $\{\theta : f_{\Theta}(\theta|\underline{X}) \geq c\}$ for some constant c .

Bayes Hypothesis test Choose H_1 if and only if $\frac{P(H_1 | \underline{X})}{P(H_0 | \underline{X})} > 1$.