

Factoradic Representation of Rational Numbers

James Philip Rowell

From ‘A Course of Pure Mathematics’ by G. H. Hardy. Chapter 1, Miscellaneous Examples.

Miscellaneous example* #2 at the end of chapter 1 in Hardy’s ‘Pure Mathematics’ presents us with a fascinating result. The theorem feels like what the ‘[basis-representation-theorem](#)’ is for integers, but this one is for rational numbers, ... beautiful!

Factorial Representation Theorem[†]

Any positive rational number can be expressed in one and only one way in the form

$$a_1 + \frac{a_2}{1 \cdot 2} + \frac{a_3}{1 \cdot 2 \cdot 3} + \dots + \frac{a_k}{1 \cdot 2 \cdot 3 \cdot \dots \cdot k},$$

where a_1, a_2, \dots, a_k are integers, and

$$0 \leq a_1, \quad 0 \leq a_2 < 2, \quad 0 \leq a_3 < 3, \quad \dots, \quad 0 < a_k < k$$

Observations that lead to the proof.

We know that any rational number[‡], say $\frac{m}{q}$, can be written as an integer part, i , PLUS a fractional part, $\frac{p}{q}$, such that $\frac{m}{q} = i + \frac{p}{q}$, where $0 \leq \frac{p}{q} < 1$ (note that i can be zero).

So if we’re trying to represent any positive rational number $\frac{m}{q}$ in terms of the theorem then the a_1 term wants to play the role of the integer part, i , and the remainder of the expression $\frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!}$ looks to be playing the roll of the rational part, $\frac{p}{q}$, such that,

$$0 \leq \frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!} < 1$$

It seemed to me a good idea to forget about the a_1 term and just focus on the a_2, a_3, \dots, a_k terms. In other words, prove the theorem for rational numbers $\frac{p}{q}$, such that $0 \leq \frac{p}{q} < 1$, then it should be trivial to extend it to ALL rational numbers by tacking the a_1 term back on at the end of the proof. Also, it started to become clear that including zero (that is, not JUST positive rational numbers) was going to simplify the task[§].

*Hardy doesn’t call them ‘Exercises’ or ‘Questions’, but that’s what they are, math exercises for the student, like calculations to perform, theorems to prove etc.

[†]...it’s not named in the text but what I’m calling the theorem!

[‡]Every variable, coefficient or constant (eg. a_1, a_k, m, n, i, p, q) in this paper is going to represent a non-negative integer. We aren’t dealing with ‘real numbers’ here, just rational numbers which we will always discuss in terms of one integer divided by another integer, like $\frac{p}{q}$.

[§]Did you notice how the theorem restricts the last term, a_k , to be strictly greater than zero, unlike all the other terms? We loosen up that restriction allow a_k to be equal to zero so all the terms are treated the same. At the very end of the proof it’s trivial to reintroduce that restriction on the a_k term.

At first glance it wasn't remotely obvious to me how I'd go about finding such an assignment of coefficients a_2, a_3, \dots, a_k for a given rational number $\frac{p}{q}$, such that $0 \leq \frac{p}{q} < 1$, let alone that it would be unique.

After playing around for a good chunk of time, and finally figuring out a way to calculate the a_i terms for a given rational number $\frac{p}{q}$, a few things started to jump out at me. For starters look at these particular rational numbers,

$$\begin{aligned}\frac{1}{2} &= \frac{1}{1 \cdot 2} = \frac{2! - 1}{2!} \\ \frac{5}{6} &= \frac{1}{1 \cdot 2} + \frac{2}{1 \cdot 2 \cdot 3} = \frac{1 \cdot 3}{1 \cdot 2 \cdot 3} + \frac{2}{1 \cdot 2 \cdot 3} = \frac{3 + 2}{6} = \frac{3! - 1}{3!} \\ \frac{23}{24} &= \frac{1}{1 \cdot 2} + \frac{2}{1 \cdot 2 \cdot 3} + \frac{3}{1 \cdot 2 \cdot 3 \cdot 4} = \frac{1 \cdot 3 \cdot 4}{1 \cdot 2 \cdot 3 \cdot 4} + \frac{2 \cdot 4}{1 \cdot 2 \cdot 3 \cdot 4} + \frac{3}{1 \cdot 2 \cdot 3 \cdot 4} = \frac{12 + 8 + 3}{24} = \frac{4! - 1}{4!}\end{aligned}$$

Good clue! It seemed to be the case that if we assign the largest possible values to the coefficients, from a_2 up to say a_k we get this rational number, $\frac{k!-1}{k!}$. This number is as close to 1 as you can get with a denominator of $k!$ without actually hitting 1. (Check it out - what happens if you add $\frac{1}{k!}$ to $\frac{k!-1}{k!}$?) This turned out to be a pretty useful observation, and it became my 'Lemma-1' in the proof below.

Also, if we assign zeros to all the coefficients then naturally we get $\frac{0}{k!}$.

So we restrict ourselves to using only a_2, a_3, \dots, a_k , then can generate the smallest rational number ($\frac{0}{k!}$) and the largest ($\frac{k!-1}{k!}$) such that $0 \leq \frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!} < 1$.

One further observation to help understand the motivation behind this proof is that by using combinatorics we can count how many possible combinations of a_i 's there are. So, we have two choices for the a_2 term (0, 1), combined with three choices for the a_3 term (0, 1, 2), combined with four choices for the a_4 term (0, 1, 2, 3), \dots combined with k choices for the a_k term (0, 1, 2, \dots , $k-1$), that gives us $2 \cdot 3 \cdot 4 \cdot \dots \cdot k = k!$ possibly different sums.

Hmmmm, the following set has $k!$ members, $\{\frac{0}{k!}, \frac{1}{k!}, \frac{2}{k!}, \dots, \frac{k!-1}{k!}\}$. So that, plus one or two other thoughts is what lead me to the proof below. I won't spoil the rest of it, to find out, go ahead and read the rest of the paper!

Lemma-1

$$\frac{1}{2!} + \frac{2}{3!} + \dots + \frac{k-1}{k!} = \frac{k! - 1}{k!}$$

Proof of Lemma-1

This equality is fairly trivial to demonstrate by induction, since $\frac{1}{2!} = \frac{2!-1}{2!}$ and,

$$\begin{aligned} & \frac{1}{2!} + \frac{2}{3!} + \dots + \frac{k-2}{(k-1)!} + \frac{k-1}{k!} \\ &= \frac{(k-1)! - 1}{(k-1)!} + \frac{k-1}{k!} \\ &= \frac{k((k-1)! - 1)}{k(k-1)!} + \frac{k-1}{k!} \\ &= \frac{k! - k + k - 1}{k!} \\ &= \frac{k! - 1}{k!} \end{aligned}$$

... thus establishing lemma-1 for all values of k. QED

Lemma-2

For integers i, k where $2 \leq i < k$,

$$\frac{1}{i!} > \frac{i}{(i+1)!} + \dots + \frac{k-1}{k!}$$

Proof of Lemma-2

$$\begin{aligned} & \frac{i}{(i+1)!} + \dots + \frac{k-1}{k!} \\ &= \left(\frac{1}{2!} + \frac{2}{3!} + \dots + \frac{k-1}{k!} \right) - \left(\frac{1}{2!} + \frac{2}{3!} + \dots + \frac{i-1}{i!} \right) \\ &= \frac{k! - 1}{k!} - \frac{i! - 1}{i!} \quad \text{(by lemma-1)} \\ &= \frac{k!}{k!} - \frac{1}{k!} - \frac{i!}{i!} + \frac{1}{i!} \\ &= \frac{1}{i!} - \frac{1}{k!} \\ &< \frac{1}{i!} \end{aligned}$$

QED

Definitions

$$\mathcal{S}_k = \left\{ \frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!} \mid 0 \leq a_2 < 2, 0 \leq a_3 < 3, \dots, 0 \leq a_k < k \right\},$$

$$\mathcal{F}_k = \left\{ \frac{0}{k!}, \frac{1}{k!}, \frac{2}{k!}, \dots, \frac{k!-1}{k!} \right\}$$

Lemma-3

$$\mathcal{S}_k = \mathcal{F}_k$$

Corollary: For integers p, k , where $k \geq 2$, every rational number $\frac{p}{k!}$, where $0 \leq \frac{p}{k!} < 1$, there is a unique sequence of integers a_2, a_3, \dots, a_k such that,

$$\frac{p}{k!} = \frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!},$$

where $0 \leq a_2 < 2, 0 \leq a_3 < 3, \dots, 0 \leq a_k < k$.

Proof of Lemma-3

It's clear that the set \mathcal{F}_k contains every rational number with denominator $k!$ where p is an integer and $0 \leq \frac{p}{k!} < 1$ and that the size of \mathcal{F}_k is $k!$. To show that the set \mathcal{S}_k is the same as \mathcal{F}_k , it suffices to show that every member of \mathcal{S}_k is also of the form $0 \leq \frac{p}{k!} < 1$, and that the size of \mathcal{S}_k is also $k!$.

The smallest member of the set \mathcal{S}_k is $\frac{0}{k!}$ and occurs when all the coefficients of the sum are zero. Furthermore, the largest member of the set occurs when all the coefficients of the sum are set to their maximum value, which gives us $\frac{k!-1}{k!}$ as shown in lemma-1.

We also note that every member of \mathcal{S}_k can be written as a rational number with $k!$ as the denominator, like so,

$$\frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_{k-1}}{(k-1)!} + \frac{a_k}{k!} = \frac{k \cdot (k-1) \cdot \dots \cdot 3 \cdot a_2}{k!} + \frac{k \cdot (k-1) \cdot \dots \cdot 4 \cdot a_3}{k!} + \dots + \frac{k \cdot a_{k-1}}{k!} + \frac{a_k}{k!}$$

Therefore any member of the set \mathcal{S}_k is of the form $0 \leq \frac{p}{k!} < 1$, where p is some integer in the range $0 \leq p \leq k! - 1$.

Furthermore, each possible assignment of values to the coefficients of $\frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!}$ produce a unique member of the set \mathcal{S}_k .

For if this weren't true and both $\frac{p}{k!} = \frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!}$ and $\frac{p}{k!} = \frac{b_2}{2!} + \frac{b_3}{3!} + \dots + \frac{b_k}{k!}$ for different coefficients a_2, a_3, \dots, a_k and b_2, b_3, \dots, b_k , then we can arrive at a contradiction as follows.

First suppose that $a_i \neq b_i$ is the first such pair of coefficients that differ from each other. In other words, $a_2 = b_2, a_3 = b_3, \dots, a_{i-1} = b_{i-1}$. Also, without loss of generality we can assume that $a_i > b_i$ and state the following equality:

$$\begin{aligned} \frac{a_i}{i!} + \frac{a_{i+1}}{(i+1)!} + \dots + \frac{a_k}{k!} &= \frac{b_i}{i!} + \frac{b_{i+1}}{(i+1)!} + \dots + \frac{b_k}{k!} \\ \Leftrightarrow \frac{a_i - b_i}{i!} &= \frac{b_{i+1} - a_{i+1}}{(i+1)!} + \dots + \frac{b_k - a_k}{k!} \end{aligned}$$

Since $a_i - b_i \geq 1$, then $\frac{a_i - b_i}{i!} \geq \frac{1}{i!}$.

Also, $\frac{i}{(i+1)!} + \dots + \frac{k-1}{k!} \geq \frac{b_{i+1} - a_{i+1}}{(i+1)!} + \dots + \frac{b_k - a_k}{k!}$ regardless of the values of the coefficients on the right side of the inequality*.

However, lemma-2 tells us,

$$\frac{a_i - b_i}{i!} \geq \frac{1}{i!} > \frac{i}{(i+1)!} + \dots + \frac{k-1}{k!} \geq \frac{b_{i+1} - a_{i+1}}{(i+1)!} + \dots + \frac{b_k - a_k}{k!},$$

demonstrating that equality between the two expressions at either end of the inequality is impossible, so our assumption that there can be a second set of coefficients to produce the same rational number $\frac{p}{k!}$ is false. Therefore any assignment of values to the coefficients of $\frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!}$ produces a unique member of the set \mathcal{S}_k .

Now we can count the number of members of the set \mathcal{S}_k , by looking at all the possible combinations of values for the coefficients a_2, a_3, \dots, a_k . There are 2 choices for the coefficient a_2 , multiplied by the 3 choices for a_3 , multiplied by the 4 choices for a_4, \dots , up to multiplying by k values that a_k can assume.

Therefore the total number of combinations of values that can be assigned to all the coefficients of $\frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!}$ is $2 \cdot 3 \cdot 4 \cdots k = k!$, which means the size of the set \mathcal{S}_k is $k!$. Recalling our previous conclusion that all members of the set \mathcal{S}_k are of the form $0 \leq \frac{p}{k!} < 1$ we can assert that $\mathcal{S}_k = \mathcal{F}_k$.

Furthermore we've shown above that there is a unique sequence of integers a_2, a_3, \dots, a_k , (with the appropriate ranges of values for each integer and $\frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!}$) for each value of $\frac{p}{k!} \in \mathcal{F}_k$. QED

Lemma-4

blah blah blah $\frac{p}{q}$ is unique for all rational numbers...

Proof Lemma-4

All rational numbers $0 \leq \frac{p}{q} < 1$ where $2 \leq q \leq k$ * are members of the set \mathcal{S}_k .

*Letting all the b 's be their maximum value, and all the a 's be zero will produce the largest numerators in each term of the sum, any other possibility will result in a smaller term for the sum.

*The list of rational numbers being $\frac{0}{2}, \frac{1}{2}, \frac{0}{3}, \frac{1}{3}, \frac{2}{3}, \frac{0}{4}, \frac{1}{4}, \frac{2}{4}, \frac{3}{4}, \frac{0}{5}, \frac{1}{5}, \frac{2}{5}, \frac{3}{5}, \frac{4}{5}, \dots, \frac{0}{k}, \frac{1}{k}, \frac{2}{k}, \dots, \frac{k-2}{k}, \frac{k-1}{k}$.

Proof of Lemma-4

Since $2 \leq q \leq k$ then

$$\begin{aligned} \frac{p}{q} &= \frac{2 \cdot 3 \cdot \dots \cdot (q-1) \cdot (q+1) \cdot \dots \cdot k}{2 \cdot 3 \cdot \dots \cdot (q-1) \cdot (q+1) \cdot \dots \cdot k} \cdot \frac{p}{q} \\ &= \frac{2 \cdot 3 \cdot \dots \cdot (q-1) \cdot p \cdot (q+1) \cdot \dots \cdot k}{2 \cdot 3 \cdot \dots \cdot (q-1) \cdot q \cdot (q+1) \cdot \dots \cdot k} \\ &= \frac{2 \cdot 3 \cdot \dots \cdot (q-1) \cdot p \cdot (q+1) \cdot \dots \cdot k}{k!} \end{aligned}$$

Therefore $\frac{p}{q}$ is an element of \mathcal{F}_k which is the same as the set \mathcal{S}_k .

QED

if $\frac{p}{q} \in \mathcal{S}_k$ then by the corollary to lemma-3 there is a unique sequence of integers, $0 \leq a_2 < 2, \quad 0 \leq a_3 < 3, \quad \dots, \quad 0 < a_k < k$ such that,

$$\begin{aligned} \frac{p}{q} &= \frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!} \\ \frac{p}{q} &= \frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!} + \frac{0}{(k+1)!} + \dots + \frac{0}{n!} \\ \frac{p}{q} &= \frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!} + \frac{a_{k+1}}{(k+1)!} + \dots + \frac{a_n}{n!} \\ 0 &\leq a_2 < 2, \quad 0 \leq a_3 < 3, \dots, 0 \leq a_k < k, a_{k+1} = 0, \dots, a_n = 0. \end{aligned}$$

Therefore, $\frac{p}{q} \in \mathcal{S}_n$ for all $n \geq k$, also demonstrating that the sum for $\frac{p}{q}$ is the same for all $n \geq k$, establishing it's unique association with $\frac{p}{q}$.

Theorem (restated)

Any positive rational number can be expressed in one and only one way in the form

$$a_1 + \frac{a_2}{1 \cdot 2} + \frac{a_3}{1 \cdot 2 \cdot 3} + \dots + \frac{a_k}{1 \cdot 2 \cdot 3 \cdot \dots \cdot k},$$

where a_1, a_2, \dots, a_k are integers, and

$$0 \leq a_1, \quad 0 \leq a_2 < 2, \quad 0 \leq a_3 < 3, \quad \dots, \quad 0 < a_k < k$$

Proof of Theorem

Thanks to Euclid we know that for all integers $j \geq 0$ and $q > 0$, there exist unique integers i and p such that,

$$j = i \cdot q + p ; \quad 0 \leq p < q$$

$$\Leftrightarrow \quad \frac{j}{q} = i + \frac{p}{q} ; \quad 0 \leq \frac{p}{q} < 1$$

Which tells us that all rational numbers $\frac{j}{q}$ can be written as an integer part, i , plus a fractional part $0 \leq \frac{p}{q} < 1$.

In our theorem, the a_1 coefficient plays the role of the integer part i , and the rest of the expression, $\frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!}$ plays the role of the fractional part $0 \leq \frac{p}{q} < 1$.

Therefore to express any rational number in the form of the theorem, first apply the Euclidean Division Theorem to $\frac{j}{q}$ and let $a_1 = i$. If there is no fractional remainder, then the theorem is trivially true, however if there is a fractional remainder $\frac{p}{q}$, then it is a member of all sets \mathcal{S}_n such that $n \geq q$.

We take for the coefficients a_2, a_3, \dots, a_k in the sum for $\frac{p}{q} \in \mathcal{S}_n$ all those for which $a_k \neq 0$ but $a_{k+1} = a_{k+2} = \dots = a_n = 0$.

By lemma-5 we know that the sum $\frac{p}{q} = \frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!}$ is uniquely associated with $\frac{p}{q}$ then clearly $\frac{j}{q} = a_1 + \frac{a_2}{2!} + \frac{a_3}{3!} + \dots + \frac{a_k}{k!}$ is uniquely associated with all rational numbers $\frac{j}{q}$.

QED

Additional Observations

While it's true that $\frac{p}{q} \in \mathcal{S}_q$, \mathcal{S}_q is not necessarily the smallest such set for which $\frac{p}{q}$ is a member.

For example, the smallest set containing $0 \leq \frac{p}{5} < 1$ is \mathcal{S}_5 however the smallest set containing $0 \leq \frac{p}{6} < 1$ is \mathcal{S}_3 .

Which is easy to see when we list the contents of a couple of sets,

$$\mathcal{S}_4 = \left\{ \frac{0}{24}, \frac{1}{24}, \frac{2}{24}, \frac{3}{24}, \frac{4}{24}, \frac{5}{24}, \frac{6}{24}, \frac{7}{24}, \frac{8}{24}, \frac{9}{24}, \frac{10}{24}, \frac{11}{24}, \frac{12}{24}, \frac{13}{24}, \frac{14}{24}, \frac{15}{24}, \frac{16}{24}, \frac{17}{24}, \frac{18}{24}, \frac{19}{24}, \frac{20}{24}, \frac{21}{24}, \frac{22}{24}, \frac{23}{24} \right\}$$

$$= \left\{ \frac{0}{24}, \frac{1}{24}, \frac{1}{12}, \frac{1}{8}, \frac{1}{6}, \frac{5}{24}, \frac{1}{4}, \frac{7}{24}, \frac{1}{3}, \frac{3}{8}, \frac{5}{12}, \frac{11}{24}, \frac{1}{2}, \frac{13}{24}, \frac{7}{12}, \frac{5}{8}, \frac{2}{3}, \frac{17}{24}, \frac{3}{4}, \frac{19}{24}, \frac{5}{6}, \frac{7}{8}, \frac{11}{12}, \frac{23}{24} \right\}$$

Which clearly doesn't contain $\frac{1}{5}$. We've established that $\frac{1}{5}$ is definitely in \mathcal{S}_5 but it's interesting to see what it looks like:

$$\frac{1}{5} = \frac{0}{2} + \frac{1}{2 \cdot 3} + \frac{0}{2 \cdot 3 \cdot 4} + \frac{4}{2 \cdot 3 \cdot 4 \cdot 5} = \frac{1}{6} + \frac{1}{30} = \frac{5+1}{30} = \frac{6}{30} = \frac{1}{5}$$

Also, $\mathcal{S}_3 = \left\{ \frac{0}{6}, \frac{1}{6}, \frac{2}{6}, \frac{3}{6}, \frac{4}{6}, \frac{5}{6} \right\} = \left\{ \frac{0}{6}, \frac{1}{6}, \frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \frac{5}{6} \right\}$, which demonstrates the claim above that \mathcal{S}_3 contains $0 \leq \frac{p}{6} < 1$.

I believe that for a given $q \geq 2$ then the smallest set for which $0 \leq \frac{p}{q} < 1$ are members is the set \mathcal{S}_k such that k is the smallest value for which q divides $k!$.

However, I'll leave that proof for another day.