

1. What is Wheelan's primary message of Chapter 1 ("What's the Point?") in *Naked Statistics: Stripping the Dread from the Data*?
 - (a) Statistics provide honest answers to complex questions
 - (b) Statistical formulas are the most important part of understanding data
 - (c) Statistics explain why relationships exist between variables
 - (d) Statistics simplify complex information to help us make better decisions
2. According to Wheelan, in *Naked Statistics: Stripping the Dread from the Data*?, what is the most common cause of inaccurate polling results?
 - (a) Incorrect calculations of standard errors
 - (b) Biased sampling methods or poorly worded questions
 - (c) Misuse of statistical software during data analysis
 - (d) Random variation in respondent answers
3. What does the following R code produce?

```
gss21 |>
  tbl_cross(
    row = incgap,
    col = sex,
    percent = "column",
    missing = "no") |>
  add_p()
```

- (a) A summary table of `sex` grouped by `incgap`, with missing values excluded and a t-test.
 - (b) A frequency table of `incgap` responses with missing values imputed and grouped by `sex`
 - (c) A cross-tabulation of `incgap` by `sex`, showing column percentages and a chi-square test of independence
 - (d) A cross-tabulation comparing the mean values of `incgap` between men and women using a t-test
4. Using 2024 GSS data, a researcher estimated how many Americans believe in life after death. Based on the 90% confidence interval from R, which interpretation is correct?

n	mean	sd	lowCI	hiCI
1283	0.83	0.38	0.81	0.85

- (a) If another survey had been conducted in 2024, there's a 90% chance the point estimate would be between .81 and .85.
- (b) About 90% of people responded to the national survey.
- (c) Because the mean is within the confidence interval, there is a .90 probability that the proportion of U.S. adults who believe in some form of life after death is .83.
- (d) The researcher is 90% confident that the proportion of U.S. adults who believe in some form of life after death is between .81 and .85.

5. If a researcher concluded that there was no difference in police use of force against Black and White men when there actually is a difference, then:
- (a) The null hypothesis was incorrectly rejected (false positive).
 - (b) The null hypothesis was incorrectly retained (false negative).
 - (c) The null hypothesis was correctly retained.
 - (d) The null hypothesis was correctly rejected.
6. Using the table, which of the following is the correct formula to find the expected frequency for U.S. liberals who oppose the death penalty in 2024?

	ideology	conservative	liberal	Total
cappun				
favor		512 (66.8%)	255 (33.2%)	767 (100.0%)
oppose		168 (32.7%)	345 (67.3%)	513 (100.0%)
Total		680 (53.1%)	600 (46.9%)	1280 (100.0%)

- (a) $(345 * 600)/1280$
 - (b) $(600 * 767)/1280$
 - (c) $(345 * 513)/1280$
 - (d) $(600 * 513)/1280$
7. In order to compare meaningfully across groups, a frequency should be converted into a/an _____.
- (a) z-score
 - (b) cumulative frequency
 - (c) raw count
 - (d) percentage
8. According to the 68-95-99.7 rule, what can we say about a person whose score is more than 2 standard deviations above the mean?
- (a) Their score is unusual and higher than most.
 - (b) Their score is likely below average.
 - (c) Their score is within the middle 68% of the distribution.
 - (d) Their score is very common.
9. What does the pipe operator (`|>`) do in dplyr?
- (a) It creates a new column in a data frame.
 - (b) It passes the result of one function into the next.
 - (c) It loads the dplyr package into memory.
 - (d) It filters rows based on a condition.

10. What is the purpose of `drop_na()` in `dplyr`?
- (a) It removes rows that contain missing values.
 - (b) It converts NA values to NULL.
 - (c) It drops columns that are entirely NA.
 - (d) It replaces missing values with zeros.
11. In a normally distributed set of test scores, a student scoring below the average will have a _____.
- (a) positive standard error
 - (b) negative z-score
 - (c) positive z-score
 - (d) negative standard error
12. Researchers calculated a 95% confidence interval of the average family size of Canadians to be from 2.7 to 3.1, based on a sample of 500 families with a standard deviation of 1.1. What change would produce a more precise estimate?
- (a) Increasing our confidence interval to 99%
 - (b) Improving our sampling strategy
 - (c) Increasing our sample size to 600
 - (d) Increasing our standard deviation to 1.3
13. Using 2024 GSS data, a researcher estimated the proportion of Americans who believe in life after death. Based on the 95% confidence interval from R, which interpretation is correct?

sex	n	mean	sd	lowCI	hiCI
male	1139	4.2	3.7	4.0	4.5
female	1174	4.0	3.5	3.8	4.2

- (a) The confidence intervals overlap, so we can't know for sure if there is a statistical difference between the average time men and women spend on leisure activities per day.
 - (b) We are 95% certain that women have less time per day than men for leisure activities, likely because of their greater time spent on housework and childcare.
 - (c) The confidence intervals do not overlap so we can conclude that the average time men and women spend on leisure activities per day are statistically significantly different.
 - (d) We cannot determine whether the null hypothesis of zero difference between the parameters will be rejected at the 0.05 level.
14. What does the `::` operator do in R?
- (a) Comments out a line of code
 - (b) Assigns a value to a variable
 - (c) Imports all functions from a package
 - (d) Specifies a function from a particular package

15. Which command installs a new package in R?
- (a) `require()`
 - (b) `library()`
 - (c) `load()`
 - (d) `install.packages()`
16. If the final grades in a statistics class were normally distributed with an average of 81.5% and a standard deviation of 1.5, about what proportion of the students' final grades were between 80% and 83%?
- (a) 50%
 - (b) 68%
 - (c) 95%
 - (d) 62%
17. National survey data shows that 69% of Canadians support mandatory childhood vaccinations. A local researcher believes that support is likely much higher in Toronto compared with the national average. Which of the following equations represents the research hypothesis?
- (a) $\mu = 0.69$
 - (b) $\mu \neq 0.69$
 - (c) $\mu > 0.69$
 - (d) $\mu < 0.69$
18. Using the U.S. General Social Survey data, a researcher tests whether the proportions who favor a law requiring a person to obtain a police permit before they could buy a gun have changed over time. Assuming an α level of .05, which of these statements best summarizes the researcher's conclusion?

t-statistic	df	p-value	95% CI Lower	95% CI Upper
1.82	3114.93	0.069	-0.00	0.06

- (a) There is a statistically significant difference in support for the police permit law between 1984 and 2024, suggesting that public opinion has changed over time.
- (b) There is a statistically significant difference in support for the police permit law between 1984 and 2024, but it was due to sampling error and cannot be interpreted.
- (c) There is not a statistically significant difference in support for the police permit law between 1984 and 2024, indicating that public opinion has remained stable.
- (d) There is a statistically significant increase in support for the police permit law from 1984 to 2024, proving that stricter gun control laws are now favored by most Americans.

19. Researchers found that couples report about 10 arguments per month on average. If a couple received a z-score of -2 for this measure, which of the following best describes the amount of arguments for this couple compared with the average?
- This couple had fewer arguments than about 5% of couples.
 - This couple had fewer arguments than about 98% of couples.
 - This couple had about 8 arguments per month.
 - This couple had fewer arguments than about 2% of couples.
20. Which of these datasets has a standard deviation of 0?
- 1, 4, 7, 10, 15
 - 0, 2, 4, 6, 8
 - 4, 4, 4, 4, 4
 - 2, 2, 5, 7, 7
21. A p-value is
- A measure that tells us the variability in a normal distribution
 - An estimate of the standard error of a sampling distribution
 - An estimate of where our population statistic lies on the true sampling distribution of the means
 - A measure of how unusual or rare our obtained statistic is compared with what is stated in the null hypothesis
22. Based on the table, which description best describes the correlation between personal income (`realinc`) and number of days of poor mental health (`mntlhlth`)?

Correlation Coefficient	t-statistic	df	p-value	95% CI Lower	95% CI Upper
-0.114	-5.04	1933	0.000	-0.16	-0.07

- Results indicate an inverse relationship ($r = -0.11$, $p < .05$), suggesting that the number of days of poor mental health decreases as personal income increases.
- Decreases in personal income was correlated with decreases in the number of days of poor mental health ($t = -5$, $p < .05$).
- There is no evidence of an association between personal income and days of poor mental health ($t = -5$).
- There is a weakly positive association between personal income and days of poor mental health ($r = -0.11$, $p < .05$).

23. Wheelan illustrates what key concept in the chapter “Inference: Why My Statistics Professor Thought I Might Have Cheated?” from the book *Naked Statistics: Stripping the Dread from the Data*?
- (a) That hypothesis testing is not useful in criminal investigations
 - (b) That unusually accurate results can raise suspicion, but statistical inference helps assess whether they are likely due to chance
 - (c) That statistical inference can be used to prove someone’s guilt beyond reasonable doubt
 - (d) That cheating can be detected by comparing mean scores across different groups

24. The level of measurement for the `hapmar` variable is _____

[1] “Happiness of marriage”

	Freq	%	% Cum.
very happy	802	59.36	59.36
pretty happy	504	37.31	96.67
not too happy	45	3.33	100.00
Total	1351	100.00	100.00

- (a) dichotomous
 - (b) nominal
 - (c) interval
 - (d) ordinal
25. What does the following code return?
- ```
gss24 |>
 group_by(sex) |>
 summarize(mean_age = mean(age, na.rm = TRUE))
```
- (a) A filtered data frame with only rows where age is not missing.
  - (b) A list of all ages grouped by sex.
  - (c) A data frame showing the average age for each sex.
  - (d) A new column named `mean_age` added to the original data frame.
26. Which of the following is a measure of the spread of a distribution?
- (a) sample size
  - (b) standard deviation
  - (c) standardized variable
  - (d) skewness
27. Based on the table, which statement best compares the relative frequency of being ‘not too happy’ between men and women?

|               | sex | male          | female        | Total         |
|---------------|-----|---------------|---------------|---------------|
| happy         |     |               |               |               |
| very happy    |     | 293 ( 20.2%)  | 386 ( 21.3%)  | 679 ( 20.8%)  |
| pretty happy  |     | 830 ( 57.1%)  | 1054 ( 58.2%) | 1884 ( 57.7%) |
| not too happy |     | 330 ( 22.7%)  | 370 ( 20.4%)  | 700 ( 21.5%)  |
| Total         |     | 1453 (100.0%) | 1810 (100.0%) | 3263 (100.0%) |

- (a) Women have a slightly higher relative frequency because more women than men reported being 'not too happy'.
  - (b) The frequency of being 'not too happy' is equal for men and women.
  - (c) Men have a slightly higher relative frequency of being 'not too happy' than women, even though fewer men reported it.
  - (d) A greater number of women reported being 'not too happy' than men, so women have a higher relative frequency.
28. A researcher compares support for universal healthcare (Support vs. Oppose) between two racial groups: Black and White respondents. Which statistical test should be used?
- (a) Chi-square test
  - (b) t-test
  - (c) Linear regression
  - (d) Pearson's correlation (r)
29. Has agreement with the statement that 'Black Americans face worse jobs, income, and housing than white Americans due to discrimination' shifted? Using a significance level of  $\alpha = 0.01$ , what conclusion should be drawn?

|                                     | year         |              | Total        |
|-------------------------------------|--------------|--------------|--------------|
|                                     | 2022         | 2024         |              |
| opinion                             |              |              |              |
| no                                  | 509 (45%)    | 522 (51%)    | 1,031 (48%)  |
| yes                                 | 619 (55%)    | 494 (49%)    | 1,113 (52%)  |
| Total                               | 1,128 (100%) | 1,016 (100%) | 2,144 (100%) |
| Pearson's Chi-squared test, p=0.004 |              |              |              |

- (a) The increase in agreement with the statement between 2022 and 2024 is statistically significant.
- (b) The small change is statistically significant, but most people still agree with the statement.
- (c) There is a statistically significant difference, so we fail to reject the null hypothesis.
- (d) The drop in agreement with the statement between 2022 and 2024 is statistically significant.

30. What will the following code do?

```
gss24 <- gss24 |>
 mutate(status = case_when(
 income >= 100000 ~ "High",
 income > 50000 & income < 100000 ~ "Medium",
 TRUE ~ "Low"
))
```

- (a) It replaces the numeric values in the income column with text labels.
- (b) It removes missing values from the income column.
- (c) It creates a new column named `status` with 3 possible values.
- (d) It filters out rows where income is less than 50,000.

31. What is the result of the following code?

```
gss24 |>
 filter(age > 20) |>
 select(age, childs)
```

- (a) rows with `age > 20` are removed
- (b) Only rows with `age > 20` and columns `age` and `childs` are returned.
- (c) The original `gss24` dataset is overwritten.
- (d) A new column is added to `gss24`

32. When examining a frequency distribution, which of the following columns generally allows you to identify the median most quickly?

- (a) percentages
- (b) cumulative percentages
- (c) frequencies
- (d) cumulative frequencies

33. What will happen if you try to use a function from a package that hasn't been loaded?

- (a) R will prompt you to install the package
- (b) R will automatically load the package
- (c) The function will run normally
- (d) You will get an error saying the function is not found

34. A sociologist wants to examine whether there is a relationship between level of education (measured in years) and annual income (in dollars) among adults in the U.S. Which statistical test is most appropriate?

- (a) Logistic regression
- (b) t-test
- (c) Chi-square test
- (d) Pearson's correlation (r)



35. Which function is used to load a package in R?

- (a) `library()`
- (b) `install.packages()`
- (c) `source()`
- (d) `load()`

36. Chi-square values cannot be \_\_\_\_\_.

- (a) outside the range of 0 to 1
- (b) positive
- (c) zero
- (d) negative

37. A respondent reports 5 poor mental health days in the past 30 days. Based on summary stats for the `mntlhlth` variable, which of the following best describes their z-score?

| freq | median | mean | sd  |
|------|--------|------|-----|
| 2279 | 0      | 4.5  | 7.9 |

- (a) The z-score is close to zero because 5 is close to the mean.
- (b) The z-score is large and positive because 5 is much higher than the mean.
- (c) The z-score is negative because 5 is below the sd.
- (d) The z-score is undefined because the median is 0.

38. Which of the following code snippets correctly uses `mutate()` to convert height from centimeters to meters?

- (a) `mutate(height = select(height / 100))`
- (b) `mutate(height <- height / 100)`
- (c) `mutate(height / 100 = height)`
- (d) `mutate(height = height / 100)`

39. A researcher summarized the `sibs` variable from the U.S. General Social Survey, a measure of how many brothers and sisters respondents had. Based on the summary table, the distribution appears to be:

| freq | median | mean | sd  |
|------|--------|------|-----|
| 3291 | 3      | 2.9  | 1.8 |

- (a) symmetrical
- (b) negatively skewed
- (c) nominal
- (d) positively skewed

40. In *Naked Statistics: Stripping the Dread from the Data?*, Wheelan explains that “most sample means will lie reasonably close to the population mean; the standard error is what defines ‘reasonably close.’ ” What does this statement imply?
- (a) The sample mean is always equal to the population mean.
  - (b) The standard error measures how much sample means tend to vary around the population mean.
  - (c) The standard error tells us how far the population mean is from the sample mean.
  - (d) The standard error increases as sample size increases.
41. Which of the following best describes the differences between the Console, R Scripts, and Quarto documents in RStudio?
- (a) R Scripts and Quarto documents both run code line-by-line, but only the Console saves your work.
  - (b) The Console and R Scripts are identical, while Quarto is used only for Markdown formatting.
  - (c) The Console is used for writing long-term code, R Scripts are for temporary testing, and Quarto is only for plotting.
  - (d) The Console runs code interactively, R Scripts store reusable code, and Quarto combines code, text, and output for dynamic reports.
42. Which of the following is a valid use of `group_by()` and `summarize()`?
- (a) `group_by(data, summarize(sex))`
  - (b) `summarize(group_by(data, age), mean(sex))`
  - (c) `data |> group_by(sex) |> summarize(freq = n())`
  - (d) `data |> summarize(mean = group_by(sex))`
43. What does the expression `5 != 3` evaluate to in R?
- (a) TRUE
  - (b) NULL
  - (c) FALSE
  - (d) NA
44. Which description best describes a correlation coefficient of  $r = -0.62$ ?
- (a) Strong negative relationship
  - (b) No relationship exists between the variables
  - (c) Weakly negative relationship
  - (d) A statistically insignificant relationship
45. In *White Logic, White Methods*, what does Gandy suggest about the use of racial statistics?
- (a) They are neutral tools most useful for academic research
  - (b) They can both expose and perpetuate discrimination
  - (c) Their use in legal proceedings increasingly lead to unfair policy outcomes
  - (d) Their use in data journalism helps mitigate online polarization

46. Which of the following statements best reflects the ASA's guidance on interpreting p-values?
- (a) A small p-value always indicates a large and important effect.
  - (b) A large p-value proves that no effect exists.
  - (c) p-values are influenced by sample size and measurement precision, and do not reflect effect size or importance.
  - (d) p-values measure the scientific or economic significance of a result.
47. Why is it considered good practice to use RStudio Projects when working on data analysis?
- (a) They ensure your code runs faster by optimizing memory usage
  - (b) They help organize your files, set a working directory, and keep your environment consistent
  - (c) They prevent syntax errors by enforcing tidyverse formatting rules
  - (d) They automatically load packages when you open RStudio
48. In the 1970s, Americans watched an average of 3.05 TV hours per day, compared with an average of 3.3 hours per day in 2024. Assuming an  $\alpha$  level of .05, what does the researcher conclude about the statistically significant difference?

| t-statistic | df      | p-value | 95% CI Lower | 95% CI Upper |
|-------------|---------|---------|--------------|--------------|
| -2.84       | 3616.79 | 0.005   | -0.43        | -0.08        |

- (a) Although there is a statistically significant difference, the direction of change is unclear.
  - (b) The confidence interval is below zero, indicating Americans watch less TV in 2024 than in the 1970s.
  - (c) The means indicate that Americans watch more TV in 2024 than in the 1970s.
  - (d) The negative t-statistic suggests a decrease, meaning Americans watch less TV in 2024 than in the 1970s.
49. According to Cohen's argument in *Citizen Scholar*, what role should descriptive statistics play in social science research?
- (a) Descriptive statistics are outdated and should be replaced by more advanced inferential methods.
  - (b) Descriptive statistics are a foundational part of scientific inquiry and should be valued as legitimate research.
  - (c) Descriptive statistics are useful predominately as a step before hypothesis testing and theory development.
  - (d) Descriptive work should be minimized in favor of studies that make strong theoretical contributions.
50. Which RStudio pane is used to write and edit R scripts and Quarto documents?
- (a) Files
  - (b) Source
  - (c) Console
  - (d) Environment

## Answer Sheet

1. (a) ☐ (b) ☐ (c) ☐ (d) ☒
2. (a) ☐ (b) ☒ (c) ☐ (d) ☐
3. (a) ☐ (b) ☐ (c) ☒ (d) ☐
4. (a) ☐ (b) ☐ (c) ☐ (d) ☒
5. (a) ☐ (b) ☒ (c) ☐ (d) ☐
6. (a) ☐ (b) ☐ (c) ☐ (d) ☒
7. (a) ☐ (b) ☐ (c) ☐ (d) ☒
8. (a) ☒ (b) ☐ (c) ☐ (d) ☐
9. (a) ☐ (b) ☒ (c) ☐ (d) ☐
10. (a) ☒ (b) ☐ (c) ☐ (d) ☐
11. (a) ☐ (b) ☒ (c) ☐ (d) ☐
12. (a) ☐ (b) ☐ (c) ☒ (d) ☐
13. (a) ☒ (b) ☐ (c) ☐ (d) ☐
14. (a) ☐ (b) ☐ (c) ☐ (d) ☒
15. (a) ☐ (b) ☐ (c) ☐ (d) ☒
16. (a) ☐ (b) ☒ (c) ☐ (d) ☐
17. (a) ☐ (b) ☐ (c) ☒ (d) ☐
18. (a) ☐ (b) ☐ (c) ☒ (d) ☐
19. (a) ☐ (b) ☒ (c) ☐ (d) ☐
20. (a) ☐ (b) ☐ (c) ☒ (d) ☐

21. (a) ☐ (b) ☐ (c) ☐ (d) ☒
22. (a) ☒ (b) ☐ (c) ☐ (d) ☐
23. (a) ☐ (b) ☒ (c) ☐ (d) ☐
24. (a) ☐ (b) ☐ (c) ☐ (d) ☒
25. (a) ☐ (b) ☐ (c) ☒ (d) ☐
26. (a) ☐ (b) ☒ (c) ☐ (d) ☐
27. (a) ☐ (b) ☐ (c) ☒ (d) ☐
28. (a) ☒ (b) ☐ (c) ☐ (d) ☐
29. (a) ☐ (b) ☐ (c) ☐ (d) ☒
30. (a) ☐ (b) ☐ (c) ☒ (d) ☐
31. (a) ☐ (b) ☒ (c) ☐ (d) ☐
32. (a) ☐ (b) ☒ (c) ☐ (d) ☐
33. (a) ☐ (b) ☐ (c) ☐ (d) ☒
34. (a) ☐ (b) ☐ (c) ☐ (d) ☒
35. (a) ☒ (b) ☐ (c) ☐ (d) ☐
36. (a) ☐ (b) ☐ (c) ☐ (d) ☒
37. (a) ☒ (b) ☐ (c) ☐ (d) ☐
38. (a) ☐ (b) ☐ (c) ☐ (d) ☒
39. (a) ☐ (b) ☒ (c) ☐ (d) ☐
40. (a) ☐ (b) ☒ (c) ☐ (d) ☐
41. (a) ☐ (b) ☐ (c) ☐ (d) ☒

42. (a) ☐ (b) ☐ (c) ☒ (d) ☐
43. (a) ☒ (b) ☐ (c) ☐ (d) ☐
44. (a) ☒ (b) ☐ (c) ☐ (d) ☐
45. (a) ☐ (b) ☒ (c) ☐ (d) ☐
46. (a) ☐ (b) ☐ (c) ☒ (d) ☐
47. (a) ☐ (b) ☒ (c) ☐ (d) ☐
48. (a) ☐ (b) ☐ (c) ☒ (d) ☐
49. (a) ☐ (b) ☒ (c) ☐ (d) ☐
50. (a) ☐ (b) ☒ (c) ☐ (d) ☐