Pet Health Record

To The Pet Owner

To live a long, healthy life, all pets require regular medical attention. This health folio provides a written record of your pet's medical background. It will be helpful if you have it available to update when you bring your pet to the veterinarian for vaccinations or medical exams.

Although brief, here are several topics that will certainly arise throughout the life of almost every pet.

Feeding

Your veterinarian may recommend a feeding program. A complete and balanced diet including protein, carbohydrates, vitamins and minerals is necessary for proper growth and disease resistance. Here are some guidelines:

- Puppies or kittens should be fed small amounts 3-4 times per day. At 6 months, reduce to twice per day and once per day for mature pets.
- Name-brand commercial pet food is carefully balanced to meet your pet's nutritional requirements. However, your veterinarian may recommend vitamin/mineral supplements.
- Table scraps disrupt your pet's balanced diet and should be avoided. Bones should never be fed.
- During pregnancy and nursing, your pet has special dietary needs. Check with your veterinarian for feeding instructions.
- · Fresh water should be available at all times.

Vaccination

A routine vaccination program significantly increases your pet's likelihood of living a normal, healthy life. Some important diseases that should be considered for prevention appear in this health record under the section "Vaccination". For maximum protection, follow your veterinarian's advice and make sure your pet receives vaccinations and boosters when scheduled.

Spay-Neuter

There are many myths about spaying or neutering that should be dispelled. First, an animal does not have to produce "at least 1 litter of offspring" to lead a normal life. Second, animals will not always gain weight after spay or neuter. Although the tendency may be there, proper nutrition and exercise will reduce that likelihood. Third, it is not inhumane to sterilize an animal. The procedure is safe and effective.

Animal shelters and pounds across the country are burdened with unwanted animals. Responsible pet owners must take the lead in reducing the number of free-roaming animals and unwanted pets.


Parasite Control

A variety of parasites affect a pet throughout its lifetime. Treatment is usually simple but requires medications that must be used carefully and according to instructions. Proper sanitation will reduce likelihood and severity of many parasite infections, but regular medical exams and treatment are necessary for best control. Among the most important parasites that affect pets and require routine treatment are heartworms, worms of the gastrointestinal tract, fleas and ticks.

Surgery

Besides spay or neuter, it is not uncommon for your pet to require surgery at some point in its life. Accidents or other injuries as well as internal disorders may require surgical procedures.

Modern anesthetics and life-support agents similar to those found in human hospitals are available to your veterinarian and minimize risks of surgery. Preand post-surgical care require attention, and you should follow the instructions of your veterinarian. Withholding food or water the day the surgery is scheduled and confinement during recovery may be recommended. Medications could be prescribed after the surgery, and sutures may require removal.



Pet Health Record		
Name		
Date of Birth		
Breed		Sex
Markings		
Owner		
Address		
City	State	Zip
Telephone /		
Breeder		
Date Acquired		

Vaccination Record

	Canine	Feline
AGE DATE Wks	Lyme Disease Distemper Adenovirus 2 Parainfluenza Parvovirus Leptospira Coronavirus Bordetella Canine/Feline	Leukemia Virus Panleukopenia Calicivirus Rhinotracheitis Pneumonitis FIP Rabies
	00000000	0000000
Wks	00000000	0000000
1 Yr	00000000	0000000
2 Yrs	00000000	0000000
3 Yrs	00000000	0000000
4 Yrs	00000000	0000000
5 Yrs	00000000	0000000
6 Yrs	00000000	0000000
7 Yrs	00000000	0000000
8 Yrs	00000000	0000000
9 Yrs	00000000	0000000
10. Yrs	00000000	0000000
11. Yrs	00000000	0000000
12. Yrs	00000000	0000000
13. Yrs		
14. Yrs	00000000	
15. Yrs	00000000	

Lab	oratory Tests	Medical / Surgical History
Hear	tworm Prevention	Date
	Test	
Date	Pos Neg Injection Comments	
	I	
	zemia/FIV	
Date	Pos Neg Comments	
	<u> </u>	
		Lyme Disease Test
		Date Pos Neg Comments
	1 Exams/Deworming	Rabies Tag Information
Date	Pos Neg Comments	Date Number
	'	

Vaccination

Rabies

Of all animal diseases, rabies is probably the most feared. The rabies virus attacks the brain and is always fatal. Most pets are exposed to rabies by bites from wild animals, particularly skunks, raccoons, bats and foxes. The disease can be transmitted to humans through the bite or scratch of an infected pet. Vaccination of all dogs and cats is the most effective means of control.

Canine Distemper

This highly contagious viral disease is found wherever dogs are found. It affects the respiratory and nervous system and is often fatal. Primary vaccination should begin at 6-12 weeks of age since dogs often contract the disease at an early age.

Canine Parvovirus (CPV)

This contagious viral disease usually causes severe diarrhea and vomiting in dogs of all ages, but is especially deadly in puppies.

Canine Coronavirus (CCV)

Coronavirus is highly contagious and can weaken dogs by causing severe diarrhea and vomiting. The disease is sometimes confused with parvovirus. The two diseases may occur simultaneously, in which case symptoms are more severe.

Canine Parainfluenza

This viral respiratory disease is often partly responsible for "kennel cough" in dogs. Infection can be severe in young puppies. Parainfluenza protection is often included in distemper-parvo vaccines.

Canine Adenovirus Type 1 and Type 2

Canine Adenovirus Type 1 infection causes infectious hepatitis which may lead to severe kidney damage. Type 2 can be a complicating factor in kennel cough. Vaccines are available that protect against both types of adenovirus.

Canine Leptospirosis

Leptospirosis is a bacterial disease carried by many wild animals. A dog can contract the disease from infected animals or by drinking contaminated water. Yearly vaccination limits your dog's chances of acquiring the disease.

Canine Bordetella

Frequently involved in kennel cough complex, this bacterial infection may occur simultaneously with distemper, adenovirus type 2 infection, parainfluenza and other respiratory infections.

Canine Borreliosis (Lyme Disease)

Borreliosis, or Lyme Disease, is an infection caused by a bacteria which is spread primarily by the painless bite of an infected tick. Symptoms of Borreliosis in the dog include fever, lethargy, muscle stiffness, depression and lack of appetite. In more severe cases, lameness occurs as a result of severe musculoskeletal or arthritic type joint pain.

Canine/Feline Giardiasis

Giardiasis in dogs, cats and humans is caused by a waterborne parasite called Giardia lamblia. The parasite is found in untreated water, i.e. puddles, ponds and creeks. Symptoms include severe diarrhea, weight loss, fever, dehydration and nausea.

Feline Panleukopenia

Feline Panleukopenia, sometimes called "feline distemper", causes severe vomiting, diarrhea and dehydration. It is the most widespread disease of cats and causes high death loss, especially among kittens. The disease is caused by a virus and is highly contagious.

Feline Rhinotracheitis

Rhinotracheitis infection is a widespread respiratory disease caused by a virus. It is most severe in small kittens and can cause profuse discharge from the eyes and nose.

Feline Calicivirus

Calicivirus infection causes a variety of symptoms including fever, excess salivation and mouth or tongue ulcers. It is usually less fatal than rhinotracheitis or panleukopenia.

Feline Pneumonitis

Respiratory infection with symptoms resembling feline viral rhinotracheitis. The disease is caused by the organism Chlamydia psittaci and can be complicated by associated bacterial infections.

Feline Leukemia

Feline Leukemia is a form of cancer in cats which is usually fatal. The disease, caused by a virus, can lead to tumor growth nearly anywhere in the body as well as a variety of other symptoms. Infected cats are unable to resist other diseases and may die from associated infections. Testing for the disease is recommended prior to initiating a vaccination program.

Feline Infectious Peritonitis (FIP)

FIP is a disease caused by a coronavirus. The virus is spread by direct cat-to-cat contact or by contact with contaminated surfaces. There are 2 manifestations of the disease, wet and dry, and both have nonspecific symptoms such as intermittent inappetence, depression, rough hair coat, weight loss, and fever. There is no cure and the disease is considered fatal.