

Amsterdam Public Health



**Ecological Momentary Assessment
In Mental Health Research:**

**A practical introduction,
with examples in R**

Jeroen Ruwaard, Lisa Kooistra & Melissa Thong

Ecological Momentary Assessment in Mental Health Research

A Practical Introduction, With Examples in R

Jeroen Ruwaard

Lisa Kooistra

Melissa Thong

2018-10-26

Contents

| | |
|--|------------|
| Preface | vii |
| I Introduction | 1 |
| 1 Introducing EMA | 3 |
| 1.1 What is EMA? | 3 |
| 1.2 Why EMA? | 4 |
| 1.3 EMA research findings: A birds-eye view | 5 |
| 1.4 What is in this manual? | 8 |
| 2 Introducing R & RStudio | 11 |
| 2.1 What are R and RStudio? | 11 |
| 2.2 Why R? | 11 |
| 2.3 Installing R & RStudio | 12 |
| 2.4 Interacting with R through the RStudio console | 12 |
| 2.5 Writing R-scripts | 13 |
| 2.6 Importing your data | 13 |
| 2.7 Extending R with Packages | 15 |
| 2.8 Getting help | 16 |
| II EMA Methods | 19 |
| 3 Study Design | 21 |
| 3.1 What is the research question | 21 |
| 3.2 Who are the prospect participants? | 21 |
| 3.3 What are the qualities of the EMA measures? | 21 |
| 3.4 Constructing the Sample Plan | 22 |
| 3.5 Power Analysis | 23 |
| 3.6 Ethical considerations | 25 |
| 4 EMA instruments | 27 |
| 4.1 General considerations | 27 |
| 4.2 Technical considerations | 28 |
| 4.3 Participant perspective | 28 |
| 4.4 Points to consider | 29 |
| 5 Data management | 31 |
| 5.1 Using RStudio-projects to manage EMA data | 31 |
| 5.2 Project directory structure | 31 |
| 5.3 The keyfile | 32 |
| 5.4 Source data | 33 |

| | |
|---|-----------|
| 5.5 Import & Prune | 33 |
| 5.6 Visual Exploration | 33 |
| 5.7 Analysis | 33 |
| III EMA Outcomes | 35 |
| 6 Mood | 37 |
| 6.1 Unidimensional mood assessment | 37 |
| 6.2 Bag-of-Items | 38 |
| 6.3 Multi-dimensional mood assessment | 39 |
| 7 Activity | 45 |
| 7.1 Accelerometry | 45 |
| 7.2 Location analysis | 47 |
| IV Analytic Approaches | 51 |
| 8 Feature Extraction and Selection | 53 |
| 8.1 Feature extraction | 53 |
| 8.2 Reliability and Validity | 55 |
| 8.3 Missing values | 55 |
| 8.4 Feature Selection | 55 |
| 9 Mixed Modeling | 57 |
| 9.1 The Mixed Model | 57 |
| 9.2 Simulating example data | 57 |
| 9.3 Fitting a mixed model in R | 58 |
| 9.4 Adding time as a predictor | 59 |
| 9.5 Adding a Two-Group Comparison | 61 |
| 9.6 Further reading | 62 |
| 10 Fitting Networks | 63 |
| 10.1 What are Networks? | 63 |
| 10.2 Fitting Networks on Single-Subject Repeated Measures Data | 63 |
| 10.3 Fitting Networks on Multiple Subjects Repeated Measures Data | 66 |
| 10.4 Discussion | 67 |
| V EMA Case Studies | 69 |
| 11 Early Warning Signs of Depression | 71 |
| 11.1 Critical Slowing Down | 71 |
| 11.2 Plotting the course of depression | 72 |
| 11.3 Mental state EMA items | 72 |
| 11.4 Running the DFA | 73 |
| 11.5 Results | 74 |
| 11.6 Discussion | 75 |
| 12 Homerange Estimation | 77 |
| 12.1 Section | 77 |
| 12.2 Spatio-temporal analysis: | 77 |
| 12.3 Section | 78 |
| 12.4 Section | 78 |

| | |
|--|------------|
| VI EMA Catalogues | 79 |
| 13 EMA Research within the APH Mental Health Consortium | 81 |
| 13.1 Depression | 82 |
| 13.2 Psychosomatic symptoms | 85 |
| 13.3 Suicidal Ideation | 88 |
| 13.4 Psychotic symptoms | 89 |
| 13.5 Symptom Networks | 90 |
| 13.6 Sleep | 91 |
| 13.7 Ambulatory Assessment | 92 |
| 14 EMA Instruments Catalogue | 95 |
| 14.1 GENEActive | 95 |
| 14.2 Ilumivu | 95 |
| 14.3 MoodBuster | 96 |
| 14.4 Movisens | 97 |
| 14.5 PsyMate | 98 |
| 14.6 RoQua | 99 |
| 15 R Packages for EMA Research | 101 |
| 15.1 Accelerometry | 102 |
| 15.2 Data management & Visual Exploration | 103 |
| 15.3 Mixed-effects modeling | 106 |
| 15.4 Power analysis | 106 |
| 15.5 Symptom Network Analysis | 107 |
| 15.6 Timeseries analysis | 110 |
| VII Closing Matters | 113 |
| Acknowledgements | 115 |

Preface

Given known limitations of retrospective self-report questionnaires, such as recall bias and poor generalisability of assessment results to real-life situations, mental health researchers increasingly adopt alternative assessment methods. One of the promising alternatives is Ecological Momentary Assessment (EMA), in which emotions and behaviours are repeatedly sampled in everyday life, through wearable electronic devices.

Repeated measurement can reveal important characteristics of the dynamics of phenomena, as illustrated by the cover of this book. With EMA, we can tap into mental health processes that were, up to very recently, unavailable to scientific research.

Conducting an EMA study, however, can be challenging. Researchers face a dazzling array of options related to the electronic wearables, outcomes selection, study design considerations, ethical and regulatory constraints, data management, statistical analysis, and study reporting. Although standards are emerging, clear guidelines for EMA research do not - at present - exist. EMA studies have unique characteristics that require specialist research skills, related to study design and statistical analysis. This manual was written to fill this gap.

The manual provides a practical introduction to EMA-research. It was written to aid beginning researchers of the Amsterdam School of Public Health (APH), who are looking for practical advice in conducting EMA studies. The manual provides an overview of EMA instruments, outcomes, methods and analytical techniques, guidelines for EMA-studies, and a catalogue of EMA research in the APH consortium.

This manual was written in Bookdown (Xie, 2016, 2018). Sources are freely available at ‘github’, via https://github.com/jruwaard/aph_ema_handbook. Please post your comments and suggestions there, or via e-mail, through aph.ema@ggzingeest.nl.

Part I

Introduction

Chapter 1

Introducing EMA

1.1 What is EMA?

Ecological Momentary Assessment (EMA) has many aliases. It is known as ‘Experience Sampling’ (Larson and Csikszentmihalyi, 1983), ‘Ambulatory Assessment’ (Ebner-Priemer and Trull, 2009) or ‘Ambulatory Self-reporting’ (Conner and Feldman Barrett, 2012), ‘Real-time Data Capturing’, the ‘Continuous Unified Electronic Diary Method’ (Ellis-Davies et al., 2012), and as the ‘Intensive-longitudinal Study Design’ (Bolger and Laurenceau, 2013). The different terms stress different aspects of EMA research. All, however, refer to research methods that involve the *repeated sampling of people’s current thoughts, emotions, behaviour, physiological states, and context, in their natural environment, typically (but not necessarily) via electronic wearable devices* (Shiffman et al., 2008).

1.1.1 Active versus Passive EMA

In EMA research, two forms of data collection are distinguished: 1) ‘Active EMA’, with which *self-report* data are collected, and 2) ‘Passive EMA’, with which *observational* data are collected. Active EMA requires participants to consciously provide information, for example by rating their current mood in response to a question that is prompted on their smartphone (see Figure 1.1). In passive EMA, information is collected through wearables or log files without active involvement of participants, for example on heart-rate, activity, smartphone use or engagement on social media (see Figure 1.2). Studies may combine active and passive EMA. A study into sleep patterns, for example, may involve both a self-report sleep diary and an accelerometer sensor (Van Der Meijden et al., 2016).




Figure 1.1: Active EMA: data are collected by prompting questions to participants, for instance by using an EMA app such as MoodBuster.




Figure 1.2: Passive EMA: data are collected automatically, for instance by accelerometers such as the GeneActive actiwatch.

1.1.2 EMA sampling

EMA is further typified by the way in which data sampling is triggered. EMA sampling may be triggered by a signal (signal-contingent sampling), an event (event-contingent sampling), or a combination of both (Conner and Lehman, 2012).

In *signal-contingent sampling*, participants respond to questions when they are prompted to do so by a signal (or ‘beeps’, as these signals are often called in the literature). In *event-contingent sampling*, study participants complete an assessment whenever a specific event occurs, such as a panic attack or alcohol consumption.

Signal-contingent sampling can follow a fixed or a random scheme. In a fixed scheme, participants are prompted at fixed time-points, for example at 9:30, 12:30, and 16:30. In a random scheme, prompts are sent at random time points, typically in pre-set intervals. For example, participants could be prompted to complete two assessments per day, one at a random time point between 10:00 and 14:00, and one at a random time point between 14:00 and 16:00. Using pre-set intervals ensures that participants do not receive several prompts within a limited time-frame (Piasecki et al., 2007). In addition, it ensures that participants are not bothered by prompts at inappropriate times (e.g., participants may not appreciate prompts after 22:00 and before 7:30).

With ‘event-contingent sampling’, the sample rate is determined by the occurrence rate of the event. One way to implement this is to simply ask study participants to complete a questionnaire whenever the event occurs. When active EMA is combined with passive EMA, it may be also be possible to trigger event-based prompts automatically based on changes in passive data, for example by triggering an EMA questionnaire automatically whenever a significant change in activity level is detected (Smyth and Stone, 2003).

1.2 Why EMA?

1.2.1 To minimize recall bias in assessment

In clinical research, self-report questionnaires are often used to assess the presence and severity of symptoms in the recent past. While useful, these retrospective self-reports are not without drawbacks, since they tap into the memory of respondents, which can be distorted (Shiffman et al., 2008; Moore et al., 2016). EMA circumvents this recall bias, by asking participants to rate their *current state*, rather than asking them to reflect on past experiences.

1.2.2 To increase the ecological validity of research findings

A key feature of EMA is the collection of data in real-world environments, as participants go about their daily activities, as opposed to data collection in controlled labs or research settings (Shiffman et al., 2008). Thus, EMA data can be expected to result in research findings that have better ecological validity and better generalization to the subject’s lived experience in real-world settings. Practical applications derived from EMA data are therefore expected to be more relevant to real-life situations.

1.2.3 To advance ideographic research

In clinical research, a distinction is often made between ideographic and nomothetic methods (Allport, 1937). Ideographic methods are those that “aim to identify patterns of behaviour within the person across a population of experiences or situations, and nomothetic methods[are] those that aim to identify patterns of behaviour across a population of individuals, rather than for any given individual” (Conner et al., 2009). The difference is important. As is increasingly recognised, group-level findings do not necessarily generalize to the individual members of the group, as shown by Figure 1.3 (adapted from Hamaker (2012)).

In contrast to more qualitative idiographic methods, such as interviews and $N = 1$ case studies, EMA allows for the collection of large amounts of quantitative data on the individual level. Thus, EMA offers a quantitative method for *idiographic* research, measuring characteristics of (unique) individuals across time and context (Shiffman et al., 2008). This allows for a better understanding of intra-individual process, and of the factors that account for variability within and between individuals.




Figure 1.3: An illustration of how group-level and individual-level processes can differ dramatically: the relationship between x on y is negative in the group (as shown by the descending regression line), but positive for individuals (marked by ellipses).

1.3 EMA research findings: A birds-eye view

The use of EMA in mental health research might seem novel but this methodology has a long track record. Already in the 1980's, early pioneers used electronic devices to elicit responses from study participants to tap into (mental) health processes in everyday life (see, e.g., Csikszentmihalyi and Larson, 2014). Recent years, however, have witnessed a large increase in EMA research. Rapid technological developments, a marked interest in the individual, and a wide recognition of the need to study health-related processes in real-life situations have all contributed to this.

EMA-based research in mental health has produced an impressive trove of findings that have supported and, sometimes, challenged existing theories on behaviour. EMA data, whether collected as self-report or via wearable device/sensor, have diagnostic, monitoring, management, or intervention applications (Patel et al. (2012); Aung et al. (2017); Evenson et al. (2015)]. Its feasibility for mental health research is evidenced by its use in observational studies and randomized controlled trials on a wide range of topics and populations. Below, we present a non-exhaustive summary of systematic reviews and meta-analyses of EMA research.

1.3.1 Active EMA

Mood disorders have been well-studied using active EMA methods (Wenze and Miller, 2010), with several reviews outlining recent findings in depression (Telford et al., 2012; Wichers et al., 2011; Burton et al., 2013), anxiety disorders (Walz et al., 2014), and depression/bipolar disorder (aan het Rot et al., 2012). The potential of EMA for use among young populations showed promising results (Dubad et al., 2018). Innovations in mobile devices has improved the feasibility and popularity of ecological momentary interventions (EMIs) for anxiety and depression (Schueller et al., 2017). A systematic review and meta-analysis of EMIs reported small to medium effects on mental health (Versluis et al., 2016).

EMA appears to be particularly well-suited to examine the role of emotions in the development and maintenance of obesity and eating disorders (Engel et al., 2016). Meta-analytic results suggest that negative affect, rather than hunger, is associated with binge eating among individuals with eating disorders (Haedt-Matt and Keel, 2011; Haedt-Matt et al., 2012). EMA methods have also significantly contributed to the understanding of the processes that drive substance use, cessation, and relapse, often in contrast with theory-driven studies largely derived from global reports collected through retrospective questionnaires (Shiffman, 2009; Swendsen, 2016).

1.3.2 Passive EMA

Objective EMA data collected passively through bio-sensors, smart devices, or context/environmental (e.g. location) has been shown to be a feasible and promising method for the longitudinal monitoring of individuals with affective disorders (Dogan et al., 2017; Kirchner and Shiffman, 2016). The potential of passive sensing via smartphone for mental health research are outlined in two reviews, encompassing the assessment of health and well-being (Cornet and Holden, 2018), and the measuring, understanding, and intervening in mental illness and maintaining mental health (Aung et al., 2017). A systematic review and meta-analysis on actigraphy reported diurnal variations in activity levels among individuals with depression (Burton et al., 2013). Compared with traditional self-reports, passive sensing was reported to be less intrusive and to result in more accurate data, while providing options for continuous monitoring and feedback.

1.3.3 Limitations

While available reviews suggest that EMA is a feasible research method that has the potential to make significant contributions to mental health research, one significant drawback of EMA highlighted in several reviews is the lack of high-quality studies. Other general limitations considered include:

- Unclear generalizability of research findings due to selective samples (Burton et al., 2013), and small sample sizes (Dogan et al., 2017)
- Unclear practice effects (Telford et al., 2012), and issues related to reactivity (Shiffman, 2009)
- Issues related to the feasibility and tolerability of prolonged and intense data periods of data collection (Wichers et al., 2011)
- Issues of privacy, ethics and informed consent (Cornet and Holden, 2018)
- Changing responsibilities of researchers and clinicians in EMA studies, e.g. in suicide ideation research (Wenze and Miller, 2010)
- Unresolved methodological issues (Dubad, 2018)

Table 1.1: A selection of reviews of EMA studies targeting various mental health conditions

| Topic | Author (Year) | Outcome |
|--------------------------|---------------------------|--|
| Anxiety disorders | Walz et al. (2014) | Provides insights to the temporal variability of symptoms, and associations between daily affect, behaviours, and situational cues. Combination of EMA and ambulatory assessment of physiological variables and treatment evaluations. |
| | Schueller et al. (2017) | Provides an overview of the distinction of EMIs from other types of treatment. Also discusses the considerations of conducting EMI research, such as design, deployment, and evaluation |
| Eating disorders | Engel et al. (2016) | An overview of studies on eating disorders, obesity, and bariatric surgery using EMA |
| Mood disorders | aan het Rot et al. (2012) | Provides an overview of EMA studies on correlates of mood, treatment effects, residual symptoms of remitted patients, paediatric populations, MDD/BD specificity, and links with neuroscience |
| | Aung et al. (2017) | Provides a conceptual review of passive sensing techniques for measuring, understanding, and treatment of mental illness. |
| | Burton et al. (2013) | Focuses on diurnal variations in activity levels among depressed individuals. |
| | Telford et al. (2012) | Identifies six themes of EMA research in MDD: methodology; positive and negative affect; cortisol secretion; antidepressant treatment; work performance; genetic risk factors |
| | Wenze and Miller (2010) | Provides an overview of EMA in mood disorder research comprising techniques used, types of population assessed, types of research questions, and a discussion of the potential of EMA in treatment settings |

| Topic | Author (Year) | Outcome |
|------------------------------------|------------------------------|--|
| | Wichers et al. (2011) | Provides an overview for the potential clinical application of EMA in the diagnostic and treatment of MDD. |
| | Versluis et al. (2016) | Provides an overview of interventions (EMI) addressing anxiety, depression, and perceived stress on positive psychological outcomes |
| | Dabad et al. (2018) | Provides an overview of the feasibility and clinical impact of mood-monitoring applications targeting young populations (10-24 years old) |
| Substance-related disorders | Shiffman (2009) | Review of processes that drive substance use, cessation, and relapse, sometimes in contrast with theory-driven studies that are largely derived from global reports collected through questionnaires |
| | Swendsen (2016) | Conceptual review of the use of mobile technologies for research on addiction and its treatment |
| Mental health/ Well-being | Cornet and Holden (2018) | Outlines the potential and challenges of passive sensing to detect status change and behaviour change following feedback on behaviour. |
| | Kirchner and Shiffman (2016) | Provides an overview of 'geographically explicit momentary assessment' (GEMA) research to enrich EMA research in mental health and well-being |
| | Dogan et al. (2017) | Provides an overview of studies that combine subjective ratings with objective EMA-data collected using smartphone-based systems |

1.4 What is in this manual?

This manual comprises six parts, in which various aspects of EMA research will be discussed.

- In **part I**, Chapter 2 introduces R, the statistical program that is an indispensable tool for the EMA researcher.
- **Part II** focuses on EMA study design (Chapter 3, EMA instruments (Chapter 4) and EMA data

management (Chapter 5).

- **Part III** details the momentary assessment of two specific outcomes: Mood (Chapter 6) and Activity (Chapter 7).
- **Part IV** introduces EMA data analysis techniques: Feature Extraction (Chapter 8), Mixed Modelling (Chapter 9), and Network Analysis (Chapter 10).
- In **part V**, the application of the preceding material is illustrated in two case studies: a study into the use EMA to detect early warnings signs of depression (Chapter 11), and a study into a new GPS-based measure of activity (Chapter 12).
- **Part VI** provides three catalogues of EMA resources. Chapter 13 lists EMA research groups within APH, chapter 14 lists EMA instruments that were found to be in use among APH researchers, and Chapter 15 summarises useful R extensions (packages) for EMA data analysis.

Chapter 2

Introducing R & RStudio

In this chapter, you will learn how to install and use two programs that are indispensable for the management and analysis of EMA data: R and RStudio.

2.1 What are R and RStudio?

R is a programming language and software environment for statistical computing and data visualisation. RStudio is a powerful user interface to R. It has many useful features that greatly simplify R-work. We strongly advise you to adopt the R/RStudio-combo.

2.2 Why R?

R, you may have been told, is for data scientists, methodologists, and scientific programmers only. It has a steep learning curve. If you are trained in SPSS, it will take time to become as productive in R as in SPSS. Why then, should you invest in R?

- Unlike SPSS, R is free. It does not eat up your budget. Why pay for something that you can get for free?
- R is cutting-edge. Methodological innovations first appear in R. Network analyses, for example (see Chapter 9), can be run in R, but not (yet) in SPSS. For some analyses, you need this alternative.
- Mastering R improves your connection to the statisticians in your team. They probably prefer R over SPSS. It is more efficient and less error-prone to all speak the same language.
- R is great for data-management. Clinical research, and especially EMA research, requires hundreds of operations on multiple raw data files. R excels at that. SPSS, frankly, does not. If you care about reproducible research (which you should), R can be a great help in putting it into practice.
- R can be used at different levels. If you want to be a basic user, that's fine. However, if you want to dive deeper, you will find that you can easily do so. You can study source code to understand a particular technique better. You can code new functions. R allows you to grow.
- R's user base is expanding every year. Chances are high that R will be the standard in your next workplace. R will look great on your CV.

You don't have to be a programmer or methodologist to use R. Yes, it takes time to master its full potential, but you should be able to run basic analyses in it within a week. This chapter will get you started.

2.3 Installing R & RStudio

Both R and RStudio are available, at no costs, for all major operating systems.

- Download R from the Comprehensive R Archive Network (CRAN), at <https://cran.r-project.org/bin/>
- Download RStudio from <http://rstudio.org>

Install R first, and RStudio second. If you install the programs in this order, RStudio will automatically find R on your computer.

If you installed R or RStudio previously, please update. This book assumes you will be working with version 3.4.2 (or higher) of R, and version 1.1.414 (or higher) of RStudio.

2.4 Interacting with R through the RStudio console

If you open RStudio, you will be presented with the interface shown in Figure 2.1. Rstudio's main window is divided in four panes (subwindows), which further contain several tabbed windows.




Figure 2.1: The RStudio Interface

Commands are sent to R starts in the bottom-left pane, named “Console”. To test this, move your cursor to the bottom line, immediately after the prompt sign (“>”). Next, type the statement below (note that ‘#’ denotes a comment line; R ignores it, so there is no immediate need to type that). To execute, press ‘Enter’.

R will execute the command and return the answer back to the console.

```
# Code snippet 2.1: R is a calculator.
1 + 1
```

Results of calculations can be saved into variables, by making use of the assignment operator (“<-”). If you type the name of a variable, R returns its value.

```
# Code snippet 2.2: Using <- to declare and set a variable.
N <- 50 + 50
N
#> [1] 100
```

To appreciate why R is such a popular tool for statistical computing, consider the following command, which, in one line, 1) uses the variable N, just created, to 2) generate 100 random numbers from the normal distribution, and 3) plot a histogram of these numbers.

```
# Code snippet 2.3: Plotting the histogram of a sample from the normal distribution.
hist(rnorm(N))
```

The plot appears in the bottom-right pane, as in Figure 2.1.

2.5 Writing R-scripts

Working in the console is a great way to interactively explore R and data, but what if you want to save a particularly useful chain of statements? For this, you can use a script file.

To create a script file, use the RStudio menu: **File > New File > R Script**. This will open a new tab in the top-left pane of RStudio, where you can edit the script.

- In the script window, type all statements that you have been entering in the console in the previous section.
- Next, select all lines in the script.
- Press **Ctrl+Enter** to run the script.

All commands in the script are executed. The commands are echoed in the console pane, and results are shown immediately, as was the case before, when you typed the commands in the console yourself.

Scripts can also be run line by line. Move the cursor to the line you want to run, and press “Ctrl+Enter”. The line is copied to the console and executed, and the cursor in the script will move to the next line, allowing you to walk through the script, step by step.

2.6 Importing your data

Something that confuses new Rstudio users, who are more familiar with SPSS, is that it is not obvious how to import data into RStudio. In SPSS, the data are in plain sight. In R, you first have to import the data.

2.6.1 Using RStudio menu's to import data

One way to load data into R is to use RStudio’s data import wizard. Follow the steps below to see how this works with data stored in a comma-separated-values (csv) format, a common data format to which many programs, including SPSS and Excell, can export data to.

- Download the example csv data file at <https://tinyurl.com/ybfafxxk> (or create a csv-version of one of your own data files).
- In RStudio’s menu, choose **File > Import Dataset > From Text (base)**.
- In the window that appears, click on **Browse** to locate the csv- file on your computer, and click **Import** in the next window (see Figure 2.2).

RStudio shows the data, in tabular view, in the top-left window, ready for analysis. You will also find a new entry in the **Environment**-tab in the top-right pane. When you click the small arrow, at the left of the name, you will see a brief summary of the contents of the data.




Figure 2.2: RStudio’s CSV import wizard.

2.6.2 Using functions to import data

While RStudio’s Data import wizard is useful, you will probably use it less over time. Most likely, you will convert to using the more efficient R commands to import data. For example, it takes only a single line to download and import the example data.

Code snippet 2.4: Importing csv-data from the internet.

```
ESMdata <- read.csv(url("https://tinyurl.com/ybfafxxk"), row.names = NULL)
```

2.6.3 Accessing your data

Since the data is now in the environment (under the name `ESMdata`), you can use it in other R commands. For example, to produce a more detailed summary of the first four columns of `ESMdata`, you type:

Code snippet 2.5: Summarising data.

```
summary(ESMdata)
      dayno      beepno      mood_relaxed      mood_down
Min.   : 1.0   Min.   : 1.00   Min.   :1.000   Min.   :-3.0000
1st Qu.: 61.0  1st Qu.: 3.00   1st Qu.:4.000   1st Qu.: 0.0000
Median :252.0  Median : 5.00   Median :4.000   Median : 0.0000
Mean   :198.9  Mean   : 5.24   Mean   :4.173   Mean   : 0.1784
3rd Qu.:303.0  3rd Qu.: 8.00   3rd Qu.:5.000   3rd Qu.: 0.0000
Max.   :366.0  Max.   :10.00   Max.   :7.000   Max.   : 3.0000
NA's    :2
```

```
      mood_irritat
Min.   :1.000
1st Qu.:1.000
Median :2.000
Mean   :2.241
3rd Qu.:3.000
Max.   :7.000
NA's   :3
```

To inspect the first 6 lines of data, type,

Code snippet 2.6: Show first 6 lines of a data frame.

```
head(ESMdata)
#>   dayno beepno mood_relaxed mood_down mood_irritat
```

```
#> 1 226    1      5      -1      1
#> 2 227    5      4      0      3
#> 3 227    6      4      0      2
#> 4 227    8      4      0      1
#> 5 227    9      4      0      2
#> 6 227   10      5      0      1
```

To view all rows of data in a spreadsheet (as in Figure 2.2, type:

```
# Code snippet 2.7: Show data as spreadsheet.
View(ESMdata)
```

To work with a specific variable in the dataset, use '\$'. For instance, to print the first 20 numbers in the `mood_relaxed` variable, type:

```
# Code snippet 2.8: Accessing a single variable in a data frame.
head(ESMdata$mood_relaxed, n = 20)
```

This allows you to apply functions to specific variables. For example, to calculate the mean of scores in `mood_relaxed`, type:

```
# Code snippet 2.19: Calculating the mean of a variable.
mean(ESMdata$mood_relaxed)
#> [1] 4.173442
```

There are many ways in which you can summarise and manipulate your data. At this point, the important milestone is that you imported and accessed data in R.

2.7 Extending R with Packages

R's attractiveness lies in the ease with which it can be extended with new functionality. Through so-called packages, which can be freely downloaded from the internet, specialised functions can be added to your workspace.

2.7.1 Installing R-packages from CRAN

Packages can be found at the CRAN website. To browse through the impressive list of available packages, see https://cran.r-project.org/web/packages/available_packages_by_name.html

If you find a package you like, you can install it via the RStudio menu system, choosing `Tools > packages`. But you can also use the console, via the `install.package` function.

A popular package, `tidyverse`, is used extensively in the examples of this manual. Package `tidyverse` comprises a set of popular packages from the creators of RStudio, that greatly simplify working with R. So, while you are at it, install this package now.

```
# Code snippet 2.10: Installing a package from CRAN.
install.package(tidyverse)
```

The `tidyverse` contains a package called 'haven', which allows you to read and write SPSS datafiles (.sav files). This is very convenient. You don't have to convert all your SPSS data to csv files. See `?read_spss` to learn how to import an SPSS-file (or use the data import wizard, by choosing 'File > Import Dataset > From SPSS', in RStudio's top-right pane).

2.7.2 Installing R-packages from GitHub

Not all packages are at CRAN. Many 'unofficial' packages are shared at a site called 'GitHub'. This book's companion R package 'emaph', for example, which contains specialised EMA functions datasets, is on GitHub.

You need package emaph to run many examples in the book, so let's install this package now.

GitHub packages can be installed via the `install_github` function, which is defined in a package called 'devtools'. So, to install 'emaph', enter the following in the console:

```
# Code snippet 2.11: Install the GitHub 'emaph' package.
install.packages("devtools")
devtools::install_github("jruwaard/emaph")
```

2.7.3 Using packages

To use packages, you have to tell R to load them. You do this with the `library` function. For example, to use package 'tidyverse' and 'emaph', type:

```
# Code snippet 2.12: Loading packages.
library(tidyverse)
library(emaph)
```

Once loaded, you can use the functions and datasets of the packages. Packge 'emaph' provides dataset 'csd', which contains the data from the 'critical slowing down'-study (Kossakowski et al., 2017; Wichers et al., 2016), in which a patient recorded his mood, for 239 days (see also Chapter 11).

To plot the irritation levels of this patient in the first six days, using the `ggplot` function from package 'ggplot2' (which is in 'tidyverse'), type:

```
# Code snippet 2.13: Using ggplot to plot EMA time series.
ggplot(data = subset(csd, dayno <= 6),
       mapping = aes(x = beepno, y = mood_irritat)) +
  geom_point() + geom_step() +
  scale_x_continuous(breaks = 1:10) +
  facet_wrap(~ dayno, nrow = 2)
```




Figure 2.3: Irritation levels of a single patient, in the first six days of an EMA study.

2.8 Getting help

R has no point-and-click menu's that you can browse through to select a statistical procedure. This is a problem for many new users. What if you want, for example, to generate random numbers from a distribution with a mean of 2 and standard deviation of 4? How to tell this to R?

2.8.1 Using ‘?’ to consult the documentation

The good thing is that you already known the name of the function to use, since we used it in the previous section: it is ‘rnorm’. To check the documentation of this function, type `?rnorm` in the console.

```
# Code snippet 2.14: Using '?' to find the documentation of a function.
?rnorm
```

This opens the documentation of the `rnorm` function in the ‘Help’-tab, in the bottom right pane, from which you learn that that the `rnorm` function accepts `mean` and `sd` (standard deviation) as additional parameters, which are 0 and 1 default, respectively (which explains why `norm(100)` worked in the previous examples). So, to generate the required numbers, you type:

```
# Code snippet 2.15: Plotting the histogram of a custom random sample
hist(rnorm(1000, mean = 2, sd = 4))
```

All functions in R are documented, and this documentation is shown in RStudio’s Help pane when you prepend `?` to the name of the function in the console.

2.8.2 Using RStudio’s global documentation index search

What if you do not know the name of a function? Suppose you want to run a t-test for independent groups. Does R have a function for that?

At the top-right of the ‘Help’ pane, RStudio has a search input field, which allows you to search through all documentation that is installed on your computer. The search field auto-completes your input. If you type a ‘t’ in this field, you will be presented with a list of functions starting with a ‘t’. In this list, you find a likely candidate: a function called `t.test`. From the documentation of this function (`?t.test`), you learn that, indeed, this is the function you were looking for.

```
# Code snippet 2.16: Running a t-test, on two simulated samples.

# generate two samples (N = 100 per group) from the normal distribution
A <- rnorm(100); B <- rnorm(100)

# the t-test should be non-significant
t.test(A, B)
#>
#> Welch Two Sample t-test
#>
#> data: A and B
#> t = -0.72282, df = 195.84, p-value = 0.4707
#> alternative hypothesis: true difference in means is not equal to 0
#> 95 percent confidence interval:
#> -0.3768889 0.1747184
#> sample estimates:
#> mean of x mean of y
#> 0.01778337 0.11886860
```

2.8.3 Learning from examples

This book contains many R code snippets. By studying these examples, you will become more familiar with R.

Some examples will introduce R language constructs and functions that are unknown to you. Learn from from these examples, by using `?` on each element that you do not understand.

2.8.4 Google

With Google, you will find many answers to your R questions. Googling for “t-test R”, for example, results in a rich set of online resources. Good resources are:

- RSeek (see <http://rseek.org/>)
- Stackoverflow: (see <https://stackoverflow.com/questions/tagged/r>)
- SearchR (see: <http://search.r-project.org/>)

2.8.5 Read books

This book does not provide a comprehensive tutorial. There is no need for that, since excellent resources are readily available. A selection is presented below.

- Many mental health researchers own a copy of Andy Field’s popular book “Discovering Statistics Using IBM SPSS Statistics” (Field, 2013). For those, Field’s R-version of this book, “Discovering Statistics Using R” (Field et al., 2012) provides a familiar companion in making the transition to R. See <https://www.discoveringstatistics.com/>
- Free manuals can be found at the official CRAN site. The manuals are dry, but complete and authoritative, since the authors are members of the R core development team. See <https://cran.r-project.org/manuals.html> (or type `help.start()` in the console).
- While at CRAN, be sure to browse the ‘contributed documentation’-section. On this page, you will find many freely available manuals contributed by the R community. See <https://cran.r-project.org/other-docs.html>

2.8.6 Online Courses

- DataCamp, an online data science education platform, offers several high-quality interactive courses in R. See <http://www.datacamp.com>
- The Try-R course at the CodeSchool website provides an alternative to DataCamp. See: <http://tryr.codeschool.com/>
- The Quick-R website provides a solid, concise, and rich introduction to R. See <https://www.statmethods.net/>

2.8.7 Learn R, in R

Package ‘swirl’ comprises a set of interactive courses that teach many aspects of the R language. See <http://swirlstats.com>

```
# Code snippet 2.18: Starting the interactive swirl-course in R.
install.packages("swirl")
library("swirl")
swirl()
```

Part II

EMA Methods

Chapter 3

Study Design

As with all scientific research, EMA studies start with mindful consideration of the study design. Issues that need to be considered are the research question(s), the hypotheses, the population of interest, and the nature of the comparison groups (Shiffman et al., 2008).

Ample information on general study design issues can be found elsewhere (see for example, the APH quality handbook, at <http://www.emgo.nl/kc/>). This chapter highlights key design aspects of EMA studies.

3.1 What is the research question

Given the plethora of new research options that emerged from the rapid development in EMA technologies, it can be tempting to dive straight into exploratory data collection, without giving much consideration to the theoretical background of the study. That, however, would be one pitfall of EMA research to avoid. Data mining is no substitute for theory. Asking participants to contribute data without a rationale is unethical. As in all scientific activities, defining the research question should be the first step.

Ask yourself what EMA could bring to your topic of interest. How is it different from traditional assessment methods? What questions does it allow you to address that you could not answer without it? For this, you could use any of the EMA advantages discussed in Chapter 1. Are you interested in real-life behaviour, in differences between participants, in changes within participants over time, or in potential causal pathways between health-related variables? What relationships do you expect to find, and why? A solid theoretical background, and clearly formulated explicit research questions and hypotheses will help to make the right choices when you have to decide on the other aspects of the study design.

3.2 Who are the prospect participants?

Given the experimental nature of EMA, studies are often piloted in healthy or sub-clinical populations. This is a recommended first step, to test the experimental procedures and to avoid unnecessary burdening of vulnerable patient populations. You should be aware, though, that results obtained in non-patient populations do not necessarily generalize to patient populations. EMA mood ratings, for example, might be much more variable in patients compared to non-patients. Pilot studies should therefore also be conducted in the target population.

3.3 What are the qualities of the EMA measures?

With the study hypotheses in place, theoretical constructs must be operationalized into quantitative measures. For this, you should consult the scientific literature on the reliability and validity of existing EMA measures.

In active EMA research, complex multi-dimensional constructs such as mood and anxiety are often measured using single-item questions, to reduce the assessment burden of participants, who are prompted several times per day. You should ask yourself (and the scientific literature) what the psychometric properties are of these measures are. How do these EMA-measures compare to scores on traditional assessments (e.g., self-report questionnaires, clinical interviews)? What is known about the variability of items scores? And what is known about the measurement errors? Surprisingly, this last question is often ignored in EMA research, even though it has been estimated that up to 30% to 50% of the observed variance in intensive repeated measures data can be attributable to measurement error (Schuurman et al., 2015).

In passive EMA research, it is important to be aware of the characteristics and limitations of the ‘data acquisition interfaces’ (Stone and Shiffman, 2002). For example, if you want to collect accelerometer data, a variety of options exist. You can collect these data via the built-in accelerometer of the smartphone of the participants, via cheap commercially available activity trackers, such as Fitbit, or via expensive wearable devices that are developed specifically for scientific research. Each option comes with specific advantages and disadvantages. Smartphone-based accelerometer data, for example, can be collected with relative ease, but these data can also be noisy and incomplete, since sample rates can often not be set and the precision of the built-in sensors varies considerably from device to device. Commercial accelerometers may have better precision and reliability (see, e.g., Evenson et al. (2015)), but manufacturers often limit access to raw data and data pre-processing algorithms, making it difficult (or even impossible) to fully analyse outcomes. Being aware of these issues when you plan the study, will help considerably in the analysis stage of your study.

3.4 Constructing the Sample Plan

An important next step is to define the EMA data sample plan. Questions that need to be answered are:

- How many days will data collection last?
- On each day, how often are participants assessed?
- How and when are participants invited for assessment?

The questions above should be answered as detailed as possible to best serve the research question and the statistical power (see below). In practice, however, it is often necessary to balance between research interests, respondent burden, and practical considerations, such as hardware limitations.

When determining the appropriate sample plan, start with mapping the expected fluctuation or patterns, based on available knowledge. For example, when an event is rare, it can be sufficient to ask participants to initiate EMA whenever the event occurs, or prompt them with an end-of-day diary. Adding more prompts in this scenario would not lead to more reliable data (Piasecki et al., 2007).

Increasing the assessment frequency and study duration will allow for a more detailed assessment of the outcome of interest. It is tempting to collect often and for a long period of time. However, this may also increase respondent burden, which may, in turn, affect study adherence and accuracy. Measurement reactivity could also occur, where the EMA-induced enhanced focus on the outcome of interest causes participants to increase or decrease on this outcome (Hufford et al., 2002; van Ballegooijen et al., 2016).

Issues related to hardware should also be considered. Electronic wearables have limited battery life and memory storage space. Memory space limitations of actigraphy watches may require participants to visit the research site. GPS-monitoring apps may have a negative impact on the battery life of the smartphone of the participants. These practical issues may also result in data loss and study drop-out.

Once all decisions related to the sampling plan are made, the procedure should be thoroughly tested. As a first step, it can be insightful to simulate the sample plan, as is done below, using the ‘sample_plan’ function, which is part of package ‘emaph’:

```
# Code snippet 3.1: simulating a signal-contingent sample plan.
library(emaph)
plan <- sample_plan(n_participants = 5,
                     n_days = 2,
```

```
times = c("09:00-11:00", "12:30", "17:00-19:00"),
plot = TRUE)
```




Figure 3.1: Generated two-day EMA sampling plan, for 5 participants

3.5 Power Analysis

The power of a statistical test is the probability that it will detect an effect when this effect, in reality, exists. It is a function of the strength of the effect size, sample size, the significance level (α), and the statistical model. Determining the power of the experiment is an important step in the design of any study - EMA studies included. Both underpowered and overpowered studies are a waste of time and resources.

Conducting a power analysis can be easy or very difficult, depending on the complexity of the experimental design and the adopted statistical technique. For simple tests, such as the t-test and ANOVA, straightforward analytical solutions exist, which are implemented in readily available tools. In R, one of those tools is package ‘pwr’.

For example, to use ‘pwr’ to calculate the power of a t.test to detect a moderate effect size ($d = 0.5$), with $n = 30$ per group, and a (two-sided) significance level α of .05, type:

```
# Code snippet 3.2: Power analysis of a t-test
# (analytical approach).
```

```
library(pwr)
pwr.t.test(d = 0.5,
            n = 30,
            sig.level = 0.05,
            type = "two.sample",
            alternative = "two.sided")
#>
#>      Two-sample t test power calculation
#>
#>      n = 30
#>      d = 0.5
#>      sig.level = 0.05
#>      power = 0.4778965
#>      alternative = two.sided
#>
```

```
#> NOTE: n is number in *each* group
```

The power is 48% - not even close to the generally adopted standard of 80%. More participants are needed to detect the hypothesized effect.

EMA study designs are often characterized by repeated measures, complex multi-level structures and the application of advanced statistical techniques. You may find that available power calculators are too limited to properly take key aspects of your design into account. If this happens, simulation techniques may help. If power is the probability that a test will detect an effect it exists, it can be determined by noting the proportion of times a statistical test reaches significance, if it is run, many times, on simulated data, in which the hypothesized effect is present. To illustrate how this works, we will calculate the power of the t-test again, through simulation:

```
# Code snippet 3.3: Power analysis of a t-test
# (simulation approach).

m1 = 0    # mean in group 1
m2 = 0.5 # mean in group 2
sd = 1    # sd (in both groups)
n = 30    # sample size, per group

# conduct the experiment many times
nsim <- 10000
p <- numeric(nsim)
for (i in 1:nsim) {

  data <- data.frame(
    outcome <- c(
      rnorm(n, m1, sd), # group 1 data
      rnorm(n, m2, sd)  # group 2 data
    ),
    group <- c(
      rep(1, n), # group 1 indicator
      rep(2, n)) # group 2 indicator
  )

  # save significance of test
  p[i] <- t.test(outcome ~ group, data)$p.value
}

# power
sum(p < 0.05) / nsim
#> [1] 0.4789
```

As can be seen, the simulation results are very close to the output of ‘pwr.t.test’.

There was no immediate need to run this simulation. We already knew that the power was 48%. The example illustrates, however, that simulation is a valid option when power calculators are too limited (or too difficult to understand...). Simulating the right data, of course, can be challenging as well, but you will find that R has packages that simplify data simulation. For example, ‘mvrnorm’ in package MASS (Venables and Ripley, 2002) can be used to generate correlated data, and package ‘simstudy’ (Goldfeld, 2018) can be used to generate complex (longitudinal) data.

3.6 Ethical considerations

All clinical studies that involve human participants need to be evaluated by a Medical Research and Ethics Committee (MERC; Dutch: ‘METC’). Recently, the committees have also been tasked to determine whether a medical device is used and to evaluate the safety and quality of the device. Researchers are therefore required to add a section in the research protocol, explaining why the software/device is or is not a medical device. The official definition of a medical device (Medical Device Act, or ‘Wet Medische Hulpmiddelen’) is as follows:

“Any instrument, apparatus or appliance, any software or material or any other article that is used alone or in combination, including any accessory and the software required for its proper operation, that is intended by the manufacturer to be used specifically for diagnostic or therapeutic purposes, and is intended by the manufacturer to be used for human beings for the purpose of:
- diagnosis, prevention, monitoring, treatment or alleviation of disease - diagnosis, monitoring, treatment, alleviation of or compensation for an injury or handicap - investigation, replacement or modification of the anatomy or of a physiological process - control of conception, and which does not achieve its principal intended action in or on the human body by pharmacological, immunological or metabolic means, but which may be assisted in its function by such means.”
(CCMO, 2018)

In short, software can be classified as a medical device when it collects patient-specific data and when it is specifically intended for one of the above-mentioned objectives. In practice, the definition of medical devices leaves a lot of room for confusion. Researchers often struggle with the question whether their assessment tools should be considered an medical device or not. For this purpose, flowcharts exist that help to determine whether an app or product should be classified as a medical device (see, e.g., Ekker and van Rest, 2013, and <http://cetool.nl/general/scanAid>). Figure 3.2 shows such a flow-chart.




Figure 3.2: Flow-chart to determine whether study software should be considered a medical device.

While planning your EMA study, you should also be mindful of the rules and regulations that apply to data collection, storage and sharing. From May 2018 onward, the European Committee has enforced the General Data Protection Regulation (GDPR, see <https://gdpr-info.eu/>; in Dutch, Algemene Verordening Gegevensbescherming - AVG), which protects the data and privacy of EU citizens. Complying to the GDPR can be a complex and time-consuming process, depending on nature of your study. Do not hesitate to consult local experts or standard guidelines provided by your organisation (see, e.g., <http://www.emgo.nl/kc/checklist-avg/>).

In EMA studies, an important GDPR-related issue concerns the Data Processing Agreement (DPA). When data processing is (partly) outsourced to a third party, a Data Processing Agreement (DPA) should be drafted, that specifies the agreements between the ‘controller’ and ‘processor’. In this context, a controller is the person or organisation that determines the why and how of data collection (for example you as a researcher). The processor processes the data on behalf of the controller, for example by providing a data-collection service, or by storing raw study data on a cloud-server.

Aspects of data processing that need to be addressed in this agreement include the context, duration and termination of agreement, the nature of the data collected, the duration of data storage, security measures taken to prevent unauthorized access, and agreements on data ownership, data sharing, the handling of personal data breaches, and liability. Most organisations offer model agreements in which all relevant issues are addressed. Third parties may offer model agreements as well. If so, however, these agreements need to be checked for compliance to local regulations.

Chapter 4

EMA instruments

Collection of EMA data has come a long way since the written diary. With technological advances, current EMA data are more likely to be collected through smart or wearable devices, connected to the internet. The various methods of EMA are developed often from distinct assessment targets (Shiffman et al., 2008). For example, a smart device will be useful for collecting data on subjective states or behaviours, whereas wearable devices are more suited to collect physiological data. Below are points to consider when deciding on a suitable EMA instrument, gleaned from the past experiences of APH researchers.

4.1 General considerations

4.1.1 Costs new development vs known player

When considering a platform for collecting EMA data, do you choose a platform developed by a known player or consider creating one of your own? Using a currently available platform has its advantages but it might not have the flexibility to meet your research needs. For example, the EMA application might not be adapted to include items specific to your research. If deciding on the creating a new platform, keep in mind of the costs required in terms of time and financial.

4.1.2 Low-tech versus high-tech

Technology has greatly increased the methods of EMA data collection. Nevertheless, it is worth considering whether your study needs to have an advanced methods of data collection. Maybe the humble written diary is sufficient for your study needs or that the you might still go ahead with the modern, more technologically advanced methods.

4.1.3 Access to a back-office

An important consideration when deciding on an EMA platform is the autonomy available to the researcher to define and decide on research items. In other words, is there access to the back office? For example, some EMA platforms do allow configuration or customization to the needs of the researcher. Also some back office offer advisory services to researchers when adapting items. For example, using EMA to collect depression data does not mean the wholesale transportation of a validated paper questionnaire onto the EMA platform. A subjective EMA survey is not similar to an online questionnaire.

4.1.4 Validation and reliability

Questions can also arise regarding the validity of EMA. Items used in subjective EMA have often not undergone rigorous psychometric testing, unlike traditional paper questionnaires. Also, EMA software are developed using algorithms which are not known or unavailable to the researcher to test for validity.

4.1.5 Data access

If planning on using a commercial EMA platform, it is important to clarify whether the researcher has access to the raw data or is the data aggregated before provision to the researcher (e.g. GENEActiv versus Fitbit). If the researcher has access to the raw data, how easily accessible is the data? Could the data be downloaded directly from the server of the EMA platform?

4.1.6 Demands on time

Subjective EMA could be burdensome to participants as data is collected multiple times a day for an extended period. In order to ensure continued participation, it is important that researchers have in place procedures and resources to manage study participants. Besides preparing a study protocol, it is also necessary to have a manual on how to operate the EMA device. Considerable time also needs to be invested for the training of participants; how EMA and the device works. The researcher might also have to factor in the time needed to contact the participant when data collection starts (e.g. the following day after start of study) to address possible problems, and after data collection (e.g. a debriefing).

4.2 Technical considerations

4.2.1 Internet access

Is access to internet or wifi an integral part of the collection process? For example, the EMA platform does not offer an app but sends an alert with a link to complete an online survey. This means that the participant needs to have online access following the alert. Otherwise, this will lead to a missed data collection. Also, if the server of the platform is down, the participant will not be able to access the online survey following an alert.

If an app is available, would the data collected be stored locally on a device before being uploaded to the server when in the vicinity of internet access or wifi?

4.2.2 Device limitations

Ensure there is adequate local storage on device, in the event that uploading of data to a central server might be hindered. EMA studies can generate a tremendous amount of data which needs adequate storage, be it in cloud or locally on a central server. If using cloud storage or from a commercial entity, This raises the following questions the ownership of the server and the data, and also the location of the server.

As EMA data could be collected using apps or in modern smartphone (e.g. GPS tracking), a consideration would be whether to use participants' own device or to provide a device for the study. If providing a dedicated study device, there are upfront costs of purchasing the devices and their accessories (e.g. battery chargers). Also, the costs of sending/returning of devices and potential loss of device needs to be factored in.

These financial factors are a non-issue if participants use their own devices. Instead there are other potential problems, namely that of compatibility of the smart devices. Currently some apps are available either only on IoS or Android. Besides considering the make of the participants device, one also needs to take into account the model. Older models might not have the feature needed for the EMA collection. Another potential issue associated with using own devices is adequate memory/storage capacity, i.e. to install the EMA app and collect the data.

Regardless of using own or a dedicated device, keep in mind that apps do have software upgrades which researchers need to be aware of and inform participants (especially if participant is using own device).

4.3 Participant perspective

Participating in an EMA study could impact negatively on the participant's in his/her social environment. If using a dedicated device, participant could have to deal with questions of the need to carry the device. This

might create feelings of stigma if the study was on a socially sensitive topic e.g.) psychiatric well-being or suicide ideation.

Or the design of the device might not be compatible with the lifestyle of the participant, e.g. a fashion-conscious female participant might be less than enamoured with the chunky design of a wearable activity sensor. Furthermore, the participants have to remember to have the device on hand during the period of data collection, sometimes even during sleep (e.g. actigraphy)

Also, in a subjective EMA study design, participants have to attend to multiple alerts. These alerts might occur at an inconvenient social moment.

4.4 Points to consider

- Internet access
- Whether a company has an app or is notification for completion of survey via an email or SMS.
- Use of participants' own device (fitness device, tablet, smartphone) or designated device – Issues of software updates, compatibility issues if using participants device, issues of storage capacity of device if data has to be stored locally (because of lack of internet access) before it can be uploaded to central system. -Issue of "stigma" of participating in an EMA study e.g. need to explain the carrying/wearing of a device that might not complement participants' look/lifestyle.

In a nutshell, before implementing an EMA study, it is advisable to do a pilot study using your own device. This will allow the researcher to experience the possible inconveniences faced by participants. And, make sure there is a Plan B!

Chapter 5

Data management

Modern EMA research generates a lot of data. Repeated self-reports, GPS-data, accelerometer data, background demographic data and traditional questionnaire data quickly add up to hundreds of megabytes of raw data. This raises the importance of proper data management. Without a proper data management plan, the EMA researcher quickly drowns in the data feed.

5.1 Using RStudio-projects to manage EMA data

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

“File | New Project...”

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

5.2 Project directory structure

```
1 project
2   |--data
3   |   |--source
4   |   |   |--keyfile.csv
5   |   |   |--GPS
6   |   |   |   |--subject1.json
7   |   |   |   |--subject2.json
8   |   |   |--accelerometer
```

```

9   |   |   |   |--subject1.bin
10  |   |   |   |--subject2.bin
11  |   |--surveys
12  |       |--demographics.sav
13  |       |--survey.sav
14  |--pruned
15      |--GPS.Rda
16      |--accelerometer.Rda
17      |--surveys.Rda
18 |--scripts
19     |--import
20     |   |--import_GPS.R
21     |   |--import_accelerometers.R
22     |   |--import_surveys.R
23     |--analysis
24     |   |--explore.R
25     |   |--test.R
26 |--output
27     |--images
28     |   |--figure_1.png
29     |   |--figure_2.png
30 |--README

```

[Template] Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

5.3 The keyfile

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Table 5.1: Sample Study Keyfile

| ID | Status | SurveyID | Watch ID | App ID |
|------|---------|--------------|----------|--------|
| P001 | QM01221 | 192.A102.83A | APC009 | |
| P002 | QM01228 | 192.A102.83B | APC010 | |

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus

nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Store personal data in another location. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

5.4 Source data

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

5.5 Import & Prune

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

5.6 Visual Exploration

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

5.7 Analysis

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Part III

EMA Outcomes

Chapter 6

Mood

Mood is a common outcome in EMA research (Myin-Germeys et al., 2016; Desmet et al., 2016). Having respondents rate their mood during the day allows researchers to assess mood fluctuation over time or reactivity to events and daily-stressors (Wenze and Miller, 2010). Often, it is studied in relation to depressive symptoms and mood disorders (aan het Rot et al., 2012). In addition, mood can be linked to other variables, such as substance abuse (Kirchner and Shiffman, 2013; Serre et al., 2015), somatic health (Engel et al., 2016; Moore et al., 2016) or activity patterns (Dunton, 2017; Marszalek et al., 2014).

The definition of mood varies across studies. Usually the concept refers to a general affective state. Following this line of reasoning, a distinction can be made between mood states (e.g. irritable, cheerful, relaxed, etc.) and discrete emotions (e.g. happy, sad, anxious, etc.), where moods are thought to be less specific and more subjective, enduring and related to context (Beedie et al., 2005; Cranford et al., 2006; Desmet et al., 2016).

Depending on the study focus and research questions, mood measurement can be operationalized in several ways. Therefore, it is vital to consider the goal of measuring mood in your own study and to choose an operationalisation that matches your hypothesis and theoretical framework. In this chapter, we will discuss the most commonly used options: 1) unidimensional mood assessment, 2) the ‘bag of items’ approach, and 3) dimensional models, namely the Circumplex model and Negative and Positive affect (NA/PA).

6.1 Unidimensional mood assessment

Perhaps the most seemingly straight-forward method to measure mood is to ask ‘face-valid’ unidimensional questions such as “How is your mood right now” (van Ballegooijen et al., 2016) or “How are you feeling right now” (van de Ven et al., 2017). Respondents usually rate these questions on a Visual Analogue Scale (VAS), aimed to indicate mood intensity. Typically, VAS scales will range from zero (low or worst mood) to 10 or 100 (good or best mood). Keep in mind that the middle of a VAS scale (e.g. 5 or 50) is generally considered a negative result, and only scores above 6 or 60 are considered acceptable or positive mood states (Groot, 2010). In order to address this issue, some researchers have proposed to use VAS-scales ranging from -1 to 1, with 0 as a neutral centre. However, such a scale implies a mood state that ranges from negative to positive, rather than absent to present. Another alternative is to use Likert scales, where the scale centre often reflects a neutral response.

Plotting data from an unidimensional item in a graph is an easy way to visually inspect within-subject change in general mood:

```
# Code snippet 6.1: plotting data over time.  
library(ggplot2)  
library(emaph)  
  
plotmood_down <- ggplot(csd, aes(x = date, y = as.numeric(mood_down))) +
```

```
geom_smooth(method = "loess", span = .05, se = FALSE, colour="dodgerblue4") +
  geom_point(size = .3, alpha = .3, position = position_jitter(height = .1),
  colour="dodgerblue2") + scale_x_date() + scale_y_continuous(breaks = 1:7) +
  xlab("Time") + ylab("Mood")
print(plotmood_down)
```




Figure 6.1: 34 weeks of mood data, from a single participant

6.1.1 Bipolar unidimensional items

Another option to assist respondents with the interpretation of one-item mood ratings, is to use a bipolar scale. These items place two opposing mood states at each end of the scale, for example by asking “Please rate your current mood on a scale of 0 to 100, on which 0 indicates happy, and 100 indicates sad” (van Rijsbergen et al., 2014). This does assume that the opposing mood states, such as happy and sad, are mutually exclusive and thus cannot occur simultaneously. The bipolar-unidimensional method was shown to be able to predict time to relapse over 5.5 years in recurrently depressed out-patients, with 6.3% of variance in time to relapse explained. This percentage was comparable to that of the HAM-D (Rijsbergen et al., 2012). Also, the scale was able to detect relapse in patients with recurrent Major Depressive Disorder (based on SCID-I interview) at a cut-off score of 55, and outperformed the HAM-D and IDS-SR. However, 47% of patients indicated by the VAS scale did not fulfil formal criteria for relapse (false positives) (van Rijsbergen et al., 2014).

6.2 Bag-of-Items

In order to make sure all constructs of interest are measured, you can also consider including a number of specific mood items in your EMA questionnaire, rather than one general unipolar item or one bipolar item. For example you can ask respondents “How depressed are you feeling right now” and “How anxious are you feeling right now (Starr and Davila, 2012). This strategy often leads to a ‘bag-of-items’ approach, where single items from various sources, such as existing questionnaires, are combined into a new EMA questionnaire. A benefit of this approach is that you can select items for which information on validity and test-retest reliability is available. A downside is that item scores can only be evaluated separately, rather than providing one overall indication of mood or well-being.

Combining data from multiple items, such as mood and loneliness, in one graph can provide respondents

with insight in the interaction between the constructs. In R two variables can easily be plotted together:

```
# Code snippet 6.2: Plotting multiple variables in one graph.
library(ggplot2)
library(emaph)

combined <- plotmood_down +
  geom_point(data=csd, aes(date, as.numeric(mood_lonely)),
             size = .3, alpha = .3, position = position_jitter(height = .1),
             colour="indianred4") +
  geom_smooth(data=csd, aes(date, as.numeric(mood_lonely)),
              method = "loess", span = .05, se = FALSE, colour="indianred2")
print(combined)
```




Figure 6.2: 34 weeks of combined mood data, from a single participant

6.3 Multi-dimensional mood assessment

Dimensional models assume that every affective state or emotion should be described by the combined score on (at least) two items, instead of just one. Over the past decades several multi-dimensional models have been specified (for an overview, see Sander and Scherer (2009)). In the context of EMA, researchers most often base their items on the Circumplex model (Russell, 1980) or the Negative/Positive affect (NA/PA) theory (Watson and Tellegen, 1985).

6.3.1 The Circumplex Model

The Circumplex Model of affect (Russell, 1980; Posner et al., 2005) argues that all mood states are a linear combination of two independent, bipolar scales: valence (ranging from unpleasant to pleasant) and arousal/activation (ranging from low to high arousal). Combining scores on these scales places the affective states in a circle on one of four quadrants (6.3). States within one quadrant are believed to be positively correlated, while states in the opposing quadrant are thought to be negatively correlated.




Figure 6.3: Russell's Circumplex model of affect.

There are several options to operationalise the Circumplex model in EMA research. For example, respondents can rate valence and arousal on two VAS scales. The most pragmatic approach is to report both scale scores separately (Asselbergs et al., 2016). Alternatively, scores can be combined to give insight into which of the four mood states (quadrants) respondents fall:

- Low aroused - unpleasant
- Low aroused - pleasant
- High aroused - unpleasant
- High aroused - pleasant

A downside of the Circumplex model is that the concepts of valence and arousal can be hard to convey to respondents, especially when translated to other languages, such as Dutch. One alternative is to adjust the scale ends, for example using “extreme sleepiness” and “extreme high energy” (Sharar et al., 2016). Another option is to use pictures or emoticons, rather than language. For this purpose, DeSmet and colleagues developed the pick-a-mood scale, which is a pictorial self-report scale (Desmet et al., 2016). The scale builds on the circumplex model and adds the third dimension “dominance” (level of experienced control over the mood state), rendering eight (instead of four) different mood states and one neutral state (6.4).




Figure 6.4: The Pick-A-Mood Circle.

6.3.2 Negative & Positive Affect

Watson and Tellegen (Watson and Tellegen, 1985) also specified the underlying theory of the Circumplex model, arguing that the diagonal quadrants represent Positive and Negative affect (PA/NA) and that these two terms are the main dimensions of affect (Watson and Clark, 1994). PA ranges from sadness to high energy, NA from calmness to distress (Watson et al., 1988). While bipolar-unidimensional assessment assumes that positive and negative affect are mutually exclusive, the PA/NA affect model assumes that these mood states can occur simultaneously. Watson and Clark (Watson and Clark, 1997) for example, showed a moderate correlation between the two constructs (.32). In order to measure Positive and Negative Affect, a designated Positive and Negative Affect Schedule was developed by Watson, Clark and Tellegen (Watson et al., 1988). Respondents are asked to indicate “to what extent you feel this way right now” on 20 affect items. The scale uses a 5-point Likert-scale, ranging from 1 (very slightly or not at all) to 5 (very) (Watson and Clark, 1994). Items include:

- Negative Affect (10): afraid, scared, nervous, jittery, irritable, hostile, guilty, ashamed, upset, distressed
- Positive Affect (10): active, alert, attentive, determined, enthusiastic, excited, inspired, interested, proud, strong

There are several short-forms available. For example, Wichers and colleagues created a 10-item short-form of the PANAS for their EMA studies. The items were based on the PANAS and their own experience with EMA (Wichers et al., 2012). Using factor analyses, the following 7-point Likert items were chosen for the questionnaire:

- Negative affect (6): insecure, lonely, anxious, low, guilty, suspicious.
- Positive affect (4): cheerful, content, energetic, enthusiastic.

PA and NA were calculated as the average score across all items and weighted for their factor loadings (Wichers et al., 2012). In R, such a factor analysis can be executed as follows:

```
# Code snippet 6.3: Performing a factor analysis.
```

```
library(dplyr)
library(tibble)
```

```
items <- csd %>%
```

```

select(c(
  "mood_enthus", "mood_cheerf",
  "mood_strong", "mood_satisfi",
  "mood_lonely", "mood_anxious",
  "mood_guilty")) %>%
scale(.) %>%
as.tibble(.) %>%
mutate_all(funs(residuals(stats::arima(., order = c(1,0,0)))))

correlations <- cor(items, use = "complete.obs")
fa = psych::fa(items,
               nfactors = 2,
               rotate = "oblimin",
               fm = "pa",
               scores = "regression")

psych::fa.diagram(fa,
                  simple = TRUE,
                  main = "")

```




Figure 6.5: Factor analysis of 7 EMA items, revealing two factors: Positive Affect (PA) and Negative Affect (NA).

```

# Code snippet 6.3: Performing a factor analysis.
library(dplyr)
library(tibble)

items2 <- csd %>%
  select(c(
    "mood_relaxed", "mood_satisfi",
    "mood_enthus", "mood_cheerf",
    "mood_strong", "mood_down",
    "mood_lonely", "mood_anxious",

```


```
"mood_guilty" )) %>%
scale(.) %>%
as.tibble(.) %>%
mutate_all(funs(residuals(stats::arima(., order = c(1,0,0)))))

require(psych)

correlations <- cor(items2, use = "complete.obs")
model <- fa(items2,
            nfactors = 2,
            rotate = "oblimin",
            fm = "pa",
            scores = "regression")

psych::fa.diagram(model,
                  simple = TRUE,
                  main = "")

colnames(model$loadings) <- c("PA", "NA")
fa.diagram(model)
```



Factor Analysis




Figure 6.6: Factor analysis of 7 EMA items, revealing two factors: Positive Affect (PA) and Negative Affect (NA).

Chapter 7

Activity

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra (Marszalek et al., 2014).

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

7.1 Accelerometry

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque.

vertical (Y), horizontal right-left (X) and horizontal front-back axis (Z).

Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

7.1.1 Data collection

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam. Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh.




Figure 7.1: One hour of raw data collected with a wrist-worn GENEActive accelerometer, sub-sampled to 10-seconds epochs (0.1 Hz)

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta.

$$1 \text{ g} = 9.81 \text{ m/s}^2$$

Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

7.1.2 Data cleaning

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam (van Hees et al., 2014, van Hees et al. (2015)).

7.1.3 Feature extraction

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt

ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

signal vector magnitude (svm):

$$\sqrt{x^2 + y^2 + z^2} \quad (7.1)$$

Euclidian Norm minus one (ENMO):

$$\sqrt{x^2 + y^2 + z^2} - g \quad (7.2)$$




Figure 7.2: One day of data of the two persons in the genea data set, summarised with ENMO, calculated from 15-minute epoch windows

7.1.4 Analysis

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

7.2 Location analysis

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

7.2.1 Global Positioning Systems (GPS)

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam. Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh.

```
# Code snippet 7.3: four-week location history.
library(emaph)
library(ggplot2)

d <- subset(locations,
            accuracy <= 100 &
            lon >= 4.80 & lon <= 5.00 &
            lat >= 52.25 & lat <= 52.50)

ggplot(d, aes(lon, lat)) +
  geom_point(alpha = .1, shape = 21, size = 3) +
  facet_wrap(~ id)
```




Figure 7.3: Four-week Location history of two people, collected with Google Timeline

7.2.2 Features

Table 7.1: Features of a GPS data set (Saeb et al., 2015).

| Name | Formula | Description |
|----------------|--|----------------------------------|
| Total Distance | $\sum (distance((lat_t, lon_t), (lat_{t-1}, lon_{t-1}))$ | Total distance between locations |

| Name | Formula | Description |
|--------------------|-----------------------------------|---|
| Location variance | $\sigma_{lon}^2 + \sigma_{lat}^2$ | Variability of location visits |
| N Places | k-means(loc, lat) | Location clusters |
| Location Entropy | $-\sum_i p_i * \log(p_i)$ | Variability in location cluster visits |
| Normalized Entropy | $\sigma_{lon}^2 + \sigma_{lat}^2$ | Entropy, controlled for number of clusters |
| Home Stay | $Entropy \log(N)$ | Percentage of time spent at home |
| Circadian Movement | lomb_scargle(lat / lon) | Consistency of location schedule based on a 24-hour period. |

Part IV

Analytic Approaches

Chapter 8

Feature Extraction and Selection

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

8.1 Feature extraction

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.




Figure 8.1: features of a series.

8.1.1 Variance

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

8.1.2 Autocorrelation

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

8.1.3 Rolling statistics

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

8.2 Reliability and Validity

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

8.3 Missing values

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra. How to prevent imputation structure to becoming a feature of your signal?

8.4 Feature Selection

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac

dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

Chapter 9

Mixed Modeling

EMA data are timeseries that are characterised by complex correlational structures, irregular sampling intervals, missing data, and substantive individual differences. Mixed models are well-suited to deal with these data. This chapter provides a brief introduction to conducting mixed models analysis of EMA data in R.

9.1 The Mixed Model

Mixed modeling can be understood as a regression technique in which separate regression functions are estimated for each cluster in the data set. In EMA data, these clusters are defined by the participants. Data from the same participant are expected to be correlated, and one way to honour this correlation is to conceptualise a separate regression for each participant. This idea, in the most simple regression model, can be expressed as:

$$Y_{ij} = \text{intercept}_i + \epsilon_{ij} \quad (9.1)$$

This models the expected value of the j-th measurement of participant i as the mean of all measurements of participant i, plus error. It defines a set of regression functions - one for each participant.

The regression functions are, however, not independent. Mixed models divide the intercepts of the individual participant regression functions into two components: 1) the intercept of the group ($\text{intercept}_{\{g\}}$; the mean intercept of all regression functions), and 2) a participant-specific component $\text{intercept}_{\{p\}}$ (i.e., the difference between the intercept of the participant and the mean intercept), i.e.:

$$\text{intercept}_i = \text{intercept}_g + \text{intercept}_p \quad (9.2)$$

The group intercept is called the ‘fixed’ effect. If we would gather more data from new participants, we would expect to find approximately the same group intercept.

The participant-specific component of the intercept is known as the ‘random’ effect. If we sample new data, we would expect a similar *variance* of the participant-specific intercept components around the group intercept. This “mixing” of fixed and random effects is what gives mixed modeling its name.

9.2 Simulating example data

To understand analysis techniques, it often helps to apply the technique to simulated data, in which parameters of interest are known.

Here, we will use the ‘sim_ema’ function from package ‘emaph’, to simulate EMA mood assessments of 100 participants, who rate their mood, three times per day, for one week. We set the mean mood (intercept_g) to 5, the variance around this mean - var(intercept_i) - to .5., and the average variance around these means within participants - the error - to 1.

As you can learn from the documentation of ‘sim_ema’ (see ‘?sim_ema’), the function expects at least two arguments: the definition of a sample plan (see ‘?sample_plan’), and a specification of the data-generating model, in the form of a list defining fixed effects, the random effects, and residual variance (i.e, the error). From these specifications, a data set is simulated (which is assigned to variable d1).

```
# Code snippet 9.1: Simulating ema data.
library(emaph)
plan <- sample_plan(n_participants = 100,
                     n_days = 7,
                     times = c("10:00-11:00",
                               "13:00-14:00",
                               "16:00-18:00"))

d1 <- sim_ema(plan,
               mm_par = list(fixed = c(intercept = 5),
                             random = c(intercept = 1),
                             error = .5),
               lim = c(0, 10))
```

Figure 9.1 shows EMA mood ratings of the first 6 participants in the simulated data set. Mean mood ratings of the participants (the red lines) vary around 5 (the grey dashed line), as specified.




Figure 9.1: Simulated EMA data of Six Participants.

9.3 Fitting a mixed model in R

Now, let’s fit a mixed model to the data, to see whether the simulation parameters are detected. For this, we will use the ‘lme’ function, from package ‘nlme’ (Pinheiro et al., 2018).

The first argument of the lme function , ‘Y ~ 1’, specifies the fixed ‘effect’ (in this case: the mean intercept). The second argument, ‘random = ~ 1 | id’ specifies the random effect: in this model, intercepts are allowed to vary between participants. The fitted model is assigned to variable fm.

```
# Code snippet 9.2: Fitting a mixed model with lme.
library(nlme)
fm <- lme(Y ~ 1, random = ~ 1 | id,
            data = d1)
```

We can now extract the fixed effects regression coefficients table, by calling the ‘summary’ function on the fitted model. The estimated intercept should be around to 5 (as this is a finite sample, we expect some deviation):

```
# Code snippet 9.3: Print fixed effects.
summary(fm)$tTable
#>           Value Std.Error   DF t-value p-value
#> (Intercept) 5.21     0.107 2000    48.8      0
```

Random effects and residual variance are shown by the ‘VarCorr’ function. Again, since we specified the data ourselves in this case, we know the ‘true’ value of these parameters: the random intercept variance should be around .5 and the residual error variance should be close to .2.

```
# Code snippet 9.4: Fitted random effects.
VarCorr(fm)
#> id = pdLogChol(1)
#>          Variance StdDev
#> (Intercept) 1.112    1.055
#> Residual    0.502    0.709
```

It can be instructive to plot the predicted values of the model, to make clear how the model ‘thinks’. As shown by Figure 9.2, the model predicts a series of straight lines, one for each participant, that vary around 5.

```
# Code snippet 9.5: Saving predicted values.
d1$predY <- predict(fm)
```




Figure 9.2: EMA ratings, of each participant in the simulated data set, as predicted by the intercept-only mixed linear model.

9.4 Adding time as a predictor

Now that we know how to fit a simple mixed model, we can consider a more complicated scenario. In the first data set, participants’ mood ratings did not change over time. Scores varied around a stable mean during the full week. Hence, there was no need to model a time effect. But suppose we would expect a systematic improvement of mood ratings over time, for instance in response to a mental health intervention?

Let’s first call ‘sim_ema’ again, with parameters that will result in data in which mood rating increase over the course of the week, 0.5 scale points per day. Let’s also assume that individual participants will vary in

the degree of mood improvement: the mean time effect will be 0.5, but this parameter is allowed to vary between participants, with a variance of 0.05.

```
# Code snippet 9.6: Simulating ema data (time effect).
d2 <- sim_ema(plan,
               mm_par = list(fixed = c(intercept = 5, time = 0.5),
                             random = c(intercept = 1, time = 0.1),
                             error = .5),
               lim = c(0, 10))
```

Figure 9.3 shows the data of the first six participants in the second data set. Both the intercept and the slope vary across the participants. Some participants improve more over time, and others improve less: the slope in this data set is a random effect.




Figure 9.3: Simulated EMA data of Six Participants (Time-varying model).

To fit the extended mixed model, time can simply be added to both the fixed and random arguments of the ‘lme’ function. Fixed effects estimated of this model should be around 5 and 0.5, since that is how we specified the data. Calling ‘summary’ on this function, we see that the fixed time effect is significant.

```
# Code snippet 9.7: A mixed model, with a random slope.
library(nlme)
fm <- lme(Y ~ 1 + time, random = ~ 1 + time | id,
            data = d2)
summary(fm)$tTable
#>           Value Std.Error   DF t-value p-value
#> (Intercept) 5.181    0.1238 1999    41.8 1.98e-275
#> time         0.445    0.0288 1999    15.5  4.33e-51
```


The random effects now has four components: the variance of the intercept, the variance of the slope, the residual error *and* the correlation between the random intercept and the random slope.

```
# Code snippet 9.8: Extracting random effects.
VarCorr(fm)
#> id = pdLogChol(1 + time)
#>          Variance StdDev Corr
#> (Intercept) 1.4390   1.200 (Intr)
#> time        0.0774   0.278 -0.123
#> Residual    0.4709   0.686
```

Model predictions clearly show how the mixed model estimated varying intercepts and slopes, that, on average, reflect the fixed effect regression formula ‘ $Y \sim 5 + 0.5 * \text{time}$ ’.

```
d2$predY <- predict(fm)

ggplot(d2, aes(x = time, y = predY, group = id)) +
  geom_line(alpha = .1, size = .6) +
  geom_smooth(aes(group = NULL), method = "lm", color = "red") +
  coord_cartesian(ylim = c(0, 10)) + theme_classic()
```



9.5 Adding a Two-Group Comparison

In data-set 1, mood ratings did not change during the week, while in data-set 2, the mood ratings increased. Suppose the two data-sets reflect the data that you collect in a two-group RCT, in which you compare the effects of a mental health intervention (data-set 2) against a waiting list condition (data-set 1). By combining the two data-sets, we can illustrate how to conduct a group comparison with ‘lme’.

Since the two data-sets are already available (in d1 and d2), the new data set can be created with just three lines of code (below). In the first line, the ‘rbind’ function is used to combine the rows of data-set 1 and 2 into a new variable: data.frame d3. The second line adds a group indicator to d3. The third line updates the id’s of the participants in the second group, to explicitly differentiate the participants in the second group from the participants in the first group.

```
# Code snippet 9.9: two-group simulation.
d3 <- rbind(d1, d2)
d3$group <- factor(c(rep(0, nrow(d1)), rep(1, nrow(d2))))
d3$id[d3$group == 1] <- d3$id[d3$group == 1] + 100
```

The effect of the intervention can be tested by adding a (fixed) ‘time * group’ interaction effect to the model. This effect, we know, is 0.5, and, as can be seen, this is what the model picks up:

```
# Code snippet 9.10: A mixed model, with two groups.
library(nlme)
fm <- lme(Y ~ 1 + time * group, random = ~ 1 + time | id,
           data = d3)
round(summary(fm)$tTable, 2)
#>          Value Std.Error DF t-value p-value
#> (Intercept) 5.24      0.12 3998   44.89    0.00
```

```
#> time      -0.01    0.02 3998   -0.51    0.61
#> group1    -0.06    0.17 198    -0.38    0.70
#> time:group1 0.46    0.03 3998   15.39    0.00
```

In Figure 9.4 below, EMA mood ratings predicted by the fitted model clearly show how the model detects 1) the fixed between-group effect, and 2) the variance in intercepts and slopes in both groups.




Figure 9.4: Predicted mood ratings

9.6 Further reading

In this chapter, we introduced mixed model analysis of EMA data. To do so, we could only touch upon the theoretical foundations of mixed models, and we deliberately used simple examples with clean simulated data. Readers, who consider the application of mixed models, are strongly advised to study additional resources.

The authoritative reference for mixed effect modeling in R is a book by Pinheiro and Bates (2000). To fully appreciate this book, however, a strong background in mathematical statistics is required. Gentle introductions in the topic can be found in XXX, YYY and ZZZ ().

Chapter 10

Fitting Networks

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

10.1 What are Networks?

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta (Borsboom and Cramer, 2013; Borsboom, 2016). Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, em tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra (Epskamp et al., 2012, Epskamp et al. (2018a)).

10.2 Fitting Networks on Single-Subject Repeated Measures Data

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem.

Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui graphicalVAR (Epskamp, 2017).

```
# Code snippet 10.1: Fitting symptom networks.
library(ggplot2)
library(graphicalVAR)
library(dplyr)

# Simulate model:
set.seed(2)
Mod <- randomGVARmodel(5, probKappaEdge = 0.8, probBetaEdge = 0.8)

# Simulate data:
d <- as.data.frame(graphicalVARsim(50, Mod$beta, Mod$kappa))

e <- tidy::gather(d %>% group_by(key) %>% mutate(t = 1:n()))
ggplot(data = e, aes(x = t, y = value)) +
  geom_point(size=.2) + geom_line() +
  facet_grid(key~.)
```




Figure 10.1: Simulated timeseries of 5 variables

10.2.1 Contemporaneous and Directed Correlations

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

```
# Code snippet 10.2: plotting symptom networks
Res <- graphicalVAR(data = d, gamma = 0,
                      nLambda = 10, verbose = FALSE)

# Show networks
plot(Res, title = FALSE)
```




Figure 10.2: Partial Contemporaneous Correlations (left) and Partial Directed Correlations (right).

10.2.2 Node Analysis

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

- Betweenness
- Closeness
- Strength

Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

- InStrength
- Outstrength

```
# Code snippet 10.3: Node Centrality plots.
```

```
library(qgraph)
centralityPlot(qgraph(Res$PCC, DoNotPlot = TRUE))
centralityPlot(qgraph(Res$PDC, DoNotPlot = TRUE))
```




Figure 10.3: Centrality plot of Partial Contemporaneous (top) and Directed (bottom) Correlations.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

10.3 Fitting Networks on Multiple Subjects Repeated Measures Data

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Code snippet 10.4: Fitting symptom networks of multiple subjects.

```
library("mlVAR")
library(methods)

Model <- mlVARsim(nPerson = 30, nNode = 5,
                   nTime = 50,    lag = 1)

fit1 <- mlVAR(Model$data,           vars = Model$vars,
              idvar = Model$idvar, verbose = FALSE,
              lags = 1,                 temporal = "correlated")
```

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

```
# Code snippet 10.5: plotting symptom networks of multiple subjects.
layout(matrix(c(1:6), ncol = 2, byrow = TRUE))

plot(Model, "contemporaneous", layout = "circle", verbose = FALSE)
plot(fit1, "contemporaneous", layout = "circle", verbose = FALSE)

plot(Model, "temporal", layout = "circle", verbose = FALSE)
plot(fit1, "temporal", layout = "circle", verbose = FALSE)

plot(Model, "between", layout = "circle")
plot(fit1, "between", layout = "circle")
```




Figure 10.4: True (left) vs estimated (right) contemporaneous (top), temporal (middle) and between-subject relationships (bottom)

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque null aut massa tempor, ac dapibus dui viverra.

10.4 Discussion

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque null aut massa tempor, ac dapibus dui viverra.

10.4.1 Pitfalls of Network Analysis

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor (Epskamp et al., 2018b; Epskamp and Fried, 2018). Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque null aut massa tempor, ac dapibus dui viverra.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet (van der Krieke et al., 2015). Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque null aut massa tempor, ac dapibus dui viverra.

10.4.2 Subsection

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet at the [<http://psychosystems.org>]. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque null aut massa tempor, ac dapibus dui viverra.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque null aut massa tempor, ac dapibus dui viverra.

Part V

EMA Case Studies

Chapter 11

Early Warning Signs of Depression

One of the promises of EMA is that it might detect signs of mental health deterioration in an early stage. Subtle changes in time series of mood variables, for example, might signal a depression relapse. If we can detect these changes, preventive interventions can be triggered to avoid the relapse.

But what changes, exactly, should we look for? What are these early warning signs?

11.1 Critical Slowing Down

Critical Slowing Down (CSD) is a concept from dynamic systems theory. In dynamic systems, state transitions are preceded by a change in which the system reacts to disturbances. In a stable state, the system quickly recovers from disturbances. Prior to a transition to a new state, however, the system takes more and more time to recover back to its current state: the variance and auto-correlation of the system increases (Scheffer et al., 2009; Dakos et al., 2008).

In this chapter, we re-analyse data from a study that aimed to demonstrate CSD in EMA-data of a single patient with a history of major depression (Groot, 2010, Kossakowski et al. (2017); Wichers et al., 2016). The patient, a 57-year old male, used EMA to monitor himself during a 239-day single-case double-blind medication reduction trial. During the experiment, he experienced a relapse, and Wichers and colleagues showed how variance and auto-correlation in the EMA data increased, several weeks prior to this relapse. The transition appeared to be preceded by CSD.

We will try to reconstruct the finding, using an alternative analysis strategy. One of the limitations of the Wicherts et al analysis is that autocorrelation was analysed at lag 1 only (i.e., only the correlation between t and $t-1$ was considered). With another analysis technique, called ‘Detrended Fluctuation Analysis’, all lags can be considered.

To conduct the analysis, we need three R packages:

- Raw EMA data of this study were published in the public domain (Kossakowski et al., 2017). We included the data in the emaph package.
- To manipulate the raw data and reconstruct the plots of the article, we are going to use several functions from the tidyverse.
- DFA is implemented in package ‘nonlinearTseries’, so we will need that as well.

```
# Code snippet 11.1: required libraries for the CSD-study re-analysis.
```

```
library(emaph)
library(tidyverse)
library(nonlinearTseries)
```

11.2 Plotting the course of depression

Let's take a look at the development of the depressive symptoms of the patient first. These were tapped with weekly assessments of the depression scale of the 'Symptom Checklist-90-Revised' (SCL-90-R; Derogatis, 1994), a well-established self-report questionnaire.

The code below reconstructs Figure 1 of the Wichers et al article. It plots the SCL90-R depression scale score over the time. Around day 120, the depression score increased considerable: the patient experienced a relapse.

```
# Code snippet 11.2: Plot depression score.
dep <- csd %>% select(dayno, scl90r_dep) %>%
  filter(!is.na(scl90r_dep)) %>% unique

# plot dep + change point (day 120 in our data)
ggplot(dep, aes(x = dayno, y = scl90r_dep, group = 1)) +
  geom_step() + scale_colour_manual(values = c("black", "red")) +
  xlab("Days") + ylab("SCL-90-R Depression")
```




Figure 11.1: SCL-90 depression score, over the study period

11.3 Mental state EMA items

Wichers and colleagues selected 13 items from the full EMA dataset, which they grouped in 5 factors: positive affect (pa; 4 items), negative affect (na; 4 items), mental unrest (mu; 3 items), suspiciousness (su; 1 item), and worrying (wo; 1 item). From these factors, an overall mental state sum score can be calculated.

```
# Code snippet 11.3: mood states calculation.

# positive affect
pa_items <- c("mood_enthus", "mood_cheerf",
             "mood_strong", "mood_satisfi")

csd$pa <- csd %>%
  select(pa_items) %>%
  rowMeans(., na.rm = TRUE)
csd$pa <- -csd$pa

# negative affect
na_items <- c("mood_lonely", "mood_anxious",
             "mood_guilty", "mood_doubt")
```

```

csd$na <- csd %>%
  select(na_items) %>%
  rowMeans(., na.rm = TRUE)

# mental unrest
mu_items <- c("mood_irritat", "pat_restl",
              "pat_agitate")
csd$mu <- csd %>%
  select(mu_items) %>%
  rowMeans(., na.rm = TRUE)

# 'single-item' states
csd$su <- csd$mood_suspic
csd$wo <- csd$pat_worry

# global mental state score
csd$ms <- rowSums(csd[c("pa", "na", "mu", "su", "wo")])

```

Rows, in which one or more of the factors have missing values, are removed from the analysis. In a full analysis, options to impute the missing values could and should be considered. However, since only 3 of the 1476 rows have missing item scores, not much is probably lost by running a simple complete-case analysis.

Code snippet 11.4: missing value removal.

```

# count number of items with missing items, per row
csd$nna <- csd %>%
  select(matches("mood_")) %>%
  is.na(.) %>% rowSums

# drop rows with missing values
csd <- csd %>% filter(nna == 0)

```

11.4 Running the DFA

Code snippet 11.5: Running the DFA.

```

# prepare result rows: one row for each day
d <- NULL
d <- csd %>%
  group_by(dayno) %>%
  summarise(ms = mean(ms))

d$ms_dfa = NA

# determine DFA, in a moving window of 32 days,
# in steps of 7 days
window <- 32
for (i in seq(window, max(csd$dayno), 7)) {

  # get the sliding window data
  w <- subset(csd, dayno > (i - window) & dayno <= i)

  # dfa: ms

```

```

dfa.analysis <- dfa(time.series = w$ms,
                      npoints = 30,
                      window.size.range = c(10, nrow(w)),
                      do.plot = FALSE)

fgn.estimation <- estimate(dfa.analysis,
                             do.plot = FALSE,
                             fit.col = "blue", fit.lwd = 2, fit.lty = 2,
                             main = "Fitting DFA to fGn")

d$ms_dfa[d$dayno == i] <- fgn.estimation
}

```

11.5 Results

We can now plot the DFA indicator over time, to see whether it peaks prior to the onset of the relapse. As can be seen, the DFA-indicator clearly peaks prior to the onset of the relapse, around day 110. Interestingly, a second peak is reached at the end of the experiment, around day 239, possibly to indicate a recovery from the relapse.

```

# Code snippet 11.6: Plot DFA results.
ggplot(na.omit(d),
       aes(x = dayno, y = ms_dfa,
            colour = factor(dayno < 120),
            group = 1)) +
  geom_point() +
  geom_line() +
  ylab("dfa (60-day window)") +
  xlim(c(0, 250)) +
  guides(color = FALSE)

```




Figure 11.2: Results of the DFA analysis.

11.6 Discussion

Our re-analysis replicates the main finding of the Wichers et al article ((Wichers et al., 2016)): several weeks prior to a depression relapse, as predicted by complex systems theory, the variance and autocorrelation in EMA mood ratings increased.

One swallow does not make summer. Yes, in this case, the DFA-indicator peaked prior to the relapse. This could be a mere coincidence. The predictive power of CSD needs to be confirmed in larger samples. Given the theoretical background, successful applications of CSD in other domains, and the present finding, these studies seem warranted.

Important additional questions remain to be answered. When it predicts a state change, what is the expected delay between this prediction and the change? Does a critical DFA-value exist? Given that critical value, are false positive and false negative rates of this prediction acceptable? These are important questions that should be answered before any clinical application of DFA can be considered.

Re-analysis of data from completed clinical studies, in which EMA data were collected, may be a first step to further explore the value of CSD. One option, for example, would be to re-analyse data from the E-COMPARED study (Kleiboer et al., 2016). In this trial, patients, who were treated for major depression, completed weekly self-report questionnaires (the Patient Health Questionnaire; PHQ-8, Kroenke et al., 2009) and daily EMA mood ratings throughout treatment, which lasted up to 16-week. Since CSD is an indicator of *any* state change (i.e., irrespective of whether the change is clinically “good” or “bad”), theory would predict a (lagged) relationship between CSD (i.e., the DFA-score) and clinically significant changes in PHQ-scores (Jacobson and Truax, 1991).

Potential clinical applications, of course, are clear. If clinically relevant changes can be predicted algorithmically from EMA ratings, automated patient feedback systems could help to prevent pending deterioration or consolidate the path towards recovery.

Chapter 12

Homerange Estimation

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

12.1 Section

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

12.2 Spatio-temporal analysis:

Package **adehabitatHR** (Calenge, 2006) was created for the study of the habitat of wild animals, using accelerometer and GPS data. Defined procedures can be used for the analysis of human data as well.

```
# code snippet 12.1
library(adehabitatHR)
```

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta.

12.3 Section

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

12.4 Section

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam vehicula augue metus, in tincidunt urna luctus sit amet. Sed ultrices, erat at laoreet semper, sem tellus hendrerit mi, eget pulvinar massa nisl ac dolor. Nunc ac tellus nec tortor interdum porta. Vestibulum hendrerit tempus condimentum. Donec a mollis sem. Aenean lectus nunc, bibendum ut orci vel, tristique pellentesque arcu. Vestibulum id laoreet neque. Phasellus at ex velit. Vestibulum scelerisque nulla ut massa tempor, ac dapibus dui viverra.

Donec sed lectus at sem ultrices commodo. Proin a viverra metus, nec scelerisque odio. Morbi viverra tristique libero vel fringilla. Sed at varius erat, id consequat nibh. Ut eget leo blandit orci posuere tincidunt ac sed erat. Aenean metus metus, eleifend ut facilisis a, fringilla ut neque. Nunc hendrerit cursus eleifend. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Vivamus enim turpis, pulvinar volutpat purus nec, lobortis imperdiet diam.

Part VI

EMA Catalogues

Chapter 13

EMA Research within the APH Mental Health Consortium

This chapter summarises recent EMA research projects within the APH Mental Health consortium, as a guide to other researchers looking for nearby EMA-expertise and research collaboration. An overview of identified projects is presented in Table 13.1. Detailed summaries are presented below.

Table 13.1: Overview of EMA research groups in the APH MH consortium.

| Focus / PI | Project | URL |
|--|--|--|
| Depression | | |
| Bockting | Imagine Your Mood Stay Fine | http://tinyurl.com/ydxeo52 http://stayfine.nl |
| Riper and Smit | ICT4Depression E-COMPARED | http://ict4depression.eu http://www.e-compared.eu |
| Penninx and Lamers | NESDA RADAR-CNS | http://nesda.nl http://radar-cns.org |
| Psychosomatic symptoms | | |
| Knoop Snoek Sprangers | Chronic Fatigue study MERITS (type 1 diabetes) FAntasTIGUE (fatigue) IMPACT (Heart failure) | http://tinyurl.com/y8wuoo http://tinyurl.com/yas8ljbo http://tinyurl.com/ybb7up http://impactonderzoek.nl |
| Psychotic symptoms | | |
| Van der Gaag | TEMSTEM VRETp | http://tinyurl.com/ybac6fcl http://psymate.eu |
| Suicidal ideation | | |
| Van Ballegooijen | CASPAR | http://tinyurl.com/y9e9evg |
| Sleep | | |
| Van Someren | | http://tinyurl.com/y8ht9lq |
| Symptom Networks | | |
| Borsboom | Psycho-systems | http://psychosystems.org |
| Ambulant Monitoring of the autonomic nervous system | | |
| De Geus | VU-AMS | http://vu-ams.nl |

13.1 Depression

13.1.1 Bockting

Prof. dr. Claudi Bockting is affiliated with the AMC psychiatry department and the Rijksuniversiteit Groningen. Her group focuses on treatment and relapse prevention of (severe) depressive disorder and related common mental health disorders. In two studies of the group, EMA measures are included.

| Aspect | Description |
|-----------------------|---|
| Project team | Claudi Bockting, PhD; Yvonne Stikkelbroek, PhD; Gerard van Rijsbergen, PhD |
| Organisation | AMC; Rijksuniversiteit Groningen |
| Project(s) | 1) ‘Imagine Your Mood’ study (IYM; 2012-2017; Slofstra et al. (2017)): This was a three-arm RCT evaluating several relapse prevention strategies in remitted depressed patients. 2) StayFine study (ongoing; http://stayfine.nl/): this RCT aims to evaluate a smartphone-based intervention to prevent relapse in young people (12-23 yrs.) with remitted anxiety or depression. |
| EMA Measures | In the IYM RCT, EMA items included mood, positive affect (PA, 5 items), negative affect (NA, 9 items) and mental imagery (2 items). Participants were prompted 10 times per day, 3 days a week, over a period of 8 weeks. Prompts were sent randomly within fixed time-intervals. The VAS mood scale used ('Please rate your current mood on a scale of 0 to 100', on which 0 indicates 'happy', and 100 indicates 'sad'), was shown to have a high positive predictive value without any false negatives at a cut-off score of 55 (van Rijsbergen et al., 2014). Compared to the HAM-D17 and the IDS-SR, the VAS also was a better predictor of current relapse status, as measured by the SCID-1 interview (variance explained for VAS: 60%; HAM-D17: 49%; IDS-SR: 34%). In the StayFine RCT, EMA will be used to monitor outcomes (for three weeks) at six assessment periods (between baseline and 36-month follow-up). Additionally, Polar watches will be used to assess activity. |
| Platforms used | Psymate, Roqua, ‘Imagine your mood’ app (custom development), Polar watches |
| Contact | https://www.amc.nl/web/research-75/person-1/prof.-dr.-c.l.h.-bockting-phd.htm http://claudibockting.com |

13.1.2 Riper and Smit

Prof. dr. Heleen Riper is affiliated with the department of clinical, neuro- and developmental psychology of the Vrije Universiteit and specialised mental health organisation GGz InGeest, Amsterdam. *Prof. dr. Jan Smit* is affiliated with the department of psychiatry of the Vrije Universiteit Medical Center (VUmc) and GGz InGeest, Amsterdam. Riper and Smit have been driving forces behind the development of MoodBuster, a research platform for the delivery of online and blended psychotherapy, which has built-in, integrated EMA functionality.

| Aspect | Description |
|---------------------|---|
| Project team | Heleen Riper, PhD, Jan Smit, PhD, and the MoodBuster consortium |
| Organisation | Vrije Universiteit Amsterdam; GGZ inGeest (and other organisations) |

| Aspect | Description |
|----------------------|--|
| Project(s) | Moodbuster was developed in the European FP7 project “ICT4Depression” (Warmerdam et al. (2012); http://www.ict4depression.eu/), and applied in the Horizon 2020 FP7 EU-project “European COMPARE Effective research on blended Depression treatment versus treatment-as-usual” (E-COMPARED; Kleiboer et al. (2016); Kemmeren et al. (2016); http://www.e-compared.eu/), MoodBuster will also be used in the EU ImpleMentAll project (http://www.implementall.eu) and in several other clinical trials that are currently (October 2018) in preparation. |
| EMA measures | In the ICT4Depression project, the MoodBuster platform included a complex set of active and passive EMA measures, including physiological sensors, accelerometers, wearables to measure sympathetic nervous system responses (chest strap and glove), and an Android EMA smartphone app. For the E-COMPARED study, the Moodbuster website and smartphone app were adapted to deliver (blended) treatment to patients with MDD, in 5 routine practice settings across Europe. A therapist portal was added, in order to allow therapists to monitor patients’ progress and send feedback messages. Smartphone- based EMA was used to asses sleep, mood, worrying, self-esteem, activities (2 items) and social contacts. Prompts were sent at a random time point between 10:00 and 22:00. At the beginning and during the final phase of treatment, patients received two additional prompts per day for one week. In the morning (around 10:00), sleep, worrying and self-esteem items were assessed. In the evening (around 22:00), these questions were repeated, along with the activity and social interaction items. Patients rated these EMA items on a visual analogue scale, ranging from 0 (low) to 10 (high). |
| Platform used | Moodbuster (http://www.moodbuster.eu) |
| Contact | https://research.vu.nl/en/persons/heleen-riper https://research.vumc.nl/en/persons/jan-smit https://research.vumc.nl/en/persons/lise-kemmeren |

13.1.3 Penninx and Lamers

Prof.dr. Brenda Penninx and Dr. Ir Femke Lamers are affiliated with the Department of Psychiatry, VU University Medical Center and GGZ InGeest. Both are involved in two EMA studies: an EMA study in the context of the Netherlands study of Depression and Anxiety (NESDA; www.nesda.nl), of which Pennix is primary Investigator, and a large international EU H2020-funded collaborative research project, in which a variety of active and passive measures are used (the ‘Remote Assessment of Disease and Relapse - Central Nervous System’-project; RADAR-CNS; <http://www.radar-cns.org>).

13.1.3.1 NESDA EMA study

| Aspect | Description |
|---------------------|---|
| Project team | Femke Lamers, PhD, Brenda Penninx, PhD (VUmc), Hariette Riese, PhD (UMCG), with the NESDA consortium |
| Organisation | VUmc, GGZ inGeest, Amsterdam |
| Project(s) | The main aim of NESDA (the Netherlands study of Depression and Anxiety) is to examine the long-term (eight-year) course and prognosis and co-morbidity of anxiety and depression, to improve quality of care and prevent chronicity (Penninx et al., 2008). Within the Wave 6 9-year follow-up of the NESDA study (2015 - 2017), a selected sample of NESDA participants with symptoms of depression and healthy controls (n = 384) was invited for an EMA diary /actigraphy study. |

| Aspect | Description |
|----------------------|---|
| EMA measures | NESDA EMA study data was collected to explore the dynamic interplay between cognitions, emotions, behaviour and environment in daily life of individuals, the feasibility of EMA in participants with affective disorders, and chronotype and diurnal patterns of activity in depressed versus non-depressed participants. EMA participants started within 31 days after the regular NESDA assessments (face-to-face interview and self-report measures). Participants could use their own smartphone (Android / iPhone) if they had sufficient data or access to WiFi for at least 80% of the two-week time period, and could borrow a phone if needed. To record the amount of physical activity during EMA monitoring, participants were also asked to wear an accelerometer (GENEActiv, 30Hz) on their non-dominant wrist, 24 hours a day, for two weeks. Recording started on the evening prior to the first EMA assessment and continued until the morning after the last assessment. Participants were prompted five times a day, for two weeks, to complete a self-report questionnaire comprising approx. 31 items. Items (7-point Likert scale, ranging from ‘not [at all]’ to ‘very’) tapped 1) valance (unpleasant to pleasant) and arousal (calm to excited), 2) current state, symptoms of depression/anxiety, physical condition, 3) social context, 4) sleep, 5) daily uplifts/hassles, 6) activities, and 7) substance use. Examples of items were: “I feel down”, “I feel cheerful” and “Where are you now” (e.g. at the neighbour’s house). In addition, there was an addendum questionnaire with one item inquiring about questionnaire burden and an open-ended question for general comments on circumstances that might influence answers. The items on valance, arousal and current state have been used in a previous study; the Uncovering the Positive Potential of Emotional Reactivity (UPPER) study (Bennik (2015)). The other items are inspired on earlier EMA studies, such as the work by Mehl and colleagues (Mehl and Conner (2012)), van Os and colleagues (Wichers et al. (2012)) and studies performed at the Interdisciplinary Center of Psychopathology and Emotion regulation (ICPE), such as the Mood and Movement in Daily life (MOOVD) study (Booij (2015)) |
| Platform used | RoQua (https://www.roqua.nl/), GENEActiv accelerometer |
| Contact | https://research.vumc.nl/en/persons/brenda-penninx |
| Information | https://research.vumc.nl/en/persons/femke-lamers https://www.nesda.nl/ |

13.1.3.2 RADAR-CNS

| Aspect | Description |
|---------------------|---|
| Project team | Femke Lamers, PhD; Brenda Penninx, PhD; Sonia Difrancesco, MSc, and the RADAR consortium. |
| Organisation | VUmc, GGZ inGeest |
| Project | The European (EU H2020-IMI) RADAR-CNS project (Remote Assessment of Disease and Relapse - Central Nervous System; http://www.radar-cns.org/) aims to study the potential of wearable devices and smartphone technology to help prevent and treat depression, multiple sclerosis and epilepsy. The project, which started in 2016, was designed to examine how remote measurement technologies can monitor and improve quality of life and psychological well-being of patients. Within this project, the VUmc/GGZ inGeest research site focuses on depression (RADAR-MDD), in collaboration with King’s College, London (KCL). |

| Aspect | Description |
|---------------------|--|
| EMA Measures | Technical goals of the project are to 1) build an end-to-end open source system with generalised data aggregation capabilities, and 2) explore (technological aspects of) big data solutions. Clinical goals of the project are to 1) assess the feasibility of continuous monitoring of patients, and 2) predict disease onset or relapse with big data prevention and risk assessment approaches. |
| Platform(s) | Data are collected during a 2-year period, in which EMA will be activated every 6 weeks for 6 consecutive days. Active EMA outcomes include (variability in) sleep quality, levels of activity, social interactions, mood, cognitive performance and stress as possible predictors of the clinical course of participants. Passive EMA measures include location and movement (GPS; actigraphy), skin temperature, galvanic skin conductance, heart rate (-variability), voice recognition, social interaction data (call/message logs), and smartphone app usage patterns. |
| Contact | In the RADAR project, an comprehensive EMA platform is developed (the RADAR RMT application; see http://thehyve.nl/cases/radar-cns and http://github.com/RADAR-base); This platform focuses on classes of data rather than specific devices, to enhance modularity and adaptability, as new devices become available. The platform will be used to integrate and control data streams from an EMA smartphone app and several wearable device (including the Empatica E4 Wristband, Pebble 2 Smartwatch, Biovotion VSM, Faros 180, and Fitbit). |
| | https://research.vumc.nl/en/persons/femke-lamers https://research.vumc.nl/en/persons/brenda-penninx http://www.radar-cns.org/ |

13.2 Psychosomatic symptoms

13.2.1 Knoop

Prof. dr Hans Knoop is professor of evidence-based psychological and behavioral interventions for medical conditions and somatic symptoms at department of Medical Psychology of the Amsterdam Medical Center (AMC). In 2015/2016, Knoop and colleagues ran an EMA study at the Expert Centre for Chronic Fatigue (ECCF; <http://nckv.nl/>).

| Aspect | Description |
|-----------------------|---|
| Project team | Margreet Worm-Smeitink, MSc; Hans Knoop, PhD, in collaboration with Judith Rosmalen, PhD, Anne van Gils, MSc, Rei Monden, PhD, and others |
| Organisation | AMC |
| Project | The CFS EMA study was designed to investigate - through timeseries analyses - whether there are differences in perpetuators of fatigue between individual patients. |
| **EMA measures | New patients attending the ECCF (n = 102) were asked to complete an e-diary, 5 times a day, for 2 weeks. The times were fixed in consultation with the participant, with a 3-hour break between each assessment. The e-diary assessed fatigue, pain, anxiety, depression, activity (physical, mental, social), patients' focus on fatigue, fatigue catastrophising, self-efficacy, fear avoidance, and social incomprehension. Participants also wore an actometer (actigraphy) during the period when the self-reports were collected. The R 'auto-var' package was used to conduct network analyses. Results are forthcoming. Identified determinants of fatigue will be used to personalize treatment of CFS-patients. |

| Aspect | Description |
|----------------------|--|
| Platform used | RoQua (https://www.roqua.nl). Participants received a link to the (web-based) e-diary via an SMS to their personal smartphone. |
| Contact Information | https://www.amc.nl/web/research-75/publications/prof.-dr.-j.a.-knoop.htm http://nkcv.nl/onderzoek/expert-centre-chronic-fatigue/ |

13.2.2 Sprangers

13.2.2.1 FAntasTIGUE

| Aspect | Description |
|----------------------|--|
| Project team | Mirjam Sprangers, PhD, in collaboration with Yvonne Goërtz, MSc; Zjala Ebadi (from July 2018), MSc, Melissa Thong, PhD, Daisy Janssen, MD, PhD, Jeanette Peters, PhD, Jan Vercoulen, PhD, Chris Burtin, MSc, Yvonne Meertens-Kerris, Arnold Coors, Jean Muris, MD, PhD, Emiel Wouters, MD, PhD, Judith Prins, PhD, and Martijn Spruit, PhD |
| Organisation | AMC (in collaboration with Ciro-Horn, Maastricht UMC, Radboud UMC, and Hasselt University) |
| Project goals | The FAntasTIGUE study examines fatigue in patients with chronic obstructive pulmonary disease (COPD; n = 400), by evaluating the course of fatigue, precipitating/perpetuating factors and hospitalisation. A secondary aim is to identify diurnal fluctuations in fatigue, by using EMA in a subsample of participants (n = 40) (Goërtz et al., 2018). |
| EMA measures | The project comprises four data collection periods (baseline, and 4, 8, and 12 month follow-up). During data collection periods, patients are provided with iPods, on which an EMA application will be pre-installed. Participants will be prompted 8 times a day, at random moments between 7:30 and 22:30, for 5 consecutive days, to answer 19 items (including nine contextual items). Measured concepts include fatigue, relaxed feeling, breathlessness, agitation, uncertainty, irritation, satisfaction, anxiety, feeling energetic, and feeling mentally fit. Items are rated on a 7-point Likert-scale, ranging from 'Not at all' to 'Very much'. In addition, participants are asked to complete a morning questionnaire soon after they awaken to assess the quality of their sleep the previous night. Participants also complete an evening questionnaire, assessing the general perception of their day just before going to bed. During the ema data collection period, patients will also be asked to wear an actigraph (activity monitor), 24 hours a day, for one week. |
| Platform used | Psymate (https://psymate.eu ; see Chapter 14) |
| Contact | https://www.amc.nl/web/research-75/publications/prof.-dr.-m.a.g.-sprangers-publications.htm |

13.2.2.2 IMPACT

| Aspect | Description |
|---------------------|---|
| Project team | Mirjam Sprangers, PhD, with Iris Hartog, MSc; Justine Netjes (until 2017), MSc; Tom Oreel, MSc; Pythia Nieuwkerk, PhD; Michael Scherer-Rath, PhD; José Henriques, MD, PhD; Hanneke van Laarhoven, MD, PhD |
| Organisation | AMC |

| Aspect | Description |
|----------------------|---|
| Project | The IMPACT project (http://www.impactonderzoek.nl/) is an NWO-funded study targeting the Quality of Life (QoL) of patients with multiple chronic morbidities, specifically those undergoing a cardiac intervention. The project aims to improve the conceptualisation of QoL and enhance the sensitivity and comprehensiveness of its measurement by taking the trait-state distinction, contextual factors, and response shift into account). Participants are cardiac patients with comorbidities who were scheduled for elective percutaneous coronary intervention (PCI) or elective coronary artery bypass graft (CABG) (N = 320). In a subsample of participants, QoL is monitored through EMA (n = 37), to monitor QoL states. The study has a longitudinal design, with three EMA data collection periods: 1) pre-treatment, 2) two weeks after treatment for PCI patients or 3 months post-treatment for CABG patients, and 3) six months post-treatment. |
| EMA measures | Participants are prompted to answer nine general and one evening questionnaire per day, for seven consecutive days. During the day, patients are beeped randomly between 7:30 and 22:30 to complete a 19-item questionnaire. Concepts measured include: positive mood (feeling energetic, relaxed feeling, cheerfulness, happy), negative mood (anxiety, sadness, irritation, worry), coronary artery disease symptoms (chest pain, tightness in chest, oppressive feeling on the chest), and general symptoms (tiredness, other types of pain, shortness of breath), and contextual items. Items are rated on a 7-point Likert-scale, ranging from 'Not at all' to 'Very much'. Patients are also asked to complete an evening questionnaire just before they go to bed. The evening questionnaire has, besides the general questionnaire, an additional set of questions from the EQ-5D, and a question on the overall health state of the day. This item is rated on a visual analogue scale from 0 (worst) to 100 (best). Data will be analysed with vector autoregressive models, using R. |
| Platform used | Psymate (https://psymate.eu); Participants are provided with iPods with a pre-installed EMA application, or , if they prefer, can use their own device. |
| Contact | https://www.amc.nl/web/research-75/publications/prof.-dr.-m.a.g.-sprangers-publications.htm |

13.2.3 Snoek

Frank Snoek is Professor of Medical Psychology, specialized in psychosocial diabetology. He heads the Diabetes Psychology Research Group (<https://www.vumc.com/branch/diabetes-psychology/>) and is consulting clinical psychologist for the VUmc Diabetes Center <https://www.vumc.nl/afdelingen/diabetescentrum/>). Snoek and co-workers use EMA to study the relationship between blood glucose variability and wellbeing.

| Aspect | Description |
|---------------------|---|
| Project team | Frank J. Snoek, PhD, Maartje de Wit, PhD, and others |
| Organisation | VUmc, AMC |
| Project | In the MERITS study ('Momentary assessment of patient Experiences in Real life of Insulin Glargine 300 in Type 1 diabetes'), Snoek and co-workers use EMA to explore whether a) blood glucose variability is associated with changes in wellbeing (mood / energy) during waking time, whether b) switching to U-300 results in less glucose variability and translates into improved mood over time within patients, and whether c) if individual differences (profiles) can be distinguished with regard to the (strength of the) association between glucose variability and changes in mood. |

| Aspect | Description |
|----------------------|--|
| EMA measures | Adult patients (N=70) with type 1 diabetes, will be (randomly) prompted to answer questions on mood (based on POMS questionnaire), diabetes distress, fear of hypoglycaemia and sleep. |
| Platform used | Ilumivu (http://ilumivu.com , see Chapter 14) |
| Contact | https://research.vumc.nl/en/persons/frank-snoek https://research.vumc.nl/en/persons/maartje-de-wit |

13.3 Suicidal Ideation

13.3.1 Van Ballegooijen

Dr. Wouter Van Ballegooijen is a post-doctoral researcher at the VU and GGZ inGeest, specialised in the use of information and communication technology in mental health care and suicide prevention. In the ‘Continuous Assessment for Suicide Prevention And Research’ study (CASPAR; Nuij et al. (2018)) Van Ballegooijen and colleagues use EMA to study pathways to suicidal behaviour.

| Aspect | Description |
|----------------------|---|
| Project team | Wouter van Ballegooijen, PhD; Chani Nuij, MSc.; Ad Kerkhof, PhD; Jan Smit, PhD; Heleen Riper, PhD |
| Organisation | Vrije Universiteit Amsterdam, VUmc, GGZ inGeest |
| Project | The ‘Continuous Assessment for Suicide Prevention And Research’ (CASPAR) study ([Nuij et al. (2018)]) aims to evaluate the feasibility of mobile safety planning and daily mobile self-monitoring in routine care treatment for patients with major depression or dysthymia and suicide risk in mental health care. Feasibility will be operationalised in terms of uptake, usage, acceptability, usability and patient satisfaction. EMA data will be used to: (a) empirically validate hypothesised psychological processes and stages of suicide pathways, (b) identify individual pathways to suicidal behaviour, and (c) profile (sub)types of suicidality. The project is expected to result in 1) an interactive smartphone-based safety plan that patients can access 24/7, 2) increased disease awareness of patients due to self-monitoring, and 3) input for the national and international field of mental health care by sharing our results and our data, ultimately contributing to more personalised interventions according to precision medicine principles, and more effective suicide prevention. |
| EMA measures | In the project, adult suicidal patients (N = 80) with major depression or dysthymia and suicide risk in mental health care will be prompted, three times a day, to answer eight self-report items (e.g. ‘I feel sad’) assessing mood, rumination, hopelessness, defeat, entrapment, burdensomeness, belongingness, impulsiveness, suicidal imagery and suicidal ideation. Items, which are based on existing questionnaires, such as the Patient Health Questionnaire (PHQ-9), are rated on a 7-point Likert-scale, ranging from ‘Not at all’ to ‘Very much’. Results are presented to patients in graphs, which patients encouraged to discuss with their clinicians. In addition to active EMA data, location data are also gathered to assess movement patterns and daily rhythms. Planned variables also include accelerometer data and smartphone usage patterns. These data are not visible to study participants. |
| Platform used | Ilumivu (http://ilumivu.com , see Chapter 14) |
| Contact | https://research.vu.nl/en/persons/wouter-van-ballegooijen https://research.vu.nl/en/persons/chani-nuij |

13.4 Psychotic symptoms

13.4.1 Van der Gaag

Prof. dr. Mark van der Gaag is professor of Clinical Psychology at the VU University in Amsterdam, and head of psychosis research at Parnassia, the Hague. He is specialised in the research and treatment of psychosis. In two recent RCTs, described below, Van der Gaag and colleagues used EMA outcomes to collect in-the-moment data.

13.4.1.1 TemStem

| Aspect | Description |
|----------------------|--|
| Team | Mark van der Gaag, PhD, Alyssa Jongeneel, MSc, David van den Berg, PhD, Dorien Scheffers, MSc |
| Organisation | Vrije Universiteit Amsterdam, VUmc, GGZ inGeest, Parnassia Psychiatric Institute |
| Project | The TemStem project focusses on people who suffer from hearing voices and are obstructed by them in their daily life. Study participants install an app that contains both an EMI and EMA function, which is designed to reduce distress and dysfunction caused by auditory verbal hallucinations (Jongeneel et al., 2018). Components of the app include 1) coping: addressing verbal working memory phonological loop with a language task, thereby blocking the hearing of voices, 2) positive reinforcement: decreasing self-reported negative self-esteem themes, 3) treatment: reducing emotional response to memories associated with voices by taxing the auditory working memory during recall of negative auditory memories (as in EMDR therapy). The TemStem study aims to explore the effect of the app on distress and dysfunction in an RCT, specifically with regard to the effect of Temstem on frequency and severity of AVH. Additional analyses will focus on the identification of working mechanisms (predictors and mediators of effects), and the usability of Temstem. |
| EMA measures | TemStem users are encouraged to fill-in nine self-report items on a daily basis. Items tap: 1) hearing voices: 6 items (e.g. "Today, the voices were disturbing"), 2) mood, 3) self-esteem, 4) the use of Temstem ("I used Temstem today"). Items, which are rated on a 7-point Likert-scale, are based on existing EMA questionnaires. Results are presented to users in separate graphs, to support users in gaining insight in the pattern of AVH over time, or after use of Temstem. Stored variables include scores of vividness of AVH pre and post use of Temstem, data on application use (duration), used function (e.g. 'Silencing' function which focuses on coping, or 'Challenging' function which is based on dual tasking), and how users feel when they hear voices. Users can choose to provide additional information on age, gender, which county in the Netherlands they are currently located, how they found the app, and why they want to use it |
| Platform used | Temstem application was developed by Reframing Studio, in collaboration with Parnassia Psychiatric Institute and TU Delft. The app is available for IOS and Android. |
| Contact | http://research.vu.nl/en/persons/m-van-der-gaag http://research.vu.nl/en/persons/alyssa-jongeneel http://research.vu.nl/en/persons/dpg-van-den-berg |

13.4.2 Van der Gaag: VRETp

| Aspect | Description |
|--------------------------|---|
| Team Organisation | Mark van der Gaag, PhD, Roos Pot-Kolder, MSc, with others Vrije Universiteit Amsterdam, VUmc, (in collaboration with UMC Groningen, Maastricht University) |
| Project | In the context of a trial exploring the effect of virtual reality exposure therapy on social participation in people with a psychotic disorder (VRETP; n = 116, Pot-Kolder et al. (2016)), EMA data were collected to assess changes in social functioning and paranoid ideation. |
| EMA measures | EMA included momentary assessment of 1) Anxiety (one item, e.g. “I feel anxious”), 2) Perceived social threat (four items, e.g. “In this company, I feel accepted”), 3) Paranoia (three items, e.g. “I feel suspicious”), and 4) Time spent with others (max. three multiple choice items inquiring about type of company (nobody, family, non-family, etc.). These items were based on previous EMA work (e.g., Collip et al. (2011)). Participants were prompted 10 times a day, during 6 days. Anxiety, threat and paranoia items were rated on a 7-point Likert scale, ranging from 1 (“not at all”) to 7 (“very”). Reports had to be completed within 15 min of the beep. To be included in the analysis, participants had to complete diary entries for at least one-third of the beeps (i.e., a minimum of 20 measurements). All 116 participants completed EMA measurements at baseline (mean number of completed self-assessments 46.1, SD 13.3), 96 participants completed the post-treatment assessment sufficiently (43.1, SD 10.1), and 87 participants completed the follow-up (43.2, SD 11.1). The trial results suggest that the addition of VR-CBT to standard treatment can reduce paranoid ideation and momentary anxiety in patients with a psychotic disorder (Pot-Kolder et al., 2018). |
| Platform used | PsyMate (http://www.psymate.eu/ , see Chapter 14). Because the PsyMate application was not finished at the time of the trial, participants were provided with a small palmtop device for the duration of the study. |
| Contact | http://research.vu.nl/en/persons/m-van-der-gaag |

13.5 Symptom Networks

13.5.1 Borsboom

The Psycho-systems group is an UvA based initiative, led by Prof. Dr. Denny Borsboom, focusing on “the development of new methodologies for psychological research using complex systems theory and network models” (<http://psychosystems.org/>).

| Aspect | Description |
|---------------------|--|
| Senior project team | UvA: Denny Borsboom, PhD; Lourens Waldorp, PhD; Han van der Maas, PhD; Sascha Epskamp, PhD; Claudia van Borkulo, PhD; Donald Robinaugh, PhD |
| Organisation | Universiteit van Amsterdam (UvA) |
| Topic | Complex systems theory and network models |
| Status | Ongoing, founded in 2013 |
| Aims | Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam (Epskamp et al., 2018c,) “temporal and contemporaneous relationships should not be overinterpreted, as these merely show predictive effects and can at most only highlight potential causal pathways.” Reproducibility Project: Psychology |
| Results | |
| Website | http://psychosystems.org/ |

13.6 Sleep

13.6.1 Van Someren

Prof. dr. Eus van Someren is affiliated with the department of sleep and cognition of the Netherlands Institute for Neuroscience (<https://herseninstituut.nl/>), the Department of Psychiatry of VUmc and GGZ Ingeest, Amsterdam. His group focusses on the study of healthy and disturbed sleep, using neuro-imaging, actigraphy and EMA.

| Aspect | Description |
|--|--|
| Team | Eus van Someren, PhD (department head), with Bart te Lindert, MSc, Wisse van der Meijden, MSc, Tessa Blanken, MSc, Michele Colombo, PhD, Kim Dekker, MSc, Jeanne Leerssen, MSc, Rick Wassing, MSc. |
| Organisation Project(s) | VUmc, GGZ inGeest, the Netherlands Institute for Neuroscience <i>Wisse van der Meijden</i> studies, among other things, the interplay between light, vigilance, and sleep. A recent study associated the post-illumination pupil response (PIPR) to blue light with multiple indices of sleep timing: a) a questionnaire on habitual lights-out time, sleep onset latency, and final wake-up time (Munich Chronotype Questionnaire, MTCQ (Roenneberg et al., 2003)); b) a one-week sleep diary on actual wake/sleep times; and c) actigraphy. Actigraphy data were obtained using either a Philips Actiwatch Spectrum or a microelectromechanical accelerometer (Move II, Movisens GmbH). Data collection was set to 30Hz. Sleep onset and final wake-up time were estimated using an algorithm that is implemented in Matlab (https://github.com/btlindert/actant-1). Participants (adolescents and young adults, n = 71) with a later sleep timing had a stronger responsiveness to blue light. The mid-sleep timing estimates from sleep diaries and actigraphy shared 94.5% of their interindividual variance (Van Der Meijden et al., 2016). |

| Aspect | Description |
|------------------|---|
| | <p><i>Bart te Lindert</i> studies the effect of the environment on sleepiness, for example by measuring light, (skin) temperature, posture, and psychological variables. A recent study focused on the effect of light intensity on Liking, Wanting and mood in insomnia disorder. In a recent study, active and passive EMA were combined. EMA prompts were sent eight times a day, for one week. Prompts were timed at quasi-random intervals between 8:00 and 22:00. In addition, participants provided input after waking up and before bedtime. The interval between prompts ranged between 16 minutes and 3 hours. EMA consisted of 22 items. Liking and Wanting (6 items each), focused on taste or smell, bodily sensation, watching or listening, interactions with other, physical activity or being busy, and receiving something. Participants indicated whether they enjoyed and wanted these categories on a 0-100 VAS scale (ranging from not to very much). Mood items were derived from the Daytime Insomnia Symptom Scale (DISS, Buysse et al. (2007)), which was developed specifically for EMA. Items focused on positive mood (5 items; relaxed, energetic, calm, happy, and efficient) and negative mood (5 items; anxious, stressed, tense, sad, and irritable). Items were scored on a 0-100 VAS-scale (ranging from very little to very much). In addition, participants wore two light sensors (Dimesimeter), one on their indoor clothes and one on their jacket. Wear-time of each sensor was derived from a build-in accelerometer. Light was sampled every minute for one week. Participants received a designated Android smartphone for the duration of EMA. Detailed information on data processing and analysis can be found in Te Lindert et al. (2018). On average, 80% of the prompts contained valid data. People with insomnia disorder ($n = 17$) had significantly lower subjective Liking and Wanting than matched controls without sleep complaints ($n = 18$). This was most apparent at low environmental light intensity. There were no overall differences between groups in Positive mood and Negative mood. Participants with insomnia did have a different diurnal profile of Positive mood and Negative mood (Te Lindert et al., 2018). The one-item Karolinska sleepiness scale is advised as a simple way to measure sleepiness in EMA. The item was validated against performance and EEG variables (Kaida et al., 2006).</p> |
| Platform(s) used | Philips Actiwatch Spectrum; Microelectromechanical accelerometer (Move II, Movisens GmbH) |
| Contact | http://research.vumc.nl/en/persons/eus-jw-van-someren http://herzeninstituut.nl/over-ons/de-organisatie/medewerkers/bart-te-lindert/ http://herzeninstituut.nl/over-ons/de-organisatie/medewerkers/wisse-van-der-meijden/ |

13.7 Ambulatory Assessment

13.7.1 De Geus

Prof. dr. Eco de Geus is head of the department of Biological Psychology and co-director of the Netherlands Twin Registry at the VU University. De Geus has been the driving force behind the development of the VU University Ambulatory Monitoring System (VU-AMS; <http://www.vu-ams.nl/vu-ams/>), a non-invasive wearable device that is used for continuous ambulatory measurement of the autonomic nervous system.

| Aspect | Description |
|--------|---|
| Team | Eco de Geus, PhD; Gonnieke Willemsen, PhD; Martin Gevonden, PhD; Denise van der Mee, MSc, Mandy Tjew-A-Sin, MSc; Cor Stoof, MSc |

| Aspect | Description |
|-------------------------|---|
| Organisation | VU |
| Project | VU-AMS was developed at the department of Biological Psychology of the Vrije Universiteit Amsterdam. The system is used worldwide by many research groups, to study stress and emotion in both laboratory and naturalistic settings (de Geus and van Doornen, 1996). VU-AMS has been used in the study of ADHD, aggression, anxiety and depressive disorders, mental, social and work-related stress, circadian rhythms, hyperventilation, migraine, sleep, and in studies linking the autonomic nervous system to metabolic and immunological risk factors (see http://www.vu-ams.nl/research/publications/). Current projects include: 1) Validation of a wristwatch-based technology, developed by Philips, to measure skin conductance responses in a laboratory (~2.5 h) and ambulatory (~22h, including the night) setting on a total of 100 subjects (van der Mee et al., ongoing, 2017 - 2021). The goal is to test whether wrist based (Philips) and palm based (VU-AMS) measured skin conductance responses accord, how these measures relate to the heart based measured pre-ejection period (PEP, VU-AMS) and whether it can be related to positive and negative affect (hourly diary prompts). The end goal for the wrist based technology is to detect sympathetic nervous system activity (measured as skin conductance responses) and present this information to the person's as an index of the current stress level, alongside a one-hour prediction of changes in stress level and cognitive functioning (Cognitive Zone Changes; http://www.ip.philips.com/licensing/program/121) 2) Ambulatory study on self-regulation among youth (Tjew A Sin et al., 2018-2019). This study is conducted at the schools as part of the neurolab project. A total of 50 students, will wear the VU-AMS device for approximately 22 hours. Autonomic nervous system activity is collected continuously over the course of a regular school day (including the night and morning thereafter) and will be related to multiple components of self-regulation, including emotion-regulation, cognitive functioning, impulsivity and inhibition |
| EMA measures | The VU-AMS device is a battery powered wearable that can record 24 to 48 hours of data (4GB storage). It measures the electrocardiogram, the impedance cardiogram, and skin conductance. With VU-AMS, the following outcomes can be collected: Heart Rate / Inter beat Interval (IBI), Heart Rate Variability (SDNN, RMSSD, IBI power spectrum: HF, LF), Respiratory Sinus Arrhythmia (RSA), Pre-Ejection Period (PEP), T-wave amplitude (TWA), Left Ventricular Ejection Time (LVET), Stroke Volume (SV) and Cardiac Output (CO), Respiration Rate (RR), Skin Conductance Level (SCL) and Skin Conductance Responses (SCRs), Movement (Hip-worn tri-axial accelerometer signals (g)). The freely available Data Analysis and Management System (DAMS) package is used for extraction and processing of VU-AMS data (http://www.vu-ams.nl/support/downloads/software/). The DAMS tool offers options for data inspection (visual inspection of raw data), automated detection of R-peaks in raw ECG signal and visual inspection of final IBI time series, event/diary-based data labelling, IBI spectral power calculation, automated scoring of RR, RSA, and PEP from the combined ECG |
| Platform(s) used | VU-AMS / DAMS (see also Chapter 14) |
| Contact | http://research.vu.nl/en/persons/jcn-de-geus https://research.vu.nl/en/persons/ahm-willemsen http://research.vu.nl/en/persons/martin-gevonden |

Chapter 14

EMA Instruments Catalogue

To conduct EMA studies, a variety apps, online applications and wearable devices are on the market. In this chapter, we list a (small) selection of instruments that we found to be in use in scientific studies within the APH consortium.

Table 14.1: EMA Instruments.

| Name | Manufacturer | URL | Description |
|------------|------------------|-------------------|---|
| GENEActive | activinsights | activinsights.com | Wrist-worn accelerometer |
| Ilumivu | Ilumivu | ilumivu.com | Android / iOS EMA app |
| MoodBuster | ICT4D Consortium | moodbuster.eu | Android EMA app |
| Movisense | MoviSence | movisense.org | Android EMA app |
| PsyMate | PsyMate | psymate.eu | Android EMA app |
| RoQua | RoQua | roqua.nl | SMS / Browser-based EMA |
| VU-AMS | VU | www.vu-ams.nl | Wearable for autonomic nervous system measurement |

14.1 GENEActive

GENEActive, sold by UK-based company Activinsights (activinsights.com), is a waterproof wrist-worn device with a high-precision, configurable 3-axial accelerometer (range: +/- 8g), an ambient light sensor, a (near-body) temperature sensor, and an event logger (a button that users can press to mark a targeted event). GENEActive was developed to accurately assess human activity in scientific studies. The device has a storage capacity of 0.5GB of raw data. At 10Hz, the device can log activity up to one month. July 2018, the price for one unit is approximately 250 euro.

GENEActive is used in a growing number of clinical studies to measure activity and sleep-wake cycles, in natural conditions, over longer periods of time. Dedicated R-packages to pre-process and analyse the raw data exist (see Chapter 15). Note, however, that no accompanying app exists with which study participants can be provided feedback about their activity. This might negatively affect wear-time and study compliance in research participants, who are accustomed to consumer activity-sampling devices, such as Fitbit, where many options for real-time feedback exist.

14.2 Ilumivu

Ilumivu)

Ilumivu (see <https://ilumivu.com/>) is an American commercial company specialized in mobile EMA application



Figure 14.1: The GENEActiv Accelerometer

development. It offers a cross-platform (Android and IOS) smartphone app (mEMA) that researchers can use to collect data from study participants. Researchers can define the assessment plan of their study in an online back-office (i.e., without any assistance from Illumivu staff).

The Ilumivu EMA toolbox library includes a rather complete set of survey elements, survey logic branching tools, and survey scheduling options. Passive EMA options include GPS tagging (of survey responses), GEO-fencing (triggering surveys at specific locations) and smartphone sensor logging (light and noise level, screen brightness, screen locked/unlocked, humidity, ambient temperature, barometric pressure, phone/SMS activity log, and basic device information). A limited number of smartwatch devices can optionally be linked to the Ilumivu app for further passive data collection. With the online backoffice, researchers can invite participants, monitor study compliance and download data (in common data formats such as CSV). The company is open for custom development, when studies require features that are not included in the default app.

Current (July 2018) subscription plans range from 3.375 dollar (basic features) to 12.000 dollar (all features), per year. Researchers, who are interested in purchasing a licence of the service, are advised to the ‘Grant Writer’s Guide’ of Ilumivu, which can be downloaded from the site of the company (at <https://ilumivu.com/pricing/writing-a-grant/>).

Ilumivu’s competitive advantage is that it has multi-platform support. It is, however, not the cheapest EMA product on the market. Being an American company, researchers should also consider EU regulations relating to personal data privacy protection, since regulations are more strict when personal data of EU citizens leave the EU.

14.3 MoodBuster

Moodbuster)

MoodBuster (<http://www.moodbuster.eu/>) is a web-based treatment platform with an integrated EMA app. The platform was developed by an international non-profit research consortium, including VU, VUmc and GGZ InGeest, in two major EU-funded research projects: ICT4Depression (see ICT4Depression.eu) and E-COMPARED (see <http://e-compared.eu>).

In the E-COMPARED trial, MoodBuster was used, in five EU-countries, to test blended treatment of major depression (Kleiboer et al., 2016). In this study, participants used the smartphone app to rate mood and various other depression-related variables, in the context of their treatment, over a period of




Figure 14.2: Ilumivu App Screenshots

up to 20 weeks. In addition, the EMA app was used in a satellite study designed to assess the effects of long-term EMA (van Ballegooijen et al., 2016). MoodBuster will also be used in the EU ImpleMentAll project (<http://www.implementall.eu>) and in several other clinical trials that are in preparation.

Currently, EMA assessment protocols are hard-coded in the app. New EMA assessments protocols can be implemented in collaboration with the MoodBuster development team. An online back-office is in development. Previous studies used an Android version of the EMA app. Cross-platform versions of the app in development.

More information on MoodBuster can be obtained from prof. dr. Heleen Riper (h.riper@vu.nl).



Figure 14.3: MoodBuster App Screenshots

14.4 Movisens

Movisens)

Movisens (<http://www.movisens.com>) is a German company that is specialised in the development of hard-

and software solutions for mobile sensing. The company sells small wearable devices that contain several high-precision sensors, including an accelerometer, gyroscope, barometer and thermometer. In addition, the company has developed an (Android) app, called MovisensXS, which can be used for active EMA research. The app can optionally be configured for smartphone logging (e.g., to log music that a study participant listens to). The wearable sensor can also be linked to the app, so that EMA questionnaires can be triggered based on targeted activity or energy expenditure patterns, such as extended periods of sedentary behaviour. Specialised software to import, pre-process and analyse raw sensor data is available for download.

Like Ilumivu, researchers can define EMA sample schedules for their study in a web-based backoffice (<https://xs.movisens.com>), using an online graphical editor. Once defined, participants can be invited to the study, through the back-office, to download the freely available Movisens App from Google Play store. The backoffice also allows researchers to monitor study compliance and download data.

MoviSensXS EMA licence costs vary from 500 to 10.000 euro's per year, depending on the required number of 'credits' which are linked to the number of EMA responses. Prospective users can test platform, without functional restrictions, with a free test account. An EMA test-study can thus be set up and started in less than a day.

A major limitation of Movisens is the lack of an iOS version of the EMA app. Study participants who own an iPhone have to be excluded from studies, or will have to be provided with an Android phone.




Figure 14.4: Movisens Sample scheme editor (left) and App Screenshots (right)

14.5 PsyMate

The PsyMate™ app (www.psymate.eu) was developed by the Department of Psychiatry and Psychology at Maastricht University in the Netherlands to assess psychological problems in daily life. The app has been validated for use in depression, bipolar disorder, and psychosis, with new scales currently being developed for a range of diseases including Parkinson's disease, pain, cardiology, hypertension, diabetes and Irritable Bowel Syndrome. It is currently used in a EU-funded project to study gene-environment interaction in schizophrenia (<http://www.eu-gei.eu/about-the-project/psymate>).

The app is free to download for iOS and Android devices on Apple and Google play stores. Uses for the app include self-monitoring of mood states, for professional support during treatment, or for research purposes. The app can be customized to address specific client needs or research projects with expertise from the PsyMate back office, which includes a working group that meets regularly to discuss and advise new projects.

Researchers have access to the raw data without having to go through the Psymate back office. A disadvantage of the Psymate app is that technical problems could arise when there are either iOS or Android updates. Communication from the Psymate back office to researchers about updates and assistance with technical problems could be a point for consideration for using this platform.




Figure 14.5: PsyMate App Screenshots

14.6 RoQua

RoQua)

RoQua (<http://www.roqua.nl/>) is a web-based Routine Outcome Monitoring system, developed and maintained by a Dutch non-profit development and service organization that is funded by several northern GGZ organisations and the Department of Psychiatry, University Medical Center Groningen. RoQUA has a sophisticated and user-friendly online back-office portal, with which researchers can define assessment protocols and invite study participants - through e-mail or SMS - to complete questionnaires online (on desktop or mobile devices). By inviting study participants several times a day to complete a questionnaire via the standard web browser of their mobile phone, active EMA can be implemented. This approach was taken in several large-scale studies, including ‘NESDA’ (<http://nesda.nl>; see also Chapter 13) and ‘HowNutsAreTheDutch’ (www.hoegekis.nl; see (Van Der Krieke et al., 2017; Krieke et al., 2016)).

At present, RoQua does not support the collection of passive EMA data. However, preliminary results have been reported with a system called ‘Physiqual’ (Blaauw et al., 2016), with which EMA data, collected with RoQUA, can be automatically combined with wearable sensor data.

14.6.1 VU-AMS

The VU University Ambulatory Monitoring System (VU-AMS; <http://www.vu-ams.nl/>), developed by the department of Biological Psychology and the Technical Department (ITM) of the Faculty of Psychology and Education, allows ambulant recording of autonomic and cardiovascular activity. VU-AMS measures heart rate, heart rate variability, Respiratory Sinus Arrhythmia, Pre-Ejection Period, Left Ventricular Ejection Time, Respiration Rate, Stroke Volume (SV) and Cardiac Output, Skin Conductance Level (SCL) and Skin Conductance Responses (SCRs) and Tri-Axial Accelerometry (of Body Movement). For the processing of VU-AMS data, a dedicated software suite called the ‘Data Analysis and Management Software’ (VU-DAMS) is available (for Windows and Mac).



Figure 14.6: Screenshots of the participant feedback web-page of the 'HowNutsAreTheDutch' project, in which data is collected by the RoQUA system




Figure 14.7: VU-AMS device

Chapter 15

R Packages for EMA Research

Many R packages exist that can help you in the management and analysis of EMA data. In this chapter, a selection of these packages are discussed. For each, we provide a summary description, a code example code, and pointers to further documentation, to give you a head start in using the packages for your work.

Table 15.1: List of R packages that are useful in EMA research.

| Category | Package | Description |
|--------------------------------------|------------------|---|
| Accelerometry | GENEActive | Import GENEActive data into R |
| | GGIR | Pre-proces and analyse raw multi-day multi-day accelerometer data. |
| | PhysicalActivity | Analyse actigraph accelerometer data. |
| Data Management & Visual Exploration | dplyr | Data transformation |
| | ggplot2 | Create graphs |
| | haven | Import and export SPSS data files |
| Mixed-effects Modeling | lubridate | Manipulate date and time variables |
| | lme4 | Fit linear and nonlinear mixed-effects models. Fast alternative to package ‘nlme’. |
| | nlme | Fit linear and nonlinear mixed effects models. Pre-dates package ‘lme4’, but is still used because it provides more advanced options to model correlational structures in the data. |
| Power Analysis | simr | Simulation-based power calculations for mixed models. |
| Simulation | simstudy | Simulate study data. |
| Spatio-temporal analysis | adehabitatHR | Developed for home range estimation of wild animals from GPS data. Useful for human data as well (see Chapter 12). |
| Symptom Networks | autovarCore | Automate the construction of vector autoregressive models. |
| | bootnet | Assess the stability of symptom networks. |
| | qgraph | Estimate and plot symptom networks. |
| Time series analysis | lomb | Calculate the Lomb-Scargle Periodogram for unevenly sampled time series. |

15.1 Accelerometry

Accelerometer data need considerable pre-processing before final analyses can be run. Raw data have to be read in from a variety of brand-specific file formats, data have to re-calibrated on a per-device basis, non-wear periods have to be detected, and summarizing measures, such as activity counts and energy-expenditure measures, have to be calculated from imputed triangular (x, y, z) data, often in several time windows (i.e., epochs).

15.1.1 Package GENERead

GENEActive, sold by Activinsights, is a wrist-worn tri-axial accelerometer that is often used in clinical research studies. With package ***GENERead*** (Fang et al., 2018), raw data can be imported into R for further processing, as illustrated below.

```
# Code snippet 15.1: Reading raw sensor data.
library(GENERead)
library(tidyr)

dat <- read.bin(system.file("binfile/TESTfile.bin", package = "GENERead"),
                verbose = FALSE, downsample = 20)
#> Processing took: 0.1 secs .
#> Loaded 1560 records (Approx 0 MB of RAM)
#> 12-05-23 16:47:50.000 (wo) to 12-05-23 16:53:01.799 (wo)

d <- as.data.frame(dat$data.out)
d <- gather(d, key = "sensor", value = "value", -timestamp)
d$timestamp <- as.POSIXct(d$timestamp,
                           origin = "1970-01-01",
                           tz = "UTC")

ggplot(d, aes(x = timestamp, y = value)) +
  geom_line() +
  facet_wrap(~sensor, scales = "free_y")
```




Figure 15.1: Raw sensor data of a GENEActive wrist-worn tri-axial accelerometer (down-sampled from 100Hz to 5Hz).

15.1.2 Package ‘GGIR’

Package *GGIR* (van Hees et al., 2018) is a packages to pre-process raw accelerometry data from three brands of wearables that are widely used in sleep and physical activity research: GENEActiv, ActiGraph and Axivity.

15.1.3 Package ‘PhysicalActivity’

Package ‘*PhysicalActivity*’ (Choi et al., 2018) provides an alternative to package ‘GGIR’, when ActiGraph data are available.

```
# Code snippet 15.2: plotting activity counts.
library(PhysicalActivity)
library(ggplot2)

data(dataSec)

d <- dataCollapser(dataSec, TS = "TimeStamp", col = "counts", by = 300)

ggplot(d, aes(x = as.POSIXct(TimeStamp), y = counts)) +
  geom_line(size = .5, alpha = .5) +
  xlab("Time") + ylab("Activity Counts")
```




Figure 15.2: Activity Counts (5-minute windows), in a Three-day Accelerometer data set.

15.2 Data management & Visual Exploration

The tidyverse is a collection of well-designed packages, authored by the team behind RStudio, that together add a consistent, modern, and efficient extension of base R functionalities. The tidyverse includes popular packages such as “*ggplot2*” (for plotting), “*haven*” (to read SPSS files), “*dplyr*” (for data manipulation), and many more (see: <http://tidyverse.org> for a full list).

15.2.1 Package *dplyr*

Package ‘*dplyr*’ (Wickham et al., 2018) implements the ‘split-apply-combine’-strategy. With ‘*dplyr*’, elementary data manipulations can be chained (using the pipe operator ‘%>%’) to elegantly implement complex data transformations.

```
# Code snippet 15.3: aggregate data by ID, through a 'pipe'.
require(dplyr)

d <- data.frame(
  c = factor(rep(1:5, each = 10)),
  score = rnorm(50)
)

b <- as_tibble(d) %>%
  group_by(c) %>%
  summarize(mean_score = mean(score))
```

A good introduction to dplyr can be found in the book ‘R for Data Science’ (Wickham and Grolemund, 2016), which can be freely accessed online (<http://r4ds.had.co.nz/>).

15.2.2 Package ‘ggplot2’

Package ***‘ggplot2’*** (Wickham, 2016) provides a collection of high-level plotting commands with which graphs can be build up in layers. It is based on ‘The Grammar of Graphics’ (Wilkinson, 2006), an influential analysis of the structure of scientific graphs. Systematic introductions are available on the the tidyverse website, and in the book ‘ggplot2: Elegant Graphics for Data Analysis’ (Wickham, 2016).

The example below illustrates how graphs are layered. In the first step, a coordinate system is set up. In step 2, all time/scores points are plotted. In step 3, a smoothed line is fitted through these points. Finally, in step 4, the graph is split on a variable ID (a subject identifier), to show individual trajectories.

```
# Code snippet 15.4: simple ggplot example.
library(ggplot2)

# simulate example data
d = data.frame(
  ID      = rep(1:4, each = 25),
  time   = rep(1:25, 4),
  score  = rnorm(100, 0, 2))

# step 1: initialise the coordinate system
g <- ggplot(d, aes(x = time, y = score)); g

# step 2: add scatterplot
g <- g + geom_point(); g

# step 3: fit a smoothed line
g <- g + geom_smooth(); g

# step 4: split plot by ID
g + facet_wrap(~ ID)
```




Figure 15.3: Plotting layers with ggplot2

15.2.3 Package ‘haven’

With package **haven** (Wickham and Miller, 2018), SPSS, STATA and SAS files can be read into R.

```
# Code snippet 15.5: read SPSS data
library(haven)

path <- system.file("examples", "iris.sav", package = "haven")
d <- read_sav(path)

attributes(d$Species)
#> $format.spss
#> [1] "F8.0"
#>
#> $class
#> [1] "labelled"
#>
#> $labels
#>   setosa versicolor virginica
#>   1          2          3
```

15.2.4 Package ‘lubridate’

EMA data analyses frequently require manipulations of datetime variables. For this, package **lubridate** (Gromelund and Wickham, 2011), which provides many functions for common date and datetime operations, can be very useful.

In the code snippet below, for example, the ‘round_date’ function is used to calculate the ENMO value from raw tri-axial accelerometer data, in 15-minute epoch windows.

```
# code snippet 15.6: rounding dates.
library(emaph)
library(lubridate)
library(ggplot2)
library(dplyr)

d <- subset(genea,
            timestamp > "2018-06-02 12:00" &
```

```

    timestamp < "2018-06-02 18:00")
d$epoch <- round_date(d$timestamp, "minute")

d <- d %>% group_by(id, epoch) %>%
  summarise(svm = sum(sqrt(x^2 + y^2 + z^2)) - 1) / length(x)

```

| id | epoch | svm |
|----|---------------------|------------|
| 1 | 2018-06-02 12:00:00 | -0.0009766 |
| 1 | 2018-06-02 12:01:00 | 0.0072235 |
| 1 | 2018-06-02 12:02:00 | 0.0023875 |
| 1 | 2018-06-02 12:03:00 | 0.0070644 |
| 1 | 2018-06-02 12:04:00 | 0.0265187 |
| 1 | 2018-06-02 12:05:00 | 0.0969168 |

To learn more about handling dates and times with ‘lubridate’, Chapter 16 of the book ‘R for Data Science’ (Wickham and Grolemund, 2016) provides a good introduction.

15.3 Mixed-effects modeling

Several R-packages for mixed-effects modelling exist. The most popular are package **nlme** (Pinheiro et al., 2018) and package ***lme4** (Bates et al., 2015). Both packages are actively used, as both provide unique functionalities.

15.3.1 nlme

Package **nlme** is introduced in Chapter 9. It is an older package (in comparison to package lme4), that is still used a lot because it provides options to model correlational structures that are not implemented (yet) in lme4.

```

# Code snippet 15.7: fit a linear mixed model, with lme.
library(nlme)
fm <- lme(distance ~ age + Sex, data = Orthodont, random = ~ 1)

fixef(fm)
#> (Intercept)      age   SexFemale
#>  17.7067130   0.6601852 -2.3210227

```

15.3.2 lme4

Package **lme4** (Bates et al., 2015) is a faster R-reimplementation of the mixed-effects model. With large data sets and complex hierarchical models, this package should probably be preferred. As can be seen below, models specifications in ‘lmer’ are different from model specifications in ‘lme’.

```

# Code snippet 15.8: fit a linear mixed model, with lme.
library(lme4)
fm <- lmer(distance ~ age + Sex + (1 | Subject), data = Orthodont)

fixef(fm)
#> (Intercept)      age   SexFemale
#>  17.7067130   0.6601852 -2.3210227

```

15.4 Power analysis

15.4.1 Package ‘simr’

With package ‘simr’ (Green and MacLeod, 2016), power of mixed-effects models can be determined via simulation. As illustrated below, the procedure requires the researcher to define the “true” parameters of a mixed model, and a single data set. Then, function ‘simPower’ can be used to simulate new datasets and tests (of a specified parameter in the model), to determine the power of the test.

```
# Code snippet 15.9: Power analysis of a two-group repeated measures design
# (simulation approach).
library(simr)

# construct design matrix
t <- 1:24
s <- 1:40
X <- expand.grid(t = t, s = s)
X$g <- c(rep(0, 24), rep(1, 24))

# fixed intercept and slope
b <- c(2, -0.1, 0, -0.5)

# random intercept variance
V1 <- 0.5

# random intercept and slope variance-covariance matrix
V2 <- matrix(c(0.5, 0.05, 0.05, 0.1), 2)

# residual standard deviation
s <- 1

model1 <- makeLmer(y ~ t * g + (1 + t | s),
                     fixef = b,
                     VarCorr = V2,
                     sigma = s,
                     data = X)

powerSim(model1,
          fixed("t:g", "lr"),
          nsim = 10,
          progress = FALSE)
#> Power for predictor 't:g', (95% confidence interval):
#>     100.0% (69.15, 100.0)
#>
#> Test: Likelihood ratio
#>     Effect size for t:g is -0.50
#>
#> Based on 10 simulations, (0 warnings, 0 errors)
#> alpha = 0.05, nrow = 960
#>
#> Time elapsed: 0 h 0 m 1 s
```

15.5 Symptom Network Analysis

When EMA is used to tap various symptoms, network analysis can reveal the dynamic interplay between these symptoms (see Chapter 10). Various packages exist to fit these networks in R. With these packages, it is relatively easy to fit a graphical network on multivariate data sets. If you are interested in conducting a network analysis, be sure to visit the Psycho-systems website, at <http://psychosystems.org>.

15.5.1 Package ‘autovarCore’

Vector autoregressive (VAR) models can be used to detect lagged relationships between multiple timeseries. In VAR, each variable is modelled as a linear function of past values (lags) of itself and of present and past values of other variables. When EMA is used to capture multiple phenomena over time, VAR can provide insight in how these phenomena interact. One challenge in VAR modelling is that many alternative models potentially exist. Package ‘**autovarCore**’ (Emerencia, 2018) was developed to help researchers to find the VAR model with the best fit to a given timeseries data set.

In the (unrealistic) example below, function ‘autovar’ is used to detect that changes in depression are positively related to past (lag 1) values of activity, in a simulated data set:

```
# Code snippet 15.11: autovar analysis.
library(autovarCore)

# simulate data
N = 100
depression <- rnorm(N)
activity <- rnorm(N)
activity_lag1 <- c(NA, activity[1:(N -1)])

depression <- depression + 0.5 * activity_lag1
d <- data.frame(depression, activity)

models_found <- autovarCore:::autovar(d, selected_column_names = c('activity', 'depression'))

# Show details for the best model found
summary(models_found[[1]]$varest$varresult$depression)
#>
#> Call:
#> lm(formula = y ~ -1 + ., data = datares)
#>
#> Residuals:
#>      Min       1Q   Median       3Q      Max
#> -2.33559 -0.48604 -0.05249  0.63636  2.28235
#>
#> Coefficients:
#>             Estimate Std. Error t value Pr(>|t|)
#> activity.l1    0.53294   0.11918   4.472 2.26e-05 ***
#> depression.l1  0.00454   0.09800   0.046  0.9632
#> depression.l2 -0.01670   0.09347  -0.179  0.8586
#> const          -0.05108   0.09678  -0.528  0.5990
#> outlier_11     1.76189   0.94114   1.872  0.0644 .
#> outlier_20     1.45376   0.94813   1.533  0.1287
#> outlier_28    -1.12760   0.95920  -1.176  0.2429
#> outlier_60     0.46033   0.96524   0.477  0.6346
#> ---
#> Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
#>
#> Residual standard error: 0.936 on 90 degrees of freedom
#> Multiple R-squared:  0.2491, Adjusted R-squared:  0.1907
#> F-statistic: 4.266 on 7 and 90 DF,  p-value: 0.0004131
```

AutovarCore is a simplified version of a more extensive package ***autovar*** [@(Emerencia, 2018), which was used in several publications (van der Krieke et al., 2015, Emerencia et al. (2016)). Further information can be found on <http://autovar.nl> and <http://autovarcore.nl>

15.5.2 Package 'qgraph'

Package ***qgraph*** (Epskamp et al., 2012) can be used to fit, visualise and analyse graphical networks.

In the example below, qgraph is used to fit a network on the ‘Critical Slowing Down’ (CSD) data set, which is included in package ‘emaph’ (see Chapter 11 and Chapter 10).

```
# Code snippet 15.12: fitting a network with qgraph.
```

```
library(qgraph)
library(emaph)

# get mood_ scores from csd data set
d <- subset(csd,
             subset = phase == "exp: db: no change",
             select = grep("mood_", names(csd))[1:5])

# Fit and plot regularized partial correlation network
g <- qgraph(cor_auto(d, detectOrdinal = FALSE),
            graph = "glasso", sampleSize = nrow(d),
            nodeNames = names(d),
            label.scale = FALSE, label.cex = .8,
            legend = TRUE, legend.cex = .5,
            layout = "spring")
```




Figure 15.4: Network of mood items from CSD data set


15.5.3 Package ‘bootnet’

In the interpretation of fitted network plots, it is important to take the stability of the network into account. Intuitively, networks that are fit on small sample data sets will be less stable than networks based on large data sets. One solution is to fit a large number of networks on subsets of the original data, through bootstrapping. In stable networks, the variance edges estimations will be small, while in unstable networks, the variance will be high. This idea is implemented in package **bootnet** (Epskamp et al., 2018a).

Below, the stability of the network that was fit in the previous example is examined with bootnet: fifty networks are fit, based on fifty bootstrapped samples. In the results plot, the red line marks the strength of the edges in the full sample, while grey confidence intervals mark the distribution of the edge weights in the bootstraps.

```
# Code snippet 15.13: a bootstrapped network.
library/bootnet

g <- estimateNetwork(d, default = "EBICglasso",
                      corArgs = list(detectOrdinal = FALSE))
results <- bootnet(g, nBoots = 50, verbose = FALSE)
plot(results, order = "mean")
```




15.6 Timeseries analysis

15.6.1 Package ‘lomb’

Disturbances in circadian rhythms have been related to depressive symptoms (see, e.g., Saeb et al., 2015). With so-called periodograms, these circadian rhythms can be detected in EMA data. Standard analysis techniques, however, expect regular timeseries, in which data are sampled at equidistant intervals. EMA data, typically, are not equidistant. One solution to this problem is to use the Lomb-Scargle periodogram procedure (Lomb, 1976), which can be applied to unevenly-sampled timeseries as well. Package **lomb** (Ruf, 1999) implements this procedure.

```
# Code snippet 15.14: calculating a Lomb-Scargle periodogram.
data(ibex, package = "lomb")
lomb::lsp(ibex[2:3])
```



Part VII

Closing Matters

Acknowledgements

We would like to thank the APH consortium, for giving us the opportunity to write this book, by granting us one of the 2017 APH mental health research funds.

We also thank the APH researchers, who we spoke with during the project, for sharing their knowledge, experience and insights in EMA research. We are also indebted to the researchers who contributed data, analyses scripts, and other materials that are presented in this book.

Finally, we like to thank the project advisors, Prof. Dr. Heleen Riper and Prof. Dr. Johannes Smit, for their valuable support and input.

Jeroen Ruwaard, Lisa Kooistra & Melissa Thong



Bibliography

- aan het Rot, M., Hogenelst, K., and Schoevers, R. A. (2012). Mood disorders in everyday life: a systematic review of experience sampling and ecological momentary assessment studies. *Clinical psychology review*, 32(6):510–23.
- Allport, G. W. (1937). Personality: A psychological interpretation.
- Asselbergs, J., Ruwaard, J., Ejdys, M., Schrader, N., Sijbrandij, M., and Riper, H. (2016). Mobile phone-based unobtrusive ecological momentary assessment of day-to-day mood: An explorative study. *Journal of medical Internet research*, 18(3):e72.
- Aung, M. H., Matthews, M., and Choudhury, T. (2017). Sensing behavioral symptoms of mental health and delivering personalized interventions using mobile technologies. *Depression and anxiety*, 34(7):603–609.
- Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1):1–48.
- Beedie, C., Terry, P., and Lane, A. (2005). Distinctions between emotion and mood. *Cognition & Emotion*, 19(6):847–878.
- Bennik, E. C. (2015). *Every dark cloud has a colored lining: The relation between positive and negative affect and reactivity to positive and negative events*. University of Groningen, Groningen.
- Blaauw, F. J., Schenk, H. M., Jeronimus, B. F., van der Krieke, L., de Jonge, P., Aiello, M., and Emerencia, A. C. (2016). Let's get Physiqual – An intuitive and generic method to combine sensor technology with ecological momentary assessments. *Journal of Biomedical Informatics*.
- Bolger, N. and Laurenceau, J.-p. (2013). Introduction to intensive methods. In *Intensive longitudinal methods: An introduction to diary and experience sampling research*. Guilford Press, New York.
- Booij, S. (2015). *Dynamics of the human stress system in depression: A combined population- and person-based approach to assess long-term changes and daily life fluctuations*. University of Groningen, Groningen.
- Borsboom, D. (2016). A network theory of mental disorders. *World Psychiatry*, 16(1):5–13.
- Borsboom, D. and Cramer, A. O. J. (2013). Network analysis: An integrative approach to the structure of psychopathology. *Annual Review of Clinical Psychology*, 9(1):91–121.
- Burton, C., McKinstry, B., Tătar, A. S., Serrano-Blanco, A., Pagliari, C., and Wolters, M. (2013). Activity monitoring in patients with depression: A systematic review. *Journal of affective disorders*, 145(1):21–28.
- Buyssse, D. J., Thompson, W., Scott, J., Franzen, P. L., Germain, A., Hall, M., Moul, D. E., Nofzinger, E. A., and Kupfer, D. J. (2007). Daytime symptoms in primary insomnia: A prospective analysis using ecological momentary assessment. *Sleep Medicine*, 8(3):198–208.
- Calenge, C. (2006). The package adehabitat for the r software: Tool for the analysis of space and habitat use by animals. *Ecological Modelling*, 197(3-4):516–519.

- CCMO (2018). Research with a medical device. Retrieved May 25 2018, from <http://www.ccmo.nl/en/research-with-a-medical-device>.
- Choi, L., Beck, C., Liu, Z., Matthews, C. E., and Buchowski, M. S. (2018). *PhysicalActivity: Process Accelerometer Data for Physical Activity Measurement*. R package version 0.2-2.
- Collip, D., Oorschot, M., Thewissen, V., Van Os, J., Bentall, R., and Myin-Germeys, I. (2011). Social world interactions: how company connects to paranoia. *Psychological medicine*, 41(5):911–921.
- Conner, T. S. and Feldman Barrett, L. (2012). Trends in ambulatory self-report: The role of momentary experience in psychosomatic medicine. *Psychosomatic Medicine*, 74(4):327–337.
- Conner, T. S. and Lehman, B. J. (2012). Getting started: Launching a study in daily life. In Mehl, M. R. and Conner, T. S., editors, *Handbook of research methods for studying daily life*., pages 89–107. Guilford Press, New York, NY, US.
- Conner, T. S., Tennen, H., Fleeson, W., and Barrett, L. F. (2009). Experience sampling methods: A modern idiographic approach to personality research. *Social and personality psychology compass*, 3(3):292–313.
- Cornet, V. P. and Holden, R. J. (2018). Systematic review of smartphone-based passive sensing for health and wellbeing. *Journal of Biomedical Informatics*, 17(1):120–132.
- Cranford, J. A., Shrout, P. E., Iida, M., Rafaeli, E., Yip, T., and Bolger, N. (2006). A procedure for evaluating sensitivity to within-person change: Can mood measures in diary studies detect change reliably? *Personality and Social Psychology Bulletin*, 32(7):917–929.
- Csikszentmihalyi, M. and Larson, R. (2014). Validity and reliability of the experience-sampling method. In *Flow and the Foundations of Positive Psychology: The Collected Works of Mihaly Csikszentmihalyi*, pages 35–54. Springer Netherlands.
- Dakos, V., Scheffer, M., van Nes, E. H., Brovkin, V., Petoukhov, V., and Held, H. (2008). Slowing down as an early warning signal for abrupt climate change. TL - 105. *Proceedings of the National Academy of Sciences of the United States of America*, 105 VN -(38):14308–14312.
- de Geus, E. and van Doornen, L. (1996). Ambulatory assessment of parasympathetic/sympathetic balance by impedance cardiography. In *Ambulatory Assessment. Computer assisted psychological and psychophysiological methods in monitoring and field studies*, pages 141–164. Hogrefe & Huber.
- Derogatis, L. R. (1994). *Symptom Checklist-90-R (SCL-90-R): Administration, scoring, and procedures manual*. Minneapolis, MN: NCS Pearson.
- Desmet, P. M. A., Vastenburg, M. H., and Romero, N. (2016). Mood measurement with Pick-A-Mood: Review of current methods and design of a pictorial self-report scale. *Journal of Design Research*, 14(3):241–279.
- Dogan, E., Sander, C., Wagner, X., Hegerl, U., and Kohls, E. (2017). Smartphone-based monitoring of objective and subjective data in affective disorders: Where are we and where are we going? systematic review. *Journal of medical Internet research*, 19(7).
- Dubad, M., Winsper, C., Meyer, C., Livanou, M., and Marwaha, S. (2018). A systematic review of the psychometric properties, usability and clinical impacts of mobile mood-monitoring applications in young people. *Psychological medicine*, 48(2):208–228.
- Dunton, G. F. (2017). Ecological momentary assessment in physical activity research. *Exercise and Sport Sciences Reviews*, 45(1):48–54.
- Ebner-Priemer, U. W. and Trull, T. J. (2009). Ecological momentary assessment of mood disorders and mood dysregulation. *Psychological Assessment*, 21(4):463–475.
- Ekker, A. and van Rest, B. (2013). *Medische apps, is certificeren nodig? In 7 stappen naar een CE-markering voor uw app*. Nictiz, Den Haag.

- Ellis-Davies, K., Sakkalou, E., Fowler, N. C., Hilbrink, E. E., and Gattis, M. (2012). CUE: The continuous unified electronic diary method. *Behavior Research Methods*, 44(4):1063–1078.
- Emerencia, A. (2018). *autovarCore: Automated Vector Autoregression Models and Networks*. R package version 1.0-4.
- Emerencia, A. C., Van Der Krieke, L., Bos, E. H., De Jonge, P., Petkov, N., and Aiello, M. (2016). Automating vector autoregression on electronic patient diary data. *IEEE Journal of Biomedical and Health Informatics*.
- Engel, S. G., Crosby, R. D., Thomas, G., Bond, D., Lavender, J. M., Mason, T., Steffen, K. J., Green, D. D., and Wonderlich, S. A. (2016). Ecological momentary assessment in eating disorder and obesity research: A review of the recent literature.
- Epskamp, S. (2017). *graphicalVAR: Graphical VAR for Experience Sampling Data*. R package version 0.2.1.
- Epskamp, S., Borsboom, D., and Fried, E. I. (2018a). Estimating psychological networks and their accuracy: A tutorial paper. *Behavior Research Methods*, 50(1):195–212.
- Epskamp, S., Borsboom, D., and Fried, E. I. (2018b). Estimating psychological networks and their accuracy: A tutorial paper. *Behavior Research Methods*, 50(1):195–212.
- Epskamp, S., Cramer, A. O. J., Waldorp, L. J., Schmittmann, V. D., and Borsboom, D. (2012). qgraph: Network visualizations of relationships in psychometric data. *Journal of Statistical Software*, 48(4):1–18.
- Epskamp, S. and Fried, E. I. (2018). A tutorial on regularized partial correlation networks. *Psychological Methods*.
- Epskamp, S., van Borkulo, C. D., van der Veen, D. C., Servaas, M. N., Isvoranu, A.-M., Riese, H., and Cramer, A. O. J. (2018c). Personalized network modeling in psychopathology: The importance of contemporaneous and temporal connections. *Clinical Psychological Science*, 6(3):416–427.
- Evenson, K. R., Goto, M. M., and Furberg, R. D. (2015). Systematic review of the validity and reliability of consumer-wearable activity trackers. *International Journal of Behavioral Nutrition and Physical Activity*, 12(1):159.
- Fang, Z., Langford, J., and Sweetland, C. (2018). *GENERead: Read GENEVA Binary Files*. R package version 2.0.2.
- Field, A. (2013). *Discovering Statistics Using IBM SPSS Statistics*. Sage Publications Ltd., 4th edition.
- Field, A., Miles, J., and Field, Z. (2012). *Discovering Statistics Using R*. SAGE Publications Ltd, 1 edition.
- Goërtz, Y. M. J., Looijmans, M., Prins, J. B., Janssen, D. J. A., Thong, M. S. Y., Peters, J. B., Burtin, C., Meertens-Kerris, Y., Coors, A., Muris, J. W. M., Sprangers, M. A. G., Wouters, E. F. M., Vercoulen, J. H., and Spruit, M. A. (2018). Fatigue in patients with chronic obstructive pulmonary disease: Protocol of the Dutch multicentre, longitudinal, observational <i>FAntasTIGUE</i> study. *BMJ Open*, 8(4):e021745.
- Goldfeld, K. (2018). *simstudy: Simulation of Study Data*. R package version 0.1.9.
- Green, P. and MacLeod, C. J. (2016). simr: An r package for power analysis of generalised linear mixed models by simulation. *Methods in Ecology and Evolution*, 7(4):493–498.
- Grolemund, G. and Wickham, H. (2011). Dates and times made easy with lubridate. *Journal of Statistical Software*, 40(3):1–25.
- Groot, P. C. (2010). Patients can diagnose too: How continuous self-assessment aids diagnosis of, and recovery from, depression. *Journal of Mental Health*, 19(4):352–362.
- Haedt-Matt, A. A. and Keel, P. K. (2011). Hunger and binge eating: A meta-analysis of studies using ecological momentary assessment. *International Journal of Eating Disorders*, 44(7):573–578.

- Haedt-Matt, A. A., Zalta, A. K., Forbush, K. T., and Keel, P. K. (2012). Experimental evidence that changes in mood cause changes in body dissatisfaction among undergraduate women. *Body image*, 9(2):216–220.
- Hamaker, E. (2012). Why researchers should think "within-person": A paradigmatic rationale. *Handbook of Research Methods for Studying Daily Life*, pages 43–61.
- Hufford, M. R., Shields, A. L., Shiffman, S., Paty, J., and Balabanis, M. (2002). Reactivity to ecological momentary assessment: An example using undergraduate problem drinkers. *Psychology of Addictive Behaviors*, 16(3):205–211.
- Jacobson, N. S. and Truax, P. (1991). Clinical significance: A statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology*, 59(1):12–19.
- Jongeneel, A., Scheffers, D., Tromp, N., Nuij, C., Delespaul, P., Riper, H., Van Der Gaag, M., and van den Berg, D. (2018). Reducing distress and improving social functioning in daily life in people with auditory verbal hallucinations: Study protocol for the 'temstem'randomised controlled trial. *BMJ open*, 8(3):e020537.
- Kaida, K., Takahashi, M., Åkerstedt, T., Nakata, A., Otsuka, Y., Haratani, T., and Fukasawa, K. (2006). Validation of the karolinska sleepiness scale against performance and eeg variables. *Clinical Neurophysiology*, 117(7):1574–1581.
- Kemmeren, L., Van Schaik, D., Riper, H., Kleiboer, A., Bosmans, J. E., and Smit, J. H. (2016). Effectiveness of blended depression treatment for adults in specialised mental healthcare: study protocol for a randomised controlled trial. *BMC psychiatry*, 16(1):113.
- Kirchner, T. R. and Shiffman, S. (2013). Ecological Momentary Assessment. In *The Wiley-Blackwell Handbook of Addiction Psychopharmacology*, pages 541–565. Wiley-Blackwell.
- Kirchner, T. R. and Shiffman, S. (2016). Spatio-temporal determinants of mental health and well-being: Advances in geographically-explicit ecological momentary assessment (gema). *Social psychiatry and psychiatric epidemiology*, 51(9):1211–1223.
- Kleiboer, A., Smit, J., Bosmans, J., Ruwaard, J., Andersson, G., Topooco, N., Berger, T., Krieger, T., Botella, C., Baños, R., Chevreul, K., Araya, R., Cerga-Pashoja, A., Cieślak, R., Rogala, A., Vis, C., Draisma, S., Schaik, A., Kemmeren, L., Ebert, D., Berking, M., Funk, B., Cuijpers, P., and Riper, H. (2016). European COMPARE Effective research on blended Depression treatment versus treatment-as-usual (E-COMPARED): Study protocol for a randomized controlled, non-inferiority trial in eight European countries. *Trials*, 17(1).
- Kossakowski, J. J., Groot, P. C., Haslbeck, J. M. B., Borsboom, D., and Wichers, M. (2017). Data from 'critical slowing down as a personalized early warning signal for depression'. *Journal of Open Psychology Data*, 5(1).
- Krieke, L. V. D., Jeronimus, B. F., Blaauw, F. J., Wanders, R. B., Emerencia, A. C., Schenk, H. M., Vos, S. D., Snippe, E., Wichers, M., Wigman, J. T., Bos, E. H., Wardenaar, K. J., and De Jonge, P. (2016). HowNutsAreTheDutch (HoeGekIsNL): A crowdsourcing study of mental symptoms and strengths. *International Journal of Methods in Psychiatric Research*.
- Kroenke, K., Strine, T. W., Spitzer, R. L., Williams, J. B., Berry, J. T., and Mokdad, A. H. (2009). The PHQ-8 as a measure of current depression in the general population. *Journal of Affective Disorders*, 114(1-3):163–173.
- Larson, R. and Csikszentmihalyi, M. (1983). The experience sampling method. In Reis, H. T., editor, *Naturalistic approaches to studying social interaction*, pages 41–56. Jossey-Bass Inc Pub, San Francisco.
- Lomb, N. R. (1976). Least-squares frequency analysis of unequally spaced data. *Astrophysics and Space Science*.

- Marszalek, J., Morgulec-Adamowicz, N., Rutkowska, I., Kosmol, A., Marszalek, J., Morgulec-Adamowicz, N., Rutkowska, I., and Kosmol, A. (2014). Using ecological momentary assessment to evaluate current physical activity, using ecological momentary assessment to evaluate current physical activity. *BioMed Research International*, 2014,:e915172.
- Mehl, M. R. and Conner, T. S., editors (2012). *Handbook of research methods for studying daily life*. Guilford Press, New York, NY, US.
- Moore, R. C., Depp, C. A., Wetherell, J. L., and Lenze, E. J. (2016). Ecological momentary assessment versus standard assessment instruments for measuring mindfulness, depressed mood, and anxiety among older adults. *Journal of Psychiatric Research*, 75:116–123.
- Myin-Germeys, I., Klippen, A., Steinhart, H., and Reininghaus, U. (2016). Ecological momentary interventions in psychiatry. *Current opinion in psychiatry*, 29(4):258–63.
- Nuij, C., van Ballegooijen, W., Ruwaard, J., de Beurs, D., Mokkenstorm, J., van Duijn, E., de Winter, R. F., O'Connor, R. C., Smit, J. H., Riper, H., and Kerkhof, A. (2018). Smartphone-based safety planning and self-monitoring for suicidal patients: Rationale and study protocol of the CASPAR (Continuous Assessment for Suicide Prevention And Research) study. *Internet Interventions*, 13:16–23.
- Patel, S., Park, H., Bonato, P., Chan, L., and Rodgers, M. (2012). A review of wearable sensors and systems with application in rehabilitation. *Journal of neuroengineering and rehabilitation*, 9(1):21.
- Penninx, B. W. J. H., Beekman, A. T. F., Smit, J. H., Zitman, F. G., Nolen, W. A., Spinhover, P., Cuijpers, P., de Jong, P. J., van Marwijk, H. W. J., Assendelft, W. J. J., van der Meer, K., Verhaak, P., Wensing, M., de Graaf, R., Hoogendoijk, W. J., Ormel, J., van Dyck, R., and NESDA Research Consortium (2008). The Netherlands Study of Depression and Anxiety (NESDA): Rationale, objectives and methods. *International journal of methods in psychiatric research*, 17(3):121–40.
- Piasecki, T. M., Hufford, M. R., Solhan, M., and Trull, T. J. (2007). Assessing clients in their natural environments with electronic diaries: Rationale, benefits, limitations, and barriers. *Psychological Assessment*, 19(1):25–43.
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., and R Core Team (2018). *nlme: Linear and Nonlinear Mixed Effects Models*. R package version 3.1-137.
- Pinheiro, J. C. and Bates, D. M. (2000). *Mixed effects models in S and S-Plus*. Springer Verlag.
- Posner, J., Russell, J. A., and Peterson, B. S. (2005). The circumplex model of affect: An integrative approach to affective neuroscience, cognitive development, and psychopathology. *Development and Psychopathology*, 17(3):715–734.
- Pot-Kolder, R., Veling, W., Geraets, C., and van der Gaag, M. (2016). Effect of virtual reality exposure therapy on social participation in people with a psychotic disorder (vretp): Study protocol for a randomized controlled trial. *Trials*, 17(1):25.
- Pot-Kolder, R. M., Geraets, C. N., Veling, W., van Beilen, M., Staring, A. B., Gijsman, H. J., Delespaul, P. A., and van der Gaag, M. (2018). Virtual-reality-based cognitive behavioural therapy versus waiting list control for paranoid ideation and social avoidance in patients with psychotic disorders: a single-blind randomised controlled trial. *The Lancet Psychiatry*, 5(3):217–226.
- Rijsbergen, G. D. V., Bockting, C. L. H., Berking, M., Koeter, M. W. J., and Schene, A. H. (2012). Can a one-item mood scale do the trick? Predicting relapse over 5-years in recurrent depression. *Plos One*, 7(10):1–5.
- Roenneberg, T., Wirz-Justice, A., and Merrow, M. (2003). Life between clocks: daily temporal patterns of human chronotypes. *Journal of biological rhythms*, 18(1):80–90.
- Ruf, T. (1999). The Lomb-Scargle periodogram in biological rhythm research: Analysis of incomplete and unequally spaced time-series. *Biological Rhythm Research*, 30:178–201.

- Russell, J. A. (1980). A circumplex model of affect. *Journal of personality and social psychology*, 39(6):1161.
- Saeb, S., Zhang, M., Karr, C. J., Schueller, S. M., Corden, M. E., Kording, K. P., and Mohr, D. C. (2015). Mobile phone sensor correlates of depressive symptom severity in daily-life behavior: An exploratory study. *Journal of Medical Internet Research*, 17(7).
- Sander, D. and Scherer, K. (2009). *Oxford companion to emotion and the affective sciences*. Oxford University Press.
- Scheffer, M., Bascompte, J., Brock, W. A., Brovkin, V., Carpenter, S. R., Dakos, V., Held, H., Van Nes, E. H., Rietkerk, M., and Sugihara, G. (2009). Early-warning signals for critical transitions.
- Schueller, S. M., Aguilera, A., and Mohr, D. C. (2017). Ecological momentary interventions for depression and anxiety. *Depression and anxiety*, 34(6):540–545.
- Schuurman, N. K., Houtveen, J. H., and Hamaker, E. L. (2015). Incorporating measurement error in n= 1 psychological autoregressive modeling. *Frontiers in psychology*, 6:1038.
- Serre, F., Fatseas, M., Swendsen, J., and Auriacombe, M. (2015). Ecological momentary assessment in the investigation of craving and substance use in daily life: A systematic review.
- Sharar, S. R., Alamdari, A., Hoffer, C., Hoffman, H. G., Jensen, M. P., and Patterson, D. R. (2016). Circumplex model of affect: A measure of pleasure and arousal during virtual reality distraction analgesia. *Games for health journal*, 5(3):197–202.
- Shiffman, S. (2009). Ecological momentary assessment (EMA) in studies of substance use. *Psychological Assessment*, 21(4):486–497.
- Shiffman, S., Stone, A. A., Hufford, M. R., Rev, A., Psychol, C., Shiffman, S., Stone, A. A., and Hufford, M. R. (2008). Ecological Momentary Assessment. *Annual Review of Clinical Psychology*, 4(1):1–32.
- Slofstra, C., Klein, N. S., Nauta, M. H., Wichers, M., Batalas, N., and Bockting, C. L. (2017). Imagine your mood: Study design and protocol of a randomized controlled micro-trial using app-based experience sampling methodology to explore processes of change during relapse prevention interventions for recurrent depression. *Contemporary clinical trials communications*, 7:172–178.
- Smyth, J. M. and Stone, A. A. (2003). Ecological momentary assessment research in behavioral medicine. *Journal of Happiness Studies*, 4(1):35–52.
- Starr, L. R. and Davila, J. (2012). Temporal patterns of anxious and depressed mood in generalized anxiety disorder: A daily diary study. *Behaviour Research and Therapy*.
- Stone, A. A. and Shiffman, S. (2002). Capturing momentary, self-report data: A proposal for reporting guidelines. *Annals of Behavioral Medicine*, 24(3):236–243.
- Swendsen, J. (2016). Contributions of mobile technologies to addiction research. *Dialogues in clinical neuroscience*, 18(2):213.
- Te Lindert, B. H., Itzhacki, J., van der Meijden, W. P., Kringselbach, M. L., Mendoza, J., and Van Someren, E. J. (2018). Bright environmental light ameliorates deficient subjective ‘liking’ in insomnia: An experience sampling study. *Sleep*, 41(4):zsy022.
- Telford, C., McCarthy-Jones, S., Corcoran, R., and Rowse, G. (2012). Experience sampling methodology studies of depression: The state of the art. *Psychological Medicine*, 42(6):1119–1129.
- van Ballegooijen, W., Ruwaard, J., Karyotaki, E., Ebert, D. D., Smit, J. H., and Riper, H. (2016). Reactivity to smartphone-based ecological momentary assessment of depressive symptoms (MoodMonitor): Protocol of a randomised controlled trial. *BMC Psychiatry*, 16(1):359.

- van de Ven, P., O'Brien, H., Henriques, R., Klein, M., Msetfi, R., Nelson, J., Rocha, A., Ruwaard, J., O'Sullivan, D., and Riper, H. (2017). ULTEMAT: A mobile framework for smart ecological momentary assessments and interventions. *Internet Interventions*, 9:74–81.
- Van Der Krieke, L., Blaauw, F. J., Emerencia, A. C., Schenk, H. M., Slaets, J. P., Bos, E. H., De Jonge, P., and Jeronimus, B. F. (2017). Temporal dynamics of health and well-being: A crowdsourcing approach to momentary assessments and automated generation of personalized feedback. *Psychosomatic Medicine*.
- van der Krieke, L., Emerencia, C. A., Bos, H. E., Rosmalen, G. J., Riese, H., Aiello, M., Sytema, S., and de Jonge, P. (2015). Ecological momentary assessments and automated time series analysis to promote tailored health care: A proof-of-principle study. *JMIR Research Protocols*, 4(3):e100.
- Van Der Meijden, W. P., Van Someren, J. L., Te Lindert, B. H., Bruijel, J., Van Oosterhout, F., Coppens, J. E., Kalsbeek, A., Cajochen, C., Bourgin, P., and Van Someren, E. J. (2016). Individual differences in sleep timing relate to melanopsin-based phototransduction in healthy adolescents and young adults. *Sleep*, 39(6):1305–1310.
- van Hees, V. T., Fang, Z., Langford, J., Assah, F., Mohammad, A., da Silva, I. C., Trenell, M. I., White, T., Wareham, N. J., and Brage, S. (2014). Autocalibration of accelerometer data for free-living physical activity assessment using local gravity and temperature: An evaluation on four continents. *Journal of Applied Physiology*, 117(7):738–744.
- van Hees, V. T., Fang, Z., Zhao, J. H., Heywood, J., Mirkes, E., Sabia, S., and Migueles, J. H. (2018). *GGIR: Raw Accelerometer Data Analysis*. R package version 1.5-24.
- van Hees, V. T., Sabia, S., Anderson, K. N., Denton, S. J., Oliver, J., Catt, M., Abell, J. G., Kivimaki, M., Trenell, M. I., and Singh-Manoux, A. (2015). A novel, open access method to assess sleep duration using a wrist-worn accelerometer. *PLoS One*, 10(11).
- van Rijsbergen, G. D., Burger, H., Hollon, S. D., Elgersma, H. J., Kok, G. D., Dekker, J., de Jong, P. J., and Bockting, C. L. (2014). How do you feel? Detection of recurrent Major Depressive Disorder using a single-item screening tool. *Psychiatry Research*, 220(1-2):287–293.
- Venables, W. N. and Ripley, B. D. (2002). *Modern Applied Statistics with S*. Springer, New York, fourth edition. ISBN 0-387-95457-0.
- Versluis, A., Verkuil, B., Spinhoven, P., van der Ploeg, M. M., and Brosschot, J. F. (2016). Changing mental health and positive psychological well-being using ecological momentary interventions: A systematic review and meta-analysis. *Journal of medical Internet research*, 18(6).
- Walz, L. C., Nauta, M. H., and aan het Rot, M. (2014). Experience sampling and ecological momentary assessment for studying the daily lives of patients with anxiety disorders: A systematic review. *Journal of Anxiety Disorders*, 28(8):925–937.
- Warmerdam, L., Riper, H., Klein, M. C., van de Ven, P., Rocha, A., Henriques, M. R., Tousset, E., Silva, H., Andersson, G., and Cuijpers, P. (2012). Innovative ict solutions to improve treatment outcomes for depression: the ict4depression project. *Annual Review of Cybertherapy and Telemedicine*, 181(1):339–343.
- Watson, D. and Clark, L. A. (1994). *The PANAS-X : Manual for the Positive and Negative Affect Schedule - expanded Form*. University of Iowa.
- Watson, D. and Clark, L. A. (1997). Measurement and mismeasurement of mood: Recurrent and emergent issues.
- Watson, D., Clark, L. A., and Tellegen, A. (1988). Development and validation of brief measures of Positive and Negative Affect : The PANAS scales. *Journal of Personality and Social Psychology*, 54(6):1063–1070.
- Watson, D. and Tellegen, A. (1985). Toward a consensual structure of mood. *Psychological bulletin*, 98(2):219.

- Wenze, S. J. and Miller, I. W. (2010). Use of ecological momentary assessment in mood disorders research. *Clinical psychology review*, 30(6):794–804.
- Wichers, M., Groot, P. C., Psychosystems, ESM, and EWS (2016). Critical slowing down as a personalized early warning signal for depression. *Psychotherapy and Psychosomatics*, 85(2):114–116.
- Wichers, M., Peeters, F., Rutten, B. P. F., Jacobs, N., Derom, C., Thiery, E., Delespaul, P., and van Os, J. (2012). A time-lagged momentary assessment study on daily life physical activity and affect. *Health Psychology*, 31(2):135–144.
- Wichers, M., Simons, C. J. P., Kramer, I. M. A., Hartmann, J. A., Lothmann, C., Myin-Germeys, I., van Bemmel, A. L., Peeters, F., Delespaul, P., and van Os, J. (2011). Momentary assessment technology as a tool to help patients with depression help themselves. *Acta Psychiatrica Scandinavica*, 124(4):262–272.
- Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. Springer.
- Wickham, H., François, R., Henry, L., and Müller, K. (2018). *dplyr: A Grammar of Data Manipulation*. R package version 0.7.6.
- Wickham, H. and Grolemund, G. (2016). *R for data science: Import, tidy, transform, visualize, and model data*. " O'Reilly Media, Inc. ".
- Wickham, H. and Miller, E. (2018). *haven: Import and Export 'SPSS', 'Stata' and 'SAS' Files*. R package version 1.1.2.
- Wilkinson, L. (2006). *The grammar of graphics*. Springer Science & Business Media.
- Xie, Y. (2016). *bookdown: Authoring Books and Technical Documents with R Markdown*. Chapman and Hall/CRC, Boca Raton, Florida. ISBN 978-1138700109.
- Xie, Y. (2018). *bookdown: Authoring Books and Technical Documents with R Markdown*. R package version 0.7.

Index

- (, 95–97, 99
Accelerometer, 45
Back-office, 27
CASPAR, 88
Chronic Fatigue, 85
Circumplex model, 39
CRAN, 15
Datamanagement, 31
 Analysis, 33
 Import and prune, 33
 Keyfile, 32
 Project directory structure, 31
 Source data, 33
 Visual exploration, 33
datetime variables, 105
E-COMPARED, 82
Ecological validity, 4
EMA, 3
 Active EMA, 3, 5, 21
 Observational EMA, *see* Passive EMA
 Passive EMA, 3, 6, 21
 Self-report EMA, *see* Active EMA
EMA instruments, 27
EMA research
 Activity patterns, 37
 Anxiety disorders, 6
 Eating disorders, 6
 Mental health/ Well-being, 6
 Mood disorders, 6, 37
 Somatic health, 37
 Substance-related disorders, 6, 37
EMA synonyms, 3
 Ambulatory Assessment, 3
 Ambulatory Self-reporting, 3
 Continuous Unified Electronic Diary Method, 3
 Experience Sampling, 3
 Intensive-longitudinal Study Design, 3
 Real-time Data Capturing, 3
FAntasTIGUE, 86
Fitbit, 28
GENEActiv, 28
GitHub, 15
GPS, 47
GraphicalVar, 63
ICT4Depression, 82
Ideographic research, 5
Imagine your Mood study, 82
IMPACT, 86
Intra-individual process, *see* Ideographic research
latitude, 47
Limitations, 6
 Feasibility, 6
 Practice effect, 6
 Privacy, 6
 Selected samples, 6
Lomb-Scargle periodogram, 110
longitude, 47
Measurement error, 21
MERITS study, 87
Methods, 21
 Ethical considerations, 25
 Power analysis, 23
 Study design, 21
Mixed Modeling, 57
Mood, 37
Mood assessment, 37
 Bag-of-Items, 38
 Bipolar unidimensional, 38
 Circumplex model, 39
 Multi-dimensional, 39
 PA/NA, 41
 Pick-a-mood scale, 39
 Unidimensional, 37
Mood definitions, 37
 Emotions, 37
 Mood states, 37
Moodbuster, 82
NESDA, 83
Nomothetic research, 5
Packages
 autovar, 108

autovarCore, 108
bootnet, 109
dplyr, 103
GGIR, 103
ggplot2, 104
haven, 105
lme4, 106
lubridate, 105
nlme, 106
PhysicalActivity, 103
qgraph, 109
simr, 106
tidyverse, 103
Positive and Negative Affect, 41
PANAS, *see* Mood assessment
Psycho-Systems, 90

R and RStudio, 11
 Console, 12
 Data access, 14
 Data import, 13
 Help, 16
 Installing, 12
 Packages, 16
 Scripts, 13
 Simulating data, 57
 Why R, 11
R packages, 15
RADAR-CNS, 84
Recall bias, 4

Sample plan, *see* Sampling
Sampling, 22
 Event-contingent, 4
 Signal-contingent, 4
Sleep, 91
SPSS, 105
StayFine study, 82
Symptom networks, 63
symptom networks, 107

TEMSTEM, 89

vector autoregressive models, 108
VU-AMS, 92