

ELM Suppression and Pedestal Structure in I-Mode Plasmas on Alcator C-Mod

by

John Reel Walk, Jr.

S.B. Physics & Mathematics (2010)
Massachusetts Institute of Technology

Submitted to the Department of Nuclear Science and Engineering
in partial fulfillment of the requirements for the degree of

Doctor of Philosophy in Applied Plasma Physics

at the

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

June 2014

© Massachusetts Institute of Technology 2014. All rights reserved.

Author
Department of Nuclear Science and Engineering
December 24, 2013

Certified by.....
Jerry W. Hughes
Research Scientist, Plasma Science and Fusion Center
Thesis Supervisor

Certified by.....
Dennis G. Whyte
Professor, Department of Nuclear Science and Engineering
Thesis Reader

Accepted by
Mujid S. Kazimi
TEPCO Professor of Nuclear Engineering
Chair, Department Committee on Graduate Students

ELM Suppression and Pedestal Structure in I-Mode Plasmas on Alcator C-Mod

by
John Reel Walk, Jr.

Submitted to the Department of Nuclear Science and Engineering
on December 24, 2013, in partial fulfillment of the
requirements for the degree of
Doctor of Philosophy in Applied Plasma Physics

Abstract

Abstract goes here.

Thesis Supervisor: Jerry W. Hughes
Research Scientist, Plasma Science and Fusion Center

Thesis Reader: Dennis G. Whyte
Professor, Department of Nuclear Science and Engineering

A dedication goes here maybe?

ACKNOWLEDGMENTS

Acknowledgements go here.

CONTENTS

1	INTRODUCTION	11
1.1	Plasmas for Fusion	11
1.1.1	Plasma Parameters	11
1.1.2	Fusion Fuels	11
1.2	Magnetic Confinement	12
1.2.1	Basic Principles	12
1.2.2	Toroidal Configurations	12
1.3	Alcator C-Mod	12
1.4	Confinement & Transport	12
1.4.1	Global Confinement	12
1.4.2	Transport Barriers	12
1.5	High-Performance Regimes	12
1.5.1	ELMy H-Mode	12
1.5.2	EDA H-Mode	12
1.5.3	I-Mode	12
1.6	Goals & Outline	12

LIST OF FIGURES

Figure 1	Binding energy per nucleon versus nuclear mass number.	12
----------	--	----

LIST OF TABLES

INTRODUCTION

1.1 PLASMAS FOR FUSION

1.1.1 Plasma Parameters

1.1.2 Fusion Fuels

Fusion collectively refers to the class of nuclear reactions merging lighter nuclei into a single heavier element. While fusion reactions for elements lighter than iron are generally exothermic, the most common and readily attainable involve isotopes of hydrogen or helium, the most promising candidates for which are shown below.



Here D and T indicate nuclei of deuterium and tritium, two heavy isotopes of hydrogen (one proton plus one and two neutrons, respectively).

Pure deuterium fuel (reactions shown in eqs. 1-2) is attractive from a research standpoint, due to the abundance and ease of use of deuterium. Deuterium is a stable nucleus, obviating the need for radiation safety on the fuel system, and is naturally occurring in relative abundance (approximately 1/6000 of hydrogen nuclei on earth are deuterium [cite](#)), allowing harvesting of deuterium fuel from seawater. However, pure-deuterium reactions suffer from low energy output per reaction [ref cross-section graphic](#) and a significantly lower reaction rate at feasible plasma conditions compared to other fuel options, setting high performance requirements for a putative DD-burning reactor.

[reaction
cross-section graphic](#)




Figure 1: Binding energy per nucleon versus nuclear mass number.

1.2 MAGNETIC CONFINEMENT

1.2.1 *Basic Principles*

1.2.2 *Toroidal Configurations*

1.3 ALCATOR C-MOD

1.4 CONFINEMENT & TRANSPORT

1.4.1 *Global Confinement*

1.4.2 *Transport Barriers*

1.5 HIGH-PERFORMANCE REGIMES

1.5.1 *ELMy H-Mode*

1.5.2 *EDA H-Mode*

1.5.3 *I-Mode*

1.6 GOALS & OUTLINE

COLOPHON

This document was typeset using `classicthesis` developed by André Miede (although aspects were changed to comply with the MIT thesis standards and the author’s personal preferences). The style was inspired by Robert Bringhurst’s seminal book on typography “*The Elements of Typographic Style*”. `classicthesis` is available for both \LaTeX and \LyX :

<http://code.google.com/p/classicthesis/>

Final Version as of December 24, 2013 (`classicthesis` version 4.1).

Hermann Zapf’s *Palatino* and *Euler* type faces (Type 1 PostScript fonts *URW Palladio L* and *FPL*) are used. The “typewriter” text is typeset in *FPL*, originally developed by Bitstream, Inc. as “Bitstream Vera”. (Type 1 PostScript fonts were made available by Malte Rosenau and Ulrich Dirr.)