# ELM Suppression and Pedestal Structure in I-Mode Plasmas on Alcator C-Mod

by

John Reel Walk, Jr.

S.B. Physics & Mathematics (2010) Massachusetts Institute of Technology

Submitted to the Department of Nuclear Science and Engineering in partial fulfillment of the requirements for the degree of

Doctor of Philosophy in Applied Plasma Physics

at the

### MASSACHUSETTS INSTITUTE OF TECHNOLOGY

June 2014

 $\hbox{$\mathbb C$}$  Massachusetts Institute of Technology 2014. All rights reserved.

uthor
Jerry W. Hughe Research Scientist, Plasma Science and Fusion Cente Thesis Superviso
Pertified by
ccepted by

# ELM Suppression and Pedestal Structure in I-Mode Plasmas on Alcator C-Mod by

John Reel Walk, Jr.

Submitted to the Department of Nuclear Science and Engineering on December 24, 2013, in partial fulfillment of the requirements for the degree of Doctor of Philosophy in Applied Plasma Physics

Abstract

Abstract goes here.

Thesis Supervisor: Jerry W. Hughes

Research Scientist, Plasma Science and Fusion Center

Thesis Reader: Dennis G. Whyte

Professor, Department of Nuclear Science and Engineering



# ACKNOWLEDGMENTS

Acknowledgements go here.

## CONTENTS

1	INT	RODUCTION 11			
	1.1	Plasmas for Fusion 11			
		1.1.1 Plasma Parameters 11			
		1.1.2 Fusion Fuels 11			
	1.2	Magnetic Confinement 12			
		1.2.1 Basic Principles 12			
		1.2.2 Toroidal Configurations 12			
	1.3	Alcator C-Mod 12			
	1.4	Confinement & Transport 12			
		1.4.1 Global Confinement 12			
		1.4.2 Transport Barriers 12			
	1.5	High-Performance Regimes 12			
		1.5.1 ELMy H-Mode 12			
		1.5.2 EDA H-Mode 12			
		1.5.3 I-Mode 12			
	1.6	Goals & Outline 12			

# LIST OF FIGURES

Figure 1 Binding energy per nucleon versus atomic mass number, with notable isotopes marked. 11

## LIST OF TABLES

INTRODUCTION

# 1

#### 1.1 PLASMAS FOR FUSION

#### 1.1.1 Plasma Parameters

#### 1.1.2 Fusion Fuels

Fusion collectively refers to the class of nuclear reactions merging lighter nuclei into a single heavier element. While fusion reactions for elements lighter than iron are generally exothermic, the most common and readily attainable involve isotopes of hydrogen or helium, the most promising candidates for which are shown below.

$$^{2}D + ^{2}D \rightarrow ^{3}T + p + 4.03 \text{ MeV}$$
 (1)

$$^{2}D + ^{2}D \rightarrow ^{3}He + n + 3.27 \text{ MeV}$$
 (2)

$$^{2}D + ^{3}He \rightarrow ^{4}He + p + 18.3 \text{ MeV}$$
 (3)


$$^{2}D + ^{3}T \rightarrow ^{4}He + n + 17.6 \text{ MeV}$$
 (4)

Here D and T indicate nuclei of deuterium and tritium, two heavy isotopes of hydrogen (one proton plus one and two neutrons, respectively).

Pure deuterium fuel (reactions shown in eqs. 1-2) is attractive from a research standpoint, due to the abundance and ease of use of deuterium. Deuterium is a stable nucleus, obviating the need for radiation safety on the fuel system, and is naturally occurring in relative abundance (approximately 1/6000 of hydrogen nuclei on earth are

reaction cross-section graphic

Figure 1: Binding energy per nucleon versus atomic mass number, with notable isotopes marked.



deuterium cite), allowing harvesting of deuterium fuel from seawater. However, pure-deuterium reactions suffer from low energy output per reaction ref cross-section graphic and a significantly lower reaction rate at feasible plasma conditions compared to other fuel options, setting high performance requirements for a putative DD-burning reactor.

- 1.2 MAGNETIC CONFINEMENT
- 1.2.1 Basic Principles
- 1.2.2 Toroidal Configurations
- 1.3 ALCATOR C-MOD
- 1.4 CONFINEMENT & TRANSPORT
- 1.4.1 Global Confinement
- 1.4.2 Transport Barriers
- 1.5 HIGH-PERFORMANCE REGIMES
- 1.5.1 ELMy H-Mode
- 1.5.2 *EDA H-Mode*
- 1.5.3 *I-Mode*
- 1.6 GOALS & OUTLINE

#### COLOPHON

This document was typeset using classicthesis developed by André Miede (although aspects were changed to comply with the MIT thesis standards and the author's personal preferences). The style was inspired by Robert Bringhurst's seminal book on typography "The Elements of Typographic Style". classicthesis is available for both LATEX and LyX:

http://code.google.com/p/classicthesis/

Final Version as of December 24, 2013 (classicthesis version 4.1).

Hermann Zapf's *Palatino* and *Euler* type faces (Type 1 PostScript fonts *URW Palladio L* and *FPL*) are used. The "typewriter" text is typeset in *FPL*, originally developed by Bitstream, Inc. as "Bitstream Vera". (Type 1 PostScript fonts were made available by Malte Rosenau and Ulrich Dirr.)