Sample limited to studies present in this tracker · Phase: all · Status: all · RT role: all · Disease: all · Country: all · Accrual: all · Source: all · Brief: all

Contents

Brief · My trials · 7 studies · October 12, 2026 at 1:39 PM

Breast; Gynecologic 2

- NRG-BR009 (OFSET)

GU 3

- NRG-GU013 (High Five)
- NRG-GU015 (ARCHER)
- S2427 (BRIGHT)

Gynecologic 4

- NRG-GY037

Head and neck; Skin 5

- A222301 (HOPE)

NSCLC 6

- A081801

Breast; Gynecologic

NRG-BR009 (OFSET)

Phase 3 · NRG, NRG Oncology · recruiting · NCT05879926

A Phase III Adjuvant Trial Evaluating the Addition of Adjuvant Chemotherapy to Ovarian Function Suppression Plus Endocrine Therapy in Premenopausal Patients With pN0-1, ER-Positive/HER2-Negative Breast Cancer and an Oncotype Recurrence Score Less Than or Equal to 25 (OFSET)

Arm 1 Ovarian function suppression (GnRH agonist of investigator's choice, or bilateral oophorectomy) plus aromatase inhibitor for 5 years.

Arm 2 Adjuvant chemotherapy of investigator's choice (NCCN- or ESMO-listed regimen), then ovarian function suppression plus aromatase inhibitor for 5 years.

Primary endpoint. Invasive breast cancer-free survival.

GU

NRG-GU013 (High Five)

Phase 3 · NRG, NRG Oncology · recruiting · NCT05946213

The Phase III 'High Five Trial' Five Fraction Radiation For High-Risk Prostate Cancer

Arm 1 Ultrahypofractionated SBRT in 5 fractions.

Arm 2 Conventional or moderate hypofractionation in 20 to 45 fractions.

Primary endpoint. Metastasis-free survival on conventional imaging.

NRG-GU015 (ARCHER)

Phase 3 · NRG, NRG Oncology · recruiting · NCT07097142

The Phase III Adaptive Radiation and Chemotherapy for Muscle Invasive Bladder Cancer Trial (ARCHER)

Arm 1 4 weeks of radiosensitizing chemotherapy plus hypofractionated radiotherapy, 55 Gy in 20 fractions.

Arm 2 4 weeks of radiosensitizing chemotherapy plus SBRT, 32.5 Gy in 5 fractions.

S2427 (BRIGHT)

Phase 2 · SWOG, National Cancer Institute (NCI) · recruiting · NCT07061964

Single Arm Phase II Study of Bladder Preservation With Immunoradiotherapy After a Clinically Meaningful Response to Neoadjuvant Therapy in Patients With Muscle Invasive Bladder Cancer (BRIGHT)

Gynecologic

NRG-GY037

Phase 3 · NRG, CCTG, National Cancer Institute (NCI) · recruiting · NCT07061977

A Phase III Study of Induction Pembrolizumab and Chemotherapy Followed by Chemoradiation and Pembrolizumab vs Chemoradiation and Pembrolizumab Both Followed by Pembrolizumab for High Risk Locally Advanced Cervical Cancer

Head and neck; Skin

A222301 (HOPE)

Phase 3 · ALLIANCE, Alliance for Clinical Trials in Oncology, Alliance · recruiting · NCT06992427

High-Dose Prophylactic Gabapentin (HOPE) to Prevent Opioid Use for Oral Mucositis Pain During Head and Neck Chemoradiotherapy: A Phase III Clinical Trial

NSCLC

A081801

Phase 3 · ALLIANCE, National Cancer Institute (NCI) · recruiting · NCT04267848

Integration of Immunotherapy Into Adjuvant Therapy for Resected NSCLC: ALCHEMIST Chemo-IO (ACCIO)

Arm B (sequential) Platinum doublet x 4 cycles, +/- PORT, then pembrolizumab 400 mg IV every 6 weeks x 8 doses (16 cycles)

Arm C (concurrent) Platinum doublet x 4 cycles with pembrolizumab 400 mg on day 1 of cycles 1 and 3, +/- PORT, then pembrolizumab 400 mg every 6 weeks x 6 doses (12 cycles), 16 total cycles of pembrolizumab