

MATH342 Review Notes

By Jack 'jryzkn's' Zhou

1 The division Algorithm

If $a, b \in \mathbb{Z}$ with $b > 0$, then $\exists!(q, r) \in \mathbb{Z}^1$ s.t.

$$a = bq + r; \quad 0 \leq r < b$$

2 Divisibility and Primes

If $a, b \in \mathbb{Z}$ with $a \neq 0$, we say that a divides b if $\exists c \in \mathbb{Z}$ s.t. $b = ac$. We write this relationship as $a \mid b$. If a does not divide b , we write $a \nmid b$.

Below are some properties of divisibility, using $a, b, c \in \mathbb{Z}$ and $m, n \in \mathbb{Z}$:

- $a \mid b \wedge b \mid c \rightarrow a \mid c$
- $c \mid a \wedge c \mid b \rightarrow c \mid (ma + nb)$

An integer n is said to have *odd* parity when $n \nmid 2$, and otherwise *even* parity when $n \mid 2$. Often times an odd number can be written in the form $n = 2k + 1$ for some integer k , and $n = 2k$ for even numbers.

A *prime* is an integer greater than 1 that is divisible by no positive integers other than 1 and itself. Otherwise, a number is said to be a *composite*. Note that by this definition. The number 2 is a prime. Often times there will be indications to exclude 2 from primes by specifying odd primes.

Here are some facts about prime numbers, where $n \in \mathbb{Z}$:

- when $n > 1$, $p \mid n$ for some prime $p \leq n$
- There are infinitely many primes
- If n is composite, then $p \mid n$ for some prime $p \leq \sqrt{n}$

The function $\pi(x)$ where x is a positive real number denotes the number of primes not exceeding x .

Dirichlet's Theorem in Arithmetic Progressions

Suppose that $a, b \in \mathbb{N}$ where $(a, b) = 1$. Then the arithmetic progression $an + b, n \in \mathbb{N}$ contains infinitely many primes.

3 Greatest Common Divisors

The *greatest common divisor* (GCD) of $a, b \in \mathbb{Z}$ is the largest divisor d such that $d \mid a$ and $d \mid b$. The GCD of a and b are often written as $\gcd(a, b)$ or (a, b) .

One way to describe the GCD is that a positive integer d is a GCD iff:

- $d \mid a$ and $d \mid b$

- if $c \in \mathbb{Z}$ s.t. $c \mid a$ and $c \mid b$, then $c \mid d$

Some facts about GCD's:

- Two integers $a, b \in \mathbb{Z}$ are said to be *relatively prime* if $(a, b) = 1$
- Suppose $d = (a, b)$, then $(\frac{a}{d}, \frac{b}{d}) = 1$
- A fraction $\frac{p}{q}$ is in lowest terms when $(p, q) = 1$
- The notion of GCD's applies to multiple values too, suppose $a_1, a_2, \dots, a_n \in \mathbb{Z}$, then

$$(a_1, a_2, \dots, a_n) = (a_1, a_2, \dots, (a_{n-1}, a_n))$$

Note that simply with the above definition, $(a_1, a_2, \dots, a_n) = 1$ means that these numbers are *mutually relatively prime*. A stronger statement is *pairwise relatively prime*, where for any pair of the numbers a_i and a_j with $i \neq j$, $(a_i, a_j) = 1$.

Bezout's Theorem

If $a, b \in \mathbb{Z}$, then $\exists m, n \in \mathbb{Z}$ s.t.

$$ma + nb = (a, b)$$

Where m, n are denoted the bezout coefficients to a, b . Furthermore, $(a, b) = 1 \Leftrightarrow ma + nb = 1$.

The set of linear combinations of a, b is the set of integer multiples of (a, b) .

Euclidean Algorithm (+Backtracking)

The Euclidean Algorithm computes (a, b) . It proceeds by heavily using a property of GCD's:

Let $b, q, r \in \mathbb{Z}$,

$$(bq + r, b) = (b, r)$$

Suppose $a = bq + r$ (by the division algorithm), we have $(a, b) = (b, r)$. As $0 \leq r < b$, the RHS will always be in lower terms: each time we apply this property, we will get smaller computations to carry out until a base case of $(b, 0)$ is reached, in which case $(b, 0) = b$.

Once the series of division algorithms are applied, the results can be used in reverse with substitution to find the bezout coefficients.

As a side note, suppose we have f_{n+1}, f_{n+2} be successive terms of the Fibonacci Sequence with $n > 1$, then the Euclidean algorithm takes exactly n divisions to show that $(f_{n+1}, f_{n+2}) = 1$.

¹The symbol $\exists!$ indicates that the existence is unique

4 Continued Fraction Expansions

Given the sequence a_0, a_1, a_2, \dots (may be infinite), a continued fraction is a fraction of the following form

$$a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots}}$$

A fraction of this form can also be written as $[a_0, a_1, a_2, \dots]$. If the sequence is infinite and has repeating elements, we denote one instance of the repeating values with a line drawn over it.

The numbers a_1, a_2, \dots, a_n are called *partial quotients* of the continued fraction, and if all a_i 's are integers, the continued fraction is said to be a *simple* continued fraction. We are only concerned with *simple* continued fractions and unless otherwise stated, all continued fractions under discussion are simple.

Finite Continued Fractions

For a rational number that can be expressed as $\frac{p}{q}$, where $p, q \in \mathbb{Z}$ and $p > q$. Its continued fraction expansion is finite. The way to generate the expansion is to consider the division algorithm applies when computing (p, q) . For each row, it will be of the form $a = q \cdot b + r$. We apply the following transformation:

$$a = bq + r \Rightarrow \frac{a}{b} = q + \frac{1}{\frac{b}{r}}$$

The $\frac{b}{r}$ term will correspond to the LHS on the next row, recall that the next row would be computing b divided by r , and thus a substitution can be made. We can continuously apply these substitutions until a continued fraction is formed.

Infinite Continued Fractions

For an irrational number n , its continued fraction expansion is computed in the following manner:

$$\alpha_0 = n$$

$$a_i = [\alpha_i] \quad \alpha_{i+1} = (\alpha_i - a_i)^{-1}$$

It's good to note that a_i is the integral component to α_i , and that $\alpha_i - a_i$ is the fractional component to α_i .

Convergents

Given a continued fraction

$$C = [a_0; a_1, a_2, \dots, a_n]$$

and $C_k = [a_0; a_1, a_2, \dots, a_k]$ with $0 < k \leq n$, C_k is called the k th convergent of C . Given C , we can compute all of the convergents of C as follows:

$$\begin{array}{lll} p_0 = a_0 & q_0 = 1 & \\ p_1 = a_0 a_1 + 1 & q_1 = a_1 & C_1 = \frac{p_1}{q_1} \\ p_k = a_k p_{k-1} + p_{k-2} & q_k = a_k q_{k-1} + q_{k-2} & C_k = \frac{p_k}{q_k} \end{array}$$