

# MATH342 Review Notes

By Jack 'jryzkn's' Zhou

## 1 The division Algorithm

If  $a, b \in \mathbb{Z}$  with  $b > 0$ , then  $\exists!(q, r) \in \mathbb{Z}^1$  s.t.

$$a = bq + r; \quad 0 \leq r < b$$

## 2 Divisibility and Primes

If  $a, b \in \mathbb{Z}$  with  $a \neq 0$ , we say that  $a$  *divides*  $b$  if  $\exists c \in \mathbb{Z}$  s.t.  $b = ac$ . We write this relationship as  $a \mid b$ . If  $a$  *does not divide*  $b$ , we write  $a \nmid b$ .

Below are some properties of divisibility, using  $a, b, c \in \mathbb{Z}$  and  $m, n \in \mathbb{Z}$ :

- $a \mid b \wedge b \mid c \rightarrow a \mid c$
- $c \mid a \wedge c \mid b \rightarrow c \mid (ma + nb)$

An integer  $n$  is said to have *odd* parity when  $n \nmid 2$ , and otherwise *even* parity when  $n \mid 2$ . Often times an odd number can be written in the form  $n = 2k + 1$  for some integer  $k$ , and  $n = 2k$  for even numbers.

A *prime* is an integer greater than 1 that is divisible by no positive integers other than 1 and itself. Otherwise, a number is said to be a *composite*. Note that by this definition. The number 2 is a prime. Often times there will be indications to exclude 2 from primes by specifying odd primes.

Here are some facts about prime numbers, where  $n \in \mathbb{Z}$ :

- when  $n > 1$ ,  $p \mid n$  for some prime  $p \leq n$
- There are infinitely many primes
- If  $n$  is composite, then  $p \mid n$  for some prime  $p \leq \sqrt{n}$

The function  $\pi(x)$  where  $x$  is a positive real number denotes the number of primes not exceeding  $x$ .

### Dirichlet's Theorem in Arithmetic Progressions

Suppose that  $a, b \in \mathbb{N}$  where  $(a, b) = 1$ . Then the arithmetic progression  $an + b, n \in \mathbb{N}$  contains infinitely many primes.

## 3 Greatest Common Divisors

The *greatest common divisor* (GCD) of  $a, b \in \mathbb{Z}$  is the largest divisor  $d$  such that  $d \mid a$  and  $d \mid b$ . The GCD of  $a$  and  $b$  are often written as  $\gcd(a, b)$  or  $(a, b)$ .

One way to describe the GCD is that a positive integer  $d$  is a GCD iff:

- $d \mid a$  and  $d \mid b$

---

<sup>1</sup>The symbol  $\exists!$  indicates that the existence is unique

- if  $c \in \mathbb{Z}$  s.t.  $c \mid a$  and  $c \mid b$ , then  $c \mid d$

Some facts about GCD's:

- Two integers  $a, b \in \mathbb{Z}$  are said to be *relatively prime* if  $(a, b) = 1$
- Suppose  $d = (a, b)$ , then  $(\frac{a}{d}, \frac{b}{d}) = 1$
- A fraction  $\frac{p}{q}$  is in lowest terms when  $(p, q) = 1$
- suppose  $(a, b) = 1$  and  $a \mid bc$ , then  $a \mid c$
- The notion of GCD's applies to multiple values too, suppose  $a_1, a_2, \dots, a_n \in \mathbb{Z}$ , then

$$(a_1, a_2, \dots, a_n) = (a_1, a_2, \dots, (a_{n-1}, a_n))$$

Note that simply with the above definition,  $(a_1, a_2, \dots, a_n) = 1$  means that these numbers are *mutually relatively prime*. A stronger statement is *pairwise relatively prime*, where for any pair of the numbers  $a_i$  and  $a_j$  with  $i \neq j$ ,  $(a_i, a_j) = 1$ .

### Bezout's Theorem

If  $a, b \in \mathbb{Z}$ , then  $\exists m, n \in \mathbb{Z}$  s.t.

$$ma + nb = (a, b)$$

Where  $m, n$  are denoted the bezout coefficients to  $a, b$ . Furthermore,  $(a, b) = 1 \Leftrightarrow ma + nb = 1$ .

The set of linear combinations of  $a, b$  is the set of integer multiples of  $(a, b)$ .

### Euclidean Algorithm (+Backtracking)

The Euclidean Algorithm computes  $(a, b)$ . It proceeds by heavily using a property of GCD's:

Let  $b, q, r \in \mathbb{Z}$ ,

$$(bq + r, b) = (b, r)$$

Suppose  $a = bq + r$  (by the division algorithm), we have  $(a, b) = (b, r)$ . As  $0 \leq r < b$ , the RHS will always be in lower terms: each time we apply this property, we will get smaller computations to carry out until a base case of  $(b, 0)$  is reached, in which case  $(b, 0) = b$ .

Once the series of division algorithms are applied, the results can be used in reverse with substitution to find the bezout coefficients.

As a side note, suppose we have  $f_{n+1}, f_{n+2}$  be successive terms of the Fibonacci Sequence with  $n > 1$ , then the Euclidean algorithm takes exactly  $n$  divisions to show that  $(f_{n+1}, f_{n+2}) = 1$ .

## 4 Continued Fraction Expansions

Given the sequence  $a_0, a_1, a_2, \dots$  (may be infinite), a continued fraction is a fraction of the following form

$$a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots}}$$

A fraction of this form can also be written as  $[a_0, a_1, a_2, \dots]$ . If the sequence is infinite and has repeating elements, we denote one instance of the repeating values with a line drawn over it.

The numbers  $a_1, a_2, \dots, a_n$  are called *partial quotients* of the continued fraction, and if all  $a_i$ 's are integers, the continued fraction is said to be a *simple* continued fraction. We are only concerned with *simple* continued fractions and unless otherwise stated, all continued fractions under discussion are simple.

### Finite Continued Fractions

For a rational number that can be expressed as  $\frac{p}{q}$ , where  $p, q \in \mathbb{Z}$  and  $p > q$ . Its continued fraction expansion is finite. The way to generate the expansion is to consider the division algorithm applies when computing  $(p, q)$ . For each row, it will be of the form  $a = q \cdot b + r$ . We apply the following transformation:

$$a = bq + r \Rightarrow \frac{a}{b} = q + \frac{1}{\frac{b}{r}}$$

The  $\frac{b}{r}$  term will correspond to the LHS on the next row, recall that the next row would be computing  $b$  divided by  $r$ , and thus a substitution can be made. We can continuously apply these substitutions until a continued fraction is formed.

### Infinite Continued Fractions

For an irrational number  $n$ , its continued fraction expansion is computed in the following manner:

$$\alpha_0 = n \\ a_i = [\alpha_i] \quad \alpha_{i+1} = (\alpha_i - a_i)^{-1}$$

It's good to note that  $a_i$  is the integral component to  $\alpha_i$ , and that  $\alpha_i - a_i$  is the fractional component to  $\alpha_i$ .

### Convergents

Given a continued fraction

$$C = [a_0; a_1, a_2, \dots, a_n]$$

and  $C_k = [a_0; a_1, a_2, \dots, a_k]$  with  $0 < k \leq n$ ,  $C_k$  is called the  $k$ th convergent of  $C$ . Given  $C$ , we can compute all of the convergents of  $C$  as follows:

$$\begin{array}{lll} p_0 = a_0 & q_0 = 1 & \\ p_1 = a_0 a_1 + 1 & q_1 = a_1 & C_1 = \frac{p_1}{q_1} \\ p_k = a_k p_{k-1} + p_{k-2} & q_k = a_k q_{k-1} + q_{k-2} & C_k = \frac{p_k}{q_k} \end{array}$$

## 5 Linear Diophantine Equations

A linear diophantine equation in two variables is an equation of the follow form

$$ax + by = c$$

where  $a, b, c \in \mathbb{Z}$  and an integer solution  $(x, y)$  is sought for. In general, an equation that is linear in the coefficients to the polynomial powers would be considered a diophantine equation. We will focus on linear diophantine equations in two variables. Much of the results we find here applies to diophantine equations of more than two variables.

Let  $d = (a, b)$ . If  $d \nmid c$ , then the diophantine equation in question has no integral solutions. If  $d \mid c$ , then there are infinitely many solutions (granted that there are no restrictions on the solution space). Indeed, once we have an initial solution  $(x_0, y_0)$ , the rest of the solutions will all be of the form

$$x = x_0 + \frac{b}{d}n, \quad y = y_0 - \frac{a}{d}n$$

for any integer  $n$ . Notice the minus sign on  $y$ .

## Systems of Linear Diophantine Equations

When there is a system of linear diophantine equations (most commonly two equations in 3 variables), the ideal approach is to substitute one equation into another, to reduce the system into a single equation. From there, solve as intended and substitute back for a full solution.

## 6 The Fundamental Theorem of Arithmetic

Every positive integer greater than 1 is *uniquely* expressible as a product of primes, with the prime factors in non decreasing order:

$$n = \prod_i p_i^{a_i}$$

The sequence of  $a_i$ 's denote the exponents to each of the prime factors. If a prime factor  $p_i \nmid n$ , then  $a_i = 0$ .

Using this prime factorization form, we can describe the GCD and LCM (least common multiple) as follows, letting  $a = \prod_i p_i^{a_i}$  and  $b = \prod_i p_i^{b_i}$ :

$$(a, b) = \prod_i p_i^{\min(a_i, b_i)} \quad [a, b] = \prod_i p_i^{\max(a_i, b_i)}$$

As  $\max(a_i, b_i) + \min(a_i, b_i) = a_i + b_i$ , we can see from above that  $(a, b) \cdot [a, b] = ab$

## 7 Congruences

Let  $m$  be a positive integer. If  $a, b \in \mathbb{Z}$ , we say that  $a$  is *congruent* to  $b$  modulo  $m$  if  $m \mid (a - b)$ . Furthermore, we can write  $a \equiv b \pmod{m}$  iff  $a = b + km$  for some integer  $k$ . Here are some properties about arithmetic in modulo  $m$ :

- $a + c \equiv b + c \pmod{m}$
- $a - c \equiv b - c \pmod{m}$
- $ab \equiv bc \pmod{m}$

The congruence relation over a modulus forms an *equivalence class*, which satisfies the following properties:

- Reflexivity:  $a \equiv a \pmod{m}$
- Symmetricity:  $a \equiv b \pmod{m} \Leftrightarrow b \equiv a \pmod{m}$
- Transitivity:  $a \equiv b \pmod{m} \wedge b \equiv c \pmod{m} \Rightarrow a \equiv c \pmod{m}$

A number in a modulo number system itself is referred to as a residue. If  $a, b$  are congruent mod  $m$ , then their residues  $a \pmod{m}$  and  $b \pmod{m}$  have to be the same as well.

A complete system of residues modulo  $m$  is a set of integers such that every integer is congruent to exactly one integer of the set. Of which, the least non-negative residues modulo  $m$  is the set

$$\{0, 1, \dots, m-1\}$$

When  $m$  is odd, the absolute least residues modulo  $m$  is the set

$$\left\{ -\frac{m-1}{2}, -\frac{m-3}{2}, \dots, -1, 0, 1, \frac{m-3}{2}, \frac{m-1}{2} \right\}$$

If  $a, b, c, m \in \mathbb{Z}^+$  with  $d = (c, m)$  and  $ac \equiv bc \pmod{m}$ , then  $a \equiv b \pmod{\frac{m}{d}}$ . In the case that  $(c, m) = 1$ , we simply have  $a \equiv b \pmod{m}$ .

If we have  $a \equiv b \pmod{m_i}$  for several values of  $i$  and each  $m_i$  are pairwise coprime, then  $a \equiv b \pmod{\prod_i m_i}$ .

## 8 Infinitality of primes of the form $4k+3$

Before the actual proof can take place, we note that primes can only either be of the form  $4k+1$  or  $4k+3$ , which are the odd numbers in mod 4.

Also we note that if two numbers  $a, b$  are both of the form  $4k+1$ , then  $ab$  will also be of the form  $4k+1$  (proof omitted for brevity).

To begin the actual proof, suppose we have a finite amount of primes of the form  $4k+3$ , Namely, all in the sequence  $\{p_i\}$  starting with  $p_0 = 3$ . We construct the following:

$$Q = 4 \prod_{i=1} p_i + 3$$

Note that  $p_0 = 3$  is omitted from the expression.

Clearly,  $Q$  is a composite value of the form  $4k+3$  by construction. It needs to have a factor that is of the form  $4k+3$ , as we have established that multiplication between numbers of the form  $4k+1$  is closed. Therefore, for some  $i$ ,  $p_i \mid Q$ . However, none of the  $p_i$ 's can divide  $Q$  by construction. Therefore,  $Q$  is another prime of the form  $4k+3$ , a contradiction to the assumption that there are only a finite amount of them.