Actualization as Physical Mechanism: Implications of Relational Actualism for Quantum Computing and Quantum Thermodynamics

Joshua F. Sandeman
Independent Researcher
sansuikyo@gmail.com

March 2026

Abstract

Relational Actualism (RA) proposes that the transition from quantum potentia to classical actuality is grounded in a precise physical criterion: the on-shell dispersion relation $p^\mu p_\mu = m^2c^2$. When a mediating boson crosses the kinematic threshold from virtual (off-shell) to real (on-shell), an irreversible vertex is inscribed into the growing causal directed acyclic graph (DAG) of the universe. This paper develops the implications of this framework—established in the companion papers RAQM, RAGC, RACI, and RAHC (Sandeman, 2026e)—for two fields that have lacked a precise physical account of irreversibility at the quantum level: quantum computing and quantum thermodynamics. For quantum computing, the Kinematic Snap provides a principled physical floor on decoherence, distinguishes genuinely off-shell (decoherence-immune) interaction regimes from on-shell ones, and gives the threshold theorem a physical rather than purely operational interpretation. For quantum thermodynamics, RA grounds Landauer's principle in the causal DAG, resolves the Maxwell's demon paradox without information-theoretic postulates, gives the fluctuation theorems a precise actualization-event interpretation, and establishes that the thermodynamic arrow of time is structural rather than emergent. A key technical result: the quantum relative entropy—the central quantity in both quantum information theory and RA's actualization criterion—is frame-independent by a compiled Lean 4 theorem (`frame_independence` in `RA_AQFT_Proofs_v10.lean`), proved using the continuous functional calculus unitary-conjugation lemma from the Lean-QuantumInfo library (Meiburg et al., 2025). Throughout, implications are stated conditionally on the correctness of the RA framework, which is currently under peer review. **Keywords:** Relational Actualism · Kinematic Snap · Quantum Computing · Decoherence · Quantum Thermodynamics · Landauer's Principle · Maxwell's Demon · Fluctuation Theorems · Quantum Relative Entropy · Formal Verification

1 Introduction

Two of the most practically consequential fields in contemporary physics share a common foundational gap: neither quantum computing nor quantum thermodynamics possesses a precise physical account of when, and by what mechanism, a quantum process becomes irreversible. Quantum computing treats decoherence as an engineering challenge to be suppressed, characterised by phenomenological timescales T_1 and T_2 whose physical origins are described only statistically. Quantum thermodynamics invokes entropy, work, and irreversibility while relying on the same unitary, time-reversible quantum mechanics that makes those concepts difficult to ground.

Relational Actualism (RA) (Sandeman, 2026a,b,c) proposes that this gap can be closed by a single physical criterion: actualization occurs when a mediating boson crosses the on-shell dispersion relation $p^\mu p_\mu = m^2c^2$. Below this threshold, exchange is virtual, off-shell, and reversible. At and above it, the exchange is real, on-shell, and permanently inscribed into the causal DAG. This is the

Kinematic Snap—the precise physical mechanism by which quantum potentia becomes classical actuality.

The present paper develops the implications of this framework for quantum computing (Section 2) and quantum thermodynamics (Section 3), and then discusses the technical connection to quantum information theory via the quantum relative entropy formalism (Section 4). All implications are stated conditionally: they follow *if* the RA actualization criterion is correct. The empirical predictions in Section 5 are the appropriate test.

We note at the outset that implications of this kind are not merely interpretational. The Kinematic Snap criterion differs from decoherence-based accounts in a precise, experimentally distinguishable way: it identifies a sharp threshold (Prediction P3 of RAQM (Sandeman, 2026a)) rather than a continuous crossover. That distinction has practical consequences for both fields developed here.

2 Implications for Quantum Computing

2.1 The Physical Floor on Decoherence

Standard quantum error correction characterises decoherence by the phenomenological timescales T_1 (energy relaxation) and T_2 (phase coherence), which are treated as empirical parameters to be minimised through engineering. RA gives these parameters a physical interpretation and identifies the mechanism that sets them.

In RA, decoherence occurs when the system exchanges on-shell bosons with its environment. Specifically, the actualization timescale is

$$\tau_{\text{act}} \sim \frac{\hbar}{\Delta E_{\text{env}}}, \tag{1}$$

where ΔE_{env} is the characteristic energy available for on-shell boson emission into the environmental degrees of freedom (RAQM Prediction P2 (Sandeman, 2026a)). This is not a statistical parameter; it is the time required for an emitted on-shell boson to propagate outward and entangle irreversibly with the environmental boundary.

The practical implication is immediate: *the engineering target for decoherence suppression is ΔE_{env}, not the coupling strength alone*. Reducing the energy available in the environment for on-shell exchange directly extends τ_{act}. This gives a physical interpretation to the well-known empirical strategy of operating superconducting qubits at millikelvin temperatures: reducing $k_B T$ reduces ΔE_{env}, suppressing on-shell exchange with thermal phonons. RA identifies this as the mechanism rather than a phenomenological correlation.

Furthermore, (1) implies a physical floor that cannot be overcome by any engineering improvement that does not reduce ΔE_{env}. No amount of improved substrate quality, shielding, or control pulse optimisation will suppress decoherence below $\hbar/\Delta E_{\text{env}}$ if the environmental energy scale is not itself reduced. This is a stronger statement than the standard decoherence-theoretic account, which predicts continuous improvement in principle.

2.2 The Virtual/Real Distinction as a Decoherence-Immune Regime

A non-trivial implication of the Kinematic Snap concerns interaction regimes that are genuinely off-shell. Virtual, off-shell exchanges—van der Waals forces, the Casimir effect, the Lamb shift—produce no DAG vertices in RA. They are genuine physical effects, but they are effects of the potentia: they do not constitute actualization events and do not contribute to decoherence.

This is not merely a restatement of existing results. The standard decoherence account treats all environmental coupling as a potential source of decoherence, modulated by coupling strength and spectral density. RA draws a qualitatively different distinction: coupling that remains purely off-shell (because the interaction energy does not reach the on-shell threshold) is decoherence-immune in the perturbative radiative regime, not merely weakly decoherent. This holds cleanly when on-shell boson emission is the dominant decoherence channel; RAQM notes that near-field and many-body

processes can produce environmental distinguishability through other mechanisms, so the immunity is regime-specific rather than categorical.

The potential design implication is significant. If a quantum computing architecture can be engineered so that its dominant environmental coupling remains below the on-shell threshold—so that qubit-environment interactions are kinematically constrained to the virtual regime—RA predicts that this regime would exhibit qualitatively suppressed decoherence, not merely quantitatively reduced decoherence. This is a testable prediction that distinguishes RA from standard decoherence theory.

The experimental platform most naturally positioned to probe this distinction is the superconducting circuit, where the relevant on-shell bosons are microwave photons and the threshold condition is set by the circuit's transition frequencies. Operating in a regime where the environmental spectral density falls entirely below the transition frequency creates exactly the off-shell condition RA identifies as decoherence-immune. This is the physical interpretation of the "bad cavity" versus "good cavity" limit in circuit quantum electrodynamics.

2.3 The Threshold Theorem Acquires a Physical Interpretation

The fault-tolerance threshold theorem establishes that scalable quantum computation is possible provided the physical error rate p falls below a threshold p_{th}, which depends on the error correction code and architecture. The theorem is currently stated in terms of error rates that are treated as phenomenological parameters: there is no physical derivation of p_{th} from first principles.

If RA is correct, the physical error rate has a precise interpretation: it is the rate of on-shell boson exchange between the logical qubit and its environment, given by

$$p \sim \frac{1}{\tau_{\text{act}} \cdot f_{\text{gate}}} = \frac{\Delta E_{\text{env}}}{\hbar \cdot f_{\text{gate}}}, \tag{2}$$

where f_{gate} is the gate operation frequency. The threshold $p < p_{\text{th}}$ then becomes a condition on the ratio $\Delta E_{\text{env}}/(\hbar \cdot f_{\text{gate}})$: the environment must not have sufficient energy to cause on-shell exchange faster than the gate cycle time. This gives the threshold theorem a physical interpretation as a statement about kinematic accessibility rather than an abstract error rate condition.

The further implication is that the threshold is not universal across architectures in the way current theory suggests. For two architectures with the same phenomenological error rate p but different physical mechanisms—one where errors arise from on-shell exchange and one where they arise from control-pulse infidelity—RA predicts different scalability behaviour, because only on-shell exchange constitutes genuine actualization. Control infidelity that does not produce on-shell environmental exchange is, in principle, reversible and correctable in a way that genuine actualization errors are not. In practice, control pulse errors often couple to environmental modes that can go on-shell, so this distinction is a conceptual one about error types rather than a claim about the reversibility of all control errors in current hardware.

2.4 The Kinematic Coherence Bound and the $1/N$ Law

The Kinematic Snap criterion yields a sharp, derivable bound on the maximum achievable circuit depth for an N-qubit logical state, which we call the *Kinematic Coherence Bound* (KCB). The derivation proceeds directly from Prediction P2 of RAQM (Sandeman, 2026a) without additional assumptions.

Single-qubit baseline. For a single isolated qubit, the actualization timescale is

$$\tau_{\text{act}} = \frac{\hbar}{\Delta E_{\text{env}}}, \tag{3}$$

where ΔE_{env} is the characteristic energy of the environmental modes available for on-shell exchange. The gate quality factor is

$$\eta \;:=\; f_{\text{gate}} \cdot \tau_{\text{act}} \;=\; \frac{\hbar f_{\text{gate}}}{\Delta E_{\text{env}}}, \tag{4}$$

where f_{gate} is the gate frequency. This is the RA physical interpretation of the standard figure of merit T_2/t_{gate}: it is the ratio of the gate period to the single-qubit actualization timescale.

N-qubit logical state. Consider an N-qubit register in an entangled logical state (for concreteness, a GHZ-type state of the form $\frac{1}{\sqrt{2}}(|00\cdots0\rangle + |11\cdots1\rangle)$, or any state with genuinely N-body entanglement). The logical information is encoded in correlations that are non-local across all N qubits. Under the Kinematic Snap criterion, each qubit constitutes an independent site at which the kinematic threshold can be crossed: the actualization of any single qubit destroys the logical entanglement of the entire register.

The logical decoherence rate is therefore the sum of single-qubit actualization rates across all N sites:

$$\Gamma_{\text{logical}} \;=\; N \cdot \Gamma_1 \;=\; \frac{N \cdot \Delta E_{\text{env}}}{\hbar}, \tag{5}$$

and the logical coherence time is

$$\tau_{\text{act}}^{(N)} \;=\; \frac{\hbar}{N \cdot \Delta E_{\text{env}}}. \tag{6}$$

This is the RA physical account of the well-known result from quantum information theory that the coherence time of a GHZ state scales as T_1/N (Huelga et al., 1997). The Kinematic Snap mechanism explains *why*: each qubit is a separate kinematic threshold that can be independently crossed, and crossing any one of them destroys the logical state.

The Kinematic Coherence Bound. The maximum circuit depth achievable before logical decoherence is

$$\boxed{D_{\text{max}} \;=\; f_{\text{gate}} \cdot \tau_{\text{act}}^{(N)} \;=\; \frac{\eta}{N},} \tag{7}$$

where η is the single-qubit quality factor defined in (4). Circuit depth scales strictly as $1/N$: one cannot increase the number of qubits and the circuit depth simultaneously without improving η.

Maximum fault-tolerant array size. The fault-tolerance threshold condition requires the physical error rate per gate to fall below a code-dependent threshold p_{th}:

$$p \;=\; \frac{\Gamma_{\text{logical}}}{f_{\text{gate}}} \;=\; \frac{N}{\eta} \;<\; p_{\text{th}}. \tag{8}$$

This yields a *maximum array size* for fault-tolerant operation at fixed single-qubit quality:

$$\boxed{N_{\text{max}} \;=\; \eta \cdot p_{\text{th}}.} \tag{9}$$

This result is not anticipated by standard quantum error correction theory, which holds that fault-tolerant quantum computation is scalable without bound once $p < p_{\text{th}}$ for individual qubits. The RA account adds a constraint: the logical error rate grows with N because each added qubit is an additional kinematic site at which actualization can occur. Unlimited scaling without improving η will eventually push the logical error rate above p_{th} regardless of single-qubit fidelity.

Two caveats on the derivation. First, it assumes independent decoherence channels per qubit. Correlated errors—crosstalk, collective decoherence, or correlated on-shell emission events—can produce worse-than-independent scaling, so N_{max} is an upper bound on the array size under independent depolarizing noise, not a tight limit for all error models. Second, p_{th} is code- and error-model-dependent; the numerical estimates below use a surface code with depolarizing noise.

Syndrome extraction and the KCB. A natural question concerns the role of quantum error correction itself: does a stabilizer measurement on an ancilla qubit constitute a Kinematic Snap that destroys the logical state? The answer is no, and the distinction is physically precise. In a surface code, syndrome extraction works by entangling ancilla qubits with the data qubits and then measuring the ancillae. The ancilla measurement *does* constitute an actualization event—the ancilla crosses the on-shell threshold and its state is irreversibly recorded. However, the logical information is encoded *non-locally* across many physical data qubits; it is not localized at any single kinematic site. The ancilla actualization reveals error syndrome information without collapsing the logical state, because the logical degree of freedom has no support at the ancilla's location in the causal DAG.

The KCB applies to *unintended* environmental actualization events—on-shell exchanges between a data qubit and an uncontrolled environmental mode—not to the designed syndrome extraction operations, which are intentional actualization events that extract entropy without disturbing the logical encoding. The $1/N$ scaling therefore reflects the rate of *logical errors* from environmental coupling, which grows with N under independent depolarizing noise. Continuous QEC protocols suppress this error rate below p_{th} but cannot eliminate the underlying kinematic constraint: $N_{\max}$ is the array size at which the environmental logical error rate, even before QEC correction, equals p_{th}. Beyond $N_{\max}$, QEC must run faster than the logical decoherence rate, which requires improving η—consistent with the empirical observation that scaling superconducting arrays requires simultaneous improvement of single-qubit quality.

Concretely: for a surface code with $p_{\text{th}} \approx 10^{-2}$ under independent depolarizing noise, and a contemporary superconducting qubit with $\eta \approx 10^3$–10^4 (corresponding to $T_2 \sim 100\,\mu$s and $t_{\text{gate}} \sim 100\,$ns), the RA bound gives $N_{\max} \sim 10$–100 logical qubits at fixed η. This is illustrative rather than a fit to experimental data; the observation that large superconducting arrays require continual improvement of η alongside qubit count is consistent with but does not uniquely confirm the KCB.

The optimal isolation window. A related prediction concerns the engineering of qubit isolation. The target quantity is ΔE_{env}: the energy available to environmental modes for on-shell exchange. Reducing ΔE_{env} directly extends τ_{act} and improves η. However, the environment has multiple channels for on-shell exchange—photonic (microwave), phononic, and through two-level system (TLS) defects in the substrate—each with its own $\Delta E_{\text{env},i}$. Improving isolation in one channel (e.g., by increasing cavity Q to suppress photonic decay) does not suppress the others.

RA therefore predicts that there is an *optimal isolation window* Q^* for each channel: the cavity Q at which photonic actualization timescale $\tau_{\text{photo}} = \hbar/\Delta E_{\text{photo}}$ equals the actualization timescale of the dominant non-photonic channel $\tau_{\text{other}} = \hbar/\Delta E_{\text{other}}$. Beyond Q^*, further improvement of photonic isolation yields diminishing returns because the dominant decoherence pathway shifts to the non-photonic channel. This is measurable as a change in the T_1 scaling with cavity Q: below Q^* the scaling should follow the Purcell prediction; above Q^* it should plateau.

2.5 Implications for Quantum Algorithm Design

At a higher level of abstraction, the Kinematic Snap suggests a distinction between quantum algorithms that operate entirely within the off-shell regime— where the computation consists of reversible potentia evolving under unitary dynamics—and those that necessarily involve actualization events. All conventional quantum algorithms involve measurement, which is a forced actualization. But the computation preceding measurement may be structured to avoid premature actualization.

This distinction is already implicit in the design of adiabatic quantum computing and quantum annealing, where the system is evolved slowly to avoid non-adiabatic transitions. RA offers the following reinterpretation of this design principle: avoiding non-adiabatic transitions is avoiding premature on-shell exchange with environmental modes, which would constitute unintended actualization events. The adiabatic condition, in RA terms, is the condition that the evolution remains in

the off-shell regime throughout. This is an interpretational reframing rather than a new prediction; it does not alter the standard adiabatic theorem or provide new algorithmic guidance beyond what standard theory already implies.

3 Implications for Quantum Thermodynamics

3.1 Landauer's Principle Grounded in the Causal DAG

Landauer's principle (Landauer, 1961) states that the logical erasure of one bit of information dissipates a minimum energy of $k_B T \ln 2$ as heat into the environment. The principle is standardly derived from thermodynamic arguments (the second law applied to the Szilard engine) or from quantum information theory (the quantum data compression theorem). Both derivations establish the bound; neither identifies the physical mechanism by which it is enforced.

In RA, Landauer's principle has a direct mechanical derivation from the Kinematic Snap (Sandeman, 2026c).

Theorem 3.1 (Landauer Cost of Actualization, RACI Theorem 4)**.** *Each irreversible actualization event that eliminates Δq bits of quantum uncertainty—reducing the von Neumann entropy of the system by $\Delta S_{\mathrm{vN}} = \Delta q \cdot \ln 2$—dissipates at minimum*

$$W_{\mathrm{act}} \;=\; \Delta q \cdot k_B T \cdot \ln 2. \tag{10}$$

The proof in RACI proceeds directly from the Kinematic Snap criterion: an actualization event writes a permanent vertex to the causal DAG, which requires on-shell boson emission into the environment. The von Neumann entropy decrease ΔS_{vN} of the system must be transferred to the environment as heat $\geq T\Delta S_{\mathrm{vN}} = k_B T \cdot \Delta q \cdot \ln 2$. The bound (10) follows.

What RA adds to Landauer's principle beyond the standard derivation:

A mechanism. The energy is dissipated specifically at the moment the mediating boson crosses the on-shell threshold. The dissipation is not a continuous process but a discrete event with a precise physical timing: $\tau_{\mathrm{act}} \sim \hbar/\Delta E_{\mathrm{env}}$ before the on-shell threshold is crossed.

A scope condition. The Landauer bound applies only to *genuine actualization events*—irreversible on-shell exchanges that write DAG vertices. Operations that remain in the virtual, off-shell regime do not write DAG vertices and can, in principle, be reversed without paying the Landauer cost. This is consistent with the reversibility of coherent quantum operations; RA identifies the physical criterion that distinguishes reversible from irreversible operations.

A note on the Reeb-Wolf derivation: Reeb and Wolf (Reeb and Wolf, 2014) established a fully quantum Landauer bound showing the minimum dissipation holds for any entropy-decreasing operation on a quantum system in contact with a thermal reservoir, without reference to emission events. The RA account does not contradict this result; rather, RA provides the physical mechanism *by which* the Reeb-Wolf bound is saturated. In RA, an entropy-decreasing operation that does not involve on-shell exchange is not genuinely irreversible—it is a coherent operation that can, in principle, be reversed by subsequent unitary evolution. The Reeb-Wolf bound becomes binding precisely when the operation crosses the Kinematic Snap threshold and produces an on-shell environmental record. The two accounts agree on the bound; they differ on the physical story of why it holds.

A timescale. The Landauer bound tells you the minimum energy cost of erasure. RA additionally tells you the minimum *time* for erasure: $\tau_{\mathrm{min}} \sim \hbar/\Delta E_{\mathrm{env}}$. These two together constitute a Landauer bound in phase space rather than merely in energy: the erasure must cost at least $\hbar$ of action in addition to $k_B T \ln 2$ of energy. This stronger version of the bound is a direct consequence of the Kinematic Snap mechanism.

3.2 The Maxwell's Demon Paradox Dissolved

The Maxwell's demon (Maxwell, 1872) paradox asks how a microscopic agent that sorts fast and slow molecules can apparently violate the second law of thermodynamics, and what physical mechanism restores it. The standard resolution via Landauer's principle (Bennett, 1982) argues that the demon's memory erasure, not the measurement, pays the thermodynamic cost. This resolution, while correct, involves an information-theoretic postulate (the equivalence of physical and logical entropy) that RA makes unnecessary.

In RA, the demon is a physical system embedded in the causal DAG. Every act of "measurement" by the demon is an actualization event: the demon's sensory apparatus exchanges on-shell bosons with the gas molecules in order to determine their velocities. Each such exchange writes a vertex to the DAG and pays the Landauer cost $W_{\text{act}} \geq k_B T \ln 2$ at the moment of measurement, not at the moment of memory erasure.

The standard resolution delays the cost to erasure; RA moves the cost to measurement. Both arrive at the same thermodynamic accounting, but through different physical pictures. The RA picture is arguably more natural: the cost is paid when the physical interaction occurs, not when a logical register is cleared.

More precisely: there is no gap in RA between the demon's information acquisition and the physical cost of that acquisition. The demon's acquisition of knowledge about a molecule's velocity *just is* the on-shell boson exchange between the demon's sensory apparatus and the molecule. The thermodynamic cost is paid at that moment. There is no separate step of "storing the information" and "later erasing it." The storage and the cost are the same physical event.

This removes the apparently paradoxical temporal gap in the standard resolution. In the standard account, the demon seemingly operates for free during the sorting phase and pays later during erasure—a picture that requires careful argument to prevent apparent violations of the second law during the sorting phase itself. In RA, the cost is paid continuously and locally during the measurement interactions, and the second law is enforced vertex by vertex as the DAG grows.

Weak measurements. The argument above applies to strong projective measurements, which constitute unambiguous actualization events in RA. Weak measurements—continuous monitoring that acquires partial information without fully collapsing the quantum state—are a more subtle case. A weak measurement that does not produce an on-shell environmental record is not, in RA, a full actualization event; it is a pre-actualization interaction that keeps the system in the reversible potentia regime. Whether and how the thermodynamic cost accumulates continuously during weak measurement, and at what point the Kinematic Snap threshold is crossed, is an open question that connects to the measurement back-action literature and constitutes a natural target for the RA framework's extension to continuous monitoring scenarios.

3.3 The Fluctuation Theorems with a Precise Actualization Criterion

The Jarzynski equality (Jarzynski, 1997) and Crooks fluctuation theorem (Crooks, 1999) relate the distribution of work performed on a system driven away from equilibrium to the free energy difference between initial and final states:

$$\langle e^{-W/k_B T} \rangle = e^{-\Delta F/k_B T} \quad \text{(Jarzynski)}, \tag{11}$$

$$\frac{P(+W)}{P(-W)} = e^{(W-\Delta F)/k_B T} \quad \text{(Crooks)}. \tag{12}$$

These theorems hold for systems driven arbitrarily far from equilibrium and have been verified in numerous experiments. They are currently derived from the assumption of microscopic reversibility (detailed balance) and the identification of work with energy exchanges during the driving protocol.

RA provides a physical decomposition of the work distribution in these theorems. An important caveat governs the following discussion: the standard work W in the Jarzynski and Crooks theorems

is defined as the total energy exchanged with the external driving agent during the protocol. If RA were to redefine W to include only on-shell exchange events—call it W_{RA}—then $W_{\mathrm{RA}} \neq W$ in general, and there is no a priori reason the standard equalities hold for the restricted quantity W_{RA} without rederivation. The analysis below therefore proceeds differently: it uses the standard W and treats RA as providing a physical decomposition of the *histories* that contribute to the work distribution, without redefining W itself.

The work distribution $P(W)$ decomposes over actualization sequences. In RA, each trajectory contributing to $P(W)$ is a causal DAG history: a specific sequence of on-shell exchange events. Virtual, off-shell exchanges occur within each trajectory but do not constitute irreversible events. The standard W for each trajectory is the total energy exchange; RA adds the physical picture that this energy is concentrated in the on-shell exchange events that write DAG vertices.

The forward and reverse protocols differ in their DAG histories. In standard statistical mechanics, detailed balance is stated over phase space trajectories. RA proposes that the more fundamental objects are causal DAG histories. The Crooks theorem, in this picture, says the ratio of probabilities of a DAG history and its time-reverse is $e^{(W-\Delta F)/k_B T}$—where W is the standard work, computed over the full trajectory including virtual exchanges.

Protocols that remain close to off-shell have fewer DAG vertices and contribute disproportionately to the low-work tail of $P(W)$. This is the physical picture of why quasi-static (nearly reversible) protocols dominate the Jarzynski average: they generate fewer irreversible actualization events per unit time.

This decomposition suggests a research program rather than a completed result: the Jarzynski equality, restated as a sum over actualization sequences with the correct weighting, may be more tractable for discrete few-body quantum systems where the DAG is a finite combinatorial object. Whether the equality holds for a suitably defined W_{RA} (restricted to on-shell contributions) is an open question whose answer would either strengthen or constrain the RA framework.

3.4 The Arrow of Time as Structural, Not Statistical

Standard statistical mechanics derives the thermodynamic arrow of time from the second law, which is in turn grounded in the initial low-entropy state of the universe. This derivation is probabilistic: it says that entropy-increasing processes are overwhelmingly more probable than entropy-decreasing ones, given typical initial conditions. The underlying microscopic dynamics remain time-reversible.

RA provides a structural account of the arrow of time that does not rely on initial conditions or probabilistic arguments. The causal DAG grows in one direction only: new vertices are added; old vertices are never removed. This is not a consequence of the second law; it is a consequence of the ontological commitment that actualization is primitive and irreversible. The past is fixed because it has been written into the DAG. The future is open because it has not.

This distinction has a precise thermodynamic consequence. In standard statistical mechanics, a process that violates the second law is *possible* (it is merely improbable). In RA, a process that removes a causal DAG vertex is not merely improbable—it is categorically excluded by the ontology. The arrow of time is not a statistical asymmetry; it is a logical one.

For quantum thermodynamics, this matters because many results in the field depend on the assumption of microscopic reversibility. RA does not violate this assumption at the level of the quantum dynamics—the Schrödinger equation is still time-reversible—but it adds an irreversible layer: the writing of DAG vertices. The interplay between these two levels (reversible quantum dynamics and irreversible DAG inscription) is precisely the interplay between pre-actualization and post-actualization regimes that RAQM characterises via Prediction P3.

4 The Quantum Information Theory Connection

4.1 Relative Entropy as the Fundamental Actualization Criterion

The quantum relative entropy $S(\rho\|\sigma) = \mathrm{Tr}[\rho(\log\rho - \log\sigma)]$ plays a central role in both quantum information theory and the RA actualization criterion. In RAGC (Sandeman, 2026b), the fundamental criterion for an actualization event is stated in terms of relative entropy with respect to the vacuum: an actualization occurs when an interaction produces a strictly irreversible increase in $S(\rho\|\sigma_0)$, where σ_0 is the Minkowski vacuum state.

This criterion is frame-independent by the following chain of results:

1. $S(U\rho U^\dagger\|U\sigma U^\dagger) = S(\rho\|\sigma)$ for any unitary U — proved using `log_unitary_conj`, which is a one-line corollary of `Matrix.cfc_conj_unitary` from the Lean-QuantumInfo library (Meiburg et al., 2025).
2. The Minkowski vacuum is Poincaré-invariant: $U\sigma_0U^\dagger = \sigma_0$ — a QFT physics input.
3. Therefore $S(U\rho U^\dagger\|\sigma_0) = S(\rho\|\sigma_0)$ — the compiled theorem `frame_independence` in `RA_AQFT_Proofs_v10.l`

This means the actualization criterion is a covariant scalar: it is the same in all inertial and uniformly accelerated frames. The Rindler thermal bath, seen by an accelerated observer, has $\Delta S_{\mathrm{rel}} = 0$—it is stationary with respect to the vacuum and does not constitute a sequence of actualization events. This is the compiled theorem `rindler_stationarity`.

4.2 The Data Processing Inequality and Causal Structure

The data processing inequality—that relative entropy can only decrease under the application of a completely positive trace-preserving (CPTP) map, $S(\mathscr{E}(\rho)\|\mathscr{E}(\sigma)) \leq S(\rho\|\sigma)$—has a natural RA interpretation. A CPTP map represents a physical process that may include actualization events (on-shell exchanges). Each such event can only reduce the distinguishability between states ρ and σ as seen from outside the system. Information is lost from the perspective of the environment when a DAG vertex is written; it is not created.

The RA formulation of the data processing inequality is therefore a statement about causal DAG evolution: each actualization step can only reduce relative entropy, not increase it. An increase in relative entropy $S(\rho\|\sigma_0)$—an actualization event—necessarily involves a physical interaction, not merely a CPTP map. This distinguishes the two directions of the inequality sharply: decreasing relative entropy corresponds to physical processes that may or may not involve actualization; increasing relative entropy (an actualization event) requires specific physical conditions (on-shell exchange).

The Petz monotonicity theorem (Petz, 1986), which establishes the data processing inequality for quantum relative entropy, is formalised in Lean-QuantumInfo (Meiburg et al., 2025) as part of the Generalised Quantum Stein's Lemma proof infrastructure. This is the mathematical substrate for the RA AQFT proof program.

4.3 Quantum Hypothesis Testing and the Actualization Criterion

The Generalized Quantum Stein's Lemma (Meiburg et al., 2025) establishes the operational significance of quantum relative entropy in the context of quantum resource theories: it determines the optimal error exponents for discriminating between a quantum state and a set of free states. We propose the following RA interpretation of this result: $S(\rho\|\sigma_0)$ quantifies how distinguishable the current state ρ is from the vacuum σ_0, which is precisely the quantity whose increase constitutes an actualization event. The GQSL itself makes no claim about RA; the connection is a proposed conceptual bridge, not a derived result.

Under this interpretation, the Stein's Lemma suggests an operational picture of the actualization threshold: an actualization event corresponds to the system crossing from a regime where ρ is statistically indistinguishable from σ_0 (in the quantum hypothesis testing sense) to one where the

distinguishability is irreversibly established. The type-I and type-II error exponents in the Stein's Lemma would then determine the precision with which this threshold can be located experimentally.

This connection suggests a natural experimental program: quantum hypothesis testing experiments that probe the vicinity of the actualization threshold could provide direct tests of the RA criterion. Specifically, the sharp threshold predicted by RAQM (Prediction P3) would manifest as a sharp transition in the hypothesis testing error exponents as the environmental coupling crosses the on-shell threshold.

5 Empirical Predictions and Experimental Touchstones

The implications developed above are conditional on the correctness of the RA framework. The following predictions, inherited from RAQM and RAGC, provide the experimental tests.

5.1 Predictions Relevant to Quantum Computing (P1–P3)

P1: Finite actualization propagation velocity. The actualization front in a spatially extended many-body quantum system propagates at a velocity bounded by the Lieb-Robinson speed (Lieb and Robinson, 1972) $v_{\mathrm{LR}} \sim J\ell/\hbar$. A decoherence event at site A should produce a time delay $\Delta t \geq |r_A - r_B|/v_{\mathrm{LR}}$ before off-diagonal coherences at site B begin to collapse. Instantaneous collapse models predict no such delay.

P2: Environment-dependent transition duration. The actualization timescale scales as $\tau_{\mathrm{act}} \sim \hbar/\Delta E_{\mathrm{env}}$. Systematically varying ΔE_{env} in a controlled superconducting circuit experiment should produce a proportional variation in transition duration. This is consistent with Minev et al. (Minev et al., 2019) and Pazourek et al. (Pazourek et al., 2015); the distinctive prediction is the *proportional* scaling.

P3: Sharp reversibility threshold. There is a sharp threshold between pre-actualization (reversible) and post-actualization (irreversible) dynamics, determined by the environmental entanglement entropy crossing a critical value. Identifying this threshold experimentally constitutes a direct measurement of the actualization criterion. Decoherence-only models predict no such sharp threshold.

5.2 Predictions Relevant to Quantum Thermodynamics

P4 (from P2): Landauer timescale. In addition to the Landauer energy bound $W_{\mathrm{act}} \geq k_B T \ln 2$, RA predicts a Landauer *timescale*: erasure cannot be completed faster than $\tau_{\mathrm{min}} \sim \hbar/\Delta E_{\mathrm{env}}$. This sets a joint energy-time bound on erasure that is stronger than the standard Landauer bound and is, in principle, measurable in fast qubit experiments.

P5: Off-shell regime is thermodynamically inert. Interactions that remain kinematically constrained to the virtual, off-shell regime do not dissipate energy in the thermodynamic sense. This is testable in carefully engineered systems where the dominant qubit-environment coupling can be tuned below the on-shell threshold, predicting a qualitative suppression of thermodynamic dissipation rather than a quantitative reduction.

6 Discussion

6.1 What RA Provides That Existing Frameworks Do Not

The implications developed in Sections 2–4 share a common structure: RA provides a *mechanism* where existing frameworks provide only *statistics*. Standard decoherence theory tells you the rate at which coherence is lost; RA tells you the physical process by which each individual coherence

loss occurs, and gives it a precise timing. Standard quantum thermodynamics invokes entropy and irreversibility; RA grounds these in the discrete, irreversible growth of a causal DAG.

This shift from statistical to mechanical description is not merely aesthetic. It has three practical consequences:

First, it gives engineering targets that are physical rather than phenomenological. The target is ΔE_{env}—a quantity with a direct physical interpretation—rather than T_2—a phenomenological timescale whose physical determinants must be inferred separately.

Second, it identifies qualitative distinctions (off-shell versus on-shell) that the statistical description treats as quantitative ones (strong coupling versus weak coupling). The difference between a system that is decoherence-immune because it is off-shell and a system that is weakly decohering because it is weakly coupled is qualitative in RA and quantitative in standard theory. That qualitative distinction may be experimentally accessible.

Third, it connects the two fields to each other and to the broader RA framework. Quantum computing and quantum thermodynamics share the relative entropy as their central quantity; RA gives that quantity a physical interpretation as the actualization criterion; and the frame-independence of that criterion is a compiled Lean theorem. The mathematical infrastructure is shared.

6.2 Scope and Limitations

The implications in this paper are derived under the assumption that the RA actualization criterion is correct. That assumption is currently under peer review in the companion paper RAQM (Sandeman, 2026a). Several implications— particularly the qualitative decoherence-immunity of the off-shell regime (Section 2) and the Landauer timescale bound (P4)—go beyond what existing experimental data require and constitute genuine predictions that may falsify the framework if tested and found wanting. This is the appropriate scientific posture; the implications are stated as conditionals, not assertions.

The Lean-verified results cited in Section 4 establish the mathematical consistency of the AQFT proof program in finite dimensions, with three axioms: the CFC unitary-conjugation lemma (sourced from Lean-QuantumInfo), the Poincaré-invariance of the vacuum (a QFT physics input), and the data processing inequality (proved in Lean-QuantumInfo). The extension to the full infinite-dimensional type III_1 algebra setting remains the primary open mathematical target.

7 Conclusion

Relational Actualism proposes that the precise physical mechanism of quantum irreversibility is the crossing of the on-shell dispersion relation by a mediating boson. This paper has developed the implications of this proposal for quantum computing and quantum thermodynamics. If correct, RA gives both fields what they currently lack: a precise physical account of when and why quantum processes become irreversible.

For quantum computing: the engineering target is the environmental energy ΔE_{env}; the off-shell regime is categorically decoherence-immune; the threshold theorem acquires a kinematic interpretation; and the distinction between reversible and irreversible errors is physically grounded.

For quantum thermodynamics: Landauer's principle has a mechanical derivation with an additional timescale bound; the Maxwell's demon pays its thermodynamic cost at measurement, not at erasure; the fluctuation theorems are restated as sums over actualization sequences; and the arrow of time is structural rather than statistical.

The shared mathematical substrate—quantum relative entropy, CPTP maps, the data processing inequality—connects both fields to the RA AQFT proof program and to the Lean-QuantumInfo library (Meiburg et al., 2025) in which key results are formally verified. The frame-independence of the actualization criterion is a compiled theorem. The remaining mathematical frontier is the extension from finite dimensions to the type III_1 von Neumann algebra setting.

The framework makes specific, discriminating predictions (Section 5) that are testable with

existing or near-future experimental platforms. Those tests are the appropriate next step.

References

C. H. Bennett. The thermodynamics of computation—a review. *International Journal of Theoretical Physics*, 21:905–940, 1982.

G. E. Crooks. Entropy production fluctuation theorem and the nonequilibrium work relation for free energy differences. *Physical Review E*, 60(3):2721–2726, 1999.

S. F. Huelga, C. Macchiavello, T. Pellizzari, A. K. Ekert, M. B. Plenio, and J. I. Cirac. Improvement of frequency standards with quantum entanglement. *Physical Review Letters*, 79(20):3865–3868, 1997.

C. Jarzynski. Nonequilibrium equality for free energy differences. *Physical Review Letters*, 78(14):2690–2693, 1997.

R. Landauer. Irreversibility and heat generation in the computing process. *IBM Journal of Research and Development*, 5(3):183–191, 1961.

E. H. Lieb and D. W. Robinson. The finite group velocity of quantum spin systems. *Communications in Mathematical Physics*, 28(3):251–257, 1972.

J. C. Maxwell. *Theory of Heat*. Longmans, Green and Co., London, 4th edition, 1872.

A. Meiburg, L. A. Lessa, and R. R. Soldati. A formalization of the generalized quantum Stein's lemma in Lean. *arXiv*:2510.08672, 2025. Lean-QuantumInfo library: `https://github.com/Timeroot/Lean-QuantumInfo`.

Z. K. Minev, S. O. Mundhada, S. Shankar, P. Reinhold, R. Gutiérrez-Jáuregui, R. J. Schoelkopf, M. Mirrahimi, H. J. Carmichael, and M. H. Devoret. To catch and reverse a quantum jump mid-flight. *Nature*, 570:200–204, 2019.

R. Pazourek, S. Nagele, and J. Burgdörfer. Attosecond chronoscopy of photoemission. *Reviews of Modern Physics*, 87(3):765–802, 2015.

D. Petz. Sufficient subalgebras and the relative entropy of states of a von Neumann algebra. *Communications in Mathematical Physics*, 105(1):123–131, 1986.

D. Reeb and M. M. Wolf. An improved Landauer principle with finite-size corrections. *New Journal of Physics*, 16(10):103011, 2014.

J. F. Sandeman. Algorithmic Causal Dynamics: A Relational Actualist Framework for Emergence, Autopoiesis, and Agentic Intelligence (RAHC). Preprint, 2026.

J. F. Sandeman. The relational universe: Interaction and the transition of the possible to the actual. Submitted to *Foundations of Physics*, 2026.

J. F. Sandeman. Relational actualism in curved spacetime: Gravity, cosmology, and the causal DAG. Preprint, 2026.

J. F. Sandeman. Relational actualism and the complexity of intelligence: Assembly theory, causal depth, and the Boltzmann brain. Preprint, 2026.