

# FONTAINE RINGS AND $p$ -ADIC $L$ -FUNCTIONS

## CONTENTS

1. One variable $p$ -adic functions	2
1.1. Functions on $\mathbf{Z}_p$	2
1.2. Distributions on $\mathbf{Z}_p$	4
1.3. Operations on the distributions	4
2. Kubota-Leopoldt $p$ -adic $L$ -functions	5
2.1. Riemann zeta function	5
2.2. Kummer congruence	6
2.3. $p$ -adic Mellin transform and Leopoldt's $\Gamma$ -transform	8
2.4. Construction of the Kubota-Leopoldt zeta function	9
2.5. The residue at $s = 1$ and the $p$ -adic zeta function	10
2.6. Dirichlet $L$ -function	11
2.7. $p$ -adic $L$ -function attaches to Dirichlet character	11
2.8. Behavior at $s = 1$ of Dirichlet $L$ -function	12
2.9. Twist by a character of conductor power of $p$	13
3. $(\varphi, \Gamma)$ -modules and $p$ -adic representations	14
3.1. The field $\tilde{\mathbf{E}}$ and its subrings.	15
3.2. The field $\tilde{\mathbf{B}}$ and its subrings	15
3.3. $(\varphi, \Gamma)$ -module and Galois representations	16
4. $(\varphi, \Gamma)$ -modules and Galois cohomology	17
4.1. The complex $C_{\varphi, \gamma}(K, V)$	17
4.2. The operator $\psi$	19
4.3. The compactness of $D(V)^{\psi=1}$	20
4.4. The module $\frac{D(V)}{\psi-1}$	22
5. Iwasawa theory and $p$ -adic representations	23
5.1. Iwasawa cohomology	23
5.2. Corestriction and $(\varphi, \Gamma)$ -modules	25
5.3. Interpretation of $D(V)^{\psi=1}$ and $\frac{D(V)}{\psi-1}$ in Iwasawa theory	26
6. De Rham representations and overconvergent representations	27
6.1. De Rham representations and crystalline representations	27
6.2. Overconvergent elements	28
6.3. Overconvergent representations	30
7. Explicit reciprocity laws and de Rham Representation	30
7.1. The Bloch-Kato exponential map and its dual	30
7.2. Explicit reciprocity law	32
7.3. Connection with the Perrin-Riou's logarithm	34
8. The $\mathbf{Q}_p(1)$ representation and Coleman's power series	34
8.1. The module $D(\mathbf{Z}_p(1))^{\psi=1}$	34
8.2. Kummer theory	35

8.3.	Multiplicative representatives	35
8.4.	Generalized Coleman's power series	36
8.5.	The map $\text{Log}_{\mathbf{Q}_p(1)}$ and $\text{Exp}_{\mathbf{Q}_p}^*$	37
8.6.	Cyclotomic units and Coates-Wiles homomorphisms	38
9.	$(\varphi, \Gamma)$ -modules and differential equations	40
9.1.	The rings $\mathbf{B}_{\max}$ and $\tilde{\mathbf{B}}_{\text{rig}}^+$	40
9.2.	The structure of $D(T)^{\psi=1}$	41
9.3.	$p$ -adic representations and differential equations	42
9.4.	The Fontaine isomorphism revisit	43
9.5.	Iwasawa algebra and power series	45
9.6.	Iwasawa algebras and differential equations	45
10.	Bloch-Kato's exponential maps: Three explicit reciprocity formulas	47
10.1.	The Bloch-Kato's exponential map and its dual revisit	47
10.2.	Perrin-Riou's big exponential map	50
10.3.	The explicit reciprocity formula	53
11.	Perrin-Riou's big regulator map	54
11.1.	Perrin-Riou's big logarithm map	54
11.2.	Cyclotomic units and Kubota-Leopoldt $p$ -adic $L$ -functions	57
	References	58

## 1. ONE VARIABLE $p$ -ADIC FUNCTIONS

In this section, we denote  $L$  a closed subfield of  $\mathbf{C}_p$ .

**1.1. Functions on  $\mathbf{Z}_p$ .** Let  $\mathcal{C}^0(\mathbf{Z}_p, L)$  be the space of continuous function from  $\mathbf{Z}_p$  to  $L$ . Since  $\mathbf{Z}_p$  is compact, every continuous function on  $\mathbf{Z}_p$  is bounded. This allows us to define a valuation  $v_{\mathcal{C}^0}$  on  $\mathcal{C}^0(\mathbf{Z}_p, L)$  by  $v_{\mathcal{C}^0}(\phi) = \inf_{x \in \mathbf{Z}_p} (\phi(x))$ , which makes  $\mathcal{C}^0(\mathbf{Z}_p, L)$  a  $L$ -Banach space.

If  $n \in \mathbf{N}$ , let  $\binom{x}{n}$  be the polynomial defined by

$$\binom{x}{n} = \begin{cases} 1 & \text{if } n = 0 \\ \frac{x(x-1)\cdots(x-n+1)}{n!} & \text{if } n \geq 1. \end{cases}$$

**Theorem 1.1.** (Mahler)  $\{\binom{x}{n}, n \in \mathbf{N}\}$  forms a Banach basis of  $\mathcal{C}^0(\mathbf{Z}_p, L)$ .

If  $h \in \mathbf{N}$ , let  $\text{LA}_h(\mathbf{Z}_p, L)$  be the space of functions from  $\mathbf{Z}_p$  to  $L$  which is analytic on  $a + p^h \mathbf{Z}_p$  for all  $a \in \mathbf{Z}_p$ , that is, if  $\phi \in \text{LA}_h(\mathbf{Z}_p, L)$ ,  $x_0 \in \mathbf{Z}_p$  and  $x \in x_0 + p^h \mathbf{Z}_p$ , then  $\phi$  can be written as the form

$$\phi(x) = \sum_{k=0}^{\infty} a_k(\phi, x_0)(x - x_0)^k,$$

where  $a_k(x_0, \phi)$  is a sequence in  $L$  such that  $\nu_p(a_k(\phi, x_0)) + kh$  tends to  $+\infty$  as  $k$  tends to  $+\infty$ . We endow  $\text{LA}_h(\mathbf{Z}_p, L)$  a valuation  $v_{\text{LA}_h}$  defined by

$$v_{\text{LA}_h}(\phi) = \inf_{x_0 \in \mathbf{Z}_p} \inf_{k \in \mathbf{N}} \nu_p(a_k(\phi, x_0)) + kh,$$

which makes  $\text{LA}_h(\mathbf{Z}_p, L)$  a  $L$ -Banach space. One can show that  $v_{\text{LA}_h}(\phi) = \inf_{a \in S} \inf_{k \in \mathbf{N}} \nu_p(\phi, a_k(a)) + kh$  where  $S$  is a representative of  $\mathbf{Z}_p/p^h \mathbf{Z}_p$ .

We denote  $\mathrm{LA}(\mathbf{Z}_p, L)$  the space of locally analytic functions on  $\mathbf{Z}_p$ . Since  $\mathbf{Z}_p$  is compact, it is an inductive limit of  $\mathrm{LA}_h(\mathbf{Z}_p, L)$ ,  $h \in \mathbf{N}$ , and we endow it with the inductive limit topology.

**Theorem 1.2.** (Amice)  $\{[\frac{n}{p^h}]!(\frac{x}{n}), n \in \mathbf{N}\}$  forms a Banach basis of  $\mathrm{LA}_h(\mathbf{Z}_p, L)$ .

**Theorem 1.3.** The function  $\phi = \sum_{n=0}^{+\infty} a_n(\phi)(\frac{x}{n}) \in \mathcal{C}^0(\mathbf{Z}_p, L)$  is in  $\mathrm{LA}(\mathbf{Z}_p, L)$  if and only if  $\liminf \frac{1}{n} \nu_p(a_n(\phi)) > 0$ .

A function  $\phi : \mathbf{Z}_p \rightarrow L$  is differentiable at  $x_0 \in \mathbf{Z}_p$  if  $\lim_{h \rightarrow 0} \frac{\phi(x_0+h) - \phi(x_0)}{h}$  exists. The limit is denoted by  $\phi'(x_0)$ . A function is said to be differentiable of order 1 if it is differentiable at all  $x_0 \in \mathbf{Z}_p$ . We say a function is differentiable of order  $k$  if its differentiation is of order  $k-1$ .

If  $r \geq 0$ , we say that  $\phi : \mathbf{Z}_p \rightarrow L$  is of class  $\mathcal{C}^r$  if there exist functions  $\phi^{(j)} : \mathbf{Z}_p \rightarrow L$  for  $0 \leq j \leq [r]$ , such that, if we define  $\varepsilon_{\phi,r} : \mathbf{Z}_p \times \mathbf{Z}_p \rightarrow L$  and  $C_{\phi,r} : \mathbf{N} \rightarrow \mathbb{R} \cup \{+\infty\}$  by

$$\varepsilon_{\phi,r}(x, y) = \phi(x+y) - \sum_{j=0}^{[r]} \phi^{(j)}(x) \frac{y^j}{j!} \quad \text{and} \quad C_{\phi,r}(h) = \inf_{x \in \mathbf{Z}_p, y \in p^h \mathbf{Z}_p} \nu_p(\varepsilon_{\phi,r}(x, y)) - rh,$$

then  $C_{\phi,r}(h)$  tends to  $+\infty$  as  $h$  tends to  $+\infty$ .

We denote  $\mathcal{C}^r(\mathbf{Z}_p, L)$  the set of functions  $\phi : \mathbf{Z}_p \rightarrow L$  of class  $\mathcal{C}^r$ . We endow  $\mathcal{C}^r(\mathbf{Z}_p, L)$  the valuation  $v_{\mathcal{C}^r}$  defined by

$$v_{\mathcal{C}^r}(\phi) = \inf \left( \inf_{0 \leq j \leq [r], x \in \mathbf{Z}_p} \nu_p\left(\frac{\phi^{(j)}(x)}{j!}\right), \inf_{x, y \in \mathbf{Z}_p} \nu_p(\varepsilon_{\phi,r}(x, y) - r\nu_p(y)) \right),$$

which makes it a  $L$ -Banach space.

**Proposition 1.4.** If  $h \in \mathbf{N}$ , and if  $r \geq 0$ , then  $\mathrm{LA}_h(\mathbf{Z}_p, L) \subset \mathcal{C}^r(\mathbf{Z}_p, L)$ . Moreover, if  $\phi \in \mathrm{LA}_h(\mathbf{Z}_p, L)$ , then

$$v_{\mathcal{C}^r}(\phi) \geq v_{\mathrm{LA}_h}(\phi) - rh.$$

*Proof.* See [Col10, proposition I.5.7]. □

If  $i \in \mathbf{N}$ , we denote  $l(i)$  the least integer  $n$  such that  $p^n > i$ . We have

$$l(0) = 0 \quad \text{and} \quad l(i) = \left\lceil \frac{\log i}{\log p} \right\rceil + 1, \text{ if } i \geq 1.$$

**Theorem 1.5.** (Mahler) The function  $\phi = \sum_{n=0}^{+\infty} a_n(\phi)(\frac{x}{n}) \in \mathcal{C}^0(\mathbf{Z}_p, L)$  is in  $\mathcal{C}^r(\mathbf{Z}_p, L)$ ,  $r \geq 0$  if and only if  $\nu_p(a_n(\phi)) - rl(n) \rightarrow +\infty$  as  $n \rightarrow +\infty$ . Moreover, the valuation  $v'_{\mathcal{C}^r}$  defined on  $\mathcal{C}^r(\mathbf{Z}_p, L)$  by the formula

$$v'_{\mathcal{C}^r}(\phi) = \inf_{n \in \mathbf{N}} \left( \nu_p(a_n(\phi)) - rl(n) \right)$$

is equivalent to the valuation  $v_{\mathcal{C}^r}$ .

*Proof.* See [Col10, proposition I.5.18]. □

**Corollary 1.6.**  $p^{[rl(n)]}(\frac{x}{n}), n \in \mathbf{N}$  forms a Banach basis of  $\mathcal{C}^r(\mathbf{Z}_p, L)$ .

↑  
you mean flow. L 1?

**1.2. Distributions on  $\mathbf{Z}_p$ .** A continuous distribution on  $\mathbf{Z}_p$  is a continuous linear function on  $\text{LA}(\mathbf{Z}_p, L)$ , that is, a linear function on  $\text{LA}(\mathbf{Z}_p, L)$  whose restriction to  $\text{LA}_h(\mathbf{Z}_p, L)$  is continuous. We denote  $\mathcal{D}(\mathbf{Z}_p, L)$  the set of continuous distributions on  $\mathbf{Z}_p$  with values in  $L$  and endow  $\mathcal{D}(\mathbf{Z}_p, L)$  with the Fréchet topology defined by the family of valuations  $v_{\text{LA}_h}$ ,  $h \in \mathbf{N}$ .

For a continuous distribution  $\mu$ , we associate it with the formal series:

$$\mathcal{A}_\mu(T) = \sum_{n=0}^{+\infty} \int_{\mathbf{Z}_p} (1+T)^x \mu = \sum_{n=0}^{+\infty} T^n \int_{\mathbf{Z}_p} \binom{x}{n} \mu,$$

which is called the Amice transform of  $\mu$ .

**Lemma 1.7.** *If  $\mu \in \mathcal{D}(\mathbf{Z}_p, L)$  and if  $\nu_p(x) > 0$ , then  $\int_{\mathbf{Z}_p} (1+z)^x \mu(x) = \mathcal{A}_\mu(z)$ .*

Let  $\mathcal{R}^+$  be the ring of power series  $f = \sum_{n=0}^{\infty} a_n T^n$  with coefficients in  $L$ , which is convergent if  $\nu_p(T) > 0$ .

*that* We say an element  $f = \sum_{n=0}^{\infty} a_n T^n \in \mathcal{R}^+$  is of order  $r$  if  $\nu_p(a_n) + rl(n)$  is bounded. We denote  $\mathcal{R}_h^+$  the subset of  $\mathcal{R}^+$  of elements of order  $r$ , and we endow  $\mathcal{R}_h^+$  the valuation  $v_r$  defined by  $v_r(f) = \inf_{n \in \mathbf{N}} \nu_p(a_n) + rl(n)$ , which makes it a  $L$ -Banach space. We endow  $\mathcal{R}^+$  the Fréchet topology defined by the family of valuations  $v_r$ .

**Theorem 1.8.** *The map  $\mu \mapsto \mathcal{A}_\mu$  is an isomorphism of Fréchet space from  $\mathcal{D}(\mathbf{Z}_p, L)$  to  $\mathcal{R}^+$ .*

*Proof.* See [Col10, Theorem II.2.2]. □

If  $r \geq 0$ , we say a continuous distribution  $\mu$  on  $\mathbf{Z}_p$  is of order  $r$  if it can be extended by continuity to  $\mathcal{C}^r$ . We denote  $\mathcal{D}_r(\mathbf{Z}_p, L)$  the set of distributions of order  $r$ , which is equipped with a valuation  $v_{\mathcal{D}_r}$  defined by

$$v_{\mathcal{D}_r}(\mu) = \inf_{f \in \mathcal{C}^r(\mathbf{Z}_p, L) - \{0\}} \left( \nu_p \left( \int_{\mathbf{Z}_p} f \mu \right) - v_{\mathcal{C}^r}(f) \right),$$

which gives  $\mathcal{D}_r(\mathbf{Z}_p, L)$  the dual topology of  $\mathcal{C}^r(\mathbf{Z}_p, L)$ .

A distribution is said to be tempered if there exist  $r \in \mathbb{R}^+$  such that it is of order  $r$ . We denote  $\mathcal{D}_{\text{temp}}(\mathbf{Z}_p, L)$  the space of tempered distributions.

**Proposition 1.9.** *The map  $\mu \mapsto \mathcal{A}_\mu$  induces an isometry from  $\mathcal{D}_r(\mathbf{Z}_p, L)$  equipped with valuation  $v_{\mathcal{D}_r}$  to  $\mathcal{R}_r^+$  equipped with valuation  $v_r$ .*

A distribution of order 0 is called the measure. By definition,  $\mathcal{D}_0(\mathbf{Z}_p, L)$  is the topological dual of the space of continuous functions. By proposition 1.9, we have a one-one correspondence from a measure to a power series of bounded coefficients.

### 1.3. Operations on the distributions.

1. Harr measure:  $\mu(\mathbf{Z}_p) = 1$  and  $\mu$  is invariant by translation. We must have  $\mu(i + p^n \mathbf{Z}_p) = \frac{1}{p^n}$  which is not bounded. Hence there exists no Harr measure on  $\mathbf{Z}_p$ .
2. Dirac measure: For  $a \in \mathbf{Z}_p$ , we define  $\delta_a$  the Dirac measure associated to  $f(a)$ . The Amice transform of  $\delta_a$  is  $\mathcal{A}_{\delta_a}(T) = (1+T)^a$ .
3. Multiplication by a function: If  $\mu$  is a distribution on  $\mathbf{Z}_p$  and  $f$  is a locally analytic function on  $\mathbf{Z}_p$ , we define the distribution  $f\mu$  by  $\int_{\mathbf{Z}_p} \phi(f\mu) = \int_{\mathbf{Z}_p} (f\phi)\mu$ .
  - Multiplication by  $x$ : We have  $x \cdot \binom{x}{n} = ((x-n) + n) \binom{x}{n} = (n+1) \binom{x}{n+1} + n \binom{x}{n}$ , hence we have

$$\mathcal{A}_{x\mu}(T) = \partial \mathcal{A}_\mu \quad \text{where } \partial = (1+T) \frac{d}{dT}.$$

- Multiplication by  $z^x$  ~~if~~ <sup>when</sup>  $\nu_p(z-1) > 0$ : By lemma 1.7, if  $\nu_p(y-1) > 0$ , and if  $\lambda$  is a continuous distribution on  $\mathbf{Z}_p$ , then  $\int_{\mathbf{Z}_p} y^x \lambda(x) = \mathcal{A}_\lambda(y-1)$ . Applying this to  $\lambda = z^x \mu$ , we obtain  $\mathcal{A}_\lambda(y-1) = \mathcal{A}_\mu(yz-1)$ . We ~~hence~~ <sup>then</sup> have the formula

$$\mathcal{A}_{z^x \mu}(T) = \mathcal{A}_\mu((1+T)z-1). \quad \text{then the}$$

4. Restriction to compact open subset: If  $X$  is a compact open subset of  $\mathbf{Z}_p$ , then characteristic function  $1_X$  is continuous on  $\mathbf{Z}_p$ . If  $\mu$  is a distribution on  $\mathbf{Z}_p$ , the measure  $1_X \mu$  is the restriction of  $\mu$  to  $X$  and is denoted by  $\text{Res}_X(\mu)$ . In particular for  $n \in \mathbf{N}$  and  $a \in \mathbf{Z}_p$ , we have  $1_{a+p^n \mathbf{Z}_p}(x) = p^{-n} \sum_{z p^n=1} z^{-a} z^x$ , hence

$$\mathcal{A}_{\text{Res}_{a+p^n \mathbf{Z}_p}(\mu)}(T) = p^{-n} \sum_{z p^n=1} z^{-a} \mathcal{A}_\mu((1+T)z-1).$$

5. Derivation of distribution: If  $\mu \in \mathcal{D}(\mathbf{Z}_p, L)$ , we define  $d\mu$  by

$$\int_{\mathbf{Z}_p} \phi(x) d\mu = \int_{\mathbf{Z}_p} \phi'(x) \mu, \quad \text{and therefore} \quad \mathcal{A}_{d\mu}(T) = \log(1+T) \cdot \mathcal{A}_\mu(T).$$

6. Actions of  $\mathbf{Z}_p^*$ ,  $\varphi$  and  $\psi$ :

- If  $a \in \mathbf{Z}_p^*$ , and if  $\mu \in \mathcal{D}(\mathbf{Z}_p, L)$ , we define  $\sigma_a(\mu) \in \mathcal{D}(\mathbf{Z}_p, L)$  by

$$\int_{\mathbf{Z}_p} \phi(x) \sigma_a(\mu) = \int_{\mathbf{Z}_p} \phi(ax) \mu, \quad \text{and therefore} \quad \mathcal{A}_{\sigma_a(\mu)}(T) = \mathcal{A}_\mu((1+T)^a - 1).$$

- $\varphi$  acts on distribution  $\mu$  by

$$\int_{\mathbf{Z}_p} \phi(x) \varphi(\mu) = \int_{\mathbf{Z}_p} \phi(px) \mu, \quad \text{and therefore} \quad \mathcal{A}_{\varphi(\mu)}(T) = \mathcal{A}_\mu((1+T)^p - 1).$$

- If  $\mu$  is a distribution on  $\mathbf{Z}_p$ , we denote  $\psi(\mu)$  the distribution on  $\mathbf{Z}_p$  defined by

$$\int_{\mathbf{Z}_p} \phi(x) \psi(\mu) = \int_{p\mathbf{Z}_p} \phi(p^{-1}x) \mu \quad \text{and therefore} \quad \mathcal{A}_{\psi(\mu)} = \psi(\mathcal{A}_\mu),$$

where  $\psi : \mathcal{K}^+ \rightarrow \mathcal{K}^+$  is defined by  $\psi(F)((1+T)^p - 1) = \frac{1}{p} \sum_{\zeta^p=1} F((1+T)\zeta - 1)$ .

The action of  $\mathbf{Z}_p^*$ ,  $\varphi$  and  $\psi$  satisfy the relations:

- (a)  $\psi \circ \phi = \text{id}$ .
- (b)  $\psi \circ \sigma_a = \sigma_a \circ \psi$  and  $\varphi \circ \sigma_a = \sigma_a \circ \varphi$  if  $a \in \mathbf{Z}_p^*$ .
- (c)  $\psi(\mathcal{A}_\mu) = 0$  if and only if  $\mu$  has support on  $\mathbf{Z}_p^*$ , and  $\mathcal{A}_{\text{Res}_{\mathbf{Z}_p^*}(\mu)} = (1 - \varphi\psi)\mathcal{A}_\mu$ .

7. Convolution of distribution: If  $\lambda$  and  $\mu$  are two distributions on  $\mathbf{Z}_p$ , we define the convolution  $\lambda * \mu$  by

$$\int_{\mathbf{Z}_p} \phi \cdot \lambda * \mu = \int_{\mathbf{Z}_p} \left( \int_{\mathbf{Z}_p} \phi(x+y) \mu(x) \right) \lambda(y).$$

Take  $\phi(x)$  the function  $x \mapsto z^x$ , where  $\nu_p(z-1) > 0$ , then we have  $\mathcal{A}_{\lambda * \mu}(z) = \mathcal{A}_\lambda(z) \mathcal{A}_\mu(z)$ . Hence we deduce  $\mathcal{A}_{\lambda * \mu} = \mathcal{A}_\lambda \cdot \mathcal{A}_\mu$ .

## 2. KUBOTA-LEOPOLDT $p$ -ADIC $L$ -FUNCTIONS

2.1. **Riemann zeta function.** Let  $\zeta(s) = \sum_{n=1}^{+\infty} n^{-s} = \prod_{p \text{ prime}} (1 - p^{-s})^{-1}$  <sup>the</sup> Riemann zeta function. Let  $\Gamma(s) = \int_{t=0}^{+\infty} e^{-ts} t^{s-1} dt$  <sup>the</sup> Gamma function, which is holomorphic on  $\text{Re}(s) > 0$  and satisfies the functional equation  $\Gamma(s+1) = s\Gamma(s)$ , ~~therefore~~ <sup>and thus</sup> it can be extended to a meromorphic function on  $\mathbb{C}$ .

Recall we have:

<sup>The</sup>

<sup>be</sup>

<sup>FONTAINE be</sup>

**Lemma 2.1.** *If  $\operatorname{Re}(s) > 1$ , then*

$$\zeta(s) = \frac{1}{\Gamma(s)} \int_0^{+\infty} \frac{1}{e^t - 1} t^s \frac{dt}{t}.$$

**Proposition 2.2.** *If  $f$  is a  $\mathcal{C}^\infty$  function on  $\mathbb{R}^+$  which decreases rapidly at infinite, then the function*

$$L(f, s) = \frac{1}{\Gamma(s)} \int_0^{+\infty} f(t) t^s \frac{dt}{t}$$

*defined on  $\operatorname{Re}(s) > 0$  admits a holomorphic extension to  $\mathbb{C}$  and if  $n \in \mathbb{N}$ , then  $L(f, -n) = (-1)^n f^{(n)}(0)$ .*

Apply the proposition to  $f_0(t) = \frac{t}{e^t - 1}$ . Let  $\sum_{n=0}^{+\infty} B_n \frac{t^n}{n!}$  be the Taylor expansion of  $f_0$  at 0, where  $B_n$  are Bernoulli number. We have, in particular

$$B_0 = 1, \quad B_1 = -\frac{1}{2}, \quad B_2 = \frac{1}{6}, \quad B_4 = \frac{-1}{30} \dots$$

Since  $f_0(t) - f_0(-t) = -t$ , we have  $B_{2k+1} = 0$  if  $k \geq 1$ .

**Theorem 2.3.**

- i) *The function  $\zeta$  has a meromorphic continuation to  $\mathbb{C}$ , which has a simple pole at  $s = 1$  with residue 1.*  
 ii) *If  $n \in \mathbb{Q}$ , then  $\zeta(-n) = (-1)^n \frac{B_{n+1}}{n+1}$ . In particular  $\zeta(-n) \in \mathbb{Q}$ .*

**2.2. Kummer congruence.** If  $a \in \mathbb{R}_+^*$ , by applying proposition 2.2 to the function  $f_a(t) = \frac{1}{e^t - 1} - \frac{a}{e^{at} - 1}$ , which is  $\mathcal{C}^\infty$  (removed the pole at  $t = 0$ ) on  $\mathbb{R}^+$  and decreases rapidly at infinity, we have

**Corollary 2.4.** *If  $a \in \mathbb{R}_+^*$ , the function  $(1 - a^{1-s})\zeta(s) = L(f_a, s)$  has an analytic continuation on  $\mathbb{C}$ , and if  $n \in \mathbb{N}$ , then  $(1 - a^{1+n})\zeta(-n) = (-1)^n f_a^{(n)}(0)$ . In particular, if  $a \in \mathbb{Q}$ , then  $(1 - a^{1+n})\zeta(-n) \in \mathbb{Q}$ .*

For a continuous distribution  $\mu$ , we associate it with the formal series:

$$\mathcal{L}_\mu(t) = \sum_{n=0}^{+\infty} \int_{\mathbf{Z}_p} e^{tx} \mu = \sum_{n=0}^{+\infty} \frac{t^n}{n!} \int_{\mathbf{Z}_p} x^n \mu,$$

which is called the Laplace transform of  $\mu$ .

We have  $\mathcal{L}_\mu(t) = \mathcal{A}_\mu(e^t - 1)$ .

**Proposition 2.5.** *If  $a \in \mathbf{Z}_p^*$ , there exists a measure  $\mu_a$  whose Laplace transform is  $f_a(t)$ . Moreover  $v_{\mathcal{O}}(\mu_a) \geq 0$  and if  $n \in \mathbb{N}$ , then  $\int_{\mathbf{Z}_p} x^n \mu_a = (-1)^n (1 - a^{1+n})\zeta(-n)$ .*

*Proof.* To show the existence of  $\mu_a$ , it suffices to prove the coefficients of series obtained by replace  $e^t$  by  $1 + T$  (Amice transform of  $\mu_a$ ) is bounded by proposition 1.9. Since  $(1 + T)^a - 1$  is of the form  $aT(1 + Tg(T))$  where  $g(T) = \sum_{n=2}^{+\infty} \frac{1}{a} \binom{a}{n} T^{n-2} \in \mathbf{Z}_p[[T]]$ , we have

$$\frac{1}{T} - \frac{a}{(1 + T)^a - 1} = \sum_{n=1}^{+\infty} (-T)^{n-1} g^n \in \mathbf{Z}_p[[T]].$$

Since the coefficients are in  $\mathbf{Z}_p$ , we have  $v_{\mathcal{O}}(\mu_a) \geq 0$ . Moreover, we have  $\int_{\mathbf{Z}_p} x^n \mu_a = \mathcal{L}_{\mu_a}^{(n)}(0) = f_a^{(n)}(0)$ .  $\square$

**Corollary 2.6.** (Kummer congruence) If  $a \in \mathbf{Z}_p^*$  and  $k \geq 1$ , if  $n_1$  and  $n_2$  are two integers  $\geq k$  such that  $n_1 \equiv n_2 \pmod{(p-1)p^{k-1}}$ , then

$$\nu_p\left((1 - a^{1+n_1})\zeta(-n_1) - (1 - a^{1+n_2})\zeta(-n_2)\right) \geq k.$$

*Proof.* Since we suppose  $n_1 \geq k$  and  $n_2 \geq k$ , we have  $\nu_p(x^{n_1}) \geq k$  and  $\nu_p(x^{n_2}) \geq k$  if  $x \in p\mathbf{Z}_p$ . On the other hand, since the order of  $(\mathbf{Z}/p^k\mathbf{Z})^*$  is  $(p-1)p^{k-1}$ , and ~~we suppose~~  $n_1 \equiv n_2 \pmod{(p-1)p^{k-1}}$ , we have  $x^{n_1} - x^{n_2} \in p^k\mathbf{Z}_p$  if  $x \in \mathbf{Z}_p^*$ . To sum up, we have  $\nu_p(x^{n_1} - x^{n_2}) \geq k$  if  $x \in \mathbf{Z}_p$  and hence  $\nu_{\mathcal{C}^0}(x^{n_1} - x^{n_2}) \geq k$ . Since  $\nu_{\mathcal{D}_0}(\mu_a) \geq 0$ , which implies

$$\nu_p((1 - a^{1+n_1})\zeta(-n_1) - (1 - a^{1+n_2})\zeta(-n_2)) = \nu_p\left(\int_{\mathbf{Z}_p} (x^{n_1} - x^{n_2})\mu_a(x)\right) \geq k.$$

□

**Proposition 2.7.** If  $a \in \mathbf{Z}_p^*$ , then

- i)  $\psi(\mu_a) = \mu_a$ .
- ii)  $\text{Res}_{\mathbf{Z}_p^*}(\mu_a) = (1 - \varphi)\mu_a$
- iii)  $\int_{\mathbf{Z}_p^*} x^n \mu_a = (1 - p^n) \int_{\mathbf{Z}_p} x^n \mu_a$  for all  $n \in \mathbf{N}$ .

*Proof.* Let  $F(T) = \psi(\frac{1}{T})$ . By definition, we have

$$\begin{aligned} F((1+T)^p - 1) &= \frac{1}{p} \sum_{\zeta^{p=1}} \frac{1}{(1+T)\zeta - 1} = \frac{-1}{p} \sum_{\zeta^{p=1}} \sum_{n=0}^{+\infty} ((1+T)\zeta)^n \\ &= - \sum_{n=0}^{+\infty} (1+T)^{pn} = \frac{1}{(1+T)^p - 1}. \end{aligned}$$

Hence we have  $\psi(\frac{1}{T}) = \frac{1}{T}$ . On the other hand, ~~we know that~~ the Amice transform of  $\mu_a$  is  $\frac{1}{T} - \frac{a}{(1+T)^a - 1} = \frac{1}{T} - a\sigma_a(\frac{1}{T})$ , and action of  $\psi$  commutes with  $\sigma_a$ . By  $\psi(\mathcal{A}_\mu) = \mathcal{A}_{\psi(\mu)}$  if  $\mu$  is a distribution, we deduce i).

ii) follows from i) since we have  $\text{Res}_{\mathbf{Z}_p^*}(\mu) = (1 - \varphi\psi)\mu$  if  $\mu$  is a distribution. iii) follows ii) and  $\int_{\mathbf{Z}_p} x^n \varphi(\mu) = \int_{\mathbf{Z}_p} (px)^n \mu$ . □

**Corollary 2.8.** Let  $a \in \mathbf{N} - \{1\}$  prime to  $p$ . Let  $k \geq 1$ . If  $n_1$  and  $n_2$  are two integers  $\geq k$  such that  $n_1 \equiv n_2 \pmod{(p-1)p^{k-1}}$ , then

$$\nu_p((1 - a^{1+n_1})(1 - p^{n_1})\zeta(-n_1) - (1 - a^{1+n_2})(1 - p^{n_2})\zeta(-n_2)) \geq k.$$

By corollary 2.8, ~~we know that~~ the function  $n \mapsto (1 - p^n)\zeta(-n)$  is continuous under  $p$ -adic topology. To have a uniform formula, we put  $q = 4$  if  $p = 2$  and  $q = p$  if  $p \neq 2$ . We denote  $\phi$  the Euler function, thus we have  $\phi(q) = 2$  if  $q = 4$  and  $\phi(q) = p - 1$  if  $p \neq 2$ .

**Theorem 2.9.** If  $i \in \mathbf{Z}/\phi(q)\mathbf{Z}$ , there exist an unique function  $\zeta_{p,i}$  continuous on  $\mathbf{Z}_p$  (resp.  $\mathbf{Z}_p - \{1\}$ ) if  $i \neq 1$  (resp.  $i = 1$ ) such that the function  $(s-1)\zeta_{p,i}$  is analytic on  $\mathbf{Z}_p$  (resp.  $i + 2\mathbf{Z}_p$  if  $p = 2$ ) and one has  $\zeta_{p,i}(-n) = (1 - p^n)\zeta(-n)$  if  $n \in \mathbf{N}$  verified  $-n \equiv i \pmod{p-1}$ .

**Remark 2.10.**  $\zeta_{p,i}$  is called the  $i$ -th branch of Kubota-Leopoldt zeta function. If  $i$  is even, then  $\zeta_{p,i}$  is identically zero since  $\zeta(-n) = 0$  if  $n \geq 2$  is even.

**2.3.  $p$ -adic Mellin transform and Leopoldt's  $\Gamma$ -transform.** We denote  $\Delta$  the group of roots of unity of  $\mathbf{Q}_p^*$ . Therefore  $\Delta$  is a cyclic group of order  $\phi(q)$  and  $\mathbf{Z}_p^*$  is disjoint union of  $\varepsilon + q\mathbf{Z}_p$  with  $\varepsilon \in \Delta$ . We denote  $\omega : \mathbf{Z}_p \rightarrow \Delta \cup \{0\}$  the function defined by  $\omega(x) = 0$  if  $x \in p\mathbf{Z}_p$ , and  $x - \omega(x) \in q\mathbf{Z}_p$ , if  $x \in \mathbf{Z}_p^*$ . If  $x \in \mathbf{Z}_p^*$ , we define  $\langle x \rangle \in 1 + q\mathbf{Z}_p$  by  $\langle x \rangle = x\omega(x)^{-1}$ .

**Proposition 2.11.** *If  $i \in \mathbf{Z}/\phi(q)\mathbf{Z}$ , the function  $x \mapsto \omega(x)^i \langle x \rangle^s$  is a locally analytic function on  $\mathbf{Z}_p$ . Moreover, we have*

- i)  $\omega(x)^i \langle x \rangle^n = x^n$  if  $n \equiv i \pmod{\phi(q)}$  and If  $x \in \mathbf{Z}_p^*$
- ii)  $\omega(x)^i \langle x \rangle^s = \lim_{\substack{n \rightarrow s \\ n \equiv i \pmod{\phi(q)}}} x^n$  for  $x \in \mathbf{Z}_p$ .

*Proof.* Note that we have  $\omega(x)^i \langle x \rangle^s = 0$  on  $p\mathbf{Z}_p$  and

$$\omega(x)^i \langle x \rangle^s = \varepsilon^i \left( \frac{x}{\varepsilon} \right)^s = \sum_{n=0}^{+\infty} \binom{s}{n} \varepsilon^{i-n} (x - \varepsilon)^n,$$

if  $x \in \varepsilon + q\mathbf{Z}_p$  and  $\varepsilon \in \Delta$ , thus the function is locally analytic.

Since the order of  $\Delta$  is  $\phi(q)$ , we have  $\omega(x)^n = \omega(x)^i$  if  $n \equiv i \pmod{\phi(q)}$ , i) and ii) follow.  $\square$

If  $i \in \mathbf{Z}/\phi(q)\mathbf{Z}$ , we defined the  $i$ -th branch of the Mellin transform of a continuous distribution  $\mu$  by the formula

$$\text{Mel}_{i,\mu}(s) = \int_{\mathbf{Z}_p} \omega(x)^i \langle x \rangle^s \mu(x) = \int_{\mathbf{Z}_p^*} \omega(x)^i \langle x \rangle^s \mu(x)$$

the second equality is because  $\omega(x) = 0$  if  $x \in p\mathbf{Z}_p$ . On the other hand, we have  $\text{Mel}_{i,\mu}(n) = \int_{\mathbf{Z}_p^*} x^n \mu$  if  $n \equiv i \pmod{\phi(q)}$ .

Let  $u$  be a topological generator of multiplicative group of  $1 + q\mathbf{Z}_p$ , and let  $\theta : 1 + q\mathbf{Z}_p \rightarrow \mathbf{Z}_p^*$  be the homomorphism which sends  $x$  to  $\frac{\log x}{\log u}$ . This homomorphism is analytic and its inverse is also. If  $f$  is a locally analytic function (resp. continuous) function on  $1 + q\mathbf{Z}_p$ , the function  $\theta^* f$  defined by  $\theta^* f(x) = f(\theta(x))$  is locally analytic (resp. continuous) on  $\mathbf{Z}_p$ .

If  $\mu$  is a distribution support on  $1 + q\mathbf{Z}_p$ , we define a distribution  $\theta_* \mu$  on  $\mathbf{Z}_p$  by the formula

$$\int_{\mathbf{Z}_p} \phi(\theta_* \mu) = \int_{1+q\mathbf{Z}_p} (\theta^* \phi) \mu.$$

In particular,  $\theta_*$  sends measure to measure.

**Lemma 2.12.** *If  $X$  is a open compact subset of  $\mathbf{Z}_p$ , If  $\alpha \in \mathbf{Z}_p^*$ , and if  $\mu$  is a continuous distribution on  $\mathbf{Z}_p$ , then*

$$\text{Res}_X(\sigma_\alpha(\mu)) = \sigma_\alpha(\text{Res}_{\alpha^{-1}X}(\mu))$$

*Proof.* Since we have  $1_X(\alpha x) = 1_{\alpha^{-1}X}(x)$  if  $X \subset \mathbf{Z}_p$ , we deduce the formula

$$\begin{aligned} \int_{\mathbf{Z}_p} \phi(x) \text{Res}_X(\sigma_\alpha(\mu)) &= \int_{\mathbf{Z}_p} 1_X(x) \phi(x) \sigma_\alpha \mu = \int_{\mathbf{Z}_p} 1_X(\alpha x) \phi(\alpha x) \mu(x) \\ &= \int_{\mathbf{Z}_p} \phi(\alpha x) (1_{\alpha^{-1}X}(x) \mu(x)) = \int_{\mathbf{Z}_p} \phi(\alpha x) \text{Res}_{\alpha^{-1}X}(\mu) \\ &= \int_{\mathbf{Z}_p} \phi(x) \sigma_\alpha(\text{Res}_{\alpha^{-1}X}(\mu)), \end{aligned}$$

USE  
MORE  
LINE'S

which proves the lemma.  $\square$



**Definition 2.13.** If  $\mu$  is a distribution on  $\mathbf{Z}_p^*$  and if  $i \in \mathbf{Z}/\phi(q)\mathbf{Z}$ , we define  $\Gamma_\mu^{(i)}$  the  $i$ -th branch of the  $\Gamma$ -transform of  $\mu$  by

$$\Gamma_\mu^{(i)} = \theta_* \text{Res}_{1+q\mathbf{Z}_p} \left( \sum_{\varepsilon \in \Delta} \varepsilon^{-i} \sigma_\varepsilon(\mu) \right) = \theta_* \left( \sum_{\varepsilon \in \Delta} \varepsilon^{-i} \sigma_\varepsilon(\text{Res}_{\varepsilon^{-1}+q\mathbf{Z}_p}(\mu)) \right),$$

where

the second equality follows from the above lemma. Moreover, it is clear that if  $\mu$  is a measure on  $\mathbf{Z}_p^*$ , then  $\Gamma_\mu^{(i)}$  is a measure on  $\mathbf{Z}_p$ , and we have  $v_{\mathcal{D}_0}(\Gamma_\mu^{(i)}) \geq v_{\mathcal{D}_0}(\mu)$ .

**Proposition 2.14.** If  $\mu$  is a continuous distribution on  $\mathbf{Z}_p^*$  and  $i \in \mathbf{Z}/\phi(q)\mathbf{Z}$ , then

$$\text{Mel}_{i,\mu}(s) = \int_{\mathbf{Z}_p^*} \omega(x)^i \langle x \rangle^s \mu(x) = \int_{\mathbf{Z}_p} u^{sy} \Gamma_\mu^{(i)}(y) = \mathcal{A}_{\Gamma_\mu^{(i)}}(u^s - 1)$$

*Proof.* The first equality is by the definition of Mellin transform and the third equality is by the definition of Amice transform. If  $y = \theta(x) = \frac{\log x}{\log u}$ , we have  $u^{sy} = \exp(s \log x) = \langle x \rangle^s$  and

$$\int_{\mathbf{Z}_p} u^{sy} \Gamma_\mu^{(i)}(y) = \int_{1+q\mathbf{Z}_p} \langle x \rangle^s \sum_{\varepsilon \in \Delta} \varepsilon^{-i} \sigma_\varepsilon(\text{Res}_{\varepsilon^{-1}+q\mathbf{Z}_p}(\mu)).$$

Using the fact that  $\omega(x) = \varepsilon^{-1}$  if  $x \in \varepsilon^{-1} + q\mathbf{Z}_p$  and  $\langle \varepsilon x \rangle = \langle x \rangle$ , we obtain

$$\int_{\mathbf{Z}_p} u^{sy} \Gamma_\mu^{(i)}(y) = \sum_{\varepsilon \in \Delta} \int_{\varepsilon^{-1}+q\mathbf{Z}_p} \omega(x)^i \langle x \rangle^s \mu(x),$$

and the proposition follows from that  $\mathbf{Z}_p^*$  is the disjoint union of  $\varepsilon + q\mathbf{Z}_p$  for  $\varepsilon \in \Delta$ .  $\square$

**Corollary 2.15.**

- i) If  $\mu$  is a continuous distribution and  $i \in \mathbf{Z}/\phi(q)\mathbf{Z}$ , the function  $\text{Mel}_{i,\mu}(s)$  is a analytic function of  $s$  and even  $u^s - 1$ .
- ii) If  $\mu$  is a measure verified  $v_{\mathcal{D}_0}(\mu) \geq 0$ , and if  $i \in \mathbf{Z}/\phi(q)\mathbf{Z}$ , then there exists  $g_{i,\mu} \in \mathcal{O}_L[[T]]$  such that  $\text{Mel}_{i,\mu}(s) = g_{i,\mu}(u^s - 1)$ .

**2.4. Construction of the Kubota-Leopoldt zeta function.** If  $i \in \mathbf{Z}/\phi(q)\mathbf{Z}$  and  $a \in \mathbf{Z}_p^*$  such that  $\langle a \rangle \neq 1$ , we define the function  $g_{a,i}$  on  $\mathbf{Z}_p$  by the formula

$$g_{a,i}(s) = \frac{1}{1 - \omega(a)^{1-i} \langle a \rangle^{1-s}} \text{Mel}_{-i,\mu_a}(-s) = \frac{1}{1 - \omega(a)^{1-i} \langle a \rangle^{1-s}} \int_{\mathbf{Z}_p^*} \omega(a)^{-i} \langle a \rangle^{-s} \mu_a.$$

By corollary 2.15,  $\text{Mel}_{-i,\mu_a}(-s)$  is an analytic function of  $s$ . On the other hand, if  $\omega(a)^{1-i} \neq 1$ , the function  $s \mapsto 1 - \omega(a)^{1-i} \langle a \rangle^{1-s}$  is a nonzero analytic function on  $\mathbf{Z}_p$  since  $\langle a \rangle^s \in 1 + q\mathbf{Z}_p$  and  $\omega(a)^{1-i} \in \Delta - \{1\}$ , therefore  $\omega(a)^{1-i} \notin 1 + q\mathbf{Z}_p$  and if  $\omega(a)^{1-i} = 1$ , the function  $1 - \langle a \rangle^{1-s}$  vanishes only at  $s = 1$ . We deduce that  $g_{a,i}$  is a function continuous on  $\mathbf{Z}_p - \{1\}$  and even on  $\mathbf{Z}_p$  if  $\omega(a)^{1-i} \neq 1$ .

Moreover, if  $-n \equiv i \pmod{\phi(q)}$ , we have  $\omega(a)^{1-i} = \omega(a)^{1+n}$  and  $\omega(x)^{-i} = \omega(x)^n$  if  $x \in \mathbf{Z}_p^*$ . Therefore

$$g_{a,i}(-n) = \frac{1}{1 - \omega(a)^{1+n} \langle a \rangle^{1+n}} \int_{\mathbf{Z}_p^*} \omega(x)^n \langle a \rangle^n \mu_a(x) = \frac{1}{1 - a^{1+n}} \int_{\mathbf{Z}_p^*} x^n \mu_a(x) = (-1)^n (1-p^n) \zeta(-n)$$

does not depend on the choice of  $a$ . If  $a$  and  $a'$  two elements of  $\mathbf{Z}_p^*$ , the function  $g_{a,i} - g_{a',i}$  is a quotient of analytic functions on  $\mathbf{Z}_p$  vanishing at infinite many points, which implies it identical zero and the function  $g_{a,i}$  is independent of choice of  $a$ . Thus we set  $\zeta_{p,i} = g_{a,i}$  for any  $a$  satisfies  $\langle a \rangle \neq 1$  and  $\omega(a)^{1-i} \neq 1$  if  $i \neq 1$  to construct Kubota-Leopoldt zeta function.

Let  $F_n = \mathbf{Q}_p(\varepsilon_{p^n})$  and  $F_\infty = \cup F_n$ . The norm  $N_{F_{n+1}/F_n}$  ~~induced~~ <sup>induces</sup> a homomorphism from  $\mu_{p^{n+1}}$  to  $\mu_{p^n}$ , where  $\mu_{p^n}$  be the set of  $p^n$ -th roots of unity in  $F_n$ . We denote the projective limit of  $\mu_{p^n}$  with respect to  $N_{F_{n+1}/F_n}$  by  $\mu_{p^\infty}$  (Tate module), which is a compact  $\mathbf{Z}_p$ -module.

The following theorem is due to Mazur and Wiles: (REF) ?

**Theorem 2.16.** *If  $i \in (\mathbf{Z}/(p-1)\mathbf{Z})^*$  is odd and if  $s \in \mathbf{Z}_p$ , then the following two conditions are equivalent:*

- i)  $\zeta_{p,i}(s) = 0$ ;
- ii) *There exists an element  $u \in \mu_{p^\infty}$  which is not killed by a power of  $p$  such that  $\sigma \in \text{Gal}(F_\infty/\mathbf{Q}_p)$  acts by the formula*

$$\sigma(u) = \omega(\chi_{\text{cycl}}(\sigma))^i \langle \chi_{\text{cycl}}(\sigma) \rangle^s \cdot u.$$

**2.5. The residue at  $s = 1$  and the  $p$ -adic zeta function.** The formal power series  $\frac{\log(1+T)}{T}$  converges on open unit disk, thus it is an Amice transform of an unique distribution  $\mu_{KL}$ . The Laplace transform of  $\mu_{KL}$  is  $\frac{t}{e^t-1} = f_0(t)$  and

$$\int_{\mathbf{Z}_p} x^n \mu_{KL} = (-1)^{n-1} n \zeta(1-n)$$

**Lemma 2.17.**  $\int_{a+p^n \mathbf{Z}_p} \mu_{KL} = \frac{1}{p^n}$

*Proof.* Since  $\int_{a+p^n \mathbf{Z}_p} \mu_{KL} = \frac{1}{p^n} \sum_{\varepsilon \in p^n=1} \varepsilon^{-a} \mathcal{A}_{\mu_{KL}}(\varepsilon-1)$  and since  $\log \varepsilon = 0$  if  $\varepsilon$  is a roots of unity of order power of  $p$ , all terms of the sum is zero except for the term corresponding to  $\varepsilon = 1$ , we get the result.  $\square$

**Proposition 2.18.** *We have*

- i)  $\psi(\mu_{KL}) = p^{-1} \mu_{KL}$
- ii)  $\text{Res}_{\mathbf{Z}_p^*}(\mu_{KL}) = (1 - p^{-1} \varphi) \mu_{KL}$
- iii)  $\int_{\mathbf{Z}_p^*} \mu_{KL} = (-1)^{n-1} n (1 - p^{n-1}) \zeta(1-n)$  if  $n \in \mathbf{N}$ .

*Proof.* i) follows from the formula  $\psi(\frac{1}{T}) = \frac{1}{T}$  (c.f. proposition 2.7) and  $\varphi(\log(1+T)) = p \log(1+T)$  and  $\psi(\varphi(a)b) = a\psi(b)$ . The rest can be deduced from proposition 2.7.  $\square$

**Theorem 2.19.** *The  $p$ -adic zeta function  $\zeta_{p,1}$  has a simple pole at  $s = 1$  with residue  $1 - \frac{1}{p}$ .*

*Proof.* According to the above, we can define the function  $\zeta_{p,i}$ , if  $i \in \mathbf{Z}/\phi(q)\mathbf{Z}$  by the formula

$$\zeta_{p,i}(s) = \frac{(-1)^{i-1}}{s-1} \text{Mel}_{1-i, \mu_{KL}}(1-s) = \frac{(-1)^{i-1}}{s-1} \int_{\mathbf{Z}_p^*} \omega^{1-i} \langle x \rangle^{1-s} \mu_{KL}(x).$$

Indeed, the function is analytic on  $\mathbf{Z}_p - \{1\}$  by above formula, and take the same value  $\zeta_{p,i}(-n) = (1 - p^n) \zeta(-n)$  if  $n \in \mathbf{N}$  satisfies  $-n \equiv i \pmod{p-1}$ . Moreover,

$$\begin{aligned} \lim_{s \rightarrow 1} (s-1) \zeta_{p,i}(s) &= \int_{\mathbf{Z}_p^*} \omega(x)^{1-i} \mu_{KL}(x) \\ &= \sum_{\alpha \in \Delta} \omega(\alpha)^{1-i} \int_{\alpha+p \mathbf{Z}_p} \mu_{KL}(x) = \begin{cases} 1 - \frac{1}{p} & \text{if } i = 1 \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

$\square$

**2.6. Dirichlet  $L$ -function.** For  $\chi$  a Dirichlet character of conductor  $D > 1$  and if  $n \in \mathbf{Z}$ , we define the Gauss sum  $G(\chi)$  by ~~the formula~~

$$G(\chi) = \sum_{a \bmod D} \chi(a) e^{2\pi i \frac{a}{D}}.$$

$\leftarrow n$  NOT used

Let

$$L(\chi, s) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s} = \prod_{p:\text{prime}} (1 - \chi(p)p^{-s})^{-1}, \quad \text{For } \operatorname{Re}(s) \geq 1,$$

$\hookrightarrow$  the Dirichlet  $L$ -function attached to  $\chi$ . By the formula

$$\chi(n) = \frac{1}{G(\chi^{-1})} \sum_{b \bmod D} \chi^{-1}(b) e^{2\pi i \frac{nb}{D}},$$

we obtain

$$L(\chi, s) = \frac{1}{G(\chi^{-1})} \sum_{b \bmod D} \chi^{-1}(b) \frac{e^{2\pi i \frac{nb}{D}}}{n^s}.$$

Using the formula  $\int_0^{+\infty} e^{-nt} t^s \frac{dt}{t} = \frac{\Gamma(s)}{n^s}$  and put  $\varepsilon_D = e^{\frac{2\pi i}{D}}$ , we obtain

$$\begin{aligned} L(\chi, s) &= \frac{1}{G(\chi^{-1})} \frac{1}{\Gamma(s)} \sum_{b \bmod D} \chi^{-1}(b) \int_0^{+\infty} \sum_{n=1}^{+\infty} \varepsilon_D^{nb} e^{-nt} t^s \frac{dt}{t} \\ &= \frac{1}{G(\chi^{-1})} \frac{1}{\Gamma(s)} \int_0^{+\infty} \sum_{b \bmod D} \frac{\chi^{-1}(b)}{\varepsilon_D^{-b} e^t - 1} t^s \frac{dt}{t}. \end{aligned}$$

In particular, proposition 2.2 implies that  $L(\chi, s)$  can be extended to a holomorphic function on  $\mathbb{C}$ . Moreover,  $L(\chi, -n) = \left(\frac{d}{dt}\right)^n \mathcal{L}_\chi(t) |_{t=0}$  where

$$\mathcal{L}_\chi(t) = \frac{-1}{G(\chi^{-1})} \sum_{b \bmod D} \frac{\chi^{-1}(b)}{\varepsilon_D^b e^t - 1}.$$

**2.7.  $p$ -adic  $L$ -function attaches to Dirichlet character.** Let  $\chi$  be a Dirichlet character of conductor  $D > 1$  prime to  $p$ . If  $\chi^{-1}(b) \neq 0$ , then  $\varepsilon_D^b$  is a roots of unity of order prime to  $p$  and distinct from 1, this implies  $\nu_p(\varepsilon_D^b - 1) = 0$ . We deduce that the power series

$$F_\chi(T) = \frac{-1}{G(\chi^{-1})} \sum_{b \bmod D} \frac{\chi^{-1}(b)}{(1+T)\varepsilon_D^b - 1} = \frac{1}{G(\chi)^{-1}} \sum_{b \bmod D} \chi^{-1}(b) \sum_{n=0}^{+\infty} \frac{\varepsilon_D^{nb}}{(\varepsilon_D^b - 1)^{n+1}} T^n$$

is of bounded coefficients (since  $\nu_p(G(\chi)G(\chi^{-1})) = \nu_p(D) = 0$ ) and hence an Amice transform of a measure  $\mu_\chi$  on  $\mathbf{Z}_p$  whose Laplace transform  $F_\chi(e^t - 1) = \mathcal{L}_\chi(t)$ . We have  $\int_{\mathbf{Z}_p} x^n \mu_\chi = \mathcal{L}_\chi^{(n)}(0) = L(\chi, -n)$  and  $v_{\mathcal{D}_0}(\mu_\chi) \geq 0$ .

**Definition 2.20.** We define the  $p$ -adic  $L$ -function associated to  $\chi$  by the Mellin transform of  $\mu_\chi$ , that is, the function  $\beta \mapsto L_p(\chi \otimes \beta)$  defined by

$$L_p(\chi \otimes \beta) = \int_{\mathbf{Z}_p^*} \beta(x) \mu_\chi(x).$$

where  $\beta$  is a locally analytic character on  $\mathbf{Z}_p^*$ . On the other hand, if  $i \in \mathbf{Z}/\phi(q)\mathbf{Z}$ , we put

$$L_{p,i}(\chi, s) = L_p(\chi \otimes (\omega^{-i}(x) \langle x \rangle^{-s})) = \int_{\mathbf{Z}_p^*} \omega^{-i} \langle x \rangle^{-s} \mu_\chi(x).$$

**Proposition 2.21.** *If  $i \in \mathbf{Z}/\phi(q)\mathbf{Z}$ , the function  $L_{p,i}(\chi, s)$  is an analytic function on  $\mathbf{Z}_p$  and we have  $L_{p,i}(\chi, -n) = (1 - \chi(p)p^n)L(\chi, -n)$  if  $n \in \mathbf{N}$  satisfying  $-n \equiv i \pmod{\phi(q)}$ .*

*Proof.* The fact that  $L_{p,i}(\chi, s)$  is an analytic function on  $\mathbf{Z}_p$  follows from corollary 2.15. On the other hand, we have

$$\sum_{\eta^p=1} \frac{1}{(1+T)\varepsilon_D\eta - 1} = p \frac{1}{(1+T)^p \varepsilon_D^{pb} - 1} ,$$

*And thus* thus we deduce the Amice transform of  $\mu_\chi$  restriction to  $\mathbf{Z}_p^*$  is

$$\frac{-1}{G(\chi)} \sum_{b \bmod D} \frac{\chi^{-1}(b)}{(1+T)\varepsilon_D^b - 1} - \frac{\chi^{-1}(b)}{(1+T)^p \varepsilon_D^{pb} - 1},$$

which can be written as  $\mathcal{A}_{\mu_\chi}(T) - \chi(p)\mathcal{A}_{\mu_\chi}((1+T)^p - 1)$ . Hence we deduce the formula

$$\mathcal{L}_{\text{Res}_{\mathbf{Z}_p^*}(\mu_\chi)}(t) = \mathcal{L}_{\mu_\chi}(t) - \chi(p)\mathcal{L}_{\mu_\chi}(pt) \quad \text{and} \quad \int_{\mathbf{Z}_p^*} x^n \mu_\chi = (1 - \chi(p))L(\chi, -n),$$

and the proposition follows.  $\square$

**2.8. Behavior at  $s = 1$  of Dirichlet  $L$ -function.** By section 2.6, we have

$$\begin{aligned} L(\chi, 1) &= \frac{1}{G(\chi^{-1})} \sum_{b \bmod D} \chi^{-1}(b) \sum_{n=0}^{+\infty} \frac{\varepsilon_D^{nb}}{n} \\ &= \frac{-1}{G(\chi^{-1})} \sum_{b \bmod D} \chi^{-1}(b) \log(1 - \varepsilon_D^b). \end{aligned}$$

We will establish the  $p$ -adic analogy of this formula by calculating  $\int_{\mathbf{Z}_p^*} x^{-1} \mu_\chi$ . To do this, we will calculate the Amice transform of  $x^{-1} \mu_\chi$  and then restrict it to  $\mathbf{Z}_p^*$ .

**Proposition 2.22.** *The Amice transform of  $x^{-1} \mu_\chi$  is*

$$\mathcal{A}_{x^{-1}\mu_\chi}(T) = \frac{-1}{G(\chi)^{-1}} \sum_{b \bmod D} \chi^{-1}(b) \log((1+T)\varepsilon_D^b - 1).$$

*Proof.* If  $\mu$  is a distribution, the relation of Amice transform of  $\mu$  and  $x^{-1}\mu$  is give by

$$(1+T) \frac{d}{dT} \mathcal{A}_{x^{-1}\mu}(T) = \mathcal{A}_\mu(T).$$

Apply the operator  $(1+T) \frac{d}{dT}$  on the right hand side of the equality in the proposition we obtain

$$\frac{-1}{G(\chi^{-1})} \sum_{b \bmod D} \chi^{-1}(b) \frac{(1+T)\varepsilon_D^b}{(1+T)\varepsilon_D^b - 1} = \frac{-1}{G(\chi^{-1})} \sum_{b \bmod D} \chi^{-1}(b) \left( \frac{1}{(1+T)\varepsilon_D^b - 1} + 1 \right)$$

which is equal to  $\mathcal{A}_{\mu_\chi}$  since  $\sum_{b \bmod D} \chi^{-1}(b) = 0$ . We deduce that the two elements have the same image by  $(1+T) \frac{d}{dT}$  and therefore differs by a locally constant function. To conclude, we must verify that the right hand side is given by a series which converges on the open unit disk. Since ~~we have~~

$$\log((1+T)\varepsilon_D^b - 1) = \log(\varepsilon_D^b - 1) + \log\left(1 + \frac{\varepsilon_D^b T}{\varepsilon_D^b - 1}\right) = \log(\varepsilon_D^b - 1) + \sum_{n=1}^{+\infty} \frac{(-1)^{n-1}}{n} \left(\frac{\varepsilon_D^b T}{\varepsilon_D^b - 1}\right)^n ,$$

and we suppose  $(D, p) = 1$ , we have  $\nu_p(\varepsilon_D^b - 1) = 0$ , and hence the series converges on open unit disk.  $\square$

**Lemma 2.23.** *The Amice transform of the restriction of  $x^{-1}\mu_\chi$  to  $\mathbf{Z}_p^*$  is defined by*

$$\begin{aligned} \mathcal{A}_{\text{Res}_{\mathbf{Z}_p^*}(x^{-1}\mu_\chi)}(T) &= \frac{-1}{G(\chi^{-1})} \sum_{b \bmod D} \chi^{-1}(b) \left( \log((1+T)\varepsilon_D^b - 1) - \frac{1}{p} \log((1+T)^p \varepsilon_D^{pb} - 1) \right) \\ &= \mathcal{A}_{x^{-1}\mu_\chi}(T) - \frac{\chi(p)}{p} \mathcal{A}_{x^{-1}\mu_\chi}((1+T)^p - 1). \end{aligned}$$

*Proof.* Use the formula for Amice transform of  $\text{Res}_{\mathbf{Z}_p^*}$ .  $\square$

By taking  $T = 0$  ~~to~~ <sup>in</sup> the above formula, we obtain

$$L_{p,1}(\chi, 1) = L_p(\chi \otimes x^{-1}) = \int_{\mathbf{Z}_p^*} x^{-1}\mu_\chi = \frac{-1}{G(\chi^{-1})} \left(1 - \frac{\chi(p)}{p}\right) \sum_{b \bmod D} \chi^{-1}(b) \log(\varepsilon_D^b - 1).$$

which differs <sup>from the</sup> complex  $L$ -function ~~case~~ by an Euler factor.

**2.9. Twist by a character of conductor power of  $p$ .** Let  $\chi$  be a Dirichlet character <sup>of</sup> conductor  $D$  prime to  $p$  and  $\beta$  be a Dirichlet character of conductor  $p^k$ . We denote  $\chi \otimes \beta$  to be the Dirichlet character of conductor  $Dp^k$  defined by  $(\chi \otimes \beta)(a) = \chi(a)\beta(a)$ , where  $\chi$  and  $\beta$  are viewed as characters mod  $Dp^k$  via the projection ~~from~~ <sup>from</sup>  $(\mathbf{Z}/Dp^k\mathbf{Z})^*$  to  $(\mathbf{Z}/D\mathbf{Z})^*$  and  $(\mathbf{Z}/p^k\mathbf{Z})^*$ .

**Lemma 2.24.** *Let  $k \geq 1$ ,  $\beta$  a Dirichlet character of conductor  $p^k$  and  $\mu$  a continuous distribution on  $\mathbf{Z}_p$ . Then we have*

$$\text{Then } \int_{\mathbf{Z}_p} \beta(x)(1+T)^x \mu(x) = \frac{1}{G(\beta)^{-1}} \sum_{c \bmod p^k} \beta^{-1}(c) \mathcal{A}_\mu((1+T)\varepsilon_{p^k}^c - 1).$$

*Proof.* We have

$$\begin{aligned} \int_{\mathbf{Z}_p} \beta(x)(1+T)^x \mu(x) &= \sum_{a \bmod p^k} \beta(a) \int_{a+p^k\mathbf{Z}_p} (1+T)^x \mu \\ &= \sum_{a \bmod p^k} \beta(a) \left( \frac{1}{p^k} \sum_{\eta^{p^k}=1} \eta^{-a} \mathcal{A}_\mu((1+T)\eta - 1) \right) \\ &= \sum_{\eta^{p^k}=1} \mathcal{A}_\mu((1+T)\eta - 1) \left( \frac{1}{p^k} \sum_{a \bmod p^k} \beta(a) \eta^{-a} \right), \end{aligned}$$

~~The~~ <sup>and the</sup> lemma follows from the identity

$$\frac{1}{p^k} \beta^{-1}(-c) G(\beta) = \frac{\beta^{-1}(c)}{G(\beta^{-1})}.$$

$\square$

**Proposition 2.25.** *If  $\mu$  is a measure on  $\mathbf{Z}_p$  with Amice transform of the form*

$$\mathcal{A}_\mu(T) = \frac{1}{G(\chi^{-1})} \sum_{b \bmod D} \chi^{-1}(b) F((1+T)\varepsilon_D^b - 1)$$

FONT

and if  $\beta$  is a Dirichlet character of conductor  $p^k$  with  $k \geq 1$ , then

$$\int_{\mathbf{Z}_p} \beta(x)(1+T)^x \mu(x) = \frac{1}{G((\chi \otimes \beta)^{-1})} \sum_{a \bmod Dp^k} (\chi \otimes \beta)^{-1}(a) F((1+T)\varepsilon_D^b \varepsilon_{p^k}^c - 1).$$

*Proof.* By the preceding lemma we have

$$\int_{\mathbf{Z}_p} \beta(x)(1+T)^x \mu(x) = \frac{-1}{G(\chi^{-1})G(\beta^{-1})} \sum_{b \bmod D} \sum_{c \bmod p^k} \chi^{-1}(b)\beta^{-1}(c) F((1+T)\varepsilon_D^b \varepsilon_{p^k}^c - 1).$$

Using the fact that every element of  $\mathbf{Z}/Dp^n\mathbf{Z}$  can be written uniquely as  $Dc + p^k b$ , where  $b \in \mathbf{Z}/D\mathbf{Z}$  and  $c \in \mathbf{Z}/p^k\mathbf{Z}$ , we have the following formula.

$$\begin{aligned} \varepsilon_{Dp^n}^a &= \varepsilon_D^b \varepsilon_{p^k}^c \\ (\chi \otimes \beta)^{-1}(a) &= \chi^{-1}(p^k)\beta^{-1}(D)\chi^{-1}(b)\beta^{-1}(c) \\ G((\chi \otimes \beta)^{-1}) &= \sum_{a \bmod Dp^k} (\chi \otimes \beta)^{-1}(a) \varepsilon_{Dp^k}^a \\ &= \chi^{-1}(p^k)\beta^{-1}(D) \left( \sum_{b \bmod D} \chi^{-1}(b)\varepsilon_D^b \right) \left( \sum_{c \bmod p^k} \beta^{-1}(c)\varepsilon_{p^k}^c \right) \\ &= \chi^{-1}(p^k)\beta^{-1}(D) G(\chi^{-1})G(\beta^{-1}) \end{aligned}$$

and the conclusion follows.  $\square$

**Proposition 2.26.** *If  $\beta$  is a non-trivial Dirichlet character of conductor  $p^n$  and if  $n \in \mathbf{N}$ , then  $L_p(\chi \otimes (x^n \beta)) = L(\chi \otimes \beta, -n)$ .*

*Proof.* By the preceding proposition and the formula for the Amice transform of  $\mu_\chi$ , we have the Amice transform of  $\beta\mu_\chi$  is

$$\frac{-1}{G((\chi \otimes \beta)^{-1})} \sum_{x \bmod Dp^n} \frac{(\chi \otimes \beta)^{-1}(x)}{(1+T)\varepsilon_{Dp^n}^x - 1}$$

and thus its Laplace transform is the function  $\mathcal{L}_{\chi \otimes \beta}(t)$ .  $\square$

### 3. $(\varphi, \Gamma)$ -MODULES AND $p$ -ADIC REPRESENTATIONS

Throughout this article,  $k$  will denote a finite field of characteristic  $p > 0$ , so if  $W(k)$  denotes the ring of Witt vectors over  $k$ , then  $F = W(k)[\frac{1}{p}]$  is a finite unramified extension of  $\mathbf{Q}_p$ . Let  $\overline{\mathbf{Q}}_p$  be the algebraic closure of  $\mathbf{Q}_p$ , let  $K$  be a totally ramified extension of  $F$ , and let  $G_K = \text{Gal}(\overline{\mathbf{Q}}_p/K)$  be the absolute Galois group of  $K$ . Let  $\mu_{p^n}$  be the group of  $p^n$ -th roots of unity; for every  $n$ , we will choose a generator  $\varepsilon^{(n)}$  of  $\mu_{p^n}$  with the additional requirement that  $(\varepsilon^{(n)})^p = \varepsilon^{(n-1)}$ . This makes  $\varprojlim \varepsilon^{(n)}$  into a generator  $\varprojlim \mu_{p^n} \simeq \mathbf{Z}_p(1)$ . We set  $K_n = K(\mu_{p^n})$  and  $K_\infty = \bigcup_{n \geq 0} K_n$ . Recall

that the cyclotomic character  $\chi : G_K \rightarrow \mathbf{Z}_p^*$  is defined by the relation:  $g(\varepsilon^{(n)}) = (\varepsilon^{(n)})^{\chi(g)}$  for all  $g \in G_K$ . The kernel of the cyclotomic character is  $H_K = \text{Gal}(\overline{\mathbf{Q}}_p/K_\infty)$ , and  $\chi$  therefore identifies  $\Gamma_K = G_K/H_K$  with an open subgroup of  $\mathbf{Z}_p^*$ .

**3.1. The field  $\tilde{\mathbf{E}}$  and its subrings.** Let  $\mathbf{C}_p$  be the completion of  $\overline{\mathbf{Q}_p}$  for the  $p$ -adic topology and let

$$\tilde{\mathbf{E}} = \varprojlim \mathbf{C}_p = \{(x^{(0)}, x^{(1)}, \dots) \mid (x^{(n+1)})^p = x^{(n)}\}$$

and let  $\tilde{\mathbf{E}}^+$  be the set of  $x \in \tilde{\mathbf{E}}$  such that  $x^{(0)} \in \mathcal{O}_{\mathbf{C}_p}$ . If  $x = (x^{(i)})$  and  $y = (y^{(i)})$  are two elements of  $\tilde{\mathbf{E}}$ , we define the sum  $x + y$  and their product  $xy$  by

$$(x + y)^{(i)} = \lim_{j \rightarrow +\infty} (x^{(i+j)} + y^{(i+j)})^{p^j} \quad \text{and} \quad (xy)^{(i)} = x^{(i)} y^{(i)},$$

which makes  $\tilde{\mathbf{E}}$  an algebraically closed field of characteristic  $p$ . If  $x = (x^{(n)}) \in \tilde{\mathbf{E}}$ , let  $\nu_E(x) = \nu_p(x^{(0)})$ . This is a valuation on  $\tilde{\mathbf{E}}$  and  $\tilde{\mathbf{E}}$  is complete for this valuation; the ring of integers of  $\tilde{\mathbf{E}}$  is  $\tilde{\mathbf{E}}^+$ . If  $\mathfrak{a}$  is an ideal of  $\mathcal{O}_{\mathbf{C}_p}$  contains  $p$  and contained in maximal ideal of  $\mathcal{O}_{\mathbf{C}_p}$ , the  $\tilde{\mathbf{E}}^+$  is identified with the projective limit of  $A_n$ , where if  $n \in \mathbf{N}$ , we put  $A_n = \mathcal{O}_{\mathbf{C}_p}/\mathfrak{a}$  and the transition amp from  $A_{n+1}$  to  $A_n$  is given by  $x \mapsto x^p$ .

Let  $\varepsilon = (1, \varepsilon^{(1)}, \dots, \varepsilon^{(n)}, \dots)$  be an element of  $\tilde{\mathbf{E}}$  such that  $\varepsilon^{(1)} \neq 1$ , this implies that  $\varepsilon^{(n)}$  is an primitive  $p^n$ -th roots of unity if  $n \geq 1$ . Let  $\bar{\pi} = \varepsilon - 1$ , we have  $\nu_E(\bar{\pi}) = \frac{p}{p-1}$  and denotes  $\mathbf{E}_{\mathbf{Q}_p}$  the subfield  $\mathbf{F}_p((\bar{\pi}))$  of  $\tilde{\mathbf{E}}$ . We denote  $\mathbf{E}$  the separable closure of  $\mathbf{E}_{\mathbf{Q}_p}$  in  $\tilde{\mathbf{E}}$  and  $\mathbf{E}^+$  (resp.  $\mathfrak{m}_{\mathbf{E}}$ ) the ring of integers (resp. the maximal ideal of  $\mathbf{E}^+$ ).

By ramification theory, if  $K$  is a finite extension of  $\mathbf{Q}_p$ , then for all  $\eta > 0$ , there exists  $n_\eta \in \mathbf{N}$  such that if  $n \geq n_\eta$ , and if  $\tau \in \Gamma_{K_n}$ , then  $\nu_p(\tau(x) - x) \geq \frac{1}{p} - \eta$ . In particular if  $\mathfrak{a}$  is an ideal of  $\mathcal{O}_{\mathbf{C}_p}$  defined by  $\mathfrak{a} = \{x \in \mathcal{O}_{\mathbf{C}_p} \mid \nu_p(x) \geq \frac{1}{p}\}$ , then  $N_{K_{n+1}/K_n}(x) - x^p \in \mathfrak{a}$  if  $n$  is large enough and  $x \in \mathcal{O}_{K_{n+1}}$ . This allows us to construct a map  $\iota_K$  from the projective limit  $\varprojlim \mathcal{O}_{K_n}$  of  $\mathcal{O}_{K_n}$  with respect to norm map to  $\tilde{\mathbf{E}}^+$  (field of norm), such that  $u = (u^{(n)})_{n \in \mathbf{N}}$  associates to  $\iota_K(u) = (x^{(n)})_{n \in \mathbf{N}}$ , where  $x^{(n)}$  is the image of  $u^{(n)}$  in  $\mathcal{O}_{\mathbf{C}_p}/\mathfrak{a}$  if  $n$  large enough. Hence we have the following proposition:

**Proposition 3.1.** *If  $K$  is a finite extension of  $\mathbf{Q}_p$ , then  $\iota_K$  induces bijection from  $\varprojlim \mathcal{O}_{K_n}$  to the ring of integers  $\mathbf{E}_K^+$  of  $\mathbf{E}_K = \mathbf{E}^{H_K}$ .*

By this proposition, one can show that  $\mathbf{E}_K$  is a finite separable extension of  $\mathbf{E}_{\mathbf{Q}_p}$  of degree  $[H_{\mathbf{Q}_p} : H_K] = [K_\infty : \mathbf{Q}_p(\mu_{p^\infty})]$  and one can identify  $\text{Gal}(\mathbf{E}/\mathbf{E}_K)$  with  $H_K$ .

**Remark 3.2.**

- i) If  $F$  is a finite unramified extension of  $\mathbf{Q}_p$  with residue field  $k_F$ , the field  $\mathbf{E}_F$  is the composition of  $k_F$  and  $\mathbf{E}_{\mathbf{Q}_p}$ , that is,  $k_F((\bar{\pi}))$ .
- ii) If  $K$  is a finite extension of  $\mathbf{Q}_p$  and  $F = K \cap \mathbf{Q}_p^{nr}$  it maximal unramified subfield, then  $\mathbf{E}_K$  is an extension of  $\mathbf{E}_F$  of degree  $[K_\infty : F_\infty]$  which is equal to  $[K_n : F_n]$  for  $n$  large enough.

**3.2. The field  $\tilde{\mathbf{B}}$  and its subrings.** Let  $\tilde{\mathbf{A}} = W(\tilde{\mathbf{E}})$  (resp.  $\tilde{\mathbf{A}}^+ = W(\tilde{\mathbf{E}}^+)$ ) the Witt vectors with coefficients in  $\tilde{\mathbf{E}}$  (resp.  $\tilde{\mathbf{E}}^+$ ). By construction, we have  $\tilde{\mathbf{A}}/p\tilde{\mathbf{A}} = \tilde{\mathbf{E}}$  (resp.  $\tilde{\mathbf{A}}^+/p\tilde{\mathbf{A}}^+ = \tilde{\mathbf{E}}^+$ ). Let

$$\tilde{\mathbf{B}} = \tilde{\mathbf{A}}[1/p] = \left\{ \sum_{k \gg -\infty} p^k [x_k] \mid x_k \in \tilde{\mathbf{E}} \right\} \quad (\text{resp. } \tilde{\mathbf{B}}^+ = \tilde{\mathbf{A}}^+[1/p] = \left\{ \sum_{k \gg -\infty} p^k [x_k] \mid x_k \in \tilde{\mathbf{E}}^+ \right\}),$$

where  $[x] \in \tilde{\mathbf{A}}$  is the Teichmüller lift of  $x \in \tilde{\mathbf{E}}$  (resp.  $\tilde{\mathbf{E}}^+$ ).

We endows  $\tilde{\mathbf{A}}$  (resp.  $\tilde{\mathbf{A}}^+$ ) the topology by taking the collection of open sets  $\{[\bar{\pi}]^k \tilde{\mathbf{A}}^+ + p^n \tilde{\mathbf{A}}\}_{k, n \geq 0}$  (resp.  $\{([\bar{\pi}]^k + p^n) \tilde{\mathbf{A}}^+\}_{k, n \geq 0}$ ) as family of neighborhoods of 0 and endow  $\tilde{\mathbf{B}} =$

$\cup_{n \in \mathbf{N}} p^{-n} \tilde{\mathbf{A}}$  (resp.  $\tilde{\mathbf{B}}^+$ ) the inductive limit topology. The action of  $G_{\mathbf{Q}_p}$  on  $\tilde{\mathbf{E}}$  can be extended by continuity to  $\tilde{\mathbf{A}}$  and  $\tilde{\mathbf{B}}$  which commutes with the Frobenius action  $\varphi$ .

For  $F$  finite unramified extension over  $\mathbf{Q}_p$ , let  $\pi = [\varepsilon] - 1$ , we define  $\mathbf{A}_F$  the closure of  $\mathcal{O}_F[[\pi, \pi^{-1}]]$  in  $\tilde{\mathbf{A}}$  by the above topology, thus

$$\mathbf{A}_F = \left\{ \sum_{k \in \mathbf{Z}} a_k \pi^k \mid a_k \in \mathcal{O}_F, \lim_{k \rightarrow -\infty} \nu_p(a_k) = +\infty \right\}$$

which is a complete discrete valuation ring with residual field  $\mathbf{E}_F$  and the Galois action and Frobenius action is defined by

$$\varphi(\pi) = (1 + \pi)^p - 1 \quad \text{and} \quad g(\pi) = (1 + \pi)^{\chi(g)} - 1 \quad g \in G_F,$$

and its fraction field  $\mathbf{B}_F = \mathbf{A}_F[\frac{1}{p}]$  is stable by actions of  $\varphi$  and  $G_F$ .

Let  $\mathbf{B}$  be the completion for the  $p$ -adic topology of the maximal unramified extension of  $\mathbf{B}_F$  in  $\tilde{\mathbf{B}}$  and  $\mathbf{A} = \mathbf{B} \cap \tilde{\mathbf{A}}$ . We have  $\mathbf{B} = \mathbf{A}[\frac{1}{p}]$  and  $\mathbf{A}$  are a complete discrete valuation ring with fractional field  $\mathbf{B}$  and residual field  $\mathbf{E}$ . We then define  $\mathbf{B}^+ = \mathbf{B} \cap \tilde{\mathbf{B}}^+$  and  $\mathbf{A}^+ = \mathbf{A} \cap \tilde{\mathbf{A}}^+$ . These rings are endowed with an action of Galois and a Frobenius deduced from those on  $\tilde{\mathbf{E}}$ .

If  $K$  is a finite extension of  $\mathbf{Q}_p$ , we put  $\mathbf{A}_K = \mathbf{A}^{H_K}$  and  $\mathbf{B}_K = \mathbf{A}_K[1/p]$ , this makes  $\mathbf{A}_K$  a complete discrete valuation ring with residual field  $\mathbf{E}_K$  and fraction field  $\mathbf{B}_K = \mathbf{A}_K[1/p]$ . On the other hand, when  $K = F$ , the definitions of  $\mathbf{A}_F$  and  $\mathbf{B}_F$  coincide with previous definitions. We put  $\mathbf{A}_F^+ = (\mathbf{A}^+)^{H_F}$  and  $\mathbf{B}_F^+ = (\mathbf{B}^+)^{H_F}$  then by using fields of norm above, we can show that  $\mathbf{A}_F^+ = \mathcal{O}_F[[\pi]]$  and  $\mathbf{B}_F^+ = F[[\pi]]$ .

If  $L$  is a finite extension of  $K$ ,  $\mathbf{B}_L$  is an unramified extension of  $\mathbf{B}_K$  of degree  $[L_\infty : K_\infty]$ . If  $L/K$  is Galois extension, then the extension  $\tilde{\mathbf{B}}_L/\tilde{\mathbf{B}}_K$  and  $\mathbf{B}_L/\mathbf{B}_K$  is Galois with Galois group  $\text{Gal}(\tilde{\mathbf{B}}_L/\tilde{\mathbf{B}}_K) = \text{Gal}(\mathbf{B}_L/\mathbf{B}_K) = \text{Gal}(\mathbf{E}_L/\mathbf{E}_K) = \text{Gal}(L_\infty/K_\infty) = H_K/H_L$ .

**Remark 3.3.**

- i) If  $\pi_K$  is a uniformizer of  $\mathbf{E}_K$ , let  $\pi_K$  be any lifting of  $\pi_K$  in  $\mathbf{A}_K$ . Then,

$$\mathbf{A}_K = \left\{ \sum_{k \in \mathbf{Z}} a_k \pi_K^k \mid a_k \in \mathcal{O}_{F'}, \lim_{k \rightarrow -\infty} \nu_p(a_k) = +\infty \right\}$$

where  $F'$  is the maximal unramified extension of  $F$  contained in  $K_\infty$ .

- ii) In the above construction, the correspondence  $R \rightarrow \tilde{R}$  is obtained by making  $\varphi$  bijective and then complete, where  $R = \{\mathbf{E}_K, \mathbf{E}, \mathbf{A}_K, \mathbf{A}, \mathbf{B}_K, \mathbf{B}\}$ .

**3.3.  $(\varphi, \Gamma)$ -module and Galois representations.** A  $p$ -adic representation  $V$  is a finite dimensional  $\mathbf{Q}_p$ -vector space with a continuous linear action of  $G_K$ . It is easy to see that there is always a  $\mathbf{Z}_p$ -lattice of  $V$  which is stable by the action of  $G_K$ , and such lattices will be denoted by  $T$  (called a  $\mathbf{Z}_p$ -representation). The main strategy due to Fontaine for studying  $p$ -adic representations of a group  $G$  is to construct topological  $\mathbf{Q}_p$ -algebras  $B$  (period rings), endowed with an action of  $G$  and some additional structures so that if  $V$  is a  $p$ -adic representation, then

$$D_B(V) = (B \otimes_{\mathbf{Q}_p} V)^G$$

is a  $B^G$ -module which inherits these structures, and so that the functor  $V \mapsto D_B(V)$  gives interesting invariants of  $V$ . We say that a  $p$ -adic representation  $V$  of  $G$  is  $B$ -admissible if we have  $B \otimes_{\mathbf{Q}_p} V \simeq B^d$  as  $B[G]$ -modules.

**Definition 3.4.** If  $K$  is a finite extension of  $\mathbf{Q}_p$ , we say



- i) A  $(\varphi, \Gamma)$ -module over  $\mathbf{A}_K$  (resp.  $\mathbf{B}_K$ ) is a  $\mathbf{A}_K$ -module of finite type (resp. a finite dimensional  $\mathbf{B}_K$ -vector space) equipped with a  $\Gamma_K$ -action and a Frobenius action  $\varphi$  which commutes with  $\Gamma_K$ .
- ii) A  $(\varphi, \Gamma)$ -module  $D$  over  $\mathbf{A}_K$  is *étale* if  $\varphi(D)$  generates  $D$  as an  $\mathbf{A}_K$ -module. A  $(\varphi, \Gamma)$ -module  $D$  over  $\mathbf{B}_K$  is *étale* if it has an  $\mathbf{A}_K$ -lattice which is *étale*, equivalently, there exists a basis  $\{e_1, \dots, e_d\}$  over  $\mathbf{B}_K$ , such that the matrix of  $\varphi$  in terms of the basis is in  $GL_d(\mathbf{A}_K)$ .

If  $K$  is a finite extension of  $\mathbf{Q}_p$  and  $V$  is a  $\mathbf{Z}_p$ -representation (resp.  $p$ -adic representation) of  $G_K$ , we put

$$D(V) = (\mathbf{A} \otimes_{\mathbf{Z}_p} V)^{H_K} \quad (\text{resp. } D(V) = (\mathbf{B} \otimes_{\mathbf{Q}_p} V)^{H_K})$$

Since the action of  $\varphi$  commutes with  $G_K$ ,  $D(V)$  is equipped with a Frobenius action  $\varphi$  which commutes with the residual action  $G_K/H_K = \Gamma_K$ . This make  $D(V)$  a  $(\varphi, \Gamma)$ -module over  $\mathbf{A}_K$  (resp.  $\mathbf{B}_K$ ).

On the other hand, if  $V$  is a  $\mathbf{Z}_p$ -representation (resp. a  $p$ -adic representation) of  $G_K$ , then  $(\mathbf{A} \otimes_{\mathbf{A}_K} D(V))^{\varphi=1}$  (resp.  $(\mathbf{B} \otimes_{\mathbf{A}_K} D(V))^{\varphi=1}$ ) is canonically isomorphic to  $V$  as a representation of  $G_K$ . In other words,  $V$  is determined by the  $(\varphi, \Gamma)$ -module  $D(V)$ .

**Theorem 3.5.** (*Fontaine*) *The correspondence*

$$V \longmapsto D(V) = (\mathbf{A} \otimes_{\mathbf{Z}_p} V)^{H_K}$$

*is an equivalence of  $\otimes$  categories from the category of  $\mathbf{Z}_p$ -representations (resp.  $p$ -adic representation) of  $G_K$  to the category of étale  $(\varphi, \Gamma)$ -module over  $\mathbf{A}_K$  (resp.  $\mathbf{B}_K$ ), and its inverse functor is*

$$D \longmapsto V(D) = (\mathbf{A} \otimes_{\mathbf{A}_K} D)^{\varphi=1}.$$

#### 4. $(\varphi, \Gamma)$ -MODULES AND GALOIS COHOMOLOGY

**4.1. The complex  $C_{\varphi, \gamma}(K, V)$ .** Let  $K$  be a finite extension of  $\mathbf{Q}_p$  such that  $\Gamma_K$  is isomorphic to  $\mathbf{Z}_p$  (i.e. contains  $\mathbf{Q}_p(\mu_p)$  if  $p \geq 3$  or three quadratic ramified extensions of  $\mathbf{Q}_2$  if  $p = 2$ ) and  $\gamma$  is a generator of  $\Gamma_K$ . If  $V$  is a  $\mathbf{Z}_p$ -representation or  $p$ -adic representation of  $G_K$  and  $f : D(V) \rightarrow D(V)$  is a  $\mathbf{Z}_p$ -linear map commutes with action of  $\Gamma$ , we denote  $C_{f, \gamma}(K, V)$  the complex

$$0 \longrightarrow D(V) \longrightarrow D(V) \oplus D(V) \longrightarrow D(V) \longrightarrow 0$$

where maps  $D(V)$  to  $D(V) \oplus D(V)$  and  $D(V) \oplus D(V)$  to  $D(V)$  respectively by

$$x \mapsto ((f-1)x, (\gamma-1)x) \quad \text{and} \quad (a, b) \mapsto (\gamma-1)a - (f-1)b$$

we denote  $Z^i(C_{f, \gamma}(K, V))$  (resp.  $B^i(C_{f, \gamma}(K, V))$ , resp.  $H^i(C_{f, \gamma}(K, V)) = \frac{Z^i(C_{f, \gamma}(K, V))}{B^i(C_{f, \gamma}(K, V))}$ ) the  $i$ -th cocycles (resp. coboundaries, resp. cohomologies) of complex  $C_{f, \gamma}(K, V)$ .

The  $C_{f, \gamma}(K, V)$  canonically and functorially identified with the Galois cohomology group  $H^i(K, V)$  (c.f. [Her98]). The following proposition gives the case of  $H^1$ .

Let  $\Lambda_K = \mathbf{Z}_p[[\Gamma_K]]$  the complete group algebra of  $\Gamma_K$ . Since  $\Gamma_K$  acts continuously on  $D(V)$ , we can view  $D(V)$  as a  $\Lambda_K$ -module. On the other hand,  $\Gamma_K$  is pro-cyclic, if  $\gamma$  is a generator of  $\Gamma_K$  and  $\gamma'$  is any element of  $\Gamma_K$ , then the element  $\frac{\gamma'-1}{\gamma-1}$  of  $\text{Frac}(\Lambda_K)$  is indeed in  $\Lambda_K$ . Moreover, the  $G_K$  action factors through  $\Gamma_K$  on  $D(V)$ , so the expression  $\frac{\sigma-1}{\gamma-1}y$  make sense if  $y \in D(V)$ ,  $\sigma \in G_K$  and  $\gamma$  is a generator of  $\Gamma_K$ .

**Proposition 4.1.**

- i) If  $(x, y) \in Z^1(C_{\varphi, \gamma}(K, V))$  and  $b \in \mathbf{A} \otimes_{\mathbf{Z}_p} V$  is a solution of  $(\varphi - 1)b = x$ , then  $\sigma \mapsto c_{x, y}(\sigma) = \frac{\sigma - 1}{\gamma - 1}y - (\sigma - 1)b$  is a cocycle of  $G_K$  with values in  $V$ .
- ii) The map sends  $(x, y) \in Z^1(C_{\varphi, \gamma}(K, V))$  to the class of  $c_{x, y}$  in  $H^1(K, V)$  induces an isomorphism  $\iota_{\varphi, \gamma}$  of  $H^1(C_{\varphi, \gamma}(K, V))$  to  $H^1(K, V)$ .

*Proof.* It clear that  $\sigma \mapsto c_{x, y}(\sigma)$  is a cocycle by definition. On the other hand, we have

$$(\varphi - 1)(c_{x, y}(\sigma)) = \frac{\sigma - 1}{\gamma - 1}((\varphi - 1)y) - (\sigma - 1)x = 0$$

since  $(\gamma - 1)x = (\varphi - 1)y$ . Hence  $c_{x, y}(\sigma) \in (\mathbf{A} \otimes_{\mathbf{Z}_p} V)^{\varphi=1} = V$ . This proves (i).

To prove (ii), suppose the image of  $c_{x, y}$  in  $H^1(K, V)$  is zero, there exist  $z \in V$  such that

$$\frac{\sigma - 1}{\gamma - 1}y - (\sigma - 1)(b + z) = 0 \quad \forall \sigma \in G_K.$$

We deduce that  $b + z$  is stable by  $H_K$  and therefore belongs to  $D(V)$ . Take  $\sigma = \gamma$ , we have  $y = (\gamma - 1)(b + z)$  and hence  $x = (\varphi - 1)(b + z)$ , which implies  $(x, y) \in B^1(C_{\varphi, \gamma}(K, V))$  and the injectivity of  $\iota_{\varphi, \gamma}$  follows.

To prove the surjectivity, let  $c \in H^1(K, V)$  and  $V'$  an extension of  $\mathbf{Z}_p$  by  $V$  corresponding to  $c$ . That is, an exact sequence

$$0 \longrightarrow V \longrightarrow V' \longrightarrow \mathbf{Z}_p \longrightarrow 0$$

such that  $e \in V'$  sends to  $1 \in \mathbf{Z}_p$  and  $\sigma(e) = e + c_\sigma$ , where  $\sigma \mapsto c_\sigma$  is the cocycle of  $G_K$  represents  $c$ . Apply functor  $D$ , we get

$$0 \longrightarrow D(V) \longrightarrow D(V') \longrightarrow D(\mathbf{Z}_p) \longrightarrow 0,$$

let  $\tilde{e} \in D(V')$  element maps to  $1 \in \mathbf{Z}_p = D(\mathbf{Z}_p)$  and  $x, y$  elements of  $D(V)$  defined by  $x = (\varphi - 1)\tilde{e}$  and  $y = (\gamma - 1)\tilde{e}$ . Since  $\gamma$  and  $\varphi$  commute,  $(x, y)$  is belongs to  $Z^1(C_{\varphi, \gamma}(K, V))$ . On the other hand,  $b = \tilde{e} - e \in \mathbf{A} \otimes_{\mathbf{Z}_p} V$  satisfies  $(\varphi - 1)b = x$ , so we have

$$c_{x, y}(\sigma) = \frac{\sigma - 1}{\gamma - 1}y - (\sigma - 1)b = (\sigma - 1)(\tilde{e} - b) = (\sigma - 1)e = c_\sigma.$$

From this, we deduce the surjectivity of  $\iota_{\varphi, \gamma}$ .  $\square$

If  $\gamma'$  is another generator of  $\Gamma_K$ , then  $\frac{\gamma - 1}{\gamma' - 1} \in \text{Frac}(\Gamma_K)$  is indeed a unit in  $\Gamma_K$  and the diagram

$$\begin{array}{ccccccc} C_{\varphi, \gamma}(K, V) : 0 & \longrightarrow & D(V) & \longrightarrow & D(V) \oplus D(V) & \longrightarrow & D(V) \longrightarrow 0 \\ & & \downarrow \frac{\gamma - 1}{\gamma' - 1} & & \downarrow \frac{\gamma - 1}{\gamma' - 1} \oplus id & & \downarrow id \\ C_{\varphi, \gamma'}(K, V) : 0 & \longrightarrow & D(V) & \longrightarrow & D(V) \oplus D(V) & \longrightarrow & D(V) \longrightarrow 0 \end{array}$$

is commutative. It hence induces via cohomology an isomorphism  $\iota_{\gamma, \gamma'}$  from  $H^1(C_{\varphi, \gamma}(K, V))$  to  $H^1(C_{\varphi, \gamma'}(K, V))$ .

Since we assume  $\Gamma_K$  is torsion free, we have  $\chi(\gamma) \in 1 + p\mathbf{Z}_p$  for  $\gamma \in \Gamma_K$ , then there exists  $k \geq 1$  such that  $\log_p(\chi(\gamma)) \in p^k\mathbf{Z}_p^*$  and we'll write  $\log_p^0(\gamma) = \log_p(\chi(\gamma))/p^k$ . The following lemma shows that  $\log_p^0(\gamma)\iota_{\varphi, \gamma}$  does not depend on the choice of generator  $\gamma$  of  $\Gamma_K$ .

**Lemma 4.2.** *If  $\gamma$  and  $\gamma'$  are two generators of  $\Gamma_K$ , then the isomorphisms  $\log_p^0(\gamma)\iota_{\varphi, \gamma}$  and  $\log_p^0(\gamma')\iota_{\varphi, \gamma'} \circ \iota_{\gamma, \gamma'}$  from  $H^1(C_{\varphi, \gamma}(K, V))$  to  $H^1(K, V)$  are equal.*

*Proof.* If  $(x, y) \in Z^1(C_{\varphi, \gamma}(K, V))$ . Let  $b$  (resp.  $b'$ ) be element of  $\mathbf{A} \otimes_{\mathbf{Z}_p} V$  verifies  $(\varphi - 1)b = x$  (resp.  $(\varphi - 1)b' = \frac{\gamma-1}{\gamma'-1}x$ ). Since  $\frac{\log_p^0(\gamma)}{\gamma-1} - \frac{\log_p^0(\gamma')}{\gamma'-1} \in \mathbf{Z}_p[[\Gamma_K]]$ , we can write the cocycle associates to  $\log_p^0(\gamma')\iota_{\varphi, \gamma} \circ \iota_{\gamma, \gamma'}(x, y) - \log_p^0(\gamma)\iota_{\varphi, \gamma}(x, y)$  as  $\sigma \mapsto (\sigma - 1)c$ , where

$$c = \left( \frac{\log_p^0(\gamma')}{\gamma' - 1} - \frac{\log_p^0(\gamma)}{\gamma - 1} \right) y - (\log_p^0(\gamma')b' - \log_p^0(\gamma)b)$$

and the relation  $(\varphi - 1)y = (\gamma - 1)x$  implies  $(\varphi - 1)c = 0$ , hence  $c \in V$  and the cocycle is indeed a coboundary, which leads to the conclusion.  $\square$

**4.2. The operator  $\psi$ .** To calculate  $H^1(C_{\varphi, \gamma}(K, V))$  we have to understand the group  $D(V)^{\varphi=1}$  and  $\frac{D(V)}{\varphi-1}$ . The problem is that the group  $\frac{D(V)}{\varphi-1}$  is too complicated to write it down. To solve this difficulty, we introduce the left inverse of  $\varphi$ .

The field  $\mathbf{B}$  is an extension of degree  $p$  of  $\varphi(B)$ , which allows up to define the operator  $\psi : \mathbf{B} \rightarrow \mathbf{B}$  by the formula  $\psi(x) = \frac{1}{p}\varphi^{-1}(\text{Tr}_{\mathbf{B}/\varphi(\mathbf{B})}(x))$ . More explicitly, one can verify that  $\{1, [\varepsilon], \dots, [\varepsilon]^{p-1}\}$  is a basis of  $\mathbf{A}$  over  $\varphi(\mathbf{A})$  (hence  $\mathbf{B}$  over  $\varphi(\mathbf{B})$ ) so we have

$$\psi\left(\sum_{i=0}^{p-1} [\varepsilon]^i \varphi(x_i)\right) = x_0 \quad x_i \in \mathbf{B} \quad \text{and} \quad \psi(\varphi(x)) = x \quad x \in \mathbf{B}.$$

The operator  $\psi$  commute with the action of  $G_K$  and  $\psi(\mathbf{A}) \subset \mathbf{A}$ .

Since  $\psi$  commutes with the action of  $G_K$ , if  $V$  is a  $\mathbf{Z}_p$ -representation or a  $p$ -adic representation of  $G_K$ , the module  $D(V)$  inherit the action of  $\psi$  and commute with  $\Gamma_K$ . That is, the unique map  $\psi : D(V) \rightarrow D(V)$  with

$$\psi(\varphi(a)x) = a\psi(x), \quad \psi(a\varphi(a)) = \psi(a)x$$

if  $a \in \mathbf{A}_K$ ,  $x \in D(V)$ .

**Proposition 4.3.** *If  $V$  is a  $\mathbf{Z}_p$ -representation or a  $p$ -adic representation of  $G_K$ , then  $\gamma - 1$  is invertible on  $D(V)^{\psi=0}$ .*

*Proof.* See [Her98].  $\square$

**Lemma 4.4.** *We have a commutative diagram of complexes*

$$\begin{array}{ccccccc} C_{\varphi, \gamma}(K, V) : 0 & \longrightarrow & D(V) & \longrightarrow & D(V) \oplus D(V) & \longrightarrow & D(V) \longrightarrow 0 \\ & & \downarrow id & & \downarrow (-\psi, id) & & \downarrow -\psi \\ C_{\psi, \gamma}(K, V) : 0 & \longrightarrow & D(V) & \longrightarrow & D(V) \oplus D(V) & \longrightarrow & D(V) \longrightarrow 0 \end{array}$$

which induces an isomorphism  $\iota$  from  $H^1(C_{\varphi, \gamma}(K, V))$  to  $H^1(C_{\psi, \gamma}(K, V))$ .

*Proof.* The commutativity of diagram follows from definition. Since  $\psi$  is surjective, the cokernel complex is 0. The kernel complex is

$$0 \longrightarrow 0 \longrightarrow D(V)^{\psi=0} \xrightarrow{\gamma-1} D(V)^{\psi=0} \longrightarrow 0,$$

which has no cohomology by proposition 4.3.  $\square$

**Notation 4.5.** We denote  $\iota_{\psi, \gamma}$  the isomorphism of  $H^1(C_{\psi, \gamma}(K, V))$  to  $H^1(K, V)$  obtained by composite  $\iota_{\varphi, \gamma}$  and  $\iota^{-1}$ .

**Remark 4.6.** The same proof as lemma 4.2 shows that  $\log_p^0(\gamma)\iota_{\psi,\gamma}$  does not depend on the generator  $\gamma$  of  $\Gamma_K$ .

**Lemma 4.7.** *The map which sends  $(x, y) \in Z^1((C_{\varphi,\gamma}(K, V))$  to the image of  $x$  in  $\frac{D(V)}{\psi-1}$  induces an exact sequence*

$$0 \longrightarrow D(V)_{\Gamma_K}^{\psi=1} \longrightarrow H^1(C_{\psi,\Gamma_K}(K, V)) \longrightarrow \left(\frac{D(V)}{\psi-1}\right)^{\Gamma_K} \longrightarrow 0$$

*Proof.*  $\bar{x} \in \frac{D(V)}{\psi-1}$  is fixed by  $\Gamma_K$  if and only if there exists  $(x, y) \in Z^1(C_{\varphi,\gamma}(K, V))$  whose image in  $\frac{D(V)}{\psi-1}$  is equal to  $\bar{x}$ . The kernel of the map is the sum of  $B^1(C_{\psi,\gamma}(K, V))$  and the set  $X$  of elements of the form  $(0, y)$  where  $y \in D(V)^{\psi=1}$ . One observes that  $X \cap B^1(C_{\varphi,\gamma}(K, V))$  is constituted by couples of the form  $(0, y)$  where  $y \in (\gamma - 1)D(V)^{\psi=1}$ .  $\square$

**Remark 4.8.** By [Her98], one can show that Herr complex indeed computes Galois cohomology group  $H^i(K, V)$ , hence we have

- $H^0(K, V) \simeq D(V)^{\psi=1, \gamma=1} \simeq D(V)^{\varphi=1, \gamma=1}$ .
- $H^2(K, V) \simeq \frac{D(V)}{(\psi-1, \gamma-1)}$ .
- $H^i(K, V) = 0$  if  $i \geq 2$ .

Similar to case of  $\varphi$ , the modules  $D(V)^{\psi=1}$  and  $\frac{D(V)}{\psi-1}$  can be interpreted naturally as Iwasawa algebra. Moreover, the module  $\frac{D(V)}{\psi-1}$  is "small" compared to  $\frac{D(V)}{\varphi-1}$ , thus we can write  $H^1(K, V)$  mainly as the submodule  $D(V)^{\psi=1}$ . More precisely, we have the following proposition whose proof would be in the following two subsections.

**Proposition 4.9.** *If  $V$  is a  $\mathbf{Z}_p$ -representation (resp. a  $p$ -adic representation) of  $G_K$ , then*

- i)  $D(V)^{\psi=1}$  is compact (resp. locally compact) and generates the  $\mathbf{A}_K$ -module  $(\mathbf{B}_K\text{-vector space}) D(V)$ .
- ii)  $\frac{D(V)}{\psi-1}$  is a free  $\mathbf{Z}_p$ -module of finite rank (resp. a finite dimensional  $\mathbf{Q}_p$ -vector space).

**Remark 4.10.** Since the  $p$ -adic representation case can be deduce from  $\mathbf{Z}_p$ -representation case by tensor  $\mathbf{Q}_p$ , we only need to treat the  $\mathbf{Z}_p$ -representation case.

**4.3. The compactness of  $D(V)^{\psi=1}$ .** The goal of this paragraph is to prove the following lemma. In particular, when  $n = 0$  and  $N = +\infty$  is equivalent to the compactness of  $D(V)^{\psi=1}$ .

**Lemma 4.11.** *If  $V$  is a  $\mathbf{Z}_p$ -representation of  $G_K$ ,  $x \in D(V)$  and  $N \in \mathbf{N} \cup \{+\infty\}$ , the set of solutions  $y \in D(V)/p^{N+1}D(V)$  of the equation  $(\psi - 1)y = x$  is compact.*

Let  $\mathbf{A}_{Q_p}^+$  is the subring  $\mathbf{Z}_p[[\pi]]$  of  $\mathbf{A}_{Q_p}$ , and let  $A = \mathbf{A}_{Q_p}^+[[\frac{p}{\pi^{p-1}}]]$ , then  $A$  is a compact subring of  $\mathbf{A}_{Q_p}$  such that elements of  $A$  can be written as  $x = \sum_{n \in \mathbf{Z}} x_n \pi^n$  where  $(x_n)_{n \in \mathbf{Z}}$  is a sequence in  $\mathbf{Z}_p$  such that we have  $\nu_p(x_n) \geq -\frac{n}{p-1}$  if  $n \leq 0$ .

If  $x \in \mathbf{A}_{Q_p}$ , let  $w_n(x) \in \mathbf{N}$  the smallest integer  $k$  such that  $x$  belongs to  $\pi^{-k}A + p^{n+1}\mathbf{A}_{Q_p}$ . If  $x$  is fixed, the sequence  $\{w_n(x)\}_{n \in \mathbf{N}}$  is increasing and we have

$$\begin{aligned} w_n(x + y) &\leq \sup(w_n(x), w_n(y)) \\ w_n(xy) &\leq \sup_{i+j=n} (w_i(x) + w_j(y)) \leq w_n(x) + w_n(y) \\ w_n(\varphi(x)) &\leq pw_n(x) \end{aligned}$$

the first two inequality follow from  $A$  is a ring and the third is because  $\frac{\varphi(\pi)}{\pi^p}$  is an unit in  $A$  (This is the reason for working with  $A$  instead of  $\mathbf{A}_{Q_p}^+$  by defining the map  $w_n$ ) and such that  $x \in \pi^{-k}A + p^{n+1}\mathbf{A}_{Q_p}$  implies  $\varphi(x) \in \varphi(\pi)^{-k}A + p^{n+1}\mathbf{A}_{Q_p} = \pi^{-pk}A + p^{n+1}\mathbf{A}_{Q_p}$ .

**Lemma 4.12.**

- i) If  $k \in \mathbf{N}$ , then  $\psi(\pi^k) \in \mathbf{A}_{Q_p}^+$  and  $\psi(\pi^{-k}) \in \pi^{-k}\mathbf{A}_{Q_p}^+$
- ii)  $\psi(A) \subset A$ .

*Proof.* ii) follows from i) and the definition of  $A$ . Since  $\varphi(\pi) = (1 + \pi)^p - 1$  is a monic polynomial of degree  $p$  in  $\pi$  and  $[\varepsilon]^i = (1 + \pi)^i$  is a monic polynomial of degree  $i$  in  $\pi$ , hence

$$\{[\varepsilon]^i \varphi(\pi)^j\}_{0 \leq i \leq p-1, j \in \mathbf{N}} \text{ forms a basis of polynomials in } \pi. \text{ Moreover, } \psi([\varepsilon]^i \varphi(\pi)^j) = \begin{cases} 0 & i \neq 0 \\ \pi^j & i = 0 \end{cases},$$

we thus deduce that  $\psi(\pi^k) \in \mathbf{A}_{Q_p}^+$  if  $k \geq 0$ . If  $k \geq 1$ , then

$$\text{Tr}_{\mathbf{A}_{Q_p}/\varphi(\mathbf{A}_{Q_p})}(\pi^{-k}) = \sum_{\zeta^p=1} ((1 + \pi)\zeta - 1)^{-k},$$

which can be written as the form  $\frac{P(\varphi(\pi))}{\varphi(\pi)^k}$ , where  $P$  is a polynomial with coefficient in  $\mathbf{Z}_p$ . Thus the conclusion follows.  $\square$

**Corollary 4.13.** If  $x \in \mathbf{A}_{Q_p}$  and  $n \in \mathbf{N}$ , then  $w_n(\psi(x)) \leq 1 + [\frac{w_n(x)}{p}] \leq 1 + \frac{w_n(x)}{p}$ .

*Proof.* Since  $\frac{\varphi(\pi)}{\pi^p}$  is an unit in  $A$  and  $\psi(\frac{x}{\varphi(\pi)^k}) = \frac{\psi(x)}{\pi^k}$ , we have

$$\psi(\pi^{-kp}A + p^{n+1}\mathbf{A}_{Q_p}) = \psi(\varphi(\pi)^{-k}A + p^{n+1}\mathbf{A}_{Q_p}) \subset \pi^{-k}A + p^{n+1}\mathbf{A}_{Q_p},$$

the conclusion follows.  $\square$

If  $U = (a_{i,j})_{1 \leq i,j \leq d} \in M_d(\mathbf{A}_{Q_p})$  and  $n \in \mathbf{N}$ , we define  $w_n(U)$  by  $w_n(U) = \sup_{i,j} w_n(a_{i,j})$ . Similarly if  $V$  is a  $\mathbf{Z}_p$ -representation of  $G_K$  and if  $e_1, \dots, e_d$  is a basis of  $D(V)$  over  $\mathbf{A}_{Q_p}$ , we put  $w_n(a) = \sup_i w_n(a_i)$  if  $a = \sum_{i=1}^d a_i e_i \in D(V)$ . Note that  $w_n$  depends on the choice of basis  $e_1, \dots, e_d$ .

**Lemma 4.14.** Let  $V$  be a  $\mathbf{Z}_p$ -representation of  $G_K$ ,  $e_1, \dots, e_d$  is a basis of  $D(V)$  over  $\mathbf{A}_{Q_p}$  and  $\Phi = (a_{i,j})$  the matrix defined by  $e_j = \sum_{i=1}^d a_{i,j} \varphi(e_i)$ . If  $x, y \in \mathbf{A}_{Q_p}$  satisfy the equation  $(\psi - 1)y = x$ , then  $w_n(y) \leq \sup \left( w_n(x), \frac{p}{p-1} (w_n(\Phi) + 1) \right)$  for all  $n \in \mathbf{N}$ .

*Proof.* Since  $\varphi(e_1), \dots, \varphi(e_d)$  is a basis of  $D(V)$  over  $\phi(D(V))$ , we can write  $x = \sum_{i=1}^d x_i \varphi(e_i)$  and  $y = \sum_{i=1}^d y_i \varphi(e_i)$ . We have  $\psi(y) = \sum_{i=1}^d \psi(y_i) e_i$  and the equation  $\psi(y) - y = x$  translate to system of equation

$$y_i = -x_i + \sum_{j=1}^d a_{i,j} \psi(y_j) \quad 1 \leq j \leq d.$$

One get the inequalities

$$w_n(y_i) \leq \sup \left( w_n(x_i), \sup_{1 \leq j \leq d} (w_n(a_{i,j}) + w_n(\psi(y_j))) \right) \leq \sup \left( w_n(x), w_n(\Phi) + \frac{w_n(y)}{p} + 1 \right)$$

for  $1 \leq i \leq d$ , which gives us the inequality

$$w_n(y) \leq \sup \left( w_n(x), w_n(\Phi) + \frac{w_n(y)}{p} + 1 \right)$$

and the conclusion follows.  $\square$

To deduce lemma 4.11. If  $n \in \mathbf{N} \cup \{+\infty\}$ , let  $X_n$  be the set of solutions of the equation  $(\psi - 1)y = x$  in  $D(V)/p^{n+1}D(V)$ . We want to show that  $X_n$  is compact. If  $n \in \mathbf{N}$ , let  $r_n = \sup(w_n(x), \frac{p}{p-1}(w_n(\Phi) + 1))$ . The set  $X_n$  is closed (since  $\psi - 1$  is continuous). By the previous lemma, the image of  $(\pi^{-r_n}A)^d$  is compact since  $A$  is. If  $N$  is finite, it suffices to take  $n = N$  to conclude. If  $N = +\infty$ , the map from  $x \in X_{+\infty}$  to the sequence of its images modulo  $p^{n+1}$  allows us to identify  $X_{+\infty}$  with the closed subset of compact set  $\prod_{n \in \mathbf{N}} X_n$ , and the conclusion follows.

#### 4.4. The module $\frac{D(V)}{\psi-1}$ .

**Lemma 4.15.** *If  $V$  be a  $\mathbf{Z}_p$ -representation of  $G_K$ , the module  $\frac{D(V)}{\psi-1}$  has no nonzero  $p$ -divisible element.*

*Proof.* Let  $x$  be a  $p$ -divisible element of  $\frac{D(V)}{\psi-1}$ . For each  $n \in \mathbf{N}$ , there exist elements  $y_n, z_n$  of  $D(V)$  such that  $x = p^n y_n + (\psi - 1)z_n$ . If we fix  $m \in \mathbf{N}$  and if  $n \geq m + 1$ , then  $z_n$  is a solution of equation  $\psi(z) - z = x \pmod{p^{m+1}}$ . Since the set of solutions is compact due to lemma 4.11, there exists a subsequence of  $\{z_n\}_{n \in \mathbf{N}}$  which converges modulo  $p^m$  for all  $m$  and we have a limit  $Z$  in  $D(V)$ . By passing to limit, we obtain  $x = (\psi - 1)Z$  and hence  $x = 0$  in  $\frac{D(V)}{\psi-1}$ .  $\square$

**Lemma 4.16.** *If  $V$  is a  $\mathbf{F}_p$ -representation of  $G_K$  and  $x \in \mathfrak{m}_{\mathbf{E}} \otimes V$ , then the series  $\sum_{n=0}^{+\infty} \varphi^n(x)$  and  $\sum_{n=1}^{+\infty} \varphi^n(x)$  converges in  $\mathfrak{m}_{\mathbf{E}} \otimes V$  and we have*

$$(\psi - 1) \left( \sum_{n=0}^{+\infty} \varphi^n(x) \right) = \psi(x) \quad \text{and} \quad (\psi - 1) \left( \sum_{n=1}^{+\infty} \varphi^n(x) \right) = x.$$

*Proof.* If  $e_1, \dots, e_d$  is a basis of  $V$  over  $F_p$  and  $x = x_1 e_1 + \dots + x_n e_d \in \mathfrak{m}_{\mathbf{E}} \otimes V$ , there exists  $r \geq 0$  such that if  $\nu_E(x_i) \geq r$  for  $1 \leq i \leq d$  implies that  $\nu_E(\varphi^n(x_i)) \geq p^n r$  tends to  $+\infty$  and hence we have  $\varphi^n(x)$  tends to 0 as  $n$  tends to  $+\infty$ . We thus deduce the convergence of the series. These formulas are consequence of the fact that  $\psi$  is a left inverse of  $\varphi$ .  $\square$

**Lemma 4.17.**

- i) *If  $V$  be a  $\mathbf{F}_p$ -representation of  $G_K$ , then  $\frac{D(V)}{\psi-1}$  is a finite dimensional  $\mathbf{F}_p$ -vector space.*
- ii) *There exists a open subgroup of  $\Gamma_K$  which acts trivially on  $\frac{D(V)}{\psi-1}$ .*

*Proof.* Let  $M = (\mathfrak{m}_{\mathbf{E}} \otimes V)^{H_K}$ , which is a lattice of  $D(V)$  fixed by  $\varphi$ . If  $x \in M$ , the series  $\sum_{n=1}^{+\infty} \varphi^n(x)$  converges in  $M$ , and by previous lemma, we have  $x = (\psi - 1)(\sum_{n=1}^{+\infty} \varphi^n(x))$ , which proves that  $(\psi - 1)D(V)$  contains  $M$ .

Since  $\psi$  is continuous, there exists  $c \in \mathbf{N}$  such that  $\psi(M) \subset \pi^{-c}M$  and since  $\psi(\pi^{-pk}x) = \pi^{-k}x$ , we have  $\psi(\pi^{-pk}M) \subset \pi^{-k-c}M$ . We deduce that if  $n \geq b = [\frac{pc}{p-1}] + 1$ , then  $\psi = 0$  in  $\frac{\pi^{-n+1}M}{\pi^{-n}M}$  and such that  $\psi - 1$  is bijective on  $\frac{\pi^{-n+1}M}{\pi^{-n}M}$ . Since  $D(V) = \bigcup_{n \in \mathbf{N}} \pi^{-n}M$ , which implies the natural map

from  $\frac{\pi^{-b}M}{\psi-1}$  to  $\frac{D(V)}{\psi-1}$  is an isomorphism.

To prove i), it suffices to note that  $(\psi - 1)M$  contained in  $M$ , which implies that  $\frac{D(V)}{\psi - 1}$  is a quotient of  $\frac{\pi^{-b}M}{\psi - 1}$ . To prove ii), this is because that  $\Gamma_K$  fixes  $M$  and hence  $\pi^k M$  for all  $k \in \mathbf{Z}$  and the action of  $\Gamma_K$  is continuous on  $D(V)$  and  $M$  is closed in  $D(V)$ , there exists an open subgroup of  $\Gamma_K$  acts trivially on  $\frac{\pi^{-b}M}{\psi - 1}$  since the module is endowed with discrete topology.  $\square$

**Corollary 4.18.** *If  $V$  be a  $\mathbf{Z}_p$ -representation of  $G_K$ , then  $\frac{D(V)}{\psi - 1}$  is a  $\mathbf{Z}_p$ -module of finite type.*

*Proof.*  $\frac{D(V)}{\psi - 1}/p \frac{D(V)}{\psi - 1} = \frac{D(V)}{(p, \psi - 1)} = \frac{D(V/p)}{\psi - 1}$  is a  $\mathbf{F}_p$ -vector space of finite type by the preceding lemma, together with lemma 4.15, we get the conclusion.  $\square$

Hence we deduce ii) of proposition 4.9 and it remains to prove that  $D(V)^{\psi=1}$  generate  $D(V)$ . We will need the following lemma.

**Lemma 4.19.** *If  $V$  be a  $\mathbf{F}_p$ -representation of  $G_K$  and  $X$  is a sub- $\mathbf{F}_p$ -vector space of  $D(V)^{\psi=1}$  of finite codimension, then  $X$  contains a basis of  $D(V)$  over  $\mathbf{E}_K$ .*

*Proof.* Let  $M = (\mathfrak{m}_{\mathbf{E}} \otimes V)^{H_K}$  as above. Note that by lemma 4.16, if  $x \in M^{\psi=0}$ , then the series  $\sum_{n=0}^{+\infty} \varphi^n(x)$  converges in  $D(V)$  to an element of  $D(V)^{\psi=1}$ . We denote it by  $eul(x)$ . Let  $e_1, \dots, e_d$  be a basis of  $M$  over  $\mathbf{E}_K^+$ . Let  $r$  the codimension of  $X$  in  $D(V)^{\psi=1}$ . If  $1 \leq i \leq d$  and  $j \geq 1$ , let  $z_{i,j} = eul(\varepsilon\varphi(\pi^j e_i))$ . If  $i$  and  $n \geq 1$  are fixed, the  $\{z_{i,j}\}_{n \leq j \leq n+r}$  form a set of  $r+1$  elements in  $D(V)^{\psi=1}$  and since  $X$  is of codimension  $r$  in  $D(V)^{\psi=1}$ , we can find elements  $\{a_{i,j}^{(n)}\}_{0 \leq j \leq r}$  of  $\mathbf{F}_p$  such that  $f_{i,n} = \sum_{j=0}^r a_{i,j}^{(n)} z_{i,j+n}$  belongs to  $X$ . Let  $\beta_{i,n} = \pi^n \sum_{j=0}^r a_{i,j}^{(n)} \pi^j$ . We have  $\lim_{n \rightarrow +\infty} (\varepsilon\varphi(\beta_{i,n}))^{-1} f_{i,n} = \varphi(e_i)$ , which implies that the determinant of  $f_{1,n}, \dots, f_{d,n}$  in the basis  $\varphi(e_1), \dots, \varphi(e_d)$  is nonzero if  $n \gg 0$  and we have  $f_{1,n}, \dots, f_{d,n}$  form a basis of  $D(V)$  over  $\mathbf{E}_K$  if  $n$  is great enough. The lemma follows.  $\square$

**Corollary 4.20.** *If  $V$  is a  $\mathbf{Z}_p$ -representation of  $G_K$ , then  $D(V)^{\psi=1}$  generates the  $\mathbf{A}_K$ -module  $D(V)$ .*

*Proof.* The snake lemma shows that the cokernel of the injective map  $D(V)^{\psi=1}/pD(V)^{\psi=1}$  to  $D(V/p)^{\psi=1}$  is identified with the  $p$ -torsion part of  $D(V)/(\psi - 1)$ . In particular, it is of finite dimension over  $\mathbf{F}_p$ . By the preceding lemma, we have  $D(V)^{\psi=1}/pD(V)^{\psi=1}$  contains a basis of  $D(V/p)$  over  $\mathbf{E}_K$ , which lifts to a basis in  $D(V)^{\psi=1}$  that generates  $D(V)$  over  $\mathbf{A}_K$ .  $\square$

## 5. IWASAWA THEORY AND $p$ -ADIC REPRESENTATIONS

**5.1. Iwasawa cohomology.** Recall that if  $n \in \mathbf{N}$ , we denote  $K_n$  the field  $K(\varepsilon^{(n)}) = K(\mu_{p^n})$ . On the other hand, if  $n \geq 1$  (resp.  $n \geq 2$  if  $p = 2$ ), the group  $\Gamma_{K_n}$  is isomorphic to  $\mathbf{Z}_p$ . We choose a generator  $\gamma_1$  of  $\Gamma_{K_1}$  and put  $\gamma_n = \gamma_1^{[K_N:K_1]}$  if  $n \geq 1$  (if  $p = 2$ , we can start from  $n = 2$ ), this makes  $\gamma_n$  a generator of  $\Gamma_{K_n}$ .

Let  $V$  be a  $p$ -adic representation of  $G_K$ . The Iwasawa cohomology groups  $H_{\text{Iw}}^i(K, V)$  are defined by  $H_{\text{Iw}}^i(K, V) = \mathbf{Q}_p \otimes_{\mathbf{Z}_p} H_{\text{Iw}}^i(K, T)$  where  $T$  is any  $G_K$ -stable lattice of  $V$  and where

$$H_{\text{Iw}}^i(K, T) = \varprojlim_{\text{cor}_{K_{n+1}/K_n}} H^i(K_n, T)$$

Each of the  $H^i(K, T)$  is a  $\mathbf{Z}_p[\Gamma_K/\Gamma_{K_n}]$ -module, and  $H_{\text{Iw}}^i(K, T)$  is then endowed with the structure of  $\mathbf{A}_K$ -module. Roughly speaking, theses cohomology groups are where Euler system live (at least locally).

If  $V$  is a  $\mathbf{Z}_p$ -representation or a  $p$ -adic representation of  $G_K$ , we endow  $\Lambda_K \otimes_{\mathbf{Z}_p} V$  the natural diagonal action of  $G_K$ . If we consider  $\Lambda_K \otimes_{\mathbf{Z}_p} V$  the space of measure of  $\Gamma_K$  with values in  $V$ , the measure  $\sigma(\mu)$  is the map sends continuous map  $f : \Gamma_K \mapsto V$  to the element

$$\int_{\Gamma_K} f(x) \sigma(\mu) = \sigma \left( \int_{\Gamma_K} f(\sigma x) \mu \right) \in V$$

If  $V$  is a  $\mathbf{Z}_p$ -representation or a  $p$ -adic representation of  $G_K$  and  $k \in \mathbf{Z}$ , we denote  $V(k)$  the twist of  $V$  by the  $k$ -th power of the cyclotomic character and if  $x \in V$ , we denote  $x(k)$  its image in  $V(k)$ .

If  $\mu \in H^m(K, \Lambda_K \otimes_{\mathbf{Z}_p} V)$  and if  $\tau \mapsto \mu_{\tau_1, \dots, \tau_m}$  is a continuous  $m$ -cocycle represents  $\mu$ , then  $\tau \mapsto (\int_{\Gamma_{K_n}} \chi(x)^k \mu_{\tau_1, \dots, \tau_m})(k)$  is a  $m$ -cycle of  $G_K$  with values in  $V(k)$  whose class  $(\int_{\Gamma_{K_n}} \chi(x)^k \mu)(k)$  in  $H^m(K_n, V(k))$  does not depend on the choice of cocycle represent  $\mu$ .

The Shapiro's lemma allows us to replace the projective limit in the definition of  $H_{\text{Iw}}^m(K, V)$  by a group cohomology.

**Proposition 5.1.** *Let  $V$  be a  $\mathbf{Z}_p$ -representation or a  $p$ -adic representation of  $G_K$ . If  $m \in \mathbf{N}$  and  $k \in \mathbf{Z}$ , the map sends  $\mu$  to  $(\dots, \int_{K_n} \chi(x)^k \mu(k), \dots)$  is an isomorphism from  $H^i(K, \Gamma_K \otimes_{\mathbf{Z}_p} V)$  to  $H_{\text{Iw}}^i(K, V(k))$ . In particular, if  $k \in \mathbf{Z}$ , the cohomology group  $H_{\text{Iw}}^m(K, V)$  and  $H_{\text{Iw}}^m(K, V(k))$  are isomorphic.*

*Proof.* The case of  $\mathbf{Q}_p$  follows from the case of  $\mathbf{Z}_p$  by tensoring  $\mathbf{Q}_p$ . If  $M$  is a  $G_{K_n}$ -module, we denote  $\text{Ind}_{K_n}^K M$  the set of continuous maps from  $G_K$  to  $M$  satisfies  $a(hx) = ha(x)$  if  $h \in G_{K_n}$ . The module  $\text{Ind}_{K_n}^K M$  is provided with a continuous action of  $G_K$ , the image  $ga$  of  $a$  by  $g \in G_K$ , is given by the formula  $(ga)(x) = a(xg)$ . If  $M$  is a  $G_K$  module, and  $a \in \text{Ind}_{K_n}^K M$ , the map sends  $x \in G_K$  to  $x^{-1}(a(x))$  is constant modulo  $G_{K_n}$ , and the map of  $\text{Ind}_{K_n}^K M$  to  $\mathbf{Z}_p[\text{Gal}(K_n/K)] \otimes M$  which sends  $a$  to  $\sum_{x \in \text{Gal}(K_n/K)} x^{-1}(ax) \delta_{x-1}$  is an isomorphism of  $G_K$ -modules. By Shapiro's lemma, we have an canonical isomorphism from  $H^i(K, \mathbf{Z}_p[\text{Gal}(K_n/K)] \otimes M)$  to  $H^i(K_n, M)$ . On the other hand, the corestriction map from  $H^i(K_{n+1}, M)$  to  $H^i(K_n, M)$  is derived from the previous isomorphism and the natural map form  $\mathbf{Z}_p[\text{Gal}(K_{n+1}/K)]$  to  $\mathbf{Z}_p[\text{Gal}(K_n/K)]$ . we thus deduce the natural map from  $H^i(K, \Lambda_K \otimes M)$  to

$$\varprojlim H^i(K, (\Lambda/\omega_n) \otimes M) = \varprojlim H^i(K_n, M).$$

It remains to show that this map is an isomorphism.

Surjectivity is obvious. To prove injectivity, it suffices to verify that the map from  $H^i(K, \Lambda_K \otimes M)$  to  $H^i(K, \Lambda/(\omega_n, p^n) \otimes M)$  is injective. Since  $\Lambda_K = \varprojlim \Lambda_K/(\omega_n, p^n)$ , it suffices to show that  $H^i(K, (\Lambda/\omega_n, p^n) \otimes M)$  satisfies the Mittag-Leffler condition (c.f. [NSK]), which is obvious since the group is finite.  $\square$

By lemma 4.7, the map  $\iota_{\psi, \gamma_n}$  identifies  $\frac{D(V)^{\psi=1}}{\gamma_n - 1}$  with a subgroup of  $H^1(K_n, V)$  if  $\Gamma_{K_n}$  is torsion free, we thus obtained a map  $h_{K_n, V}^1 : D(V)^{\psi=1} \rightarrow H^1(K_n, V)$ . Explicitly, if  $y \in D(V)^{\psi=1}$ , then  $(\varphi - 1)y \in D(V)^{\psi=0}$  and since  $\gamma_n - 1$  is invertible on  $D(V)^{\psi=0}$ , there exist  $x_n \in D(V)^{\psi=0}$  satisfies  $(\gamma_n - 1)x_n = (\varphi - 1)y$  (i.e.  $(x_n, y) \in Z_{\varphi, \gamma_n}^1(K_n, V)$ ). On the other hand, lemma 4.2 implies that the image  $\iota_{\psi, n}(y)$  and  $\log_p^0(\gamma_n) \iota_{\varphi, \gamma_n}(x_n, y)$  in  $H^1(K_n, V)$  does not depend on the choice of  $\gamma$ .

By lemma 5.3 below, we have  $\text{cor}_{K_{n+1}/K_n} \circ h_{K_{n+1}, V}^1 = h_{K_n, V}^1$ . On the other hand, if  $\Gamma_{K_n}$  is no longer torsion free, we define  $h_{K_n, V}^1$  by the relation  $\text{cor}_{K_{n+1}/K_n} \circ h_{K_{n+1}, V}^1 = h_{K_n, V}^1$ . By



this way, we associate every element in  $D(V)^{\psi=1}$  to a collection of Galois cohomology class  $h_{K_n, V}^1(y) \in H^1(K_n, V)$  for  $n \geq 1$ . The main result of this section is:

**Theorem 5.2.** (Fontaine) *Let  $V$  be a  $\mathbf{Z}_p$ -representation or a  $p$ -adic representation of  $G_K$ .*

- i) *If  $y \in D(V)^{\psi=1}$ , then  $(\dots, h_{K_n, V}^1(y), \dots) \in H_{\text{Iw}}^1(K, V)$ .*
- ii) *The map  $\text{Log}_{V^*(1)}^* : D(V)^{\psi=1} \rightarrow H_{\text{Iw}}^1(K, V)$  defined by above is an isomorphism.*

**5.2. Corestriction and  $(\varphi, \Gamma)$ -modules.** i) of theorem 5.2 is a consequence of the following lemma.

**Lemma 5.3.** *If  $n \geq 1$ , let*

$$T_{\gamma, n} : H^1(C_{\varphi, \gamma_n}(K_n, V)) \rightarrow H^1(C_{\varphi, \gamma_{n-1}}(K_{n-1}, V))$$

*the map induced by  $(x, y) \in Z^1(C_{\varphi, \gamma_n}(K_n, V))$  to  $(\frac{\gamma_n-1}{\gamma_{n-1}-1}x, y) \in Z^1(C_{\varphi, \gamma_{n-1}}(K_{n-1}, V))$ . Then the diagram*

$$\begin{array}{ccc} H^1(C_{\varphi, \gamma_n}(K_n, V)) & \xrightarrow{T_{\gamma, n}} & H^1(C_{\varphi, \gamma_{n-1}}(K_{n-1}, V)) \\ \downarrow \iota_{\varphi, \gamma_n} & & \downarrow \iota_{\varphi, \gamma_{n-1}} \\ H^1(K_n, V) & \xrightarrow{\text{cor}_{K_n/K_{n-1}}} & H^1(K_{n-1}, V) \end{array}$$

*is commutative.*

*Proof.* Recall that if  $G$  is a group,  $M$  is a  $G$ -module and  $H$  a subgroup of finite index of  $G$ , the corestriction map  $\text{cor} : H^1(H, N) \rightarrow H^1(G, M)$  can be written in the following way: let  $X \subset G$  is a system of representatives of  $G/H$  and, if  $g \in G$ , let  $\tau_g$  is the permutation of  $X$  defined by  $\tau_g(x)H = gxH$  if  $x \in X$ . If  $c \in H^1(H, M)$  and  $h \mapsto c_h$  is a cocycle which represents  $c$ , then

$$g \mapsto \sum_{x \in X} \tau_g(x)(c_{\tau_g(x)^{-1}gx})$$

is a cocycle of  $G$  with values in  $M$  whose class in  $H^1(G, M)$  does not depend on the choice of  $X$  and is equal to  $\text{cor}(c)$ .

If  $N$  is a  $G$ -submodule of  $M$  such that the image of  $c$  in  $H^1(H, N)$  is trivial (i.e. there exists  $b \in N$  such that we have  $c_h = (h-1)b$  for all  $h \in H$ ), then  $\text{cor}(c)$  is the class of the cocycle  $g \mapsto (g-1)(\sum_{x \in X} xb)$ .

In particular, we put  $G = G_{K_{n-1}}$ ,  $H = G_{K_n}$  and, if  $\tilde{\gamma}_{n-1}$  is a lift of  $\gamma_{n-1}$  in  $G_{K_{n-1}}$ , we take  $X = \{1, \tilde{\gamma}_{n-1}, \dots, \tilde{\gamma}_{n-1}^{p-1}\}$ . Take  $N = \text{Frac}(\mathbf{Z}_p[[G_{K_{n-1}}]]) \otimes_{\mathbf{Z}_p[[G_{K_{n-1}}]]} (A \otimes_{\mathbf{Z}_p} V)$ . If  $(x, y) \in Z^1(C_{\varphi, \gamma}(K_n, V))$  and if  $b \in \mathbf{A} \otimes T$ , the cocycle  $c_{x, y}$  is given by the formula  $c_{x, y}(\tau) = (\tau-1)c$ , where  $c = \frac{y}{\tilde{\gamma}_n-1} - b \in N$ . It follows that  $\text{cor}_{K_n/K_{n-1}}(\iota_{\varphi, \gamma_n}(x, y))$  is represented by the cocycle

$$\tau \mapsto (\sigma-1)\left(\sum_{i=0}^{p-1} \tilde{\gamma}_{n-1}^i c\right) = (\sigma-1)\left(\frac{y}{\tilde{\gamma}_{n-1}-1} - \sum_{i=0}^{p-1} \tilde{\gamma}_{n-1}^i b\right)$$

and since

$$(\varphi-1)\left(\sum_{i=0}^{p-1} \tilde{\gamma}_{n-1}^i b\right) = \sum_{i=0}^{p-1} \tilde{\gamma}_{n-1}^i ((\varphi-1)b) = \frac{\tilde{\gamma}_{n-1}^p - 1}{\tilde{\gamma}_{n-1} - 1} b = \frac{\gamma_n - 1}{\gamma_{n-1} - 1} x,$$

we see that this cocycle is just  $\iota_{\varphi, \gamma_{n-1}}(T_{\gamma, n}(x, y))$ , and the conclusion follows.  $\square$

**Remark 5.4.** One can also hide the explicit calculation by noting that, if  $n \geq 1$ , the diagram

$$\begin{array}{ccccccc} C_{\varphi, \gamma_n}(K_n, V) : 0 & \longrightarrow & D(V) & \longrightarrow & D(V) \oplus D(V) & \longrightarrow & D(V) \longrightarrow 0 \\ & & \downarrow \frac{\gamma_n-1}{\gamma_{n-1}-1} & & \downarrow (\frac{\gamma_n-1}{\gamma_{n-1}-1}, id) & & \downarrow id \\ C_{\varphi, \gamma_{n-1}}(K_n, V) : 0 & \longrightarrow & D(V) & \longrightarrow & D(V) \oplus D(V) & \longrightarrow & D(V) \longrightarrow 0 \end{array}$$

is commutative and functorial on  $V$  and induces a homomorphism of cohomology group from  $H^*(K_n, \cdot)$  to  $H^*(K_{n-1}, \cdot)$  which coincides with the corestriction map at  $* = 0$  and hence is corestriction map.

**5.3. Interpretation of  $D(V)^{\psi=1}$  and  $\frac{D(V)}{\psi-1}$  in Iwasawa theory.** We turn to prove ii) of theorem 5.2. The lemma 5.3 implies that the map  $(\iota_{\psi, \gamma_n})_{n \in \mathbf{N}}$  induces an isomorphism from the projective limit of  $H^1(C_{\psi, \gamma_n}(K_n, V))$  with respect to the map  $T_{\gamma, n}$  to  $H_{\text{Iw}}^1(K, V)$ . On the other hand, lemma 4.7, implies by passing to the projective limit, that we have an exact sequence:

$$0 \longrightarrow \varprojlim \frac{D(V)^{\psi=1}}{\gamma_n-1} \longrightarrow \varprojlim H^1(C_{\psi, \gamma_n}(K_n, V)) \longrightarrow \varprojlim (\frac{D(V)}{\psi-1})^{\Gamma_{K_n}}$$

The projective limit of  $\frac{D(V)^{\psi=1}}{\gamma_n-1}$  is by the natural maps induced by the identity on  $D(V)^{\psi=1}$  and that of  $(\frac{D(V)}{\psi-1})^{\gamma_n=1}$  with respect to the map

$$\frac{\gamma_{n+1}-1}{\gamma_n-1} : (\frac{D(V)}{\psi-1})^{\gamma_n=1} \rightarrow (\frac{D(V)}{\psi-1})^{\gamma_{n-1}=1}.$$

Hence ii) of theorem 5.2 is followed by the following proposition:

**Proposition 5.5.** *If  $V$  is a  $\mathbf{Z}_p$ -representation of  $G_K$ , then*

- i) *The natural map from  $D(V)^{\psi=1}$  to  $\varprojlim \frac{D(V)^{\psi=1}}{\gamma_n-1}$  is an isomorphism.*
- ii)  *$\varprojlim (\frac{D(V)}{\psi-1})^{\gamma_n=1} = 0$*

*Proof.* i) Let  $(x_n)_{n \in \mathbf{N}} \in \varprojlim \frac{D(V)^{\psi=1}}{\gamma_n-1}$ . The compactness of  $D(V)^{\psi=1}$  [c.f. proposition 4.9 i)] implies that the sequence  $x_n$  admits a accumulation points  $x \in D(V)^{\psi=1}$  and the image of  $x \in \varprojlim \frac{D(V)^{\psi=1}}{\gamma_n-1}$  is by construction  $(x_n)_{n \in \mathbf{N}}$ . The natural map from  $D(V)^{\psi=1}$  to  $\varprojlim \frac{D(V)^{\psi=1}}{\gamma_n-1}$  is hence surjective.

By the compactness of  $D(V)^{\psi=1}$  and the fact that if  $x \in D(V)$ , then  $(\gamma_n - 1)x$  tend to 0 when  $n$  tend to  $+\infty$  implies that if  $U$  is open in  $D(V)$  fixed by  $\Gamma$ , then there exist  $n_U \in \mathbf{N}$  such that  $(\gamma_n - 1)D(V)^{\psi=1} \subset U$  if  $n \geq n_U$ . This implies that  $\bigcap_{n \in \mathbf{N}} (\gamma_n - 1)(D(V)^{\psi=1}) = \{0\}$  and we prove the injectivity.

ii)  $\frac{D(V)}{\psi-1}$  is a free  $\mathbf{Z}_p$ -module of finite rank [c.f. proposition 4.9 ii)], the sequence  $(\frac{D(V)}{\psi-1})^{\gamma_n=1}$  is stationary since it is increasing. One can deduce the fact that there exists  $n_0 \in \mathbf{N}$  such that  $\frac{\gamma_n-1}{\gamma_{n-1}-1}$  is multiplication by  $p$  on  $(\frac{D(V)}{\psi-1})^{\gamma_n=1}$  if  $n \geq n_0$ , which proves the statement since  $\frac{D(V)}{\psi-1}$  has no  $p$ -divisible element [c.f. lemma 4.15].  $\square$

**Remark 5.6.** We have  $H^2(K_n, V) \cong H^2(C_{\psi, \gamma_n}(K_n, V)) = \frac{D(V)}{(\psi-1, \gamma_n-1)}$ . We deduce that if  $V$  is a  $\mathbf{Z}_p$ -representation, then  $H_{\text{Iw}}^2(K, V)$  is a projective limit of  $\frac{D(V)}{(\psi-1, \gamma_n-1)}$  since  $\frac{D(V)}{\psi-1}$  is a  $\mathbf{Z}_p$ -module of finite type which  $\Gamma_K$  acts continuously by ii) of lemma 4.17, the natural map from  $\frac{D(V)}{\psi-1}$  to

the projective limit of  $\frac{D(V)}{(\psi-1, \gamma_n-1)}$  is an isomorphism, this proves that  $\frac{D(V)}{\psi-1}$  is identified with  $H_{\text{Iw}}^2(K, V)$ .

By above proposition, one can summarize the the above results as follows:

**Corollary 5.7.** *The complex of  $\mathbf{Q}_p \otimes_{\mathbf{Z}_p} \Lambda_K$ -modules*

$$0 \longrightarrow D(V) \xrightarrow{1-\psi} D(V) \longrightarrow 0$$

*computes the Iwasawa cohomology of  $V$ .*

There is a natural projection map  $\text{pr}_{K_n, V} : H_{\text{Iw}}^i(K, V) \rightarrow H^i(K_n, V)$  and when  $i = 1$  it is of course equal to the composition of:

$$H_{\text{Iw}}^1(K, V) \longrightarrow D(V)^{\psi=1} \xrightarrow{h_{K_n, V}^1} H^1(K_n, V).$$

The  $H_{\text{Iw}}^i(K, V)$  have been studied in detail by Perrin-Riou, who proved the following

**Proposition 5.8.** *If  $V$  is a  $p$ -adic representation of  $G_K$ , then*

- i) *The torsion submodule of  $H_{\text{Iw}}^1(K, V)$  is a  $\mathbf{Q}_p \otimes_{\mathbf{Z}_p} \Lambda_K$ -module isomorphic to  $V^{H_K}$  and  $H_{\text{Iw}}^1(K, V)/V^{H_K}$  is a free  $\mathbf{Q}_p \otimes_{\mathbf{Z}_p} \Lambda_K$ -module whose rank is  $[K : \mathbf{Q}_p] \dim_{\mathbf{Q}_p} V$ .*
- ii)  *$H_{\text{Iw}}^2(K, V)$  is isomorphic to  $V(-1)^{H_K}$  as  $\mathbf{Q}_p \otimes_{\mathbf{Z}_p} \Lambda_K$ -module. In particular, it is torsion.*
- iii)  *$H_{\text{Iw}}^i(K, V) = 0$  when  $i \neq 1, 2$ .*

*Proof.* See [Per94, 3.2.1]. □

## 6. DE RHAM REPRESENTATIONS AND OVERCONVERGENT REPRESENTATIONS

**6.1. De Rham representations and crystalline representations.** Recall  $\tilde{\mathbf{A}}^+ = W(\tilde{\mathbf{E}}^+)$  the ring of Witt vectors with coefficients in  $\tilde{\mathbf{E}}^+$  and if  $x \in \tilde{\mathbf{E}}^+$ . We define the homomorphism  $\theta : \tilde{\mathbf{A}}^+ \rightarrow \mathcal{O}_{\mathbf{C}_p}$  by

$$\theta\left(\sum_{k \geq 0} p^k [x_k]\right) = \sum_{k \geq 0} p^k x_k^{(0)}$$

One can show that this is a surjective map and  $\ker(\theta : \tilde{\mathbf{A}}^+ \rightarrow \mathcal{O}_{\mathbf{C}_p})$  is generated by  $\omega = \pi/\varphi^{-1}(\pi)$ .

We can extend  $\theta$  to a homomorphism from  $\tilde{\mathbf{B}}^+ = \tilde{\mathbf{A}}^+[\frac{1}{p}]$  to  $\mathbf{C}_p$ , and we denote  $\mathbf{B}_{\text{dR}}^+$  the ring  $\varprojlim \tilde{\mathbf{B}}^+ / (\ker \theta)^n$ , thus  $\theta$  can be extended by continuity to a homomorphism from  $\mathbf{B}_{\text{dR}}^+$  to  $\mathbf{C}_p$ . This makes  $\mathbf{B}_{\text{dR}}^+$  a discrete valuation ring with maximal ideal  $\ker \theta$  and residue field  $\mathbf{C}_p$ . The action of  $G_{\mathbf{Q}_p}$  on  $\tilde{\mathbf{A}}^+$  extend by continuity to an action of  $G_{\mathbf{Q}_p}$  on  $\mathbf{B}_{\text{dR}}^+$ . The series  $\log[\varepsilon] = \sum_{n=1}^{+\infty} \frac{(-1)^{n-1}}{n} \pi^n$  converges in  $\mathbf{B}_{\text{dR}}^+$  to an element which we denote it by  $t$ , which is a generator of  $\ker \theta$  with an  $G_{\mathbf{Q}_p}$  action defined by  $\sigma(t) = \chi(\sigma)t$  where  $\sigma \in G_{\mathbf{Q}_p}$ . This element can be viewed as  $p$ -adic analogy of  $2\pi i$ .

We put  $\mathbf{B}_{\text{dR}} = \mathbf{B}_{\text{dR}}^+[t^{-1}]$ , this makes  $\mathbf{B}_{\text{dR}}$  a field with filtration defined by  $\text{Fil}^i \mathbf{B}_{\text{dR}} = t^i \mathbf{B}_{\text{dR}}^+$ . This filtration is stable by the action of  $G_K$ .

Let  $K$  is a finite extension of  $\mathbf{Q}_p$  and  $V$  be a  $p$ -adic representation of  $G_K$ . We say  $V$  is de Rham if the  $\mathbf{B}_{\text{dR}}$ -admissible, which is equivalent to the  $K$ -vector space  $\mathbf{D}_{\text{dR}}(V) = (\mathbf{B}_{\text{dR}} \otimes_{\mathbf{Q}_p} V)^{G_K}$  is of dimension  $d = \dim_{\mathbf{Q}_p}(V)$ . On the other hand,  $\mathbf{D}_{\text{dR}}(V)$  is endowed with a filtration induced by  $\mathbf{B}_{\text{dR}}$ . We have  $\text{Fil}^i \mathbf{D}_{\text{dR}}(V) = \mathbf{D}_{\text{dR}}(V)$  if  $i \ll 0$  and  $\text{Fil}^i \mathbf{D}_{\text{dR}}(V) = \{0\}$  if  $i \gg 0$ .

The ring  $\mathbf{B}_{\text{cris}}^+$  is defined by

$$\mathbf{B}_{\text{cris}}^+ = \left\{ \sum_{n \geq 0} a_n \frac{\omega^n}{n!} \mid a_n \in \tilde{\mathbf{B}}^+ \text{ is a sequence converges to } 0 \right\},$$

and  $\mathbf{B}_{\text{cris}}^+[\frac{1}{t}]$ . The ring  $\mathbf{B}_{\text{cris}}$  is a subring of  $\mathbf{B}_{\text{dR}}$  stable under  $G_{\mathbf{Q}_p}$  containing  $t$  and the action of  $\varphi$  on  $\tilde{\mathbf{B}}^+$  is extended by continuity to an action of  $\mathbf{B}_{\text{cris}}^+$ . In particular, we have  $\varphi(t) = pt$ .

We say  $V$  is crystalline if it is  $\mathbf{B}_{\text{cris}}$ -admissible, which is equivalent to the  $F = K \cap \mathbf{Q}_p^{ur}$ -vector space  $\mathbf{D}_{\text{cris}}(V) = (\mathbf{B}_{\text{cris}} \otimes V)^{G_K}$  is of dimension  $d = \dim_{\mathbf{Q}_p}(V)$ . The action of  $\varphi$  on  $\mathbf{B}_{\text{cris}}$  commutes with the action of  $G_{\mathbf{Q}_p}$ , which endows  $\mathbf{D}_{\text{cris}}$  a natural semi-linear action of  $\varphi$ .

We have  $(\mathbf{B}_{\text{dR}} \otimes_{\mathbf{Q}_p} V)^{G_K} = \mathbf{D}_{\text{dR}}(V) = K \otimes_F \mathbf{D}_{\text{cris}}(V)$ , thus the crystalline representation is de Rham and  $K \otimes_F \mathbf{D}_{\text{cris}}(V)$  is a filtered  $K$ -vector space. Hence if  $V$  is de Rham (resp. crystalline) and  $k \in \mathbf{Z}$ , so is  $V(k)$ , and we have  $\mathbf{D}_{\text{dR}}(V(k)) = t^{-k} \mathbf{D}_{\text{dR}}(V)$  (resp.  $\mathbf{D}_{\text{cris}}(V(k)) = t^{-k} \mathbf{D}_{\text{cris}}(V)$ ).

**6.2. Overconvergent elements.** Every element  $x$  of  $\tilde{\mathbf{B}}$  can be written uniquely as the form  $\sum_{k \gg -\infty} p^k [x_k]$ , where  $x_k$  is element of  $\tilde{\mathbf{E}}$  and the series converges in  $\mathbf{B}_{\text{dR}}^+$  if and only if the series  $\sum_{k \gg -\infty} p^k x_k^{(0)}$  converges in  $\mathbf{C}_p$ , which is equivalent to  $k + \nu_E(x_k)$  tends to  $+\infty$  as  $k$  tends to  $+\infty$ . More generally, if  $n \in \mathbf{N}$ ,  $\varphi^{-n}(x)$  converges if and only if  $k + p^{-n} \nu_E(x_k)$  tends to  $+\infty$  as  $k$  tends to  $+\infty$ .

For  $r \geq 0$ , we set

$$\tilde{\mathbf{B}}^{\dagger, r} = \{x \in \tilde{\mathbf{B}} \mid \lim_{k \rightarrow +\infty} \nu_E(x_k) + \frac{pr}{p-1}k = +\infty\}.$$

This makes  $\tilde{\mathbf{B}}^{\dagger, r}$  into an intermediate ring between  $\tilde{\mathbf{B}}^+$  and  $\tilde{\mathbf{B}}$ . We denote  $\tilde{\mathbf{B}}^{\dagger} = \cup_{r \geq 0} \tilde{\mathbf{B}}^{\dagger, r}$ , which is a subfield of  $\tilde{\mathbf{B}}$  with action of  $G_K$  and  $\varphi$ . On the other hand, we have a well-defined injection map  $\varphi^{-n} : \tilde{\mathbf{B}}^{\dagger, r_n} \rightarrow \mathbf{B}_{\text{dR}}^+$ , where  $r_n = p^{n-1}(p-1)$ .

We denote  $\tilde{\mathbf{A}}^{\dagger, r} = \tilde{\mathbf{B}}^{\dagger, r} \cap \tilde{\mathbf{A}}$ , that is, the subring of elements  $x = \sum_{k=0}^{+\infty} p^k [x_k]$  of  $\tilde{\mathbf{A}}$  such that  $\nu_E(x_k) + \frac{pr}{p-1}k$  tends to  $+\infty$  as  $k$  tends to  $+\infty$ . We have  $\tilde{\mathbf{B}}^{\dagger, n} = \tilde{\mathbf{A}}^{\dagger, n}[\frac{1}{p}]$ .

By putting  $\mathbf{B}^{\dagger} = \mathbf{B} \cap \tilde{\mathbf{B}}^{\dagger}$ ,  $\mathbf{A}^{\dagger, n} = \mathbf{A} \cap \tilde{\mathbf{A}}^{\dagger, r}$  and  $\mathbf{B}^{\dagger, r} = \mathbf{B} \cap \tilde{\mathbf{B}}^{\dagger, r}$ , we define a subring  $\mathbf{B}^{\dagger}$  of  $\mathbf{B}$  fixed by  $\varphi$  and  $G_{\mathbf{Q}_p}$ , and if  $n \in \mathbf{N}$ , subrings  $\mathbf{A}^{\dagger, r}$  and  $\mathbf{B}^{\dagger, r}$  of  $\mathbf{B}$  are fixed by  $G_{\mathbf{Q}_p}$ . By construction,  $\varphi^{-n}(\mathbf{B}^{\dagger, r_n})$  is naturally identified with a subring of  $\mathbf{B}_{\text{dR}}^+$ . Finally, if  $K$  is a finite extension of  $\mathbf{Q}_p$ , we set  $\mathbf{B}_K^{\dagger} = (\mathbf{B}^{\dagger})^{H_K}$ ,  $\mathbf{A}_K^{\dagger, r} = (\mathbf{A}^{\dagger, r})^{H_K}$  and  $\mathbf{B}_K^{\dagger, r} = (\mathbf{B}^{\dagger, r})^{H_K}$ .

Let  $e_K$  the ramification index of  $K_{\infty}$  over  $F_{\infty}$  and  $F' \subset K_{\infty}$  be the maximal unramified extension of  $\mathbf{Q}_p$  contained in  $K_{\infty}$ . Let  $\bar{\pi}_K$  be a uniformizer of  $\mathbf{E}_K = k_{F'}((\bar{\pi}_K))$  and  $\bar{P}_K \in E_{F'}$  be a minimal polynomial of  $\bar{\pi}_K$  and  $\delta = \nu_E(\bar{P}'(\bar{\pi}_K))$ . Choose  $P_K \in \mathbf{A}_{F'}$  such that it modulo  $p$  is  $\bar{P}_K$ . By Hensel's lemma, there exists a unique  $\pi_K \in \mathbf{A}_K$  such that  $P_K(\pi_K) = 0$  and  $\pi_K = \bar{\pi}_K$  modulo  $p$ . In particular, if  $K = F'$ , one can take  $\pi_K = \pi$ .

The terminology "overconvergent" can be explained by the following proposition:

**Proposition 6.1.** *If  $r \geq r(K)$ , then the map  $f \mapsto f(\pi_K)$  from  $\mathcal{B}_{F'}^{e_K r}$  to  $\mathbf{B}_K^{\dagger, r}$  is an isomorphism, where  $\mathcal{B}_{F'}^{\alpha}$  is the set of power series  $f(T) = \sum_{k \in \mathbf{Z}} a_k T^k$  such that  $a_k$  is a bounded sequence of elements of  $F'$ , and such that  $f(T)$  is holomorphic on the  $p$ -adic annulus  $\{p^{-1/\alpha} \leq \|T\| < 1\}$ .*

*Proof.* See lemma II.2.2 [CC98]. □

**Proposition 6.2.** *If  $K$  is a finite extension of  $\mathbf{Q}_p$ , then  $\mathbf{B}_K^{\dagger}$  is an extension of  $\mathbf{B}_{\mathbf{Q}_p}^{\dagger}$  of degree  $[\mathbf{B}_K : \mathbf{B}_{\mathbf{Q}_p}] = [K_{\infty} : \mathbf{Q}_p(\mu_{p^{\infty}})]$  and there exists  $a(K) \in \mathbf{N}$  such that if  $n \geq a(K)$ , then  $\varphi^{-n}(\mathbf{B}_K^{\dagger, r_n}) \subset K_n[[t]]$ , where  $r_n = p^{n-1}(p-1)$ .*

*Proof.* In the case  $K$  is unramified over  $\mathbf{Q}_p$ , one can follow proposition 6.1 i) using the fact that  $K_n[[t]]$  is closed in  $\mathbf{B}_{\text{dR}}^+$  and the formula

$$\varphi^{-n}(\pi) = \varphi^{-n}([\varepsilon] - 1) = [\varepsilon^{p^{-n}}] - 1 = \varepsilon^{(n)} \exp(t/p^n) - 1 \in K_n[[t]].$$

For general case, by remark 3.2, there exists  $\omega = (\omega^{(n)})_{n \in \mathbf{N}} \in \varprojlim \mathcal{O}_{K_n}$  such that  $\omega^{(n)}$  is a uniformizer of  $\mathcal{O}_{K_n}$  if  $n$  large enough and then  $\bar{\pi}_K = \iota_K(\omega)$  is a uniformizer of  $\mathbf{E}_K$  such that it is totally ramified of degree  $e_K$  over  $\mathbf{E}_{F'}$ . Let  $\bar{P}(X) = X^{e_K} + \bar{a}_{e_K-1}X^{e_K-1} + \dots + \bar{a}_0 \in \mathbf{E}_{F'}[X]$  be the minimal polynomial of  $\bar{\pi}_K$  over  $\mathbf{E}_{F'}$  and let  $\delta = \nu_E(\bar{P}'(\bar{\pi}_K))$ . If  $0 \leq i \leq e_K - 1$ , let  $a_i \in \mathcal{O}_F[[t]] \subset \mathbf{A}_F$  whose reduction modulo  $p$  is  $\bar{a}_i$  and let  $P(X) = X^{e_K} + a_{e_K-1}X^{e_K-1} + \dots + a_0 \in \mathbf{A}_F[X]$ . By Hensel's lemma, the equation  $P(X) = 0$  has a unique solution  $\pi_K$  in  $A_K$  whose reduction modulo  $p$  is  $\bar{\pi}_K$  and we can write it in the form

$$(1) \quad \pi_K = [\bar{\pi}_K] + \sum_{i=1}^{+\infty} p^i [\alpha_i],$$

where  $\alpha_i$  are elements of  $\tilde{\mathbf{E}}$  verify  $\nu_E(\alpha_i) \geq -i\delta$ . In particular,  $\pi_K \in \mathbf{A}_K^{\dagger,r}$  if  $\frac{p}{p-1}r \geq \delta$ , hence we have  $\mathbf{A}_K^{\dagger,r} = \mathbf{A}_F^{\dagger,r}[\pi_K]$  if  $\frac{p}{p-1}r \geq \delta$ . Thus it suffices to prove it when  $n$  large enough, then  $\pi_{K,n} = \varphi^{-n}(\pi_K) \in K_n[[t]]$ .

Let  $P_n$  (resp.  $Q_n$ ) be polynomial obtained by the map  $\theta \circ \varphi^{-n}$  (resp.  $\varphi^{-n}$ ) apply on the coefficients of  $P$ , which is a polynomial with coefficients in  $\mathcal{O}_{F_n}$  (resp.  $F_n[[t]]$ ) with  $\theta(\pi_{K,n})$  (resp.  $\pi_{K,n}$ ) as a root. On the other hand, by definition of  $\iota_K$  (c.f. 3.2), we have  $\nu_p(\omega^{(n)} - \bar{\pi}_K^{(n)}) \geq \frac{1}{p}$  if  $n$  large enough and formula (1) shows that  $\nu_p(\theta(\pi_{K,n}) - \bar{\pi}_K^{(n)}) \geq (1 - \frac{\delta}{p^n})$ . Then we have  $\nu_p(P_n(\omega^{(n)})) \geq \frac{1}{p}$  if  $n$  large enough and

$$\nu_p(P'_n(\omega^{(n)})) = \frac{1}{p^n} \nu_E(P'(\bar{\pi}_K)) = \frac{\delta}{p^n} < \frac{1}{2p}$$

if  $n$  large enough. By Hensel's lemma, the equation  $P_n(X) = 0$  has a unique solution in  $\mathbf{C}_p$  close to  $\omega^{(n)}$  and hence belongs to  $\mathcal{O}_{K_n}$  since  $\omega^{(n)}$  and the coefficients of  $P_n$  do. We deduce that  $\theta(\pi_{K,n})$  belongs to  $K_n$ . By using the Hensel's lemma again, one can show that  $Q_n$  has a unique solution in  $\mathbf{B}_{\text{dR}}^+$  whose image by  $\theta$  is  $\theta(\pi_{K,n})$  and thus belongs to  $K_n[[t]]$ .  $\square$

We endow  $\mathbf{B}_{Q_p}$  the differential operator  $\partial$  defined by continuity and the derivation  $\partial\pi = 1 + \pi$ . We therefore have  $\partial = [\varepsilon] \frac{d}{d\pi} = \frac{d}{dt}$  (Note that  $t \notin \mathbf{B}_{Q_p}$ ). The derivation can be extended uniquely to a maximal unramified extension of  $\mathbf{B}_{Q_p}$  in  $\tilde{\mathbf{B}}$ , hence by continuity to a derivation  $\partial$  from  $\mathbf{B}$  to  $\mathbf{B}$ .

**Lemma 6.3.** *If  $K$  is a finite extension of  $\mathbf{Q}_p$ , there exists  $m(K) \in \mathbf{Z}$  such that, if  $n \geq m(K)$  and  $x$  in  $\mathbf{B}_K^{\dagger,rn}$ , then*

- i)  $\partial x \in \mathbf{B}_K^{\dagger,rn}$ .
- ii)  $\varphi^{-n}(\partial x) = p^n \partial(\varphi^{-n}(x))$ .

*Proof.* If  $K = \mathbf{Q}_p$ , explicit calculation using proposition 6.1 i), shows that we can take  $m(K) = 1$ . For the general case, let  $\alpha$  be a generator of  $\mathbf{B}_K^\dagger$  over  $\mathbf{B}_{\mathbf{Q}_p}^\dagger$  and  $P$  be its minimal polynomial. The identity,

$$0 = \partial(P(\alpha)) = P'(\alpha)\partial\alpha + \partial P(\alpha),$$

where  $\partial P$  is the polynomial obtained by applying  $\partial$  on the coefficients of  $P$ , shows that  $\partial\alpha = -\frac{\partial P(\alpha)}{P'(\alpha)} \in \mathbf{B}_K^\dagger$ . It is then possible to take  $m(K)$  any integer such that  $\mathbf{B}_K^{\dagger, m(K)}$  contains  $\partial\alpha$  and  $\alpha$ .

For ii), it suffices to note that  $\varphi^{-n} \circ \partial$  is  $p^n \partial \circ \varphi^{-n}$  are two derivations of  $\mathbf{B}_K^{\dagger, r_n}$  coincides on  $\mathbf{B}_{\mathbf{Q}_p}^{\dagger, r_n}$  by

$$\begin{aligned}\varphi^{-n} \circ \partial([\varepsilon]) &= \varphi^{-n}([\varepsilon]) = \varepsilon^{(n)} \exp(p^{-n}t) \\ p^n \partial \circ \varphi^{-n}([\varepsilon]) &= p^n \frac{d}{dt}(\varepsilon^{(n)} \exp(p^{-n}t)) = \varepsilon^{(n)} \exp(p^{-n}t).\end{aligned}$$

□

### 6.3. Overconvergent representations.

**Definition 6.4.** If  $V$  is a  $p$ -adic representation of  $G_K$ , we set

$$\mathbf{D}^\dagger(V) = (\mathbf{B}^\dagger \otimes_{\mathbf{Q}_p} V)^{H_K} \quad \text{and} \quad \mathbf{D}^{\dagger, r}(V) = (\mathbf{B}^{\dagger, r} \otimes_{\mathbf{Q}_p} V)^{H_K}$$

We have  $\dim_{\mathbf{B}_K^\dagger} \mathbf{D}^\dagger(V) \leq \dim_{\mathbf{Q}_p} V$  and we say that  $V$  is overconvergent if the equality holds, which is equivalent to  $D(V)$  has a basis over  $\mathbf{B}_K$  made up of elements of  $\mathbf{D}^\dagger(V)$ .

**Proposition 6.5.**

- i) Every  $p$ -adic representations of  $G_K$  is overconvergent.
- ii) There exists  $r(V)$  such that  $D(V)^{\psi=1} \subset \mathbf{D}^{\dagger, r(V)}(V)$ .
- iii) If  $V$  is overconvergent and  $n \in \mathbf{N}$ , then  $\gamma_n - 1$  admits a continuous inverse on  $\mathbf{D}^\dagger(V)^{\psi=0}$ . Moreover, there exists  $n_2(V)$  such that if  $n \geq n_2(V)$ , then

$$(\gamma_n - 1)^{-1}(\mathbf{D}^{\dagger, r_n}(V)^{\psi=0}) \subset \mathbf{D}^{\dagger, r_{n+1}}(V)^{\psi=0}$$

*Proof.* i), iii) see [CC98]. ii) follows from lemma 4.14. □

## 7. EXPLICIT RECIPROCITY LAWS AND DE RHAM REPRESENTATION

**7.1. The Bloch-Kato exponential map and its dual.** Let  $K$  be a finite extension of  $\mathbf{Q}_p$  and  $V$  a  $p$ -adic representation of  $G_K$ . We have fundamental exact sequence

$$0 \longrightarrow \mathbf{Q}_p \longrightarrow \mathbf{B}_{\text{cris}}^{\varphi=1} \longrightarrow \mathbf{B}_{\text{dR}}/\mathbf{B}_{\text{dR}}^+ \longrightarrow 0$$

(c.f. [Col98, proposition III 3.5]). Tensoring this exact sequence with  $V$  and taking the invariant under the action of  $G_K$ , we obtain:

$$0 \longrightarrow V^{G_K} \longrightarrow \mathbf{D}_{\text{cris}}(V)^{\varphi=1} \longrightarrow ((\mathbf{B}_{\text{dR}}/\mathbf{B}_{\text{dR}}^+) \otimes V)^{G_K} \longrightarrow H_e^1(K, V) \longrightarrow 0$$

where we denote  $H_e^1(K, V)$  the kernel of the natural map from  $H^1(K, V)$  to  $H^1(K, \mathbf{B}_{\text{cris}}^{\varphi=1} \otimes V)$ . We denote the isomorphism induced by connecting homomorphism

$$\exp_{K, V} : \frac{\mathbf{D}_{\text{dR}}(V)}{\text{Fil}^0 \mathbf{D}_{\text{dR}}(V) + \mathbf{D}_{\text{cris}}(V)^{\varphi=1}} \longrightarrow H_e^1(K, V) \subset H^1(K, V)$$

the Bloch-Kato exponential of  $V$  over  $K$  and we denote its inverse by

$$\log_{K, V} : H_e^1(K, V) \longrightarrow \frac{\mathbf{D}_{\text{dR}}(V)}{\text{Fil}^0 \mathbf{D}_{\text{dR}}(V) + \mathbf{D}_{\text{cris}}(V)^{\varphi=1}}$$

the Bloch-Kato logarithm of  $V$  over  $K$ . Moreover, if  $V$  is de Rham and  $k \gg 0$ , then  $\exp_{K, V(k)}$  is an isomorphism from  $\mathbf{D}_{\text{dR}}(V(k))$  to  $H^1(K, V(k))$ .

The choice of  $t$  gives an isomorphism from  $\mathbf{D}_{\mathrm{dR}}(\mathbf{Q}_p(1)) = t^{-1}K$  to  $K$ . If  $V$  is a  $p$ -representation of  $G_K$ , the couple  $[\cdot, \cdot]_{\mathbf{D}_{\mathrm{dR}}(V)}$  is defined by composing the maps

$$\mathbf{D}_{\mathrm{dR}}(V) \otimes \mathbf{D}_{\mathrm{dR}}(V^*(1)) \cong \mathbf{D}_{\mathrm{dR}}(V \otimes V^*(1)) \longrightarrow \mathbf{D}_{\mathrm{dR}}(\mathbf{Q}_p(1)) \cong K \xrightarrow{\mathrm{Tr}_{K/\mathbf{Q}_p}} \mathbf{Q}_p$$

is non-degenerate, hence  $\mathbf{D}_{\mathrm{dR}}(V^*(1))$  can be naturally identified with the dual of  $\mathbf{D}_{\mathrm{dR}}(V)$ . Similarly, via the cup product

$$H^1(K, V) \times H^1(K, V^*(1)) \rightarrow H^2(K, \mathbf{Q}_p(1)) = \mathbf{Q}_p,$$

$H^1(K, V^*(1))$  is naturally identified with the dual of  $H^1(K, V)$ . This allows us to view the map  $\exp_{K, V^*(1)}^*$  as transpose of the map  $\exp_{K, V^*(1)} : \mathbf{D}_{\mathrm{dR}}(V^*(1)) \rightarrow H^1(K, V^*(1))$  as a map from  $H^1(K, V)$  to  $\mathbf{D}_{\mathrm{dR}}(V)$ , whose image is contained in  $\mathrm{Fil}^0(\mathbf{D}_{\mathrm{dR}}(V))$ . If  $V$  is de Rham and  $k \gg 0$ , the map  $\exp_{K, V^*(1+k)}^*$  is an isomorphism from  $H^1(K, V(-k))$  to  $\mathbf{D}_{\mathrm{dR}}(V(-k))$ .

If  $x \in K_\infty$  and  $n \in \mathbf{N}$ , then  $\frac{1}{p^n} \mathrm{Tr}_{K_m/K_n}(x)$  does not depend on the choice of integer  $m \geq n+1$  such that  $x$  belongs to  $K_m$ . We denote  $T_n$  the above  $\mathbf{Q}_p$ -linear map from  $K_\infty$  to  $K_n$ . If  $n \geq 1$  and  $x \in K_n$ , then  $T_n(x) = p^{-n}x$ . We have

$$T_m = \mathrm{Tr}_{K_n/K_m} \circ T_n \quad \text{if } n \geq m.$$

We also denote  $T_n$  the map from  $K_\infty((t))$  to  $K_n((t))$  defined by  $T_n(\sum_{k=0}^{+\infty} a_k t^k) = \sum_{k=0}^{+\infty} T_n(a_k) t^k$ .

**Proposition 7.1.**

- i)  $K_\infty((t))$  is dense in  $\mathbf{B}_{\mathrm{dR}}^{H_K}$  and  $T_n$  can be extended to a  $\mathbf{Q}_p$ -linear map from  $\mathbf{B}_{\mathrm{dR}}^{H_K}$  to  $K_n((t))$ .
- ii) If  $F \in \mathbf{B}_{\mathrm{dR}}^{H_K}$ , then  $\lim_{n \rightarrow +\infty} p^n T_n(F) = F$ .

*Proof.* See [Col98], proposition V.4.5. □

Let  $V$  be a de Rham representation of  $G_K$ , we have  $\mathbf{B}_{\mathrm{dR}} \otimes_{\mathbf{Q}_p} V \cong \mathbf{B}_{\mathrm{dR}} \otimes_K \mathbf{D}_{\mathrm{dR}}(V)$  and  $H^1(K, \mathbf{B}_{\mathrm{dR}} \otimes V) = H^1(K, \mathbf{B}_{\mathrm{dR}} \otimes \mathbf{D}_{\mathrm{dR}}(V)) = H^1(K, \mathbf{B}_{\mathrm{dR}}) \otimes \mathbf{D}_{\mathrm{dR}}(V)$ . Since  $K \cong H^1(K, \mathbf{B}_{\mathrm{dR}})$  via  $x \mapsto x \cup \log \chi$ , we get an isomorphism

$$\mathbf{D}_{\mathrm{dR}}(V) \rightarrow H^1(K, \mathbf{B}_{\mathrm{dR}} \otimes V); \quad x \mapsto x \cup \log \chi$$

**Proposition 7.2.** *If  $V$  is a de Rham representation, the map sends  $x \in \mathbf{D}_{\mathrm{dR}}(V)$  to a cocycle  $\tau \mapsto x \log \chi(\tau) \in \mathbf{D}_{\mathrm{dR}}(V) \subset \mathbf{B}_{\mathrm{dR}} \otimes V$  induces an isomorphism from  $\mathbf{D}_{\mathrm{dR}}(V)$  to  $H^1(K, \mathbf{B}_{\mathrm{dR}} \otimes V)$  and the map  $\exp_{V^*(1)}^*$  is the composition of the inverse of the above isomorphism and the natural map from  $H^1(K, V)$  to  $H^1(K, \mathbf{B}_{\mathrm{dR}} \otimes V)$ .*

*Proof.* See [Kato93] proposition 1.4. of chapter II. □

We define the map  $\mathrm{pr}_{K_n} : \mathbf{B}_{\mathrm{dR}}^{H_K} \rightarrow K_n((t))$  by the formula  $\mathrm{pr}_{K_n}(x) = \frac{1}{[K_m:K_n]} \mathrm{Tr}_{K_m/K_n}(x)$  if  $x \in K_\infty$  and  $m \geq n$  such that  $x \in K_m$  and there exists  $a'(K) \geq 1$  such that one has  $p^n T_n = \mathrm{pr}_{K_n}$  if  $n \geq a'(K)$ . From (ii) of proposition 7.1, we can show that  $\lim_{n \rightarrow +\infty} \mathrm{pr}_{K_n} x = x$  if  $x \in \mathbf{B}_{\mathrm{dR}}^{H_K}$ .

If  $V$  is a de Rham representation, the natural map from  $\mathbf{B}_{\mathrm{dR}}^{H_K} \otimes_K \mathbf{D}_{\mathrm{dR}}(V)$  to  $(\mathbf{B}_{\mathrm{dR}} \otimes_{\mathbf{Q}_p} V)^{H_K}$  is an isomorphism and we can extend the map  $T_n$  and  $\mathrm{pr}_{K_n}$  for  $n \in \mathbf{N}$  by linearity to  $\mathbf{B}_{\mathrm{dR}}^{H_K} \otimes_K \mathbf{D}_{\mathrm{dR}}(V)$ . On the other hand, if  $F \in K_\infty((t)) \otimes \mathbf{D}_{\mathrm{dR}}(V)$ , we can write  $F$  uniquely as the form  $\sum_{k \gg -\infty} t^k d_k$ , where  $d_k \in K_\infty \otimes \mathbf{D}_{\mathrm{dR}}(V)$ . We denote  $\partial_{V(-k)}(F)$  the element  $t^k d_k$  of  $K_\infty \otimes \mathbf{D}_{\mathrm{dR}}(V(-k))$ .

**Proposition 7.3.** *Let  $V$  is a  $p$ -adic representation of  $G_K$  and  $n, m \in \mathbf{N}$  be two integers. If  $c \in H^1(K_m, V(-k))$ , there exists cocycle  $\tau \mapsto c_\tau$  on  $\Gamma_{K_m}$  with values in  $(\mathbf{B}_{\text{dR}} \otimes V(-k))^{H_K}$  which has the same image as  $c$  in  $H^1(K_m, \mathbf{B}_{\text{dR}} \otimes V(-k))$ . Moreover, if  $V$  is de Rham. then*

$$\exp_{V^*(1+k)}^*(c) = \partial_{V(-k)} \circ \text{pr}_{K_m} \left( \frac{1}{\log_p(\chi(\gamma))} c_\gamma \right)$$

for all  $\gamma \in \Gamma_{K_m}$  such that  $\log_p(\chi(\gamma)) \neq 0$

*Proof.* Since  $H^1(K_\infty, \mathbf{B}_{\text{dR}} \otimes V)$  is zero (c.f. [Col98] theorem IV.3.1), the inflation map from  $H^1(\Gamma_{K_m}, (\mathbf{B}_{\text{dR}} \otimes V)^{H_K})$  to  $H^1(K_m, \mathbf{B}_{\text{dR}} \otimes V)$  is an isomorphism, hence we have the existence of cocycle  $\tau \mapsto c_\tau$ . On the other hand, if  $V$  is de Rham, the map  $\tau \mapsto \partial_{V(-k)} \circ \text{pr}_{K_m}(c_\tau)$  is a cocycle on  $\Gamma_{K_m}$  with values in  $\mathbf{D}_{\text{dR}}(V(-k))$  which  $\Gamma_{K_m}$  acts trivially. It is of the form  $\tau \mapsto d \log_p \chi(\tau)$ , where  $d \in \mathbf{D}_{\text{dR}}(V(-k))$  and if  $c$  is zero, which implies  $\tau \mapsto c_\tau$  is a coboundary, hence  $d = 0$ . One can deduce that  $\partial_{V(-k)} \circ \text{pr}_{K_m} \left( \frac{1}{\log_p \chi(\gamma)} c_\gamma \right) \in \mathbf{D}_{\text{dR}}(V(-k))$  does not depend on  $\gamma \in \Gamma_{K_m}$  such that  $\chi(\gamma) \neq 0$  and the choice of cocycle  $\tau \mapsto c_\tau$  representing  $c$ , which provides us a natural map from  $H^1(K, V(-k))$  to  $\mathbf{D}_{\text{dR}}(V(-k))$  coincides with  $\exp_{V^*(1+k)}^*$  by proposition 7.2.  $\square$

**7.2. Explicit reciprocity law.** Let  $V$  be a de Rham representation of  $G_K$  and let  $n(V) \geq n_1(V)$  smallest integer satisfies  $r_{n(V)} \geq r_V$  (c.f. proposition 6.5). If  $\mu \in H_{\text{Iw}}^1(K, V)$ , then  $\text{Exp}_{V^*(1)}^*(\mu) \in D(V)^{\psi=1}$ . On the other hand,  $D(V)^{\psi=1} \subset \mathbf{D}^{\dagger, r_V}(V)$ . If  $n \geq n(V)$ , we can view  $\varphi^{-n}(\text{Exp}_{V^*(1)}^*(\mu))$  as an element in  $\mathbf{B}_{\text{dR}} \otimes V$ . Since  $\varphi^{-n}(\text{Exp}_{V^*(1)}^*(\mu))$  is an element of  $\mathbf{B}_{\text{dR}} \otimes V$  fixed by  $H_K$ , we can consider its image under  $T_m$ .

**Theorem 7.4.** *Let  $V$  be a de Rham representation and  $m \in \mathbf{N}$ .*

- i) *If  $n \geq \sup(m, n(V))$  and  $\mu \in H_{\text{Iw}}^1(K, V)$ , then  $T_m(\varphi^{-n}(\text{Exp}_{V^*(1)}^*(\mu)))$  is an element in  $K_m((t)) \otimes_K \mathbf{D}_{\text{dR}}(V)$  independent of  $n$ , we denote it by  $\text{Exp}_{V^*(1), K_m}^*(\mu)$ .*
- ii) *If  $\mu \in H_{\text{Iw}}^1(K, V)$ , then*

$$\text{Exp}_{V^*(1), K_m}^*(\mu) = \sum_{k \in \mathbf{Z}} \exp_{V^*(1+k)}^* \left( \int_{\Gamma_{K_m}} \chi(x)^{-k} \mu \right).$$

- iii) *There exists  $m(V) \geq n(V)$  such that if  $m \geq m(V)$  and  $\mu \in H_{\text{Iw}}^1(K, V)$ , then*

$$\text{Exp}_{V^*(1), K_m}^*(\mu) = p^{-m} \varphi^{-m}(\text{Exp}_{V^*(1)}^*(\mu)).$$

**Remark 7.5.**

- i) The image of  $H^1(K_m, V(-k))$  by  $\exp_{V^*(1+k)}^*$  is contained in  $\text{Fil}^0 \mathbf{D}_{\text{dR}}(V(-k)) = \text{Fil}^0(t^K \mathbf{D}_{\text{dR}}(V))$  which is zero if  $k \ll 0$ . Hence the series in ii) converges in  $\mathbf{B}_{\text{dR}} \otimes \mathbf{D}_{\text{dR}}(V)$ .
- ii) We have a map  $\mu \in H_{\text{Iw}}^1(K, V) \mapsto \int_{\Gamma_{K_n}} \chi^k \mu \in H^1(G_{K_n}, V(k))$ , thus  $\exp_{V^*(1+k)}^* \left( \int_{\Gamma_{K_n}} \chi^{-k} \mu \right) \in t^k K_n \otimes_K \mathbf{D}_{\text{dR}}(V)$ .
- iii) For  $n \geq n(V)$ , we have  $\varphi^{-n}(\mathbf{D}^{\dagger, r_n}(V)) \subset K_n((t)) \otimes \mathbf{D}_{\text{dR}}(V)$ .

*Proof.* Given that  $T_r = \text{Tr}_{K_m/K_r} \circ T_m$  if  $r \leq m$  and if  $L_1 \subset L_2$  are two finite extension of  $K$ , then the diagram

$$\begin{array}{ccc} H^1(L_2, V) & \xrightarrow{\exp_{V^*(1)}^*} & L_2 \otimes \mathbf{D}_{\text{dR}}(V) \\ \downarrow \text{cor}_{L_2/L_1} & & \downarrow \text{Tr}_{L_2/L_1} \otimes \text{id} \\ H^1(L_1, V) & \xrightarrow{\exp_{V^*(1)}^*} & L_1 \otimes \mathbf{D}_{\text{dR}}(V) \end{array}$$



is commutative. Thus, to prove i) and ii), it suffices to prove them for  $m$  large enough. We can therefore suppose that  $m \geq n(V) + 1$ ,  $\text{pr}_{K_m} = p^m \text{T}_m$  and  $\log_p^0(\gamma_m) = \frac{\log_p(\chi(\gamma_m))}{p^m}$ .

Denote  $y$  the element  $\text{Exp}_{V^*(1)}^*(\mu)$  in  $D(V)^{\psi=1}$  and if  $i \in \mathbf{Z}$ , denote  $y(i)$  the image of  $y$  in  $D(V(i))^{\psi=1} = D(V)^{\psi=1}$  (same as set but different as Galois module by twist  $\chi^i$ ). By construction of  $\text{Exp}_{V^*(1)}^*$  (indeed its inverse),  $\int_{\Gamma_{K_n}} \chi(x)^{-k} \mu$  is represented by the cocycle

$$\sigma \mapsto c'_\sigma = \log_p^0(\gamma_m) \left( \frac{\sigma - 1}{\gamma_m - 1} y(-k) - (\sigma - 1)b \right),$$

where  $b \in \mathbf{A} \otimes V$  is a solution of the equation  $(\varphi - 1)b = (\gamma_m - 1)^{-1}((\varphi - 1)y)(-k)$ .

By definition of  $n(V)$ , we have  $y \in \mathbf{D}^{\dagger, r_n(V)}(V) \subset \mathbf{D}^{\dagger, r_{m-1}}(V)$  and  $(\varphi - 1)y \in \mathbf{D}^{\dagger, r_m}(V)$ , which implies that  $(\gamma_m - 1)^{-1}(\varphi - 1)y(-k) \in \mathbf{D}^{\dagger, r_{m+1}}(V)$  by proposition 6.5, and the same argument as lemma 4.14 implies that  $b \in \mathbf{A}^{\dagger, r_m} \otimes V$ . Since we suppose that  $n \geq \sup(m, n(V))$ , we have  $\varphi^{-n}(b)$  and  $\varphi^{-n}(y)$  are both in  $\mathbf{B}_{\text{dR}}^+ \otimes V$  and  $c'_\sigma = \varphi^{-n}(c'_\sigma)$  is a cocycle with values in  $\mathbf{B}_{\text{dR}} \otimes V$  which differs from the cocycle

$$\sigma \mapsto c_\sigma = \frac{\log_p \chi(\gamma_m)}{p^m} \frac{\sigma - 1}{\gamma_m - 1} \varphi^{-n}(y(-k))$$

by a coboundary  $\sigma \mapsto \frac{\log_p \chi(\gamma_m)}{p^m} (\sigma - 1) \varphi^{-n}(b)$ . Since  $y$  is fixed by  $H_K$ , the cocycle  $\sigma \mapsto c_\sigma$  has values in  $(\mathbf{B}_{\text{dR}} \otimes V)^{H_K}$  which allows us to use proposition 7.3 to calculate it and we obtain

$$\text{exp}_{V^*(1+k)}^* \left( \int_{\Gamma_{K_m}} \chi(x)^{-k} \mu \right) = \frac{1}{\log_p \chi(\gamma_m)} \partial_{V(-k)}(\text{pr}_{K_m}(c_{\gamma_m})) = \frac{1}{p^m} \partial_{V(-k)}(\text{pr}_{K_m}(\varphi^{-n}(y)))$$

and since  $\frac{1}{p^m} \text{pr}_{K_m} = \text{T}_m$  and  $\text{T}_m(x) = \sum_{k \in \mathbf{Z}} \partial_{V(-k)}(\text{T}_m(x))$  if  $x \in (\mathbf{B}_{\text{dR}} \otimes V)^{H_K}$ , we deduce i) and ii).

To prove iii), it suffices to show that if  $m$  is large enough, then  $\varphi^{-m}(\text{Exp}_{V^*(1)}^*(\mu)) \in K_m((t)) \otimes_K \mathbf{B}_{\text{dR}}(V)$ . We need the following lemma:

**Lemma 7.6.** *Let  $d$  be an integer  $\geq 1$ . If  $U \in GL_d(\mathbf{B}_{\text{dR}}^{H_K})$  and there exists  $n \in \mathbf{N}$  such that  $U^{-1}\gamma(U) \in GL_d(K_n((t)))$ , then there exists  $m \in \mathbf{N}$  such that  $U \in GL_d(K_m((t)))$ .*

*Proof.* Let  $A = U^{-1}\gamma(U)$ . If  $m \geq n$ , let  $U_m = \text{pr}_{K_m}(U)$ . Using the fact that  $\text{pr}_{K_m}$  is  $K_n((t))$ -linear if  $m \geq n$ , we obtain, by applying  $\text{pr}_{K_m}$  to the identity  $UA = \gamma(U)$ , the relation  $U_m A = \gamma(U_m)$ . On the other hand, since  $\lim_{m \rightarrow +\infty} U_m = U$ , there exists  $m \geq n$  such that  $U_m$  is invertible. Subtract  $A$  by the above identity, We have  $UU_m^{-1}$  is fixed by  $\gamma$  and therefore belongs to  $GL_d(K)$ . We hence deduce that  $U$  belongs to  $GL_d(K_m((t)))$ .  $\square$

Let  $e_1, \dots, e_d$  be a basis of  $\mathbf{D}^{\dagger, r_n(V)}(V)$  over  $\mathbf{B}_K^{\dagger, r_n(V)}$  which contains  $D(V)^{\psi=1}$  and  $f_1, \dots, f_d$  a basis of  $\mathbf{B}_{\text{dR}}(V)$  over  $K$ . Let  $A = (a_{i,j}) \in GL_d(\mathbf{B}_K^{\dagger})$ ,  $B = (b_{i,j}) \in GL_d(\mathbf{B}_K^{\dagger})$  and if  $m \geq n(V)$ ,  $C^{(m)} = (c_{i,j}^{(m)}) \in GL_d(\mathbf{B}_{\text{dR}}^{H_K})$  the matrices defined by

$$\gamma(e_i) = \sum_{j=1}^d a_{i,j} e_j, \quad \varphi(e_i) = \sum_{j=1}^d b_{i,j} e_j \quad \text{and} \quad \varphi^{-m}(e_i) = \sum_{j=1}^d c_{i,j}^{(m)} f_j.$$

The relation  $\gamma \circ \varphi^{-m} = \varphi^{-m} \circ \gamma$  and  $\varphi^{-m} = \varphi^{-(m+1)} \circ \varphi$  is translated to

$$\gamma(C^{(m)}) = C^{(m)} \varphi^{-m}(A) \quad \text{and} \quad C^{(m)} = C^{(m+1)} \varphi^{-(m+1)}(C^{-1})$$

since  $f_1, \dots, f_d$  is fixed by  $\gamma$ . There exists  $n_0 \geq n(V)$  such that  $A$  and  $B$  belongs to  $GL_d(\mathbf{B}_K^{\dagger, r_{n_0}})$ . Since there exists  $m_0 \in \mathbf{N}$  such that  $\varphi^{-m}(\mathbf{B}_K^{\dagger, r_m}) \in K_m[[t]]$ , if  $m \geq m_0$ . By above relations and lemma 7.6, there exists  $m(V) \geq \sup(n_0, m_0) = m_1$  such that  $C^{(m_1)} \in GL_d(K_{m(V)}((t)))$ , which implies that  $C^{(m)} \in GL_d(K_{m(V)}((t)))$  for  $m \geq m(V)$  by second relation. Since  $x \in D(V)^{\psi=1}$  is of the form  $\sum_{i=1}^d x_i e_i$  where  $x \in \mathbf{B}^{\dagger, r_{n(V)}}$  and  $\varphi^{-m}(\mathbf{B}^{\dagger, r_{n(V)}}) \subset K_m[[t]]$  if  $m \geq m(V)$  by the choice of  $m(V)$ , we have the inclusion  $\varphi^{-m}(D(V)^{\psi=1}) \subset K_m((t)) \otimes_K \mathbf{D}_{\text{dR}}(V)$  if  $m \geq m(V)$ . This proves iii).  $\square$

**7.3. Connection with the Perrin-Riou's logarithm.** Our Goal in this paragraph is to compare  $\text{Exp}_{V^*(1)}^*$  and Perrin-Riou's logarithm constructed in [Col98]. Let's recall the construction of logarithm map.

**Proposition 7.7.** *Let  $V$  be a de Rham representation. Let  $W$  be the finite dimensional  $\mathbf{Q}_p$ -vector space  $\cup_{n \in \mathbf{N}} (\mathbf{B}_{\text{cris}}^{\varphi=1} \otimes V)^{G_{K_n}}$ . Let  $\mu \in H_{\text{Iw}}^1(K_n, V)$  such that  $\int_{\Gamma_{K_n}} \mu \in H_e^1(K, V)$  for all  $n \in \mathbf{N}$  and  $\tau \rightarrow \mu_\tau$  a continuous cocycle represent  $\mu$ . Finally, if  $n \gg 0$ , let  $c_n$  be the unique element of  $(\mathbf{B}_{\text{cris}}^{\varphi=1} \otimes V)/W$  verifies  $(1 - \tau)c_n = \int_{K_n} \mu_\tau$  for all  $\tau \in G_{K_n}$ .*

- i) *The sequence  $p^n c_n$  converges in  $(\mathbf{B}_{\text{cris}}^{\varphi=1} \otimes V)/W$  to an element of  $(\mathbf{B}_{\text{cris}}^{\varphi=1} \otimes V)^{G_K}/W$  denoted by  $\text{Log}_V(\mu)$ .*
- ii) *If  $n \in \mathbf{N}$ , then*

$$t \frac{d}{dt} T_n(\text{Log}_V(\mu)) = \sum_{k \in \mathbf{N}} \exp_{V^*(1+k)}^* \left( \int_{\Gamma_{K_n}} \chi(x)^{-k} \mu \right).$$

*Proof.* See [Col98, Theorem VI.3.1 and Theorem VII.1.1].  $\square$

**Remark 7.8.**

- i) There exists  $k_0 \in \mathbf{N}$  such that the condition  $\int_{\Gamma_{K_n}} \mu \in H_e^1(K, V)$  for all  $n \in \mathbf{N}$  holds automatically if we replace  $V$  by  $V(k)$  for  $k \geq k_0$ .
- ii) The operator  $\frac{d}{dt}$  annihilates  $K_\infty \otimes \mathbf{D}_{\text{dR}}(V)$  and hence  $W$ , which explains why we don't need to pass to quotient  $W$  in formula (ii).

The connection between  $\text{Log}_V$  and  $\text{Exp}_{V^*(1)}^*$  in the case  $V$  is de Rham is by:

**Theorem 7.9.** *Let  $V$  be a de Rham representation of  $G_K$ . There exists  $m(V) \geq n(V)$  such that if  $m \geq m(V)$  and  $\mu \in H_{\text{Iw}}^1(K, V)$  such that  $\int_{\Gamma_{K_n}} \mu \in H_e^1(K_n, V)$  for all  $n \in \mathbf{N}$ , then*

$$p^{-m} \varphi^{-m}(\text{Exp}_{V^*(1)}^*(\mu)) = t \frac{d}{dt} (T_m(\text{Log}_V(\mu)))$$

*Proof.* Given ii) of proposition 7.7, it is immediately followed by theorem 7.4.  $\square$

**Remark 7.10.** It is possible that the theorem is empty, that is there exists no nonzero element in  $H_{\text{Iw}}^1(K, V)$  satisfies the assumptions in proposition 7.7, but as we note above, if we replace  $V$  by  $V(k)$  for  $k \gg 0$ , then the assumptions of the theorem is verified for all elements of  $H_{\text{Iw}}^1(K, V)$ .

## 8. THE $\mathbf{Q}_p(1)$ REPRESENTATION AND COLEMAN'S POWER SERIES

**8.1. The module  $D(\mathbf{Z}_p(1))^{\psi=1}$ .** The module  $\mathbf{Z}_p(1)$  is just  $\mathbf{Z}_p$  with the action of  $G_{\mathbf{Q}_p}$  defined by  $g \in G_{\mathbf{Q}_p}$ ,  $x \in \mathbf{Z}_p(1)$ ,  $g(x) = \chi(g)x$ . We shall study the exponential map

$$\text{Exp}_{\mathbf{Q}_p}^* : H_{\text{Iw}}^1(\mathbf{Q}_p, \mathbf{Z}_p(1)) \rightarrow D(\mathbf{Z}_p(1))^{\psi=1}.$$

Note that  $D(\mathbf{Z}_p(1)) = (\mathbf{A} \otimes \mathbf{Z}_p(1))^{H_{\mathbf{Q}_p}} = \mathbf{A}_{\mathbf{Q}_p}(1)$ , with usual actions of  $\varphi$  and  $\psi$ , and for  $\gamma \in \Gamma$ ,  $\gamma(f(\pi)) = \chi(\gamma)f((1 + \pi)^{\chi(\gamma)} - 1)$ , for all  $f(\pi) \in \mathbf{A}_{\mathbf{Q}_p}(1)$ .

**Proposition 8.1.**  $(\mathbf{A}_{\mathbf{Q}_p})^{\psi=1} = \mathbf{Z}_p \cdot \frac{1}{\pi} \oplus (\mathbf{A}_{\mathbf{Q}_p}^+)^{\psi=1}$ .

*Proof.* Note that we have  $\psi(\mathbf{A}_{\mathbf{Q}_p}^+) \subset \mathbf{A}_{\mathbf{Q}_p}^+$ ,  $\psi(\frac{1}{\pi}) = \frac{1}{\pi}$  and  $\nu_E(\psi(x)) \geq [\frac{\nu_E(x)}{p}]$  if  $x \in \mathbf{E}_{\mathbf{Q}_p}^+$ . These facts imply that  $\psi - 1$  is bijective on  $\mathbf{E}_{\mathbf{Q}_p}/\pi^{-1}\mathbf{E}_{\mathbf{Q}_p}^+$  and hence it is also bijective on  $\mathbf{A}_{\mathbf{Q}_p}/\pi^{-1}\mathbf{A}_{\mathbf{Q}_p}^+$ . Thus  $\psi(x) = x$  implies  $x \in \pi^{-1}\mathbf{A}_{\mathbf{Q}_p}^+$ .  $\square$

**8.2. Kummer theory.** We define the Kummer map  $\kappa : K^* \rightarrow H^1(K, \mathbf{Q}_p(1))$  as follows: For  $a \in K^*$ , we choose  $x$  any element in  $\tilde{\mathbf{E}}$  satisfying  $x^{(0)} = a$ , then  $\tau \mapsto (1 - \tau)(\frac{\log[x]}{t})(1)$  is a 1-cocycle on  $G_K$  with values in  $\mathbf{Q}_p(1)$  whose image in  $H^1(K, \mathbf{Q}_p(1))$  is defined to be  $\kappa(a)$ .

Recall that  $\varepsilon = (1, \varepsilon^{(1)}, \dots) \in \mathbf{E}_{\mathbf{Q}_p}^+$ ,  $\varepsilon^{(1)} \neq 1$ . Let  $F_n = \mathbf{Q}_p(\varepsilon^{(n)})$  and  $\kappa_n : F_n^* \rightarrow H^1(F_n, \mathbf{Q}_p(1))$  be the Kummer maps defined above. Since  $\text{cor}_{F_{n+1}/F_n} \circ \kappa_{n+1} = \kappa_n \circ \text{N}_{F_{n+1}/F_n}$ , which induces a map

$$\kappa : \varprojlim F_n^* \rightarrow H_{\text{Iw}}^1(\mathbf{Q}_p, \mathbf{Q}_p(1)).$$

We have

$$H_{\text{Iw}}^1(\mathbf{Q}_p, \mathbf{Z}_p(1)) = \mathbf{Z}_p \cdot \kappa(\pi) \oplus \kappa(\varprojlim \mathcal{O}_{F_n}^*).$$

**8.3. Multiplicative representatives.** Recall  $\mathbf{B}$  is a extension of degree  $p$  of  $\varphi(\mathbf{B})$  (totally ramified since residual extension is purely inseparable). Define the multiplicative map  $\text{N} : \mathbf{B} \rightarrow \mathbf{B}$  by the formula  $\text{N}(x) = \varphi^{-1}(\text{N}_{\mathbf{B}/\varphi(\mathbf{B})}(x))$ . This is an multiplicative analogy of  $\psi$ .

**Lemma 8.2.** *If  $x \in \mathbf{E}^*$  and  $U_x$  denote the set  $y \in \mathbf{A}$  whose reduction modulo  $p$  is  $x$ , then  $\text{N}$  is a contractible map of  $U_x$  for the  $p$ -adic topology.*

*Proof.* Note that  $\text{N}$  induces the identity on  $\mathbf{E}$  and thus the fixes  $U_x$ . On the other hand, if  $y \equiv 1 \pmod{p^k}$ , we have

$$\text{N}(y) \equiv 1 + \varphi^{-1} \text{Tr}_{\mathbf{B}/\varphi(\mathbf{B})}(y - 1) = 1 + p\psi(y - 1) \pmod{p^{2k}},$$

which implies in particular  $\text{N}(y) - 1 \in p^{k+1}\mathbf{A}$ . We deduce that if  $y_1, y_2$  two elements of  $U_x$  verified  $y_1 - y_2 \in p^k\mathbf{A}$ , then  $\text{N}(y_1) - \text{N}(y_2) = \text{N}(y_2)(\text{N}(y_2^{-1}y_1) - 1) \in p^{k+1}\mathbf{A}$ , which proves the lemma.  $\square$

**Corollary 8.3.**

- i) *If  $x \in \mathbf{E}$ , there exists an unique element  $\hat{x} \in \mathbf{A}$  whose image modulo  $p$  is  $x$  and  $\text{N}(\hat{x}) = \hat{x}$ .*
- ii) *If  $x$  and  $y$  are two elements in  $\mathbf{E}$ , the  $\widehat{xy} = \hat{x}\hat{y}$ .*

*Proof.* i) follows from the above lemma if  $x \neq 0$  and completeness of  $U_x$  for the  $p$ -adic topology. On the other hand,  $\text{N}(p^k\mathbf{A}) \subset p^{p^k}\mathbf{A}$ , this proves that 0 is the only element of  $y$  of  $p\mathbf{A}$  verified  $\text{N}(y) = y$  and the uniqueness follows. ii) follows from the uniqueness of i).  $\square$

**Remark 8.4.** There are two multiplicative maps from  $\mathbf{E}$  to  $\tilde{\mathbf{A}}$ , namely the map  $x \rightarrow \hat{x}$  and the Techmuller map  $[x]$ . We have  $\hat{x} \neq [x]$  unless  $x \in \overline{\mathbf{F}}_p$ .

**Lemma 8.5.** *Let  $K$  be a finite extension of  $\mathbf{Q}_p$  and  $d = [\mathbf{B}_K : \mathbf{B}_{\mathbf{Q}_p}] = [K_\infty : \mathbf{Q}_p(\mu_{p^\infty})]$ . If  $n(K)$  is the smallest integer  $n \geq 2$  such that there exist  $e_1, \dots, e_d \in \tilde{\mathbf{A}}_K^{\dagger, r_n}$  such that  $\varphi(e_1), \dots, \varphi(e_d)$  form a basis of  $\tilde{\mathbf{A}}_K^{\dagger, r_{n+1}}$  over  $\tilde{\mathbf{A}}_{\mathbf{Q}_p}^{\dagger, r_{n+1}}$  and if  $n \geq n(K)$ , then  $\text{N}(\tilde{\mathbf{A}}_K^{\dagger, r_{n+1}}) \subset \tilde{\mathbf{A}}_K^{\dagger, r_n}$ .*

*Proof.* By definition of  $n(K)$ , if  $n \geq n(K)$  and  $x \in \mathbf{A}_K^{\dagger, r_{n+1}}$ , we can write  $x$  as the form  $x = \sum_{i=1}^d x_i \varphi(e_i)$  where  $x_i \in \mathbf{A}_{\mathbf{Q}_p}^{\dagger, r_{n+1}}$ . On the other hand, we can write  $x_i$  of the form  $x_i = \sum_{j=0}^{p-1} x_{i,j} [\varepsilon]^j$  where  $x_{i,j} = \varphi(\psi([\varepsilon]^{-j} x_i))$  and corollary 4.13 and proposition 6.1 show that we have  $x_{i,j} \in \varphi(\mathbf{A}_{\mathbf{Q}_p}^{\dagger, r_n})$ . We hence deduce the coordinate  $y_j = \sum_{i=1}^d x_{i,j} \varphi(e_i)$  of  $x$  in basis  $1, [\varepsilon], \dots, [\varepsilon]^{p-1}$  of  $\mathbf{B}$  over  $\varphi(\mathbf{B})$  belongs to  $\mathbf{A}_K^{\dagger, r_{n+1}} \cap \varphi(\mathbf{B}) = \varphi(\mathbf{A}_K^{\dagger, r_n})$ . On the other hand,  $N_{\mathbf{B}/\varphi(\mathbf{B})}$  is the determinant of the multiplication by  $x$  in  $\mathbf{B}$  considered as a vector space of dimension  $p$  over  $\varphi(\mathbf{B})$ , therefore the determinant of the matrix

$$\begin{pmatrix} y_0 & [\varepsilon]^p y_{p-1} & \cdots & [\varepsilon]^p y_1 \\ y_1 & y_0 & \ddots & \vdots \\ \vdots & \vdots & \ddots & [\varepsilon]^p y_{p-1} \\ y_{p-1} & y_{p-2} & \cdots & y_0 \end{pmatrix}.$$

We deduce that  $N_{\mathbf{B}/\varphi(\mathbf{B})}$  belongs to  $\varphi(\mathbf{A}_K^{\dagger, r_n})$ . Together with the relation  $N = \varphi^{-1} \circ N_{\mathbf{B}/\varphi(\mathbf{B})}$ , we complete the proof.  $\square$

**Corollary 8.6.** *If  $x \in \mathbf{E}_K^+$ , then  $\hat{x} \in \mathbf{A}_K^{\dagger, r_{n(K)}}$ . Moreover, if  $K$  is a unramified extension over  $\mathbf{Q}_p$  and  $x \in \mathbf{E}_K^+$ , then  $\hat{x} \in \mathbf{A}_K^+ = \mathbf{A}_K \cap \mathbf{A}_{\mathbf{Q}_p}^+ = \mathcal{O}_K[[\pi]]$ .*

*Proof.* Let  $v \in \mathbf{A}_K^+$  whose image in  $\mathbf{E}_K$  is  $x$  and let  $n \geq n(K)$  such that  $v \in \mathbf{A}_K^{\dagger, r_n}$ . Let  $(v_k)_{k \in \mathbf{N}}$  the sequence of elements in  $\mathbf{A}_K$  defined by  $v_0 = v$  and  $v_k = N(v_{k-1})$  if  $k \geq 1$ . By lemma 8.2, the sequence tends to  $\hat{x}$  in  $\mathbf{A}_K$  as  $k$  tends to  $+\infty$ . On the other hand, lemma 8.5, implies that  $v_k \in \mathbf{A}_K^{\dagger, r_n}$  for  $k \in \mathbf{N}$  and since  $\mathbf{A}_K^{\dagger, r_n}$  is relatively compact in  $\mathbf{A}_K^{\dagger, r_{n+1}}$ , which implies  $\hat{x} \in \mathbf{A}_K^{\dagger, r_{n+1}}$  and the result follows by using the lemma 8.5 by descending  $\mathbf{A}_K^{\dagger, r_{n+1}}$  to  $\mathbf{A}_K^{\dagger, r_{n(K)}}$ .

In the case where  $K$  is unramified over  $\mathbf{Q}_p$ , the reduction modulo  $p$  induces a surjection from  $\mathbf{A}_K^+$  to  $\mathbf{E}_K^+$  and since  $\mathbf{A}_K^+$  is a closed subring of  $\mathbf{A}_K$  fixed by  $N$ , similar proof shows that  $v \in \mathbf{A}_K^+$  implies that  $\hat{x} \in \mathbf{A}_K^+$ .  $\square$

**8.4. Generalized Coleman's power series.** Let's recall the construction of Coleman's power series.

**Proposition 8.7.** *Let  $F$  be a finite unramified extension of  $\mathbf{Q}_p$ . If  $u = (u^{(n)})_{n \in \mathbf{N}}$  is an element of the projective limit  $\varprojlim \mathcal{O}_{F_n}^*$  of  $\mathcal{O}_{F_n}^*$  with respect to the norm map, there exists a unique power series  $\text{Col}_u(T)$  in  $\mathcal{O}_F[[T]]^*$  such that we have  $\text{Col}_u^{\varphi^{-n}}(\varepsilon^{(n)} - 1) = u^{(n)}$  for all  $n \in \mathbf{N}$ .*

*Proof.* See [Co79].  $\square$

**Lemma 8.8.** *If  $K$  is a finite extension of  $\mathbf{Q}_p$  and  $n \geq n(K)$ , then the diagram*

$$\begin{array}{ccc} \mathbf{A}_K^{\dagger, r_{n+1}} & \xrightarrow{N} & \mathbf{A}_K^{\dagger, r_n} \\ \varphi^{-(n+1)} \downarrow & & \downarrow \varphi^{-n} \\ K_{n+1}[[t]] & \xrightarrow{N_{K_{n+1}/K_n}} & K_n[[t]] \end{array}$$

*is commutative.*

*Proof.* By definition,  $N_{\mathbf{B}/\varphi(\mathbf{B})}$  (resp.  $N_{K_{n+1}/K_n}(\varphi^{-(n+1)}(x))$ ) is the determinant of the multiplication by  $x$  (resp.  $\varphi^{-(n+1)}(x)$ ) over  $\mathbf{B}$  (resp.  $K_{n+1}[[t]]$ ) considered as a  $\varphi(\mathbf{B})$ -vector space

(resp.  $K_n[[t]]$ -module) and the commutativity of the diagram follows from the fact  $\varphi^{(n+1)}$  is a ring homomorphism and  $\varphi^{-n} \circ N = \varphi^{-(n+1)} \circ N_{\mathbf{B}/\varphi(\mathbf{B})}$ .  $\square$

Denote the map  $\theta_n$  the homomorphism  $\theta \circ \varphi^{-n}$  from  $\mathbf{B}^{\dagger, r_n}$  to  $\mathbf{C}_p$ .

**Lemma 8.9.** *If  $u = (u^{(n)}) \in \varprojlim \mathcal{O}_{K_n}$  and  $n \geq n(K)$ , then  $\theta_n(\widehat{\iota_K(u)}) = u^{(n)}$ .*

*Proof.* By the preceeding lemma,  $(\theta_n(\widehat{\iota_K(u)}))_{n \geq n(K)}$  belongs to  $\varprojlim \mathcal{O}_{K_n}$ . On the other hand, since  $[\iota_K(u)] - \widehat{\iota_K(u)} \in \widetilde{\mathbf{A}}_K^{\dagger, r_{n(K)}} \cap p\widetilde{\mathbf{A}}$ , which implies that if  $n \geq n(K)$ , then  $\nu_p(\theta_n([\iota_K(u)] - \widehat{\iota_K(u)})) \geq 1 - \frac{1}{p^{n-n(K)}}$  and since  $\nu_p(\theta_n([\iota_K(u)] - u^{(n)})) \geq \frac{1}{p}$  if  $n$  large enough, we show that  $(\theta_n(\widehat{\iota_K(u)}))_{n \geq n(K)}$  has same image as  $u$  in  $\mathbf{E}_K^+$  (c.f. proposition 3.1), so it is equal.  $\square$

**Proposition 8.10.** *Let  $K$  be a finite extension of  $\mathbf{Q}_p$ ,  $F = K_\infty \cap \mathbf{Q}_p^{ur}$  and  $e_K = [K_\infty : F_\infty]$ .*

- i) *If  $e_K = 1$  and  $u \in \varprojlim \mathcal{O}_{K_n}$ , then  $\widehat{\iota_K(u)} = \text{Col}_u(\pi)$ .*
- ii) *When  $e_K \geq 2$ , there exist Laurent series  $f_0, \dots, f_{e_K-1} \in \mathcal{O}_F((T))$  converges in the annulus  $0 < \nu_p(x) < \frac{1}{(p-1)p^{n(K)-1}}$  such that, if  $n \geq n(K)$ , then  $(u^{(n)})^{e_K} + f_{e_K-1}^{\varphi^{-n}}(\varepsilon^{(n)} - 1)(u^{(n)})^{e_K-1} + \dots + f_0^{\varphi^{-n}}(\varepsilon^{(n)} - 1) = 0$ .*

*Proof.* By corollary 8.6,  $\widehat{\iota_K(u)} \in \mathbf{A}_F^+$  if  $u \in \varprojlim \mathcal{O}_{K_n}$ . In particular, there exists  $f \in \mathcal{O}_F[[T]]$  such that  $\widehat{\iota_K(u)} = f(\pi)$ . On the other hand, by applying lemma 8.9 to the map  $\theta_n$ , we obtained  $u^{(n)} = f^{\varphi^{-n}}(\varepsilon^{(n)} - 1)$ , which shows  $f = \text{Col}_u$  by the characterization of  $\text{Col}_u$ .

ii) By corollary 8.6,  $\widehat{\iota_K(u)} \in \mathbf{A}_K^{\dagger, r_{n(K)}}$ . On the other hand,  $\mathbf{A}_K^{\dagger, r_{n(K)}}$  is of dimension  $e_K$  over  $\mathbf{A}_F^{\dagger, r_{n(K)}}$  (by the definition of  $n(K)$ ); so we can find elements  $\tilde{f}_0, \dots, \tilde{f}_{e-1} \in \mathbf{A}_F^{\dagger, r_{n(K)}}$  such that we have  $\widehat{\iota_K(u)}^{e_K} + \tilde{f}_{e-1}\widehat{\iota_K(u)}^{e_K-1} + \dots + \tilde{f}_0 = 0$ , by lemma 8.9, we obtain the result.  $\square$

### 8.5. The map $\text{Log}_{\mathbf{Q}_p(1)}$ and $\text{Exp}_{\mathbf{Q}_p}^*$ .

**Lemma 8.11.** *If  $u \in \mathbf{E}_K$ , the sequence  $(\varphi^{-n}(\widehat{\iota_K(u)}))^{p^n}$  converge in to  $[\iota_K(u)]$  in  $\widetilde{\mathbf{A}}$  and  $\mathbf{B}_{\text{dR}}^+$ .*

*Proof.* Since  $\widehat{\iota_K(u)} \in \mathbf{A}^{\dagger, r_{n(K)}}$  with image  $\iota_K(u)$  in  $\mathbf{E}$ , it can be written as the form  $[\iota_K(u)] + \sum_{k=1}^{+\infty} p^k[x_k]$ , where  $x_k$  are elements of  $\widetilde{\mathbf{E}}$  satisfying  $\nu_E(x_k) \geq -kp^{n(K)}$ . We have the formula  $v_n = \varphi^{-n}(\widehat{\iota_K(u)}) = [\iota_K(u)^{p^{-n}}] + \sum_{k=1}^{+\infty} p^k[x_k^{p^{-n}}]$  and the congruence  $v_n^{p^n} \equiv [\iota_K(u)] \pmod{p^{n+1}\widetilde{\mathbf{A}}}$ , thus it converges in  $\widetilde{\mathbf{A}}$ .

Let  $\alpha$  an element in  $\widetilde{\mathbf{E}}^+$  verified  $\nu_E(\alpha) = \frac{p-1}{p}$ , thus  $(\frac{p}{[\alpha]})^i$  tends to 0 in  $\mathbf{B}_{\text{dR}}^+$  as  $i$  tends to  $+\infty$ . If  $n \geq n(K) + 1$ , the above formula shows that  $v_n$  belongs to the subring  $A$  (c.f. section 4.4) of  $\mathbf{B}_{\text{dR}}^+$  of elements of the form  $y = \sum_{i=0}^{+\infty} y_i(\frac{p}{[\alpha]})^i$ , where  $y_i$  are elements in  $\mathbf{A}$  and we have  $v_n - [\iota_K(u)^{p^{-n}}] \in \frac{p}{[\alpha]}A$ . We deduce that  $v_n^{p^n}$  tends to  $[\iota_K(u)]$  in  $A$  and a fortiori in  $\mathbf{B}_{\text{dR}}^+$ .  $\square$

**Proposition 8.12.** *Let  $K$  be a finite extension of  $\mathbf{Q}_p$  and  $u \in \varprojlim \mathcal{O}_{K_n}$ .*

- i)  $\text{Log}_{\mathbf{Q}_p(1)}(\kappa(u)) = t^{-1} \log[\iota_K(u)]$
- ii) *If  $n \geq n(K)$ , then  $T_n(\text{Log}_{\mathbf{Q}_p(1)}(\kappa(u))) = t^{-1} \log \varphi^{-n}(\widehat{\iota_K(u)})$ .*
- iii)  $\text{Exp}_{\mathbf{Q}_p}^*(\kappa(u)) = \widehat{\iota_K(u)}^{-1} \partial \widehat{\iota_K(u)}$ , where  $\partial$  is the derivation  $(1 + \pi) \frac{d}{d\pi}$  (see 6.2).

*Proof.* By construction of Kummer map, if  $u_n$  any element in  $\tilde{\mathbf{E}}^+$  satisfying  $u_n^{(0)} = u^{(n)}$ , then  $\tau \mapsto (1 - \tau)(\frac{\log[u_n]}{t})(1)$  is a 1-cocycle on  $G_{K_n}$  with values in  $\mathbf{Q}_p(1)$  whose image in  $H^1(K_n, \mathbf{Q}_p(1))$  is equal to  $\kappa_n(u^{(n)})$ . Since we suppose that  $u^{(n)} \in \mathcal{O}_{K_n}^*$ , we have  $\log[u_n] \in \mathbf{B}_{\text{cris}}$  and  $\frac{\log[u_n]}{t}(1) \in \mathbf{B}_{\text{cris}}^{\varphi=1} \otimes \mathbf{Q}_p(1)$ , proving that  $\kappa_n(u^{(n)}) \in H_e^1(K_n, \mathbf{Q}_p(1))$ . Hence we deduce the formula

$$\text{Log}_{\mathbf{Q}_p(1)}(\kappa(u)) = \lim_{n \rightarrow +\infty} p^n \frac{\log[u_n]}{t}(1) = t^{-1} \lim_{n \rightarrow +\infty} \log([u_n]^{p^n}).$$

Finally, we have  $\nu_p(\theta([ \iota_K(u)^{p^{-n}} ]) - \theta([u_n])) \geq \frac{1}{p}$  if  $n$  large enough, therefore  $[u_n]^{p^n}$  tends to  $[ \iota_K(u) ]$  as  $n$  tends to  $+\infty$ . We complete i).

By i) and lemma 8.11, we have

$$\text{T}_n(\text{Log}_{\mathbf{Q}_p(1)}(\kappa(u))) = t^{-1} \lim_{m \rightarrow +\infty} \text{T}_n(p^m \log(\varphi^{-m}(\widehat{\iota_K(u)}))).$$

On the other hand, if  $m \geq n$ , we have  $\text{T}_n = \text{Tr}_{K_m[[t]]/K_n[[t]]} \circ T_m$  and since  $\varphi^{-m}(\widehat{\iota_K(u)}) \in K_m[[t]]$  and the restriction of  $T_m$  on  $K_m[[t]]$  is multiplication by  $p^{-m}$ , we obtain the formula

$$\begin{aligned} \text{T}_n(\text{Log}_{\mathbf{Q}_p(1)}(\kappa(u))) &= t^{-1} \lim_{m \rightarrow +\infty} \text{Tr}_{K_m[[t]]/K_n[[t]]}(\log(\varphi^{-m}(\widehat{\iota_K(u)}))) \\ &= t^{-1} \lim_{m \rightarrow +\infty} \log(\text{N}_{K_m[[t]]/K_n[[t]]}(\varphi^{-m}(\widehat{\iota_K(u)}))) \end{aligned}$$

and this completes ii) by using lemma 8.8.

Note that  $t^{-1}$  is a generator of  $\mathbf{D}_{\text{dR}}(\mathbf{Q}_p(1))$ . ii) and theorem 7.9 implies that if  $n$  is large enough, we have

$$\begin{aligned} \varphi^{-n}(\text{Exp}_{\mathbf{Q}_p}^*(\kappa(u))) &= t^{-1} \left( t \frac{d}{dt} (p^n \log \varphi^{-n}(\widehat{\iota_K(u)})) \right) \\ &= p^n \frac{\frac{d}{dt}(\varphi^{-n}(\widehat{\iota_K(u)}))}{\varphi^{-n}(\widehat{\iota_K(u)})} \\ &= \varphi^{-n}(\widehat{\iota_K(u)})^{-1} \frac{\partial \widehat{\iota_K(u)}}{\partial \iota_K(u)} \quad \text{by lemma 6.3,} \end{aligned}$$

which complete iii). □

### 8.6. Cyclotomic units and Coates-Wiles homomorphisms.

**Example 8.13.** Let  $K = \mathbf{Q}_p$ ,  $V = \mathbf{Q}_p(1)$  and  $u = (\frac{\zeta_{p^n}-1}{\zeta_{p^n}})_{n \geq 1} \in \varprojlim \mathcal{O}_{F_n}^*$ . Then its Coleman's power series is  $\text{Col}_u(T) = \frac{1+T}{T}$ . By iii) of proposition 8.12, we have  $\text{Exp}_{\mathbf{Q}_p}^*(\kappa(u)) = (\frac{\partial \text{Col}_u(T)}{\partial \text{Col}_u(T)})(\pi) = \frac{1}{\pi}$ . On the other hand, since

$$\varphi^{-1}(\pi)^{-1} = ([\varepsilon^{\frac{1}{p}}] - 1)^{-1} = \frac{1}{\varepsilon^{(1)} \exp(t/p) - 1},$$

by iii) of theorem 7.4, we have

$$\begin{aligned}
 \text{Exp}_{\mathbf{Q}_p, F_1}^*(\kappa(u)) &= \frac{1}{p} \text{Tr}_{F_1/\mathbf{Q}_p} \varphi^{-1} \left( \frac{1}{\pi} \right) \\
 &= \frac{1}{p} \sum_{\zeta^p=1, \zeta \neq 1} \frac{1}{\zeta \exp(t/p)} \\
 &= \frac{-1}{t} \left( \frac{t}{1 - \exp(t)} - \frac{t/p}{1 - \exp(t/p)} \right) \\
 &= \sum_{k=1}^{+\infty} (1 - p^{-k}) \zeta(1-k) \frac{(-t)^{k-1}}{(k-1)!}.
 \end{aligned}$$

Thus by ii) of theorem 7.4,

$$\exp_{\mathbf{Q}_p(1+k)^*}^* \left( \int_{\Gamma_{F_1}} \chi^{-k} \kappa(\mu) \right) = \begin{cases} 0 & \text{if } k \leq 0 \\ (1 - p^{-k}) \zeta(1-k) \frac{(-t)^{k-1}}{(k-1)!} & \text{if } k \geq 1 \end{cases}.$$

**Example 8.14.** Let  $K = \mathbf{Q}_p$ ,  $V = \mathbf{Q}_p(1)$  and  $u = (\frac{\zeta_{p^n}^a - 1}{\zeta_{p^n} - 1})_{n \geq 1} \in \varprojlim \mathcal{O}_{F_n}^*$ , where  $a \in \mathbf{Z}$ . Then its Coleman's power series is  $\text{Col}_u(T) = \frac{(1+T)^a - 1}{T}$ . By iii) of proposition 8.12, we have  $\text{Exp}_{\mathbf{Q}_p}^*(\kappa(u)) = (\frac{\partial \text{Col}_u(T)}{\partial u(T)})(\pi) = \frac{a(1+\pi)^a}{(1+T)^{a-1}} - \frac{1+T}{T}$ . On the other hand, since

$$\varphi^{-1}(\pi)^{-1} = ([\varepsilon^{\frac{1}{p}}] - 1)^{-1} = \frac{1}{\varepsilon^{(1)} \exp(t/p) - 1},$$

by iii) of theorem 7.4, we have

$$\begin{aligned}
 \text{Exp}_{\mathbf{Q}_p, F_1}^*(\kappa(u)) &= \frac{1}{p} \text{Tr}_{F_1/\mathbf{Q}_p} \varphi^{-1} (\text{Exp}_{\mathbf{Q}_p}^*(\kappa(u))) \\
 &= a - 1 + \frac{1}{p} \sum_{\zeta^p=1, \zeta \neq 1} \frac{a}{\zeta \exp at/p - 1} - \frac{1}{\zeta \exp t/p} \\
 &= a - 1 + \frac{-1}{t} \left( \frac{at}{1 - \exp(at)} - \frac{at/p}{1 - \exp(at/p)} - \frac{t}{1 - \exp(t)} + \frac{t/p}{1 - \exp(t/p)} \right) \\
 &= \sum_{k=1}^{+\infty} (1 - p^{-k}) (a^k - 1) \zeta(1-k) \frac{(-t)^{k-1}}{(k-1)!}.
 \end{aligned}$$

Thus by ii) of theorem 7.4,

$$\exp_{\mathbf{Q}_p(1+k)^*}^* \left( \int_{\Gamma_{F_1}} \chi^{-k} \kappa(\mu) \right) = \begin{cases} 0 & \text{if } k \leq 0 \\ (a^k - 1)(1 - p^{-k}) \zeta(1-k) \frac{(-t)^{k-1}}{(k-1)!} & \text{if } k \geq 1 \end{cases}.$$

**Example 8.15.** Let  $K = \mathbf{Q}_p(\zeta_d)$ ,  $V = \mathbf{Q}_p(1)$  and  $\varepsilon$  is a Dirichlet character of conductor  $d \geq 1$  prime to  $p$ . Set  $u = (\frac{-1}{G(\varepsilon^{-1})} \sum_{b \bmod d} \frac{\varepsilon^{-1}(b)}{\zeta_d^b \zeta_{p^n} - 1})_{n \geq 1} \in \varprojlim \mathcal{O}_{K_n}^*$ , then we have

$$\text{Col}_u(T) = \frac{-1}{G(\varepsilon^{-1})} \sum_{b \bmod d} \frac{\varepsilon^{-1}(b)}{\zeta_d^b (1+T) - 1}$$

and thus

$$\mathrm{Exp}_{\mathbf{Q}_p}^*(\kappa(u)) = \frac{-1}{G(\varepsilon^{-1})} \sum_{b \bmod d} \frac{\varepsilon^{-1}(b)}{\zeta_d^b(1 + \pi) - 1}.$$

Hence we have,

$$\begin{aligned} \mathrm{Exp}_{\mathbf{Q}_p, K_1}^*(\kappa(u)) &= p^{-1} \mathrm{Tr}_{K_1/K} \varphi^{-1}(\mathrm{Exp}_{\mathbf{Q}_p}^*(\kappa(u))) \\ &= p^{-1} \frac{-1}{G(\varepsilon^{-1})} \sum_{z^p=1, z \neq 1} \sum_{b \bmod d} \varepsilon^{-1}(b) \frac{1}{\zeta_d^{b/p} z \exp(t/p) - 1} \\ &= \frac{-1}{G(\varepsilon^{-1})} \sum_{b \bmod d} \varepsilon^{-1}(b) \left( \frac{1}{1 - \zeta_d^b \exp(t)} - p^{-1} \frac{1}{1 - \zeta_d^{b/p} \exp(t/p)} \right) \\ &= \sum_{b \bmod d} \frac{\varepsilon(b) \exp(bt)}{1 - \exp(dt)} - p^{-1} \varepsilon(p) \frac{\varepsilon(b) \exp(bt/p)}{1 - \exp(dt/p)} \\ &= \sum_{k=1}^{+\infty} (1 - \varepsilon(p) p^{-k}) L(1 - k, \varepsilon) \frac{t^{k-1}}{(k-1)!} \end{aligned}$$

and thus

$$\exp_{\mathbf{Q}_p^*(1+k)}^* \left( \int_{\Gamma_{K_1}} \chi^{-k} \kappa(\mu) \right) = \begin{cases} 0 & \text{if } k \leq 0 \\ (1 - \varepsilon(p) p^{-k}) L(1 - k, \varepsilon) \frac{t^{k-1}}{(k-1)!} & \text{if } k \geq 1 \end{cases}.$$

These allow us to define a homomorphism  $\mathrm{CW}_{k,n}$  from  $H_{\mathrm{Iw}}^1(K, V)$  to  $K_n \otimes \mathbf{D}_{\mathrm{dR}}(V)$  by putting

$$\mathrm{CW}_{k,n}(\mu) = \partial_k(\mathrm{T}_{K_n}(\mathrm{Log}_V(\mu))).$$

for each  $n \in \mathbf{N}$  and  $k \in \mathbf{Z}$ , the homomorphism is a generalization of Coates-Wiles homomorphism and we have the following theorem by proposition 7.7.

**Theorem 8.16.** *If  $\mu \in H_{\mathrm{Iw}}^1(K, V)$ , if  $n \in \mathbf{N}$  and  $k \in \mathbf{Z}$ , then*

$$\mathrm{CW}_{k,n}(\mu) = -\exp^* \left( \int_{\Gamma_{K_n}} \chi(x)^{-k} \mu \right).$$

**Remark 8.17.** The map

$$\varprojlim \mathcal{O}_{\mathbf{Q}_p(\mu_{p^n})}^* \rightarrow H_{\mathrm{Iw}}^1(\mathbf{Q}_p, \mathbf{Q}_p(1)) \rightarrow \mathbf{Q}_p, \quad u \mapsto \exp_{\mathbf{Q}_p^*(1+k)}^* \left( \int_{\Gamma_{F_1}} \chi^{-k} \kappa(\mu) \right)$$

is just the Coates-Wiles homomorphism.

## 9. $(\varphi, \Gamma)$ -MODULES AND DIFFERENTIAL EQUATIONS

**9.1. The rings  $\mathbf{B}_{\max}$  and  $\tilde{\mathbf{B}}_{\mathrm{rig}}^+$ .** The ring  $\mathbf{B}_{\max}^+$  is defined by

$$\mathbf{B}_{\max}^+ = \left\{ \sum_{n \geq 0} a_n \frac{\omega^n}{p^n} \mid a_n \in \tilde{\mathbf{B}}^+ \text{ is a sequence converges to } 0 \right\},$$

and  $\mathbf{B}_{\max} = \mathbf{B}_{\max}^+[\frac{1}{t}]$ . It is closely related to  $\mathbf{B}_{\mathrm{cris}}$  but tends to be more amenable. One could replace  $\omega$  by any generator of  $\ker(\theta)$  in  $\tilde{\mathbf{A}}^+$ . The ring  $\mathbf{B}_{\max}$  injects canonically into  $\mathbf{B}_{\mathrm{dR}}$  and, in particular, it is endowed with the induced Galois action and filtration, as well as with a continuous Frobenius  $\varphi$ , extending the map  $\varphi : \tilde{\mathbf{B}}^+ \rightarrow \tilde{\mathbf{B}}^+$ . Note that  $\varphi$  does not extend continuously to  $\mathbf{B}_{\mathrm{dR}}$ . We sets  $\tilde{\mathbf{B}}_{\mathrm{rig}}^+ = \cap_{n=0}^{+\infty} \varphi^n(\mathbf{B}_{\max}^+)$ .



We recall a representation  $V$  of  $G_K$  is crystalline if it is  $\mathbf{B}_{\text{cris}}$ -admissible, which is equivalent to  $\mathbf{B}_{\text{max}}$ -admissible or  $\tilde{\mathbf{B}}_{\text{rig}}^+[\frac{1}{t}]$ -admissible (because  $\cap_{n=0}^{+\infty} \varphi^n(\mathbf{B}_{\text{max}}^+) = \cap_{n=0}^{+\infty} \varphi^n(\mathbf{B}_{\text{cris}}^+)$  and the periods of crystalline representations live in finite dimensional  $F$ -vector subspaces of  $\mathbf{B}_{\text{max}}$ , stable by  $\varphi$  and so in fact in  $\cap_{n=0}^{\infty} \varphi^n(\mathbf{B}_{\text{max}}^+[\frac{1}{t}])$ ; that is, the  $F$ -vector space

$$\mathbf{D}_{\text{cris}}(V) = (\mathbf{B}_{\text{cris}} \otimes_{\mathbf{Q}_p} V)^{G_K} = (\mathbf{B}_{\text{max}} \otimes_{\mathbf{Q}_p} V)^{G_K} = (\tilde{\mathbf{B}}_{\text{rig}}^+[\frac{1}{t}] \otimes_{\mathbf{Q}_p} V)^{G_K}$$

is of dimension  $d = \dim_{\mathbf{Q}_p}(V)$ . Then  $\mathbf{D}_{\text{cris}}(V)$  is endowed with a Frobenius  $\varphi$  induced by that of  $\mathbf{B}_{\text{max}}$  and  $(\mathbf{B}_{\text{dR}} \otimes_{\mathbf{Q}_p} V)^{G_K} = \mathbf{D}_{\text{dR}}(V) = K \otimes_F \mathbf{D}_{\text{cris}}(V)$  so that a crystalline representation is also de Rham and  $K \otimes_F \mathbf{D}_{\text{cris}}(V)$  is a filtered  $K$ -vector space.

If  $V$  is a  $p$ -adic representation, we say  $V$  is Hodge-Tate, with Hodge Tate weights  $h_1, \dots, h_d$ , if we have a decomposition  $\mathbf{C}_p \otimes_{\mathbf{Q}_p} V \cong \bigoplus_{j=1}^d \mathbf{C}_p(h_j)$ . We say that  $V$  is positive if its Hodge-Tate weights are all negative. By using the map  $\theta : \mathbf{B}_{\text{dR}}^+ \rightarrow \mathbf{C}_p$ , it is easy to see that a de Rham representation is Hodge-Tate and that the Hodge-Tate weights of  $V$  are those integers  $h$  such that  $\text{Fil}^{-h} \mathbf{D}_{\text{dR}}(V) \neq \text{Fil}^{-h+1} \mathbf{D}_{\text{dR}}(V)$ .

**9.2. The structure of  $D(T)^{\psi=1}$ .** Recall in section 3, we introduce  $(\varphi, \Gamma)$ -modules and their relation with Galois representation. Let us now set  $K = F$  (i.e. we are working in an unramified extension of  $\mathbf{Q}_p$ ). We say that a  $p$ -adic representation  $V$  of  $G_F$  is of finite height if  $D(V)$  has a basis over  $\mathbf{B}_F$  made up of elements of  $D^+(V) = (\mathbf{B}^+ \otimes_{\mathbf{Q}_p} V)^{H_F}$ . A result of [Col99, proposition III.2] shows that  $V$  is of finite height if and only if  $D(V)$  has a sub- $\mathbf{B}_F^+$ -module which is free of rank  $d$ , and stable by  $\varphi$ . Let us recall the main result of [Col99, theorem 1] regarding crystalline representation of  $G_F$ :

**Theorem 9.1.** *If  $V$  is a crystalline representation of  $G_F$ , then  $V$  is of finite height.*

Let  $V$  be a crystalline representation of  $G_F$  and let  $T$  denote a  $G_F$  stable lattice of  $V$ . The following proposition is proved in [Ber04, proposition II.1.1]

**Proposition 9.2.** *If  $T$  is a lattice in a positive crystalline representation  $V$ , then there exists a unique sub- $\mathbf{A}_F^+$ -module  $\mathbf{N}(T)$  of  $D^+(T)$ , which satisfies the following conditions:*

1.  $\mathbf{N}(T)$  is an free  $\mathbf{A}_F^+$ -module of rank  $d = \dim_{\mathbf{Q}_p} V$ ;
2. the action of  $\Gamma_F$  preserves  $\mathbf{N}(T)$  and is trivial on  $\mathbf{N}(T)/\pi \mathbf{N}(T)$ ;
3. there exists an integer  $r \geq 0$  such that  $\pi^r D^+(T) \subset \mathbf{N}(T)$ .

Moreover,  $\mathbf{N}(T)$  is stable by  $\varphi$ , and the  $\mathbf{B}_F^+$ -module  $\mathbf{N}(V) = \mathbf{B}_F^+ \otimes_{\mathbf{A}_F^+} \mathbf{N}(T)$  is the unique sub- $\mathbf{B}_F^+$ -module of  $D^+(V)$  satisfying the corresponding conditions.

The  $\mathbf{A}_F^+$ -module  $\mathbf{N}(T)$  is called the Wach module associated to  $T$ .

Notice that  $\mathbf{N}(T(-1)) = \pi \mathbf{N}(T) \otimes e_{-1}$ . When  $V$  is no longer positive, we can therefore defined  $\mathbf{N}(T)$  as  $\pi^{-h} \mathbf{N}(T(-h)) \otimes e_h$  for  $h$  large enough so that  $V(-h)$  is positive. Using the results of [Ber04, III.4], one can show that:

**Proposition 9.3.** *If  $T$  is a lattice in a crystalline representation  $V$  of  $G_F$ , whose Hodge-Tate weights are in  $[a, b]$ , then  $\mathbf{N}(T)$  is the unique sub- $\mathbf{A}_F^+$ -module of  $D^+(T)[1/\pi]$  which is free of rank  $d$ , stable by  $\Gamma_F$  with the action of  $\Gamma_F$  being trivial on  $\mathbf{N}(T)/\pi \mathbf{N}(T)$  and such that  $\mathbf{N}(T)[1/\pi] = D^+(T)[1/\pi]$ .*

Finally, we have  $\varphi(\pi^b \mathbf{N}(R)) \subset \pi^b \mathbf{N}(T)$  and  $\pi^b \mathbf{N}(T)/\varphi^*(\pi^b)$  is killed by  $q^{b-a}$ , where  $q = \varphi(\omega)$ . The construction  $T \mapsto \mathbf{N}(T)$  gives a bijection between Wach modules over  $\mathbf{A}_F^+$  which are lattices in  $\mathbf{N}(V)$  and Galois lattices  $T$  in  $V$ .

Indeed  $D(V)^{\psi=1}$  is not very far from being included in  $\mathbf{N}(V)$ :

**Theorem 9.4.** *If  $V$  is a crystalline representation of  $G_F$ , whose Hodge-Tate weights are in  $[a; b]$ , then  $D(V)^{\psi=1} \subset \pi^{a-1}\mathbf{N}(V)$ . In addition, if  $V$  has no quotient isomorphic to  $\mathbf{Q}_p(a)$ , then actually  $D(V)^{\psi=1} \subset \pi^a\mathbf{N}(V)$ .*

*Proof.* See [Ber03, Theorem A.3].  $\square$

**9.3.  $p$ -adic representations and differential equations.** In this paragraph, we recall some of the results of [Ber02], which allow us to recover  $\mathbf{D}_{\text{cris}}(V)$  from the  $(\varphi, \Gamma)$ -module associated to  $V$ . Let  $\mathcal{H}_{F'}^\alpha$  be the set of power series  $f(T) = \sum_{k \in \mathbf{Z}} a_k T^k$  such that  $a_k$  is a sequence (not necessarily bounded) of elements of  $F'$ , and such that  $f(T)$  is holomorphic on the  $p$ -adic annulus  $\{p^{-1/\alpha} \leq |T| < 1\}$ .

For  $r \geq r(K)$  (c.f. proposition 6.1), define  $\tilde{\mathbf{B}}_{\text{rig}, K}^{\dagger, r}$  as the set of  $f(\pi_K)$  where  $f(T) \in \mathcal{H}_{F'}^{e_K r}$ . Obviously,  $\mathbf{B}_K^{\dagger, r} \subset \tilde{\mathbf{B}}_{\text{rig}, K}^{\dagger, r}$  and the second ring is the completion of the first one for the natural Fréchet topology. If  $V$  is a  $p$ -adic representation, let

$$\mathbf{D}_{\text{rig}}^{\dagger, r}(V) = \tilde{\mathbf{B}}_{\text{rig}, K}^{\dagger, r} \otimes_{\mathbf{B}_K^{\dagger, r}} \mathbf{D}^{\dagger, r}(V) \quad \text{and} \quad \mathbf{D}_{\text{rig}}^{\dagger}(V) = (\tilde{\mathbf{B}}_{\text{rig}}^{\dagger})^{H_K} \otimes_{\mathbf{B}_K^{\dagger}} \mathbf{D}^{\dagger}(V).$$

One of the main technical tools of [Ber02] is the construction of a large ring  $\tilde{\mathbf{B}}_{\text{rig}}^{\dagger}$ , which contains  $\tilde{\mathbf{B}}_{\text{rig}}^{\dagger}$  and  $\tilde{\mathbf{B}}^{\dagger}$ . This ring is a bridge between  $p$ -adic Hodge theory and the theory of  $(\varphi, \Gamma)$ -modules.

As a consequence of the two above inclusions, we have:

$$\mathbf{D}_{\text{cris}}(V) \subset (\tilde{\mathbf{B}}_{\text{rig}}^{\dagger}[\frac{1}{t}] \otimes_{\mathbf{Q}_p} V)^{G_K} \quad \text{and} \quad \mathbf{D}_{\text{rig}}^{\dagger}(V)[\frac{1}{t}] \subset (\tilde{\mathbf{B}}_{\text{rig}}^{\dagger}[\frac{1}{t}] \otimes_{\mathbf{Q}_p} V)^{H_K}.$$

One of the main result of [Ber02] is:

**Theorem 9.5.** *If  $V$  is a  $p$ -adic representation of  $G_K$  then  $\mathbf{D}_{\text{cris}}(V) = (\mathbf{D}_{\text{rig}}^{\dagger}(V)[\frac{1}{t}])^{\Gamma_K}$ . If  $V$  is positive, then  $\mathbf{D}_{\text{cris}}(V) = \mathbf{D}_{\text{rig}}^{\dagger}(V)^{\Gamma_K}$ .*

*Proof.* See [Ber02, theorem 3.6].  $\square$

Note that  $\tilde{\mathbf{B}}_{\text{rig}, K}^{\dagger, r}$  is the completion of  $\mathbf{B}_K^{\dagger, r}$  for the ring's natural Fréchet topology and that  $\mathbf{B}_{\text{rig}, K}^{\dagger}$  is the union of the  $\tilde{\mathbf{B}}_{\text{rig}, K}^{\dagger, r}$ . Similarly, there is a natural Fréchet topology on  $\tilde{\mathbf{B}}_{\text{rig}}^{\dagger, r}$ ,  $\tilde{\mathbf{B}}_{\text{rig}}^{\dagger, r}$  is the completion of  $\tilde{\mathbf{B}}_{\text{rig}}^{\dagger, r}$  for that topology and  $\tilde{\mathbf{B}}_{\text{rig}}^{\dagger} = \cup_{r \geq 0} \tilde{\mathbf{B}}_{\text{rig}}^{\dagger, r}$ . Actually, one can show that  $\tilde{\mathbf{B}}_{\text{rig}}^{\dagger} \subset \tilde{\mathbf{B}}_{\text{rig}}^{\dagger, r}$  for any  $r$  and there is an exact sequence

$$0 \longrightarrow \tilde{\mathbf{B}}^{\dagger} \longrightarrow \tilde{\mathbf{B}}_{\text{rig}}^{\dagger} \oplus \tilde{\mathbf{B}}_{\text{rig}}^{\dagger, r} \longrightarrow \tilde{\mathbf{B}}_{\text{rig}}^{\dagger, r} \longrightarrow 0$$

which one can take as providing a definition of  $\tilde{\mathbf{B}}_{\text{rig}}^{\dagger, r}$ .

Recall that if  $n \geq 0$  and  $r_n = p^{n-1}(p-1)$ , then there is a well-defined injective map  $\varphi^{-n} : \tilde{\mathbf{B}}_{\text{rig}}^{\dagger, r_n} \rightarrow \mathbf{B}_{\text{dR}}^+$  (c.f. section 6.2), and the map extends to an injective map  $\varphi^{-n} : \tilde{\mathbf{B}}_{\text{rig}}^{\dagger, r_n} \rightarrow \mathbf{B}_{\text{dR}}^+$  (see [Ber02, corollary 2.13]).

Let  $\mathbf{B}_{\text{rig}, F}^+$  be the set of  $f(\pi)$  where  $f(T) = \sum_{k \geq 0} a_k T^k$  with  $a_k \in F$ , and such that  $f(T)$  is holomorphic on the  $p$ -adic open unit disk. Set  $\mathbf{D}_{\text{rig}}^+(V) = \mathbf{B}_{\text{rig}, F}^+ \otimes_{\mathbf{B}_F^+} D^+(V)$ . One can show the following refinement of theorem 9.5:

**Proposition 9.6.** *We have  $\mathbf{D}_{\text{cris}}(V) = (\mathbf{D}_{\text{rig}}^+(V)[1/t])^{\Gamma_F}$  and if  $V$  is positive then  $\mathbf{D}_{\text{cris}}(V) = \mathbf{D}_{\text{rig}}^+(V)^{\Gamma_F}$ .*

Indeed if  $\mathbf{N}(V)$  is the Wach module associated to  $V$ , then  $\mathbf{N}(V) \subset D^+(V)$  when  $V$  is positive and it is shown in [Ber03, II.2] that under that hypothesis,  $\mathbf{D}_{\text{cris}}(V) = (\mathbf{B}_{\text{rig},F}^+ \otimes_{\mathbf{B}_F^+} \mathbf{N}(V))^{\Gamma_F}$ .

**9.4. The Fontaine isomorphism revisited.** The purpose of this section is to recall the constructions in section 4.2 and extend them a little bit. Let  $V$  be a  $p$ -adic representation of  $G_K$ . Recall in section 4.2, we construct a map  $h_{K,V}^1 : D(V)^{\psi=1} \rightarrow H^1(K, V)$ , and when  $\Gamma_K$  is torsion free, it gives rise to an exact sequence:

$$0 \longrightarrow D(V)_{\Gamma_K}^{\psi=1} \xrightarrow{h_{K,V}^1} H^1(K, V) \longrightarrow \left(\frac{D(V)}{\psi-1}\right)^{\Gamma_K} \longrightarrow 0$$

We shall extend  $h_{K,V}^1$  to a map  $h_{K,V}^1 : \mathbf{D}_{\text{rig}}^{\dagger}(V)^{\psi=1} \rightarrow H^1(K, V)$ .

**Lemma 9.7.** *If  $r$  is large enough and  $\gamma \in \Gamma_K$  then*

$$1 - \gamma : \mathbf{D}_{\text{rig}}^{\dagger,r}(V)^{\psi=0} \rightarrow \mathbf{D}_{\text{rig}}^{\dagger,r}(V)^{\psi=0}$$

*is an isomorphism*

*Proof.* We first show that  $1 - \gamma$  is injective. By theorem 9.5, an element in the kernel of  $1 - \gamma$  would be in  $\mathbf{D}_{\text{cris}}(V)$  and therefore in  $\mathbf{D}_{\text{cris}}(V)^{\psi=0}$ , which is obviously 0.

To prove surjectivity. Recall that by iii) of proposition 6.5, if  $r$  is large enough and  $\gamma \in \Gamma_K$  then  $1 - \gamma : \mathbf{D}_{\text{rig}}^{\dagger,r}(V)^{\psi=0} \rightarrow \mathbf{D}_{\text{rig}}^{\dagger,r}(V)^{\psi=0}$  is an isomorphism whose inverse is uniformly continuous for the Fréchet topology of  $\mathbf{D}_{\text{rig}}^{\dagger,r}(V)$ .

In order to show the surjectivity of  $1 - \gamma$  it is therefore enough to show that  $\mathbf{D}_{\text{rig}}^{\dagger,r}(V)^{\psi=0}$  is dense in  $\mathbf{D}_{\text{rig}}^{\dagger,r}(V)^{\psi=0}$  for the Fréchet topology. For  $r$  large enough,  $\mathbf{D}_{\text{rig}}^{\dagger,r}(V)$  has a basis in  $\varphi(\mathbf{D}_{\text{rig}}^{\dagger,r/p}(V))$  so that

$$\begin{aligned} \mathbf{D}_{\text{rig}}^{\dagger,r}(V)^{\psi=0} &= (\mathbf{B}_K^{\dagger,r})^{\psi=0} \cdot \varphi(\mathbf{D}_{\text{rig}}^{\dagger,r/p}(V)) \\ \mathbf{D}_{\text{rig}}^{\dagger,r}(V)^{\psi=0} &= (\tilde{\mathbf{B}}_{\text{rig},K}^{\dagger,r})^{\psi=0} \cdot \varphi(\mathbf{D}_{\text{rig}}^{\dagger,r/p}(V)). \end{aligned}$$

The fact that  $\mathbf{D}_{\text{rig}}^{\dagger,r}(V)^{\psi=0}$  is dense in  $\mathbf{D}_{\text{rig}}^{\dagger,r}(V)^{\psi=0}$  for the Fréchet topology will therefore follow from the density of  $(\mathbf{B}_K^{\dagger,r})^{\psi=0}$  in  $(\tilde{\mathbf{B}}_{\text{rig},K}^{\dagger,r})^{\psi=0}$ . The last statement follows from the facts that by definition  $\mathbf{B}_K^{\dagger,r/p}$  is dense in  $\tilde{\mathbf{B}}_{\text{rig},K}^{\dagger,r/p}$  and that

$$(\mathbf{B}_K^{\dagger,r})^{\psi=0} = \bigoplus_{i=1}^{p-1} [\varepsilon]^i \varphi(\mathbf{B}_K^{\dagger,r/p}) \quad \text{and} \quad (\tilde{\mathbf{B}}_{\text{rig},K}^{\dagger,r})^{\psi=0} = \bigoplus_{i=1}^{p-1} [\varepsilon]^i \varphi(\tilde{\mathbf{B}}_{\text{rig},K}^{\dagger,r/p}).$$

□

**Lemma 9.8.** *The following maps are all surjective and the kernel is  $\mathbf{Q}_p$*

$$1 - \varphi : \tilde{\mathbf{B}}^{\dagger} \rightarrow \tilde{\mathbf{B}}^{\dagger}, \quad 1 - \varphi : \tilde{\mathbf{B}}_{\text{rig}}^+ \rightarrow \tilde{\mathbf{B}}_{\text{rig}}^+ \quad \text{and} \quad 1 - \varphi : \tilde{\mathbf{B}}_{\text{rig}}^{\dagger} \rightarrow \tilde{\mathbf{B}}_{\text{rig}}^{\dagger}$$

*Proof.* Since  $\tilde{\mathbf{B}}_{\text{rig}}^+ \subset \mathbf{B}_{\text{rig}}^+$  and  $\tilde{\mathbf{B}}^{\dagger} \subset \mathbf{B}_{\text{rig}}^{\dagger}$  it is enough to show that  $(\tilde{\mathbf{B}}_{\text{rig}}^{\dagger})^{\varphi=1} = \mathbf{Q}_p$ . If  $x \in (\tilde{\mathbf{B}}_{\text{rig}}^{\dagger})^{\varphi=1}$ , then [Ber02, proposition 3.2] shows that actually  $x \in (\tilde{\mathbf{B}}_{\text{rig}}^+)^{\varphi=1}$ , and therefore  $x \in (\tilde{\mathbf{B}}_{\text{rig}}^+)^{\varphi=1} = (\mathbf{B}_{\text{max}}^+)^{\varphi=1} = \mathbf{Q}_p$  by [Col98, proposition III 3.5].

The surjectivity of  $1 - \varphi : \tilde{\mathbf{B}}_{\text{rig}}^+ \rightarrow \tilde{\mathbf{B}}_{\text{rig}}^+$  results from the surjectivity of  $1 - \varphi$  on the first two spaces since by [Ber02, lemma 2.18], one can write  $\alpha \in \tilde{\mathbf{B}}_{\text{rig}}^+$  as  $\alpha = \alpha^+ + \alpha^-$  with  $\alpha^+ \in \tilde{\mathbf{B}}_{\text{rig}}^+$  and  $\alpha^- \in \tilde{\mathbf{B}}^{\dagger}$ .

The surjectivity of  $1 - \varphi : \tilde{\mathbf{B}}_{\text{rig}}^+ \rightarrow \tilde{\mathbf{B}}_{\text{rig}}^+$  follows from the facts that  $1 - \varphi : \mathbf{B}_{\text{max}}^+ \rightarrow \mathbf{B}_{\text{max}}^+$  is surjective ([Col98, proposition III 3.1]) and that  $\tilde{\mathbf{B}}_{\text{rig}}^+ = \cap_{n=0}^{\infty} \varphi^n(\mathbf{B}_{\text{max}}^+)$ .

The surjectivity of  $1 - \varphi : \tilde{\mathbf{B}}^\dagger \rightarrow \tilde{\mathbf{B}}^\dagger$  follows from the facts that  $1 - \varphi : \tilde{\mathbf{B}} \rightarrow \tilde{\mathbf{B}}$  is surjective (it is surjective on  $\tilde{\mathbf{A}}$  as can be seen by reducing modulo  $p$  and using the fact that  $\tilde{\mathbf{E}}$  is algebraically closed) and that if  $\beta \in \tilde{\mathbf{B}}$  is such that  $(1 - \varphi)\beta \in \tilde{\mathbf{B}}^\dagger$ , then  $\beta \in \tilde{\mathbf{B}}^\dagger$ .

If  $x = \sum_{i=0}^{+\infty} p^i[x_i] \in \tilde{\mathbf{A}}$ , let us set  $w_k(x) = \inf_{i \leq k} \nu_E(x_i) \in \mathbb{R} \cup \{+\infty\}$ . The definition of  $\tilde{\mathbf{B}}^{\dagger,r}$  shows that  $x \in \tilde{\mathbf{B}}^{\dagger,r}$  if and only if  $\lim_{k \rightarrow +\infty} w_k(x) + \frac{pr}{p-1}k = +\infty$ . A short computation shows that  $w_k(\varphi(x)) = pw_k(x)$  and that  $w_k(x+y) \geq \inf(w_k(x), w_k(y))$  with equality if  $w_k(x) \neq w_k(y)$ .

It is then clear that

$$\lim_{k \rightarrow +\infty} w_k((1 - \varphi)x) + \frac{pr}{p-1}k = +\infty \implies \lim_{k \rightarrow +\infty} w_k(x) + \frac{p(r/p)}{p-1}k = +\infty$$

and so if  $x \in \tilde{\mathbf{A}}$  is such that  $(1 - \varphi)x \in \tilde{\mathbf{B}}^{\dagger,r}$  then  $x \in \tilde{\mathbf{B}}^{\dagger,r/p}$  and likewise for  $x \in \tilde{\mathbf{B}}$  by multiplication by a suitable power of  $p$ . This shows the second fact.  $\square$

**Proposition 9.9.** *If  $y \in \mathbf{D}_{\text{rig}}^\dagger(V)^{\psi=1}$  and  $\Gamma_K$  is torsion free, there exists  $b \in \tilde{\mathbf{B}}_{\text{rig}}^\dagger \otimes_{\mathbf{Q}_p} V$  such that  $(\gamma - 1)(\varphi - 1)b = (\varphi - 1)y$  and the formula*

$$h_{K,V}^1(y) = \log_p^0(\gamma)[\sigma \mapsto \frac{\sigma - 1}{\gamma - 1}y - (\sigma - 1)b]$$

*then defines a map  $h_{K,V}^1 : \mathbf{D}_{\text{rig}}^\dagger(V)_{\Gamma_K}^{\psi=1} \mapsto H^1(K, V)$  which does not depend either on the choice of generator  $\gamma$  of  $\Gamma_K$  or on the particular solution  $b$ , and if  $y \in D(V)^{\psi=1} \subset \mathbf{D}_{\text{rig}}^\dagger(V)^{\psi=1}$ , then  $h_{K,V}^1(y)$  coincides with the cocycle constructed in section 4.2.*

*Proof.* Our construction closely follows section 4.2; to simplify the notations, we may assume that  $\log_p^0(\gamma) = 1$ . The fact that  $h_{K,V}^1$  is independent of the choice of  $\gamma$  is same as lemma 4.2.

Let us start by showing the existence of  $b \in \tilde{\mathbf{B}}_{\text{rig}}^\dagger \otimes_{\mathbf{Q}_p} V$ . If  $y \in \mathbf{D}_{\text{rig}}^\dagger(V)^{\psi=1}$ , then  $(\varphi - 1)y \in \mathbf{D}_{\text{rig}}^\dagger(V)^{\psi=0}$ . By lemme 9.7, there exists  $x \in \mathbf{D}_{\text{rig}}^\dagger(V)^{\psi=0}$  such that  $(\gamma - 1)x = (\varphi - 1)y$ . By lemma 9.8, there exists  $b \in \tilde{\mathbf{B}}_{\text{rig}}^\dagger \otimes_{\mathbf{Q}_p} V$  such that  $(\varphi - 1)b = x$ .

Recall that we define  $h_{K,V}^1(y) \in H^1(K, V)$  by the formula:

$$h_{K,V}^1(y)(\sigma) = \frac{\sigma - 1}{\gamma - 1}y - (\sigma - 1)b.$$

Notice that, a priori,  $h_{K,V}^1(y) \in H^1(K, \tilde{\mathbf{B}}_{\text{rig}}^\dagger \otimes_{\mathbf{Q}_p} V)$ , but

$$\begin{aligned} (\varphi - 1)h_{K,V}^1(y)(\sigma) &= \frac{\sigma - 1}{\gamma - 1}(\varphi - 1)y - (\sigma - 1)(\varphi - 1)b \\ &= \frac{\sigma - 1}{\gamma - 1}(\gamma - 1)x - (\sigma - 1)x \\ &= 0, \end{aligned}$$

so that  $h_{K,V}^1(y)(\sigma) \in (\mathbf{B}_{\text{rig}}^\dagger)^{\varphi=1} \otimes_{\mathbf{Q}_p} V = V$ . In addition, two different choices of  $b$  differ by an element of  $(\tilde{\mathbf{B}}_{\text{rig}}^\dagger)^{\varphi=1} \otimes_{\mathbf{Q}_p} V = V$ , and therefore give rise to two cohomologous cocycles.

It is clear that if  $y \in D(V)^{\psi=1} \subset \mathbf{D}_{\text{rig}}^\dagger(V)^{\psi=1}$ , then  $h_{K,V}^1$  coincide with the cocycle constructed in section 4.2, as can be seen by their identical construction, and it is immediate that if  $y \in (\gamma - 1)\mathbf{D}_{\text{rig}}^\dagger(V)$ , then  $h_{K,V}^1(y) = 0$ .  $\square$

**Lemma 9.10.** *We have  $\text{cor}_{K_{n+1}/K_n} \circ h_{K_{n+1},V}^1 = h_{K_n,V}^1$ .*

*Proof.* Same as lemma 5.3.  $\square$

**9.5. Iwasawa algebra and power series.** Given a finite unramified extension  $F$  of  $\mathbf{Q}_p$ , denote by  $\Lambda(\Gamma_F)$  (resp.  $\Lambda(\Gamma_F^1)$ ) the Iwasawa algebra  $\mathbf{Z}_p[[\Gamma_F]]$  (resp.  $\mathbf{Z}_p[[\Gamma_F^1]]$ ).

Let

$$\mathcal{H} = \{f \in \mathbf{Q}_p[[\Delta]][[X]] \mid f \text{ converges on the open unit disk}\},$$

and define  $\mathcal{H}(\Gamma_F)$  to be the set of  $f(\gamma - 1)$  with  $f(X) \in \mathcal{H}$  and  $\gamma$  a topological generator of  $\Gamma$ . We may identify  $\Lambda(\Gamma_F) \otimes \mathbf{Q}_p$  with the subring of  $\mathcal{H}(\Gamma_F)$  consisting of power series with bounded coefficients. Note that  $\mathcal{H}(\Gamma)$  may be identified with the continuous dual of the space of locally analytic functions on  $\Gamma_F$ , with multiplication corresponding to convolution, implying that its definition is independent of the choice of generator  $\gamma_F$  (c.f. section 1.2).

The action of  $\Gamma_F$  on  $\mathbf{B}_{\text{rig},\mathbf{Q}_p}^+$  gives an isomorphism of  $\mathcal{H}(\Gamma_F)$  with  $(\mathbf{B}_{\text{rig},\mathbf{Q}_p}^+)^{\psi=0}$  via the Mellin transform [Per01, corollary B.2.8]

$$\begin{aligned} \mathfrak{M} : \mathcal{H}(\Gamma_F) &\rightarrow (\mathbf{B}_{\text{rig},\mathbf{Q}_p}^+)^{\psi=0} \\ f(\gamma - 1) &\mapsto f(\gamma - 1)(\pi + 1). \end{aligned}$$

In particular,  $\Lambda(\Gamma_F)$  corresponds to  $(\mathbf{A}_{\mathbf{Q}_p}^+)^{\psi=0}$  under  $\mathfrak{M}$ . Similarly, we define  $\mathcal{H}(\Gamma_F^1)$  as the subring of  $\mathcal{H}(\Gamma_F)$  defined by power series over  $\mathbf{Q}_p$ , rather than  $\mathbf{Q}_p[[\Delta]]$ . Then,  $\mathcal{H}(\Gamma_F^1)$  (resp.  $\Lambda(\Gamma_F^1)$ ) corresponds to  $(1 + \pi)\varphi(\mathbf{B}_{\text{rig},\mathbf{Q}_p}^+)$  (resp.  $(1 + \pi)\varphi(\mathbf{A}_{\mathbf{Q}_p}^+)$ ) under  $\mathfrak{M}$ .

**9.6. Iwasawa algebras and differential equations.** By [Ber02, proposition 2.24], we have maps  $\varphi^{-n} : \widetilde{\mathbf{B}}_{\text{rig}}^{\dagger,r_n} \rightarrow \mathbf{B}_{\text{dR}}^+$  whose restriction to  $\mathbf{B}_{\text{rig},F}^+$  satisfied  $\varphi^{-n}(\mathbf{B}_{\text{rig},F}^+) \subset F_n[[t]]$  and which can be characterized by the fact that  $\pi$  maps to  $\varepsilon^{(n)} \exp(t/p^n) - 1$ .

Recall if  $z \in F_n((t)) \otimes_F \mathbf{D}_{\text{cris}}(V)$ , we denote the constant coefficient of  $z$  by  $\partial_V(z) \in F_n \otimes_F \mathbf{D}_{\text{cris}}(V)$ .

**Lemma 9.11.** *If  $y \in (\mathbf{B}_{\text{rig},F}^+[1/t] \otimes_F \mathbf{D}_{\text{cris}}(V))^{\psi=1}$ , then for any  $m \geq n \geq 0$ , the element*

$$p^{-m} \text{Tr}_{F_m/F_n} \partial_V(\varphi^{-m}(y)) \in F_n \otimes \mathbf{D}_{\text{cris}}(V)$$

*does not depend on  $m$  and we have*

$$p^{-m} \text{Tr}_{F_m/F_n} \partial_V(\varphi^{-m}(y)) = \begin{cases} p^{-n} \partial_V(\varphi^{-n}(y)) & \text{if } n \geq 1 \\ (1 - p^{-1} \varphi^{-1}) \partial_V(y) & \text{if } n = 0 \end{cases}$$

*Proof.* Recall that if  $y = t^{-l} \sum_{k=0}^{+\infty} a_k \pi^k \in \mathbf{B}_{\text{rig},F}^+[1/t] \otimes_F \mathbf{D}_{\text{cris}}(V)$ , then

$$\varphi^{-m}(y) = p^{ml} t^{-l} \sum_{k=0}^{+\infty} \varphi^{-m}(a_k) (\varepsilon^{(m)} \exp(t/p^m) - 1)^k,$$

and that by the definition of  $\psi$ ,  $\psi(y) = y$  means that:

$$\varphi(y) = \frac{1}{p} \sum_{\zeta^p=1} y(\zeta(1+T) - 1).$$

The lemma then follows from the fact that if  $m \geq 2$ , then the conjugates of  $\varepsilon^{(m)}$  under  $\text{Gal}(F_m/F_{m-1})$  are the  $\zeta \varepsilon^{(m)}$ , where  $\zeta^p = 1$ , while if  $m = 1$ , then the conjugates of  $\varepsilon^{(1)}$  under  $\text{Gal}(F_1/F)$  are the  $\zeta$ , where  $\zeta^p = 1$  but  $\zeta \neq 1$ .  $\square$

Recall that since  $F$  is an unramified extension of  $\mathbf{Q}_p$ ,  $\Gamma_F \simeq \mathbf{Z}_p^*$  and that  $\Gamma_{F_n} = \text{Gal}(F_\infty/F_n)$  is the set of elements  $\gamma \in \Gamma_F$  such that  $\chi(\gamma) \in 1 + p^n \mathbf{Z}_p$ .

The Iwasawa algebra of  $\Gamma_F$  is  $\Lambda_{\mathcal{O}_F} = \mathbf{Z}_p[[\Gamma_F]] \cong \mathbf{Z}_p[\Delta_F] \otimes_{\mathbf{Z}_p} \mathbf{Z}_p[[\Gamma_{F_1}]]$ , and we set  $\mathcal{H}(\Gamma_F) = \mathbf{Q}_p[\Delta_F] \otimes_{\mathbf{Q}_p} \mathcal{H}(\Gamma_F^1)$  where  $\mathcal{H}(\Gamma_F^1)$  is the set of  $f(\gamma - 1)$  with  $\gamma \in \Gamma_F^1$  and where  $f(X) \in \mathbf{Q}_p[[X]]$  is convergent on the  $p$ -adic open unit disk. We define  $\nabla_i \in \mathcal{H}(\Gamma_F)$  by

$$\nabla_i = \frac{\log(\gamma)}{\log_p(\chi(\gamma))} - i.$$

We will also use the operator  $\nabla_0/(\gamma_n - 1)$ , where  $\gamma_n$  is a topological generator of  $\Gamma_F^n$ . It is defined by the formula

$$\frac{\nabla_0}{\gamma_n - 1} = \frac{\log(\gamma_n)}{\log_p(\chi(\gamma_n))(\gamma_n - 1)} = \frac{1}{\log_p(\chi(\gamma_n))} \sum_{i \geq 1} \frac{(1 - \gamma_n)^{i-1}}{i},$$

or equivalently by

$$\frac{\nabla_0}{\gamma_n - 1} = \lim_{\eta \in \Gamma_F^n, \eta \rightarrow 1} \frac{\eta - 1}{\gamma_n - 1} \frac{1}{\log_p(\chi(\eta))}.$$

It is easy to see that  $\nabla_0/(\gamma_n - 1)$  acts on  $F_n$  by  $1/\log_p(\chi(\gamma_n))$ .

The algebra  $\mathcal{H}(\Gamma_F)$  acts on  $\mathbf{B}_{\text{rig},F}^+$  and one can easily check that

$$\nabla_i = t \frac{d}{dt} - i = \log(1 + \pi) \partial - i, \quad \text{where } \partial = (1 + \pi) \frac{d}{d\pi}.$$

In particular,  $\nabla_0 \mathbf{B}_{\text{rig},F}^+ \subset t \mathbf{B}_{\text{rig},F}^+$  and if  $i \geq 1$ , then

$$\nabla_{i-1} \circ \cdots \circ \nabla_0 \subset t^i \mathbf{B}_{\text{rig},F}^+.$$

**Lemma 9.12.** *If  $n \geq 1$ , then  $\nabla_0/(\gamma_n - 1)(\mathbf{B}_{\text{rig},F}^+)^{\psi=0} \subset (t/\varphi^n(\pi))(\mathbf{B}_{\text{rig},F}^+)^{\psi=0}$  so that if  $i \geq 1$ , then*

$$\nabla_{i-1} \circ \cdots \circ \nabla_1 \circ \frac{\nabla_0}{\gamma_n - 1} (\mathbf{B}_{\text{rig},F}^+)^{\psi=0} \subset \left(\frac{t}{\varphi^n(\pi)}\right)^i (\mathbf{B}_{\text{rig},F}^+)^{\psi=0}.$$

*Proof.* Since  $\nabla_i = t \cdot d/dt - i$ , the second claim follows easily from the first one. By the standard properties of  $p$ -adic holomorphic functions, what we need to do is to show that if  $x \in (\mathbf{B}_{\text{rig},F}^+)^{\psi=0}$ , then

$$\frac{\nabla_0}{\gamma_n - 1} x(\varepsilon^{(m)} - 1) = 0$$

for all  $m \geq n + 1$ .

On the other hand, up to a scalar factor, one has for  $m \geq n + 1$ :

$$\frac{\nabla_0}{\gamma_n - 1} x(\varepsilon^{(m)} - 1) = \text{Tr}_{F_m/F_n} x(\varepsilon^{(m)} - 1),$$

which can be seen from the fact that

$$\frac{\nabla_0}{\gamma_n - 1} = \lim_{\eta \in \Gamma_F^n, \eta \rightarrow 1} \frac{\eta - 1}{\gamma_n - 1} \cdot \frac{1}{\log_p(\chi(\eta))}.$$

On the other hand, the fact  $\psi(x) = 0$  implies that for every  $m \geq 2$ ,  $\text{Tr}_{F_m/F_{m-1}} x(\varepsilon^{(m)} - 1) = 0$ . This completes the proof.  $\square$

Finally, let us point out that the actions of any element of  $\mathcal{H}(\Gamma_F)$  and  $\varphi$  commute. Since  $\varphi(t) = pt$ , we also see that  $\partial \circ \varphi = p\varphi \circ \partial$ .

We will henceforth assume that  $\log_p(\chi(\gamma_n)) = p^n$ , and in addition  $\nabla_0/(\gamma_n - 1)$  acts on  $F_n$  by  $p^{-n}$ .

## 10. BLOCH-KATO'S EXPONENTIAL MAPS: THREE EXPLICIT RECIPROCITY FORMULAS

In this section, we explain the results of Berger in [Ber03] on explicit reciprocity formulas when  $V$  is a crystalline representation of an unramified field.

Recall  $H_K = \text{Gal}(\overline{\mathbf{Q}_p}/K_\infty)$ , let  $\Delta_K$  be the torsion subgroup of  $\Gamma_K = G_K/H_K = \text{Gal}(K_\infty/K)$  and let  $\Gamma_K^1 = \text{Gal}(K_\infty/K(\mu_p))$ , so that  $\Gamma_K \simeq \Delta_K \times \Gamma_K^1$ . Let  $\Gamma_K = \mathbf{Z}_p[[\Gamma_K]]$  and  $\mathcal{H}(\Gamma_K) = \mathbf{Q}_p[\Delta_K] \otimes_{\mathbf{Q}_p} \mathcal{H}(\Gamma_K^1)$  where  $\mathcal{H}(\Gamma_K^1)$  is the set of  $f(\gamma_1 - 1)$  with  $\gamma_1 \in \Gamma_{K_1}$  and where  $f(T) \in \mathbf{Q}_p[[T]]$  is a power series which converges on the  $p$ -adic unit disk.

When  $F$  is an unramified extension of and  $V$  is a crystalline representation of  $G_F$ , Perrin-Riou has constructed in [Per94] a period map  $\Omega_{V,h}$  which interpolates the  $\exp_{F,V(k)}$  as  $k$  runs over the positive integers. It is crucial ingredient in the construction of  $p$ -adic  $L$ -funtions, and is a vast generalization of Coleman's isomorphism.

The main result of [Per94] is the construction, for a crystalline representation of  $V$  of  $G_F$  of a family of maps (parameterized by  $h \in \mathbf{Z}$ ):

$$\Omega_{V,h} : (\mathcal{H}(\Gamma_F) \otimes_{\mathbf{Q}_p} \mathbf{D}_{\text{cris}}(V))^{\Delta=0} \rightarrow \mathcal{H}(\Gamma_F) \otimes_{\Lambda_F} H_{\text{Iw}}^1(F, V)/V^{H_F},$$

whose main property is that they interpolate Bloch-Kato's exponential map. More precisely, if  $h, j \gg 0$ , then the diagram:

$$\begin{array}{ccc} (\mathcal{H}(\Gamma_F) \otimes_{\mathbf{Q}_p} \mathbf{D}_{\text{cris}}(V(j)))^{\Delta=0} & \xrightarrow{\Omega_{V(j),h}} & \mathcal{H}(\Gamma_F) \otimes_{\Gamma_F} H_{\text{Iw}}^1(F, V(j))/V(j)^{H_F} \\ \Xi_{n,V(j)} \downarrow & & \downarrow \text{pr}_{F_n, V(j)} \\ F_n \otimes_F \mathbf{D}_{\text{cris}}(V) & \xrightarrow{(h+j-1)! \times \exp_{F_n, V(j)}^*} & H^1(F_n, V(j)). \end{array}$$

is commutative where  $\Delta$  and  $\Xi$  are two maps whose definition is rather technical (see section 10.2 for a precisely definition).

Using the inverse of Perrin-Riou's map, one can then associate to an Euler system a  $p$ -adic  $L$ -function. For example, if one starts with  $V = \mathbf{Q}_p(1)$ , then Perrin-Riou's map is the inverse of the Coleman isomorphism and one recovers Kubota-Leopoldt  $p$ -adic  $L$ -functions (See section 11.2).

The goal of this section is to give formulas for  $\exp_{K,V}$ ,  $\exp_{K,V^*(1)}^*$  and  $\Omega_{V,h}$  in terms of the  $(\varphi, \Gamma)$ -module associated to  $V$ .

**10.1. The Bloch-Kato's exponential map and its dual revisit.** Recall in section 7.1, we defined the Bloch-Kato's exponential map and its dual. The goal of this paragraph is to compute Bloch-Kato's exponential map and its dual in terms of the  $(\varphi, \Gamma)$ -module of  $V$ . Let  $h \geq 1$  be an integer such that  $\text{Fil}^{-h} \mathbf{D}_{\text{cris}}(V) = \mathbf{D}_{\text{cris}}(V)$ .

Recall that we have seen that  $\mathbf{D}_{\text{cris}}(V) = (\mathbf{D}_{\text{rig}}^+[1/t])^{\Gamma_F}$  and by [Ber04, II.3], there is an isomorphism

$$\mathbf{B}_{\text{rig}, F}^+[1/t] \otimes_F \mathbf{D}_{\text{cris}}(V) = \mathbf{B}_{\text{rig}, F}^+[1/t] \otimes_F \mathbf{D}_{\text{rig}}^+(V).$$

If  $y \in \mathbf{B}_{\text{rig},F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V)$ , then the fact that  $\text{Fil}^{-h} \mathbf{D}_{\text{cris}}(V) = \mathbf{D}_{\text{cris}}(V)$  implies by result of [Ber04, II.3] that  $t^h y \in \mathbf{D}_{\text{rig}}^+(V)$ , so that if

$$y = \sum_{i=0}^d y_i \otimes d_i \in (\mathbf{B}_{\text{rig},F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V))^{\psi=1},$$

then

$$\nabla_{h-1} \circ \cdots \nabla_0(y) = \sum_{i=0}^d t^h \partial^h y_i \otimes d_i \in \mathbf{D}_{\text{rig}}^+(V)^{\psi=1}.$$

One can apply the operator  $h_{F_n,V}^1$  to  $\nabla_{h-1} \circ \cdots \nabla_0(y)$ , then we have:

**Theorem 10.1.** *If  $y \in (\mathbf{B}_{\text{rig},F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V))^{\psi=1}$ , then*

$$h_{F_n,V}^1(\nabla_{h-1} \circ \cdots \nabla_0(y)) = (-1)^{h-1} (h-1)! \begin{cases} \exp_{F_n,V}(p^{-n} \partial_V(\varphi^{-n}(y))) & \text{if } n \geq 1 \\ \exp_{F,V}((1-p^{-1}\varphi^{-1})\partial_V(y)) & \text{if } n = 0 \end{cases}$$

*Proof.* Because the diagram

$$\begin{array}{ccc} F_{n+1} \otimes_F \mathbf{D}_{\text{cris}}(V) & \xrightarrow{\exp_{F_{n+1},V}} & H^1(F_{n+1}, V) \\ \text{Tr}_{F_{n+1}/F_n} \otimes id \downarrow & & \downarrow \text{cor}_{F_{n+1}/F_n} \\ F_n \otimes \mathbf{D}_{\text{cris}}(V) & \xrightarrow{\exp_{F_n,V}} & H^1(F_n, V) \end{array}$$

is commutative, it is enough to prove the theorem under the assumption that  $\Gamma_F^n$  is torsion free. Let us set  $y_h = \nabla_{h-1} \circ \cdots \nabla_0(y)$ . Since we are assuming for simplicity that  $\chi(\gamma_n) = p^n$ , the cocycle  $h_{F_n,V}^1(y_h)$  is defined by:

$$h_{F_n,V}^1(y_h)(\sigma) = \frac{\sigma-1}{\gamma_n-1} y_h - (\sigma-1) b_{n,h}$$

where  $b_{n,h}$  is a solution of the equation  $(\gamma_n-1)(\varphi-1)b_{n,h} = (\varphi-1)y_h$ . In lemma 9.12 above, we prove that

$$\nabla_{i-1} \circ \cdots \nabla_1 \circ \frac{\nabla_0}{\gamma_n-1} (\mathbf{B}_{\text{rig},F}^+)^{\psi=0} \subset \left(\frac{t}{\varphi^n(\pi)}\right)^i (\mathbf{B}_{\text{rig},F}^+)^{\psi=0}.$$

It is then clear that if one sets

$$z_{n,h} = \nabla_{h-1} \circ \cdots \nabla_1 \circ \frac{\nabla_0}{\gamma_n-1} (\varphi-1)y,$$

then

$$z_{n,h} \in \left(\frac{t}{\varphi^n(\pi)}\right)^h (\mathbf{B}_{\text{rig},F}^+)^{\psi=0} \otimes_F \mathbf{D}_{\text{cris}}(V) \subset \varphi^n(\pi^{-h}) \mathbf{D}_{\text{rig}}^+(V)^{\psi=0} \subset \mathbf{D}_{\text{rig}}^+(V)^{\psi=0}.$$

Let  $q = \varphi(\pi)/\pi$ . By lemma 10.2 below, there exists an element  $b_{n,h} \in \varphi^{n-1}(\pi^{-h}) \widetilde{\mathbf{B}}_{\text{rig}}^+ \otimes_{\mathbf{Q}_p} V$  such that

$$(\varphi - \varphi^{n-1}(q^h))(\varphi^{n-1}(\pi^h) b_{n,h}) = \varphi^n(\pi^h) z_{n,h},$$

so that  $(1-\varphi)b_{n,h} = z_{n,h}$  with  $b_{n,h} \in \varphi^{n-1}(\pi^{-h}) \widetilde{\mathbf{B}}_{\text{rig}}^+ \otimes_{\mathbf{Q}_p} V$ .

If we set  $w_{n,h} = \nabla_{h-1} \circ \cdots \nabla_1 \circ \frac{\nabla_0}{\gamma_n-1} y$ , then  $w_{n,h}$  and  $b_{n,h} \in \mathbf{B}_{\text{max}} \otimes_{\mathbf{Q}_p} V$  and the cocycle  $h_{F_n,V}^1(y_h)$  is then given by the formula  $h_{F_n,V}^1(y_h)(\sigma) = (\sigma-1)(w_{n,h} - b_{n,h})$ . Now  $(\varphi-1)b_{n,h} = z_{n,h}$  and  $(\varphi-1)w_{n,h} = z_{n,h}$  as well, so that  $w_{n,h} - b_{n,h} \in \mathbf{B}_{\text{max}}^{\varphi=1} \otimes_{\mathbf{Q}_p} V$ .



We can also write

$$h_{F_n, V}^1(y_h)(\sigma) = (\sigma - 1)(\varphi^{-n}(w_{n, h}) - \varphi^{-n}(b_{n, h})).$$

Since we know that  $b_{n, h} \in \varphi^{n-1}(\pi^{-h})\mathbf{B}_{\max}^+ \otimes_{\mathbf{Q}_p} V$ , we have  $\varphi^{-n}(b_{n, h}) \in \mathbf{B}_{\mathrm{dR}}^+ \otimes_{\mathbf{Q}_p} V$ .

The definition of Bloch-Kato exponential gives rise to the following construction: if  $x \in \mathbf{D}_{\mathrm{dR}}(V)$  and  $\tilde{x} \in \mathbf{B}_{\max}^{\varphi=1} \otimes_{\mathbf{Q}_p} V$  is such that  $x - \tilde{x} \in \mathbf{B}_{\mathrm{dR}}^+ \otimes_{\mathbf{Q}_p} V$  then  $\exp_{K, V}(x)$  is the class of the cocycle  $g \mapsto g(\tilde{x}) - \tilde{x}$ .

The theorem therefore follow from the fact that:

$$\varphi^{-n}(w_{n, h}) - (-1)^{h-1}(h-1)!p^{-n}\partial_V(\varphi^{-n}(y)) \in \mathbf{B}_{\mathrm{dR}}^+ \otimes_{\mathbf{Q}_p} V,$$

since we already know that  $\varphi^{-n}(b_{n, h}) \in \mathbf{B}_{\mathrm{dR}}^+ \otimes_{\mathbf{Q}_p} V$ .

In order to show this, first notice that

$$\varphi^{-n}(y) - \partial_V(\varphi^{-n}(y)) \in tF_n[[t]] \otimes_F \mathbf{D}_{\mathrm{cris}}(V).$$

We can therefore write

$$\frac{\nabla_0}{\gamma_n - 1} \varphi^{-n}(y) = p^{-n} \partial_V(\varphi^{-n}(y)) + tz_1$$

and a simple recurrence shows that

$$\nabla_{i-1} \circ \cdots \circ \frac{\nabla_0}{1 - \gamma_n} \varphi^{-n}(y) = (-1)^{i-1}(i-1)!p^{-n} \partial_V(\varphi^{-n}(y)) + t^i z_i,$$

with  $z_i \in F_n[[t]] \otimes_F \mathbf{D}_{\mathrm{cris}}(V)$ . By taking  $i = h$ , we see that

$$\varphi^{-n}(w_{n, h}) - (-1)^{h-1}(h-1)!p^{-n} \partial_V(\varphi^{-n}(y)) \in \mathbf{B}_{\mathrm{dR}}^+ \otimes_{\mathbf{Q}_p} V,$$

Since we choose  $h$  such that  $t^h \mathbf{D}_{\mathrm{cris}}(V) \subset \mathbf{B}_{\mathrm{dR}}^+ \otimes_{\mathbf{Q}_p} V$ . □

**Lemma 10.2.** *If  $\alpha \in \tilde{\mathbf{B}}_{\mathrm{rig}}^+$ , then there exists  $\beta \in \tilde{\mathbf{B}}_{\mathrm{rig}}^+$  such that*

$$(\varphi - \varphi^{n-1}(q^h))\beta = \alpha.$$

*Proof.* By [Ber02, proposition 2.19], the ring  $\tilde{\mathbf{B}}^+$  is dense in  $\tilde{\mathbf{B}}_{\mathrm{rig}}^+$  for the Fréchet topology. Hence, if  $\alpha \in \tilde{\mathbf{B}}_{\mathrm{rig}}^+$ , then there exists  $\alpha_0 \in \tilde{\mathbf{B}}^+$  such that  $\alpha - \alpha_0 = \varphi^n(\pi^h)\alpha_1$  with  $\alpha_1 \in \tilde{\mathbf{B}}_{\mathrm{rig}}^+$ .

The map  $\varphi - \varphi^{n-1}(q^h) : \tilde{\mathbf{B}}^+ \rightarrow \tilde{\mathbf{B}}^+$  is surjective because  $\varphi - \varphi^{n-1}(q^h) : \tilde{\mathbf{A}}^+ \rightarrow \tilde{\mathbf{A}}^+$  is surjective, as can be seen by reducing modulo  $p$  using the fact that  $\tilde{\mathbf{E}}$  is algebraically closed and that  $\tilde{\mathbf{E}}^+$  is its ring of integers.

One can therefore write  $\alpha_0 = (\varphi - \varphi^{n-1}(q^h))\beta_0$ . Finally by lemma 9.8, there exists  $\beta \in \tilde{\mathbf{B}}_{\mathrm{rig}}^+$  such that  $\alpha_1 = (\varphi - 1)\beta_1$ , so that  $\varphi^n(\pi^h)\alpha_1 = (\varphi - \varphi^{n-1}(q^h))(\varphi^{n-1}(\pi^h)\beta_1)$ . □

**Theorem 10.3.** *If  $y \in (\mathbf{D}_{\mathrm{rig}}^\dagger(V))^{\psi=1}$  and  $y \in \mathbf{D}_{\mathrm{rig}}^+(V)[1/t]$  (so that in particular  $y \in (\mathbf{B}_{\mathrm{rig}, F}^+[1/t] \otimes_F \mathbf{D}_{\mathrm{cris}}(V))^{\psi=1}$ ), then*

$$\exp_{F_n, V^*(1)}^*(h_{F_n, V}^1(y)) = \begin{cases} p^{-n} \partial_V(\varphi^{-n}(y)) & \text{if } n \geq 1 \\ (1 - p^{-1} \varphi^{-1}) \partial_V(y) & \text{if } n = 0 \end{cases}$$

*Proof.* Since the following diagram

$$\begin{array}{ccc} H^1(F_{n+1}, V) & \xrightarrow{\exp_{F_{n+1}, V^*(1)}^*} & F_{n+1} \otimes \mathbf{D}_{\text{cris}}(V) \\ \text{cor}_{F_{n+1}/F_n} \downarrow & & \downarrow \text{Tr}_{F_{n+1}/F_n} \otimes \text{id} \\ H^1(F_n, V) & \xrightarrow{\exp_{F_n, V^*(1)}^*} & F_n \otimes \mathbf{D}_{\text{cris}}(V) \end{array}$$

is commutative, we only need to prove the theorem when  $\Gamma_F^n$  is torsion free by lemma 10.1. We then have (assuming that  $\chi(\gamma_n) = p^n$  for simplicity) :

$$h_{F_n, V}^1(y)(\sigma) = \frac{\sigma - 1}{\gamma_n - 1} y - (\sigma - 1)b,$$

where  $(\gamma_n - 1)(\varphi - 1)b = (\varphi - 1)y$ . Recall that  $\tilde{\mathbf{B}}_{\text{rig}}^\dagger = \cup_{r \geq 0} \tilde{\mathbf{B}}_{\text{rig}}^{\dagger, r}$ . Since  $b \in \tilde{\mathbf{B}}_{\text{rig}}^\dagger \otimes_{\mathbf{Q}_p} V$ , there exists  $m \gg 0$  such that  $b \in \tilde{\mathbf{B}}_{\text{rig}}^{\dagger, r_m} \otimes_{\mathbf{Q}_p} V$  and that the map  $\varphi^{-m}$  embeds  $\tilde{\mathbf{B}}_{\text{rig}}^{\dagger, r_m}$  into  $\mathbf{B}_{\text{dR}}^+$ . we can then write

$$h^1(y)(\sigma) = \frac{\sigma - 1}{\gamma_n - 1} \varphi^{-m}(y) - (\sigma - 1)\varphi^{-m}(b),$$

and  $\varphi^{-m}(b) \in \mathbf{B}_{\text{dR}}^+ \otimes_{\mathbf{Q}_p} V$ . In addition,  $\varphi^{-m}(y) \in F_m((t)) \otimes_F \mathbf{D}_{\text{cris}}(V)$  and  $\gamma_n - 1$  is invertible on  $t^k F_m \otimes_F \mathbf{D}_{\text{cris}}(V)$  for every  $k \neq 0$ . This shows that the cocycle  $h_{F_n, V}^1$  is cohomologous in  $H^1(F_n, \mathbf{B}_{\text{dR}} \otimes_{\mathbf{Q}_p} V)$  to

$$\sigma \mapsto \frac{\sigma - 1}{\gamma_n - 1} (\partial_V(\varphi^{-m}(y)))$$

which is itself cohomologous (since  $\gamma_n - 1$  is invertible on  $F_m^{\text{Tr}_{F_m/F_n}=0}$ ) to

$$\sigma \mapsto \frac{\sigma - 1}{\gamma_n - 1} (p^{n-m} \text{Tr}_{F_m/F_n} \partial_V(\varphi^{-m}(y))) = \sigma \mapsto p^{-n} \log_p(\chi(\bar{\sigma})) p^{n-m} \text{Tr}_{F_m/F_n} \partial_V(\varphi^{-m}(y)).$$

It follows from this and proposition 7.2 and lemma 9.11 that

$$\exp_{F_n, V^*(1)}^*(h_{F_n, V}^1(y)) = p^{-m} \text{Tr}_{F_m/F_n} \partial_V(\varphi^{-m}(y)) = \begin{cases} p^{-n} \partial_V(\varphi^{-n}(y)) & \text{if } n \geq 1 \\ (1 - p^{-1} \varphi^{-1}) \partial_V(y) & \text{if } n = 0. \end{cases}$$

□

**10.2. Perrin-Riou's big exponential map.** By using the results of the previous paragraphs, we can give a uniform formula for the image of an element  $y \in (\mathbf{B}_{\text{rig}, F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V))^{\psi=1}$  in  $H^1(F_n, V(j))$  under the composition of the following maps:

$$(\mathbf{B}_{\text{rig}, F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V))^{\psi=1} \xrightarrow{\nabla_{h-1} \circ \dots \circ \nabla_0} \mathbf{D}_{\text{rig}}^\dagger(V)^{\psi=1} \xrightarrow{\otimes e_j} \mathbf{D}_{\text{rig}}^\dagger(V(j))^{\psi=1} \xrightarrow{h_{F_n, V(j)}^1} H^1(F_n, V(j))$$

Here  $e_j$  is a basis of  $\mathbf{Q}_p(j)$  such that  $e_{j+k} = e_j \otimes e_k$  so that if  $V$  is a  $p$ -adic representation, then we have compatible isomorphisms of  $\mathbf{Q}_p$ -vector spaces  $V \rightarrow V(j)$  given by  $v \mapsto v \otimes e_j$ .

**Theorem 10.4.** *If  $y \in (\mathbf{B}_{\text{rig}, F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V))^{\psi=1}$ , and  $h \geq 1$  is an integer such that  $\text{Fil}^{-h} \mathbf{D}_{\text{cris}}(V) = \mathbf{D}_{\text{cris}}(V)$ , then for all  $j$  with  $h + j \geq 1$ , we have :*

$$h_{F_n, V(j)}^1(\nabla_{h-1} \circ \dots \circ \nabla_0(y) \otimes e_j) = (-1)^{h+j-1} (h + j - 1)! \times \begin{cases} \exp_{F_n, V(j)}(p^{-n} \partial_{V(j)}(\varphi^{-n}(\partial^{-j} y \otimes t^{-j} e_j))) & \text{if } n \geq 1 \\ \exp_{F, V(j)}((1 - p^{-1} \varphi^{-1}) \partial_{V(j)}(\partial^{-j} y \otimes t^{-j} e_j)) & \text{if } n = 0 \end{cases}$$

while if  $h + j \leq 0$ , then we have:

$$\exp_{F_n, V^*(1-j)}^*(h_{F_n, V}^1(\nabla_{h-1} \circ \cdots \circ \nabla_0(y) \otimes e_j)) = \frac{1}{(-h-j)!} \begin{cases} p^{-n} \partial_{V(j)}(\varphi^{-n}(\partial^j y \otimes t^{-j} e_j)) & \text{if } n \geq 1 \\ (1 - p^{-1} \varphi^{-1}) \partial_{V(j)}(\partial^{-j} y \otimes t^{-j} e_j) & \text{if } n = 0 \end{cases}$$

*Proof.* If  $h + j \geq 1$ , then we have the following commutative diagram:

$$\begin{array}{ccc} \mathbf{D}_{\text{rig}}^+(V)^{\psi=1} & \xrightarrow{\otimes e_j} & \mathbf{D}_{\text{rig}}^+(V(j))^{\psi=1} \\ \nabla_{h-1} \circ \cdots \circ \nabla_0 \uparrow & & \uparrow \nabla_{h+j-1} \circ \cdots \circ \nabla_0 \\ (\mathbf{B}_{\text{rig}, F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V))^{\psi=1} & \xrightarrow{\partial^{-j} \otimes t^{-j} e_j} & (\mathbf{B}_{\text{rig}, F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V(j)))^{\psi=1}. \end{array}$$

and the theorem is then a straightforward consequence of theorem 10.1 applied to  $\partial^{-j} y \otimes t^{-j} e_j$ ,  $h + j$  and  $V(j)$ .

On the other hand, if  $h + j \leq 0$ , and  $\Gamma_F^n$  is torsion free, then theorem 10.3 shows that

$$\exp_{F_n, V^*(1-j)}^*(h_{F_n, V(j)}^1(\nabla_{h-1} \circ \cdots \circ \nabla_0(y) \otimes e_j)) = p^{-n} \partial_{V(j)}(\varphi^{-n}(\nabla_{h-1} \circ \cdots \circ \nabla_0(y) \otimes e_j))$$

in  $\mathbf{D}_{\text{cris}}(V(j))$ , and a short computation involving Taylor series shows that

$$p^{-n} \partial_{V(j)}(\varphi^{-n}(\nabla_{h-1} \circ \cdots \circ \nabla_0(y) \otimes e_j)) = (-h-j)!^{-1} p^{-n} \partial_{V(j)}(\varphi^{-n}(\partial^{-j} y \otimes t^{-j} e_j)).$$

Finally, to get the case  $n = 0$ , one just needs to use the corresponding statement of theorem 10.3 or equivalently corestrict.  $\square$

**Remark 10.5.** The notation  $\partial^{-j}$  is not injective on  $\mathbf{B}_{\text{rig}, F}^+$  (it is surjective by integration) but it can be checked that it leads to no ambiguity in the formulas above.

We will now use the above result to give a construction of Perrin-Riou's exponential map. If  $f \in \mathbf{B}_{\text{rig}, F}^+ \otimes \mathbf{D}_{\text{cris}}(V)$ , we define  $\Delta(f)$  to be the image of  $\oplus_{k=0}^h \partial^k(f)(0)$  in  $\oplus_{k=0}^h (\mathbf{D}_{\text{cris}}(V))/(1 - p^k \varphi)(k)$ . There is then an exact sequence of  $\mathbf{Q}_p \otimes_{\mathbf{Z}_p} \Lambda_F$ -modules (cf [Per94, section 2.2]):

$$0 \longrightarrow \oplus_{k=0}^h t^k \mathbf{D}_{\text{cris}}(V)^{\varphi=p^{-k}} \longrightarrow (\mathbf{B}_{\text{rig}, F}^+ \otimes \mathbf{D}_{\text{cris}}(V))^{\psi=1} \xrightarrow{1-\varphi} (\mathbf{B}_{\text{rig}, F}^+)^{\psi=0} \otimes_F \mathbf{D}_{\text{cris}}(V) \xrightarrow{\Delta} \oplus_{k=0}^h \frac{\mathbf{D}_{\text{cris}}(V)}{1 - p^k \varphi}(k) \longrightarrow 0.$$

If  $f \in ((\mathbf{B}_{\text{rig}, F}^+)^{\psi=0} \otimes_F \mathbf{D}_{\text{cris}}(V))^{\Delta=0}$ , then by the above exact sequence there exists

$$y \in (\mathbf{B}_{\text{rig}, F}^+ \otimes \mathbf{D}_{\text{cris}}(V))^{\psi=1}$$

such that  $f = (1 - \varphi)y$ , and since  $\nabla_{h-1} \circ \cdots \circ \nabla_0$  kills  $\oplus_{k=0}^{h-1} t^k \mathbf{D}_{\text{cris}}(V)^{\varphi=p^{-k}}$  we see that  $\nabla_{h-1} \circ \cdots \circ \nabla_0(y)$  does not depend upon the choice of such  $y$  unless  $\mathbf{D}_{\text{cris}}(V)^{\varphi=p^{-h}} \neq 0$ .

**Definition 10.6.** Let  $h \geq 1$  be an integer such that  $\text{Fil}^{-h} \mathbf{D}_{\text{cris}}(V) = \mathbf{D}_{\text{cris}}(V)$  and such that  $\mathbf{D}_{\text{cris}}(V)^{\varphi=p^{-h}} = 0$ . One deduce from the above construction a well-defined map

$$\Omega_{V, h} : ((\mathbf{B}_{\text{rig}, F}^+)^{\psi=0} \otimes_F \mathbf{D}_{\text{cris}}(V))^{\Delta=0} \rightarrow \mathbf{D}_{\text{rig}}^+(V)^{\psi=1}$$

given by  $\Omega_{V, h}(f) = \nabla_{h-1} \circ \cdots \circ \nabla_0(y)$ , where  $y \in (\mathbf{B}_{\text{rig}, F}^+ \otimes \mathbf{D}_{\text{cris}}(V))^{\psi=1}$  is such that  $f = (1 - \varphi)y$ .

If  $\mathbf{D}_{\text{cris}}(V)^{\varphi=p^{-h}} \neq 0$  then we get a map

$$\Omega_{V, h} : ((\mathbf{B}_{\text{rig}, F}^+)^{\psi=0} \otimes_F \mathbf{D}_{\text{cris}}(V))^{\Delta=0} \rightarrow \mathbf{D}_{\text{rig}}^+(V)^{\psi=1} / V^{G_F = \chi^h}.$$

**Theorem 10.7.** *If  $V$  is a crystalline representation and  $h \geq 1$  is such that we have  $\mathrm{Fil}^{-h} \mathbf{D}_{\mathrm{cris}}(V) = \mathbf{D}_{\mathrm{cris}}(V)$ , then the map*

$$\Omega_{V,h} : ((\mathbf{B}_{\mathrm{rig},F}^+)^{\psi=0} \otimes_F \mathbf{D}_{\mathrm{cris}}(V))^{\Delta=0} \rightarrow \mathbf{D}_{\mathrm{rig}}^+(V)^{\psi=1} / V^{H_F}$$

*which takes  $f \in ((\mathbf{B}_{\mathrm{rig},F}^+)^{\psi=0} \otimes_F \mathbf{D}_{\mathrm{cris}}(V))^{\Delta=0}$  to  $\nabla_{h-1} \circ \cdots \circ \nabla_0((1-\varphi)^{-1}f)$  is well defined and coincides with Perrin-Riou's exponential map.*

*Proof.* The map  $\Omega_{V,h}$  is well defined because as we seen above the kernel of  $1-\varphi$  is killed by  $\nabla_{h-1} \circ \cdots \circ \nabla_0$ , except for  $t^h \mathbf{D}_{\mathrm{cris}}(V)^{\varphi=p^{-h}}$ , which is mapped to copies of  $\mathbf{Q}_p(h) \in V^{H_F}$ .

The fact that  $\Omega_{V,h}$  coincides with Perrin-Riou's exponential map follows directly from theorem 10.4 above applied to those  $j$ 's for which  $h+j \geq 1$ , and the fact that by [Per94, theorem 3.2.3], the  $\Omega_{V,h}$  are uniquely determined by the requirement that they satisfy the following diagram for  $h, j \gg 0$ :

$$\begin{array}{ccc} (\mathcal{H}(\Gamma_F) \otimes_{\mathbf{Q}_p} \mathbf{D}_{\mathrm{cris}}(V(j))^{\Delta=0}) & \xrightarrow{\Omega_{V(j),h}} & \mathcal{H}(\Gamma_F) \otimes_{\Gamma_F} (H_{\mathrm{Iw}}^1(F, V(j)/V(j)^{H_F})) \\ \Xi_{n,V(j)} \downarrow & & \downarrow \mathrm{pr}_{F_n, V(j)} \\ F_n \otimes_F \mathbf{D}_{\mathrm{cris}}(V) & \xrightarrow{(h+j-1)! \exp_{F_n, V(j)}} & H^1(F_n, V(j)). \end{array}$$

Here  $\Xi_{n,V(j)}(g) = p^{-n}(\varphi \otimes \varphi)^{-n}(f)(\varepsilon^{(n)} - 1)$  where  $f$  is such that

$$(1-\varphi)f = g(\gamma-1)(1+\pi) \in (\mathbf{B}_{\mathrm{rig},F}^+ \otimes_F \mathbf{D}_{\mathrm{cris}}(V))^{\psi=0}$$

and the  $\varphi$  on the left of  $\varphi \otimes \varphi$  is the Frobenius on  $\mathbf{B}_{\mathrm{rig},F}^+$  while the  $\varphi$  on the right is the Frobenius on  $\mathbf{D}_{\mathrm{cris}}(V)$ .

Note that by theorem 5.2, we have an isomorphism  $D(V)^{\psi=1} \simeq H_{\mathrm{Iw}}^1(F, V)$  and therefore we get a map  $\mathcal{H}(\Gamma_F) \otimes_{\Lambda_F} H_{\mathrm{Iw}}^1(F, V) \rightarrow \mathbf{D}_{\mathrm{rig}}^+(V)^{\psi=1}$ . On the other hand, there is a map

$$\mathcal{H}(\Gamma_F) \otimes_{\mathbf{Q}_p} \mathbf{D}_{\mathrm{cris}}(V(j)) \rightarrow (\mathbf{B}_{\mathrm{rig},F}^+ \otimes_F \mathbf{D}_{\mathrm{cris}}(V))^{\psi=0}$$

which sends  $\sum f_i(\gamma-1) \otimes d_i$  to  $\sum f_i(\gamma-1)(1+\pi) \otimes d_i$ . These two maps allow us to compare the diagram above with the formulas given by theorem 10.4.  $\square$

**Remark 10.8.** By the above remarks, if  $V$  is a crystalline representation and  $h \geq 1$  is such that  $\mathrm{Fil}^{-h} \mathbf{D}_{\mathrm{cris}}(V) = \mathbf{D}_{\mathrm{cris}}(V)$  and  $\mathbf{Q}_p(h) \not\subset V$ , then the map

$$\Omega_{V,h} : ((\mathbf{B}_{\mathrm{rig},F}^+)^{\psi=0} \otimes_F \mathbf{D}_{\mathrm{cris}}(V))^{\Delta=0} \rightarrow \mathbf{D}_{\mathrm{rig}}^+(V)^{\psi=1}$$

which takes  $f \in ((\mathbf{B}_{\mathrm{rig},F}^+)^{\psi=0} \otimes_F \mathbf{D}_{\mathrm{cris}}(V))^{\Delta=0}$  to  $\nabla_{h-1} \circ \cdots \circ \nabla_0((1-\varphi)^{-1}f)$  is well defined, without having to kill the  $\Lambda_F$ -torsion of  $H_{\mathrm{Iw}}^1(F, V)$ .

**Remark 10.9.** It is clear from theorem 10.4 that we have:

$$\Omega_{V,h}(x) \otimes e_j = \Omega_{V(j),h+j}(\partial^j x \otimes t^{-j} e_j) \quad \text{and} \quad \nabla_h \circ \Omega_{V,h}(x) = \Omega_{V,h+1}(x)$$

and following Perrin-Riou, one can use these formulas to extend the definition of  $\Omega_{V,h}$  to all  $h \in \mathbf{Z}$  by tensoring all  $\mathcal{H}(\Gamma_F)$ -modules with the field of fractions of  $\mathcal{H}(\Gamma_F)$

**10.3. The explicit reciprocity formula.** Recall we have a map  $\mathcal{H}(\Gamma_F) \rightarrow (\mathbf{B}_{\text{rig}, \mathbf{Q}_p}^+)^{\psi=0}$  which sends  $f(\gamma-1)$  to  $f(\gamma-1)(1+\pi)$ , this map is a bijection and its inverse is the Mellin transform so that if  $g(\pi) \in (\mathbf{B}_{\text{rig}, \mathbf{Q}_p}^+)^{\psi=0}$ , then  $g(\pi) = \mathfrak{M}(g)(1+\pi)$ . If  $f, g \in (\mathbf{B}_{\text{rig}, \mathbf{Q}_p}^+)^{\psi=0}$  then we define  $f * g$  by the formula  $\mathfrak{M}(f * g) = \mathfrak{M}(f)\mathfrak{M}(g)$ . Let  $[-1] \in \Gamma_F$  be the element such that  $\chi([-1]) = -1$ , and let  $\iota$  be the involution of  $\Gamma_F$  which sends  $\gamma$  to  $\gamma^{-1}$ . The operator  $\partial^j$  on  $(\mathbf{B}_{\text{rig}, \mathbf{Q}_p}^+)^{\psi=0}$  corresponds to  $\text{Tw}_j$  on  $\Gamma_F$  ( $\text{Tw}_j$  is defined by  $\text{Tw}_j(\gamma) = \chi(\gamma^j)\gamma$ ). We will make use of the facts that  $\iota \circ \partial^j = \partial^{-j} \circ \iota$  and  $[-1] \circ \partial^j = (-1)^j \partial^j \circ [-1]$ .

If  $V$  is a crystalline representation, then the natural maps

$$\mathbf{D}_{\text{cris}}(V) \otimes_F \mathbf{D}_{\text{cris}}(V^*(1)) \longrightarrow \mathbf{D}_{\text{cris}}(\mathbf{Q}_p(1)) \xrightarrow{\text{Tr}_{F/\mathbf{Q}_p}} \mathbf{Q}_p$$

allow us to define a perfect pairing  $[\cdot, \cdot]_V : \mathbf{D}_{\text{cris}}(V) \times \mathbf{D}_{\text{cris}}(V^*(1))$  which we extend by linearity to

$$[\cdot, \cdot]_V : (\mathbf{B}_{\text{rig}, F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V))^{\psi=0} \times (\mathbf{B}_{\text{rig}, F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V^*(1)))^{\psi=0} \rightarrow (\mathbf{B}_{\text{rig}, \mathbf{Q}_p}^+)^{\psi=0}$$

by the formula  $[f(\pi) \otimes d_1, g(\pi) \otimes d_2]_V = (f * g)(\pi)[d_1, d_2]_V$ .

We can also define a semi-linear pairing (with respect to  $\iota$ )

$$\langle \cdot, \cdot \rangle_V : \mathbf{D}_{\text{rig}}^+(V)^{\psi=1} \times \mathbf{D}_{\text{rig}}^+(V)^{\psi=1} \rightarrow (\mathbf{B}_{\text{rig}, \mathbf{Q}_p}^+)^{\psi=0}$$

by the formula

$$\langle \cdot, \cdot \rangle_V = \varprojlim_{\tau \in \Gamma_F / \Gamma_F^n} \sum \langle \tau^{-1}(h_{F_n, V}^1(y_1)), h_{F_n, V^*(1)}^1(y_2) \rangle_{F_n, V} \cdot \tau(1 + \pi)$$

where the pairing  $\langle \cdot, \cdot \rangle_{F_n, V}$  is given by the cup product:

$$\langle \cdot, \cdot \rangle_{F_n, V} : H^1(F_n, V) \times H^1(F_n, V^*(1)) \rightarrow H^2(F_n, \mathbf{Q}_p(1)) \cong \mathbf{Q}_p.$$

The pairing  $\langle \cdot, \cdot \rangle_V$  satisfies the relation  $\langle \gamma_1 x_1, \gamma_2 x_2 \rangle_V = \gamma_1 \iota(\gamma_2) \langle x_1, x_2 \rangle_V$ . Perrin-Riou's explicit reciprocity formula is then:

**Theorem 10.10.** *If  $x_1 \in (\mathbf{B}_{\text{rig}, F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V))^{\psi=0}$  and  $x_2 \in (\mathbf{B}_{\text{rig}, F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V^*(1)))^{\psi=0}$ , then for every  $h$ , we have*

$$(-1)^h \langle \Omega_{V, h}(x_1), [-1] \cdot \Omega_{V^*(1), 1-h}(x_2) \rangle_V = -[x_1, \iota(x_2)]_V.$$

*Proof.* By the theory of  $p$ -adic interpolation, it is enough to prove that if  $x_i = (1 - \varphi)y_i$  with  $y_1 \in (\mathbf{B}_{\text{rig}, F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V))^{\psi=1}$  and  $y_2 \in (\mathbf{B}_{\text{rig}, F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V^*(1)))^{\psi=1}$ , then for all  $j \gg 0$ ;

$$(\partial^{-j}(-1)^h \langle \Omega_{V, h}(x_1), [-1] \cdot \Omega_{V^*(1), 1-h}(x_2) \rangle_V)(0) = -(\partial^{-j}[x_1, \iota(x_2)]_V)(0).$$

The above formula is equivalent to:

$$(1) \quad (-1)^{h+j} \langle h_{F, V(j)}^1 \Omega_{V(j), h+j}(\partial^{-j} x_1 \otimes t^{-j} e_{-j}), h_{F, V^*(1-j)}^1 \Omega_{V^*(1-j), 1-h-j}(\partial^j x_2 \otimes t^j e_{-j}) \rangle_{F, V(j)} \\ = [\partial_{V(j)}(\partial^{-j} x_1 \otimes t^{-j} e_j), \partial_{V^*(1-j)}(\partial^j x_2 \otimes t^j e_{-j})]_{V(j)}.$$

By combining theorems 10.4 and 10.7 with remark 10.9, we see that for  $j \gg 0$ :

$$h_{F, V(j)}^1 \Omega_{V(j), h+j}(\partial^{-j} x_1 \otimes t^{-j} e_j) = (-1)^{h+j-1} \exp_{F, V(j)}((h+j-1)!(1-p^{-1}\varphi^{-1})\partial_{V(j)}(\partial^{-j} y_1 \otimes t^{-j} e_j)),$$

and that

$$h_{F, V^*(1-j)}^1 \Omega_{V^*(1-j), 1-h-j}(\partial^j x_2 \otimes t^j e_{-j}) \\ = (\exp_{F, V^*(1-j)}^*)^{-1} (h+j-1)!^{-1} ((1-p^{-1}\varphi^{-1})\partial_{V^*(1-j)}(\partial^j y_2 \otimes t^j e_{-j})).$$

Using the fact that by definition, if  $x \in \mathbf{D}_{\text{cris}}(V(j))$  and  $y \in H^1(F, V(j))$  then

$$[x, \exp_{F, V^*(1-j)}^* y]_{V(j)} = \langle \exp_{F, V(j)} x, y \rangle_{F, V(j)},$$

we see that

$$\begin{aligned} (2) \quad & \langle h_{F, V(j)}^1 \Omega_{V(j), h+j}(\partial^{-j} x_1 \otimes t^{-j} e_j), h_{F, V^*(1-j)}^1 \Omega_{V^*(1-j), 1-h-j}(\partial^j x_2 \otimes t^j e_{-j}) \rangle_{F, V(j)} \\ &= (-1)^{h+j-1} [(1-p^{-1}\varphi^{-1})\partial_{V(j)}(\varphi^{-j} y_1 \otimes t^{-j} e_j), (1-p^{-1}\varphi^{-1})\partial_{V^*(1-j)}(\partial^j y_2 \otimes t^j e_{-j})]_{V(j)}. \end{aligned}$$

It is easy to see that under  $[\cdot, \cdot]$ , the adjoint of  $(1-p^{-1}\varphi^{-1})$  is  $1-\varphi$  and that if  $x_i = (1-\varphi)y_i$ , then

$$\begin{aligned} \partial_{V(j)}(\partial^{-j} x_1 \otimes t^{-j} e_j) &= (1-\varphi)\partial_{V(j)}(\partial^{-j} y_1 \otimes t^{-j} e_j), \\ \partial_{V^*(1-j)}(\partial^j x_2 \otimes t^j e_{-j}) &= (1-\varphi)\partial_{V^*(1-j)}(\partial^j y_2 \otimes t^j e_{-j}), \end{aligned}$$

So that (2) implies (1), and this proves the theorem.  $\square$

## 11. PERRIN-RIOU'S BIG REGULATOR MAP

Let  $F$  be a finite unramified extension over  $\mathbf{Q}_p$  and  $V$  a continuous  $p$ -adic representation of  $G_F$ , which is crystalline with Hodge-Tate weights  $\geq 0$  and with no quotient isomorphic to the trivial representation. In [Per95], Perrin-Riou construct a big logarithm map

$$\mathcal{L}_{F,V}^{\Gamma_F} : H_{\text{Iw}}^1(F, V) \longrightarrow \mathcal{H}(\Gamma_F) \otimes_{\mathbf{Q}_p} \mathbf{D}_{\text{cris}}(V)$$

which interpolates the values of Bloch-Kato's dual exponential and logarithm maps for  $V(j)$ ,  $j \in \mathbf{Z}$ , over each  $F_n$ .

In this section, we follow [LZ11, Appendix B] to adapt Berger's explicit formulas to construct Perrin-Riou's big logarithm and use it to calculate Kubota-Leopoldt  $p$ -adic  $L$ -function.

**11.1. Perrin-Riou's big logarithm map.** Let  $V$  be a positive crystalline representation of  $\text{Gal}(F_{\infty}/F)$  and  $x \in \mathcal{H}(\Gamma_F) \otimes_{\Lambda_F} H_{\text{Iw}}^1(F, V)$ . We write  $x_j$  for the image of  $x$  in  $H_{\text{Iw}}^1(F, V(-j))$ , and  $x_{j,n}$  for the image of  $x_j$  in  $H^1(F_n, V(-j))$ . If we identify  $x$  with its image in  $D(V)^{\psi=1}$ , then  $x_j$  corresponds to the element  $x \otimes e_{-j} \in D(V)^{\psi=1} \otimes e_{-j} = D(V(-j))^{\psi=1}$ .

Since  $V$  is positive, we may interpret  $x$  as an element of the module  $(\mathbf{B}_{\text{rig}, F}^+[1/t] \otimes \mathbf{D}_{\text{cris}}(V))^{\psi=1}$ .

We shall assume:

$$(3) \quad x \in (\mathbf{B}_{\text{rig}, F}^+ \otimes_{\mathbf{B}_F^+} \mathbf{N}(V))^{\psi=1} \subset (\mathbf{B}_{\text{rig}, F}^+[1/t] \otimes_F \mathbf{D}_{\text{cris}}(V))^{\psi=1}.$$

The condition is satisfied if  $V$  has no quotient isomorphic to  $\mathbf{Q}_p$  (c.f. theorem 9.4).

Recall in section 6.2, we define  $\partial$  denote the differential operator  $(1+\pi)\frac{d}{d\pi}$  (or  $\frac{d}{dt}$ ) on  $\mathbf{B}_{\text{rig}, F}^+$  and we have a map

$$\partial_V \circ \varphi^{-n} : \mathbf{B}_{\text{rig}, F}^+[1/t] \otimes_F \mathbf{D}_{\text{cris}}(V) \rightarrow F_n \otimes_F \mathbf{D}_{\text{cris}}(V)$$

which sends  $\pi^k \otimes d$  to the constant coefficient of  $(\zeta_n \exp(t/p^n) - 1)^k \otimes \varphi^{-n}(d) \in F_n((t)) \otimes_F \mathbf{D}_{\text{cris}}(V)$ .

For  $m \in \mathbf{Z}$ , define  $\Gamma^*(m)$  the leading term of Taylor series expansion of  $\Gamma(x)$  at  $x = m$ ; thus

$$\Gamma^*(m) = \begin{cases} j! & \text{if } n \geq 0 \\ \frac{(-1)^{-j-1}}{(-j-1)!} & \text{if } n \leq -1 \end{cases}$$

**Proposition 11.1.** *Define*

$$R_{j,n}(x) = \frac{1}{j!} \times \begin{cases} p^{-n} \partial_{V(-j)}(\varphi^{-n}(\partial^j x \otimes t^j e_{-j})) & \text{if } n \geq 1 \\ (1 - p^{-1} \varphi^{-1}) \partial_{V(-j)}(\partial^j x \otimes t^j e_{-j}) & \text{if } n = 0 \end{cases}$$

Then we have

$$R_{j,n}(x) = \begin{cases} \exp_{F_n, V^*(1-j)}^*(x_{j,n}) & \text{if } j \geq 0 \\ \log_{F_n, V(-j)}(x_{j,n}) & \text{if } j \leq -1 \end{cases}$$

*Proof.* This result is essentially a minor variation on theorem 10.4. The case  $j \geq 0$  is immediate from theorem 10.1 applied with  $V$  replaced by  $V(-j)$  and  $x$  by  $x \otimes e_{-j}$ , using the formula

$$\partial_{V(-j)}(\varphi^{-n}(x \otimes e_{-j})) = \frac{1}{j!} \partial_{V(-j)}(\varphi^{-n}(\partial^j x \otimes t^j e_{-j})).$$

For the formula for  $j \leq -1$ , we choose  $h$  such that  $\text{Fil}^{-h} \mathbf{D}_{\text{cris}}(V) = \mathbf{D}_{\text{cris}}(V)$ . The element  $\partial^j x \otimes t^j e_{-j}$  lies in  $(\mathbf{B}_{\text{rig}, F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V(-j)))^{\psi=1}$ . Applying theorem 10.1 with  $V, h$  and  $x$  replaced by  $V(-j)$ ,  $h - j$ , and  $\partial^j x \otimes t^{-j} e_j$ , we see that

$$\Gamma^*(j+1) R_{j,n}(x) = \Gamma^*(j-h+1) \log_{F_n, V(-j)}[(\nabla_0 \circ \cdots \circ \nabla_{h-1} x)_{j,n}].$$

For  $x \in \mathcal{H}(\Gamma_F) \otimes_{\Lambda_F} H_{\text{Iw}}^1(F, V)$ , we have

$$(\nabla_r x)_{j,n} = (j-r) x_{j,n},$$

so we have

$$(\nabla_0 \circ \cdots \circ \nabla_{h-1} x)_{j,n} = (j)(j-1) \cdots (j-h+1) x_{j,n}$$

as require.  $\square$

For  $\omega$  a finite order character on  $\Gamma_F$  of conductor  $n$ , we denote

$$G(\omega) = \sum_{\sigma \in \Gamma_F / \Gamma_{F_n}} \omega(\sigma) \zeta_{p^n}^\sigma.$$

the Gauss sum of  $\omega$ .

**Proposition 11.2.** *If  $x$  is as above, and  $\mathcal{L}_V^{\Gamma_F}(x)$  is the unique element of  $\mathcal{H}(\Gamma_F) \otimes_F \mathbf{D}_{\text{cris}}(V)$  such that  $\mathcal{L}_V^{\Gamma_F}(x) \cdot (1 + \pi) = (1 - \varphi)x$ , then for any  $j \in \mathbf{Z}$  we have*

$$(1 - \varphi) \partial_{V(-j)}(\varphi^{-n}(\partial^j x \otimes t^j e_{-j})) = \mathcal{L}_V^{\Gamma_F}(x)(\chi^j) \otimes t^j e_{-j},$$

while for any finite order character  $\omega$  of  $\Gamma_F$  of conductor  $n \geq 1$ , we have

$$\left( \sum_{\sigma \in \Gamma_F / \Gamma_F^n} \omega(\sigma)^{-1} \sigma \right) \cdot \partial_{V(-j)}(\varphi^{-n}(\partial^j x \otimes t^j e_{-j})) = G(\omega) \varphi^{-n}(\mathcal{L}_V^{\Gamma_F}(x)(\chi^j \omega) \otimes t^j e_{-j}).$$

*Proof.* We note that

$$\mathcal{L}_{V(-j)}^{\Gamma_F}(\partial^j x \otimes t^j e_{-j}) = \text{Tw}_j(\mathcal{L}_V^{\Gamma_F}(x)) \otimes t^j e_{-j},$$

so it suffices to prove the result for  $j = 0$ . Suppose we have  $x = \sum_{k \geq 0} v_k \pi^k$  where  $v_k \in \mathbf{D}_{\text{cris}}(V)$ .

Then

$$\partial_V(\varphi^{-n}(x)) = \sum_{k \geq 0} \varphi^{-n}(v_k)(\zeta_{p^n} - 1)^k.$$

On the other hand,

$$\partial_V(\varphi^{-n}((1-\varphi)x)) = \sum_{k \geq 0} \varphi^{-n}(v_k)(\zeta_{p^n} - 1)^k - \sum_{k \geq 0} \varphi^{1-n}(v_k)(\zeta_{p^{n-1}} - 1)^k.$$

Applying the operator  $e_\omega = \sum_{\sigma \in \Gamma_F/\Gamma_F^n} \omega(\sigma)\sigma$ , we have for  $n \geq 1$

$$e_\omega \cdot \partial_V(\varphi^{-n}(x)) = e_\omega \cdot \partial_V(\varphi^{-n}((1-\varphi)x)),$$

since  $e_\omega$  is zero on  $F_{n-1}((t))$ .

However, since the map  $\partial_V \circ \varphi^{-n}$  is a homomorphism of  $\Gamma_F$ -modules, we have

$$\begin{aligned} e_\omega \cdot \partial_V(\varphi^{-n}((1-\varphi)x)) &= e_\omega \cdot \partial_V(\varphi^{-n}(\mathcal{L}_V^{\Gamma_F}(x) \cdot (1+\pi))) \\ &= \varphi^{-n}(\mathcal{L}_V^{\Gamma_F}(x)) \cdot e_\omega \partial_V(\varphi^{-n}(1+\pi)) \\ &= G(\omega) \varphi^{-n}(\mathcal{L}_V^{\Gamma_F}(x)(\omega)). \end{aligned}$$

This completes the proof of the proposition for  $j = 0$ .  $\square$

**Definition 11.3.** Let  $x \in H_{\text{Iw}}^1(F, V)$ . If  $\eta$  is any continuous character of  $\Gamma_F$ , denote by  $x_\eta$  the image of  $x$  in  $H_{\text{Iw}}^1(F, V(\eta^{-1}))$ . If  $n \geq 0$ , denote by  $x_{\eta,n}$  the image of  $x_\eta$  in  $H^1(F_n, V(\eta^{-1}))$ .

Thus  $x_{\chi^j,n} = x_{j,n}$  in the previous notation. The next lemma is valid for arbitrary de Rham representations of  $G_F$  (with no restriction on Hodge-Tate weights):

**Lemma 11.4.** *For any finite-order character  $\omega$  factoring through  $\Gamma_F/\Gamma_F^n$ , with values in a finite extension  $E/F$ , we have*

$$\sum_{\sigma \in \Gamma_F/\Gamma_F^n} \omega(\sigma)^{-1} \exp_{F_n, V^*(1)}^*(x_{0,n})^\sigma = \exp_{F_n, V(\omega^{-1})^*(1)}^*(x_{\omega,0})$$

and

$$\sum_{\sigma \in \Gamma/\Gamma_n} \omega(\sigma)^{-1} \log_{F_n, V}(x_{0,n})^\sigma = \log_{F_n, V(\omega^{-1})}(x_{\omega,0})$$

where we identify  $\mathbf{D}_{\text{dR}}(V(\omega^{-1})) \cong (E \otimes_F F_n \otimes_F \mathbf{D}_{\text{cris}}(V))^{\Gamma=\omega}$ .

*Proof.* This follows from the compatibility of the maps  $\exp^*$  and  $\log$  with the corestriction maps (c.f. Theorem 10.1 and 10.3).  $\square$

Combining the three results above, we obtain:

**Theorem 11.5.** *Let  $j \in \mathbf{Z}$  and let  $x$  satisfies (3). Let  $\eta$  be a continuous character of  $\Gamma_F$  of the form  $\chi^j \omega$ , where  $\omega$  is a finite-order character of conductor  $n$ .*

i) *If  $j \geq 0$ , we have*

$$\mathcal{L}_V^{\Gamma_F}(x)(\eta) = j! \times \begin{cases} (1-p^j\varphi)(1-p^{-1-j}\varphi^{-1})^{-1} \left( \exp_{F, V(\eta^{-1})^*(1)}^*(x_{\eta,0}) \otimes t^{-j}e_j \right) & \text{if } n = 0 \\ G(\omega)^{-1} p^{n(1+j)} \varphi^n \left( \exp_{F, V(\eta^{-1})^*(1)}^*(x_{\eta,0}) \otimes t^{-j}e_j \right) & \text{if } n \geq 1. \end{cases}$$

ii) *If  $j \leq -1$ , we have*

$$\mathcal{L}_V^{\Gamma_F}(x)(\eta) = \frac{(-1)^{-j-1}}{(-j-1)!} \times \begin{cases} (1-p^j\varphi)(1-p^{-1-j}\varphi^{-1})^{-1} \left( \log_{F, V(\eta^{-1})}(x_{\eta,0}) \otimes t^{-j}e_j \right) & \text{if } n = 0 \\ G(\omega)^{-1} p^{n(1+j)} \varphi^n \left( \log_{F, V(\eta^{-1})}(x_{\eta,0}) \otimes t^{-j}e_j \right) & \text{if } n \geq 1. \end{cases}$$



In both case, we assume that  $(1 - p^{-1-j}\varphi^{-1})$  is invertible on  $\mathbf{D}_{\text{cris}}(V)$  when  $\eta = \chi^j$ .

**11.2. Cyclotomic units and Kubota-Leopoldt  $p$ -adic  $L$ -functions.** The relation between Coleman's power series and the Perrin-Riou's big logarithm map is given by the following diagram:

$$\begin{array}{ccc} \varprojlim \mathcal{O}_{F_n}^* & \xrightarrow{\kappa} & H_{\text{Iw}}^1(F, \mathbf{Z}_p(1)) \\ \text{Col} \downarrow & & \downarrow \mathcal{L}_{F, \mathbf{Q}_p(1)}^\Gamma \\ \mathcal{O}_F[[\pi]]^* & & \\ (1 - \frac{\varphi}{p}) \log \downarrow & & \downarrow \\ \mathcal{O}_F[[\pi]]^{\psi=0} & \longrightarrow & \mathcal{H}(\Gamma) \otimes_{\mathbf{Q}_p} \mathbf{D}_{\text{cris}}(F, \mathbf{Q}_p(1)) \end{array}$$

If we identify  $\mathbf{D}_{\text{cris}}(F, \mathbf{Q}_p(1))$  with  $F$  via the basis vector  $t^{-1} \otimes e_1$ , then the bottom map sends  $f \in \mathcal{O}_F[[\pi]]^{\psi=0}$  to  $\nabla_0 \cdot \mathfrak{M}^{-1}(f)$ , where  $\nabla_0 = \frac{\log \gamma}{\log \chi(\gamma)}$  for any non-identity element  $\gamma \in \Gamma_1$  and  $\mathfrak{M}$  is the Mellin transform defined in section 9.5. Thus the image of the bottom map is precisely  $\nabla_0 \cdot \Lambda_{\mathcal{O}_F}(\Gamma) \subset \mathcal{H}_F(\Gamma)$ ; and if we define

$$h_F(u) = \nabla_0^{-1} \cdot \mathcal{L}_{F, \mathbf{Q}_p(1)}^\Gamma(\kappa(u)) \in \Lambda_{\mathcal{O}_F}(\Gamma),$$

then we have

$$\mathfrak{M}(h_F(u)) = (1 - \frac{\varphi}{p}) \log \text{Col}_u(u).$$

By calculation in section 8.6, we can use theorem 11.5 to calculate the Kubota-Leopoldt  $p$ -adic  $L$ -functions.

**Example 11.6.** (Kubota-Leopoldt  $p$ -adic zeta-function) Let  $K = \mathbf{Q}_p$ ,  $V = \mathbf{Q}_p(1)$  and

$$u = (\frac{\zeta_{p^n} - 1}{\zeta_{p^n}})_{n \geq 1} \in \varprojlim \mathcal{O}_{\mathbf{Q}_p(\mu_{p^n})}^*.$$

Then by calculation in section 8.6, we have

$$\begin{aligned} h_F(u)(\chi^k) &= \chi^k(\nabla_0^{-1}) \cdot k! \cdot \frac{(1 - p^k \varphi)}{(1 - p^{-1-k} \varphi^{-1})} \left( \exp_{\mathbf{Q}_p, V^*(1-j)}^*(u_{k,0}) \otimes t^{-k} e_k \right) \\ &= \frac{1}{k} k! \cdot \frac{(1 - p^k \varphi)}{(1 - p^{-1-k} \varphi^{-1})} \left( (1 - p^{-k}) \zeta(1 - k) \frac{t^{k-1}}{(k-1)!} \otimes t^{-k} e_k \right) \\ &= (1 - p^{k-1}) \zeta(1 - k) t^{-1} \end{aligned}$$

and for  $\omega$  a finite order character of  $\Gamma$  of conductor  $n$ , we have

$$\begin{aligned} h_F(u)(\chi^k \omega) &= \chi^k(\nabla_0^{-1}) \cdot k! \cdot G(\omega)^{-1} p^{n(1+k)} \varphi^n \left( \exp_{\mathbf{Q}_p, V(\eta^{-1})^*(1)}^*(u_{\eta,0}) \otimes t^{-k} e_k \right) \\ &= \frac{1}{k} k! \cdot G(\omega)^{-1} p^{n(1+k)} \varphi^n \left( p^{-(n+1)k} G(\omega) L(1 - k, \omega) \frac{t^{k-1}}{(k-1)!} \otimes t^{-k} e_k \right) \\ &= L(1 - k, \omega) t^{-1} \end{aligned}$$

**Example 11.7.** (Kubota-Leopoldt  $p$ -adic  $L$ -function) Let  $K = \mathbf{Q}_p(\zeta_d)$ ,  $V = \mathbf{Q}_p(1)$  and  $\varepsilon$  is a Dirichlet character of conductor  $d \geq 1$  prime to  $p$ . Set  $u = (\frac{-1}{G(\varepsilon^{-1})} \sum_{0 \leq a \leq d-1} \varepsilon(a)^{-1} \frac{\zeta_d^a \zeta_{p^n}}{\zeta_d^a \zeta_{p^n} - 1})_{n \geq 1}$ .

Then by calculation in section 8.6, we have

$$\begin{aligned} h_F(u)(\chi^k) &= \chi^k(\nabla_0^{-1}) \cdot k! \cdot \frac{(1-p^k\varphi)}{(1-p^{-1-k}\varphi^{-1})^{-1}} \left( \exp_{K,V^*(1-j)}^*(u_{k,0}) \otimes t^{-k}e_k \right) \\ &= \frac{1}{k} k! \cdot \frac{(1-p^k\varphi)}{(1-p^{-1-k}\varphi^{-1})^{-1}} \left( (1-\varepsilon(p)p^{-k})L(1-k, \varepsilon) \frac{t^{k-1}}{(k-1)!} \otimes t^{-k}e_k \right) \\ &= (1-\varepsilon(p)p^{k-1})L(1-k, \varepsilon)t^{-1} \end{aligned}$$

and for  $\omega$  a finite order character of  $\Gamma$  of conductor  $n$ , we have

$$\begin{aligned} h_F(u)(\chi^k\omega) &= \chi^k(\nabla_0^{-1}) \cdot k! \cdot G(\omega)^{-1}p^{n(1+k)}\varphi^n \left( \exp_{\mathbf{Q}_p,V(\eta^{-1})^*(1)}^*(u_{\eta,0}) \otimes t^{-k}e_k \right) \\ &= \frac{1}{k} k! \cdot G(\omega)^{-1}p^{n(1+k)}\varphi^n \left( p^{-(n+1)k}G(\omega)L(1-k, \omega\varepsilon) \frac{t^{k-1}}{(k-1)!} \otimes t^{-k}e_k \right) \\ &= L(1-k, \omega\varepsilon)t^{-1} \end{aligned}$$

## REFERENCES

- [Ber02] L. Berger, *Representations  $p$ -adiques et quations differentielles*, Invent. Math. **148** (2002), 219-284.
- [Ber03] L. Berger, *Bloch and Kato's exponential map: three explicit formulas* Doc. Math. Extra Vol. **3** (2003): 99-129.
- [Ber04] L. Berger, *Limites de representations cristallines* Compositio Math. **140** (2004): 14731498.
- [BCDDPR] M. Bertolini, F. Castella, H. Darmon, S. Dasgupta, K. Prasanna and V. Rotger  *$p$ -adic  $L$ -functions and Euler systems: a tale in two trilogies*, Automorphic forms and Galois representations, London Mathematical Society Lecture Notes Series **414** (2014).
- [CC98] F. Cherbonnier and P. Colmez, *Representations  $p$ -adiques surconvergentes*, Invent. Math. **133** (1998), 581-611.
- [CC99] F. Cherbonnier and P. Colmez, *Thorie d'Iwasawa des representations  $p$ -adiques d'un corps local*, J. Amer. Math. Soc. **12** (1999), 241-268.
- [Co79] R. Coleman, *Division values in local fields*, Invent. Math. **53** (1979), 91-116.
- [Col98] P. Colmez, *Thorie d'Iwasawa des Representations de de Rham d'un Corps Local*, Annals Math. **148** (1998), 485-571.
- [Col99] P. Colmez, *Representations cristallines et representations de hauteur finie*, J. Reine Angew. Math. **514** (1999), 119-143.
- [Col02] P. Colmez, *Arithmetique de la fonction zeta*, Journees mathematiques **X-UPS** (2002), 37-164.
- [Col04] P. Colmez, *Fontaine rings and  $p$ -adic  $L$ -functions*, Course given at Tsinghua University during the fall of 2004.
- [Col10] P. Colmez, *Fonctions d'une variable  $p$ -adique*, Astrisque. **330** (2010), 13-59.
- [Her98] L. Herr, *Sur la cohomologie galoisienne des corps  $p$ -adiques*, Bulletin de la Socit Mathematique de France. **126** (1998), 563-600.
- [Kato93] K. Kato, *Lectures on the approach to Iwasawa theory for Hasse-Weil  $L$ -functions via  $\mathbf{B}_{dR}$* , in Arithmetic Algebraic Geometry, Lecture Notes in math. **1553** (1993).
- [LLZ11] A. Lei, D. Loeffler and S.L. Zerbes, *Coleman maps and the  $p$ -adic regulator*, Algebra & Number Theory. **5** (2011), 1095-1131.
- [LZ11] D. Loeffler and S.L. Zerbes, *Iwasawa theory and  $p$ -adic  $L$ -functions over  $\mathbf{Z}_p^2$ -extensions*, to appear in Int. J. Number Theory.
- [NSK] J. Neukirch, A. Schmidt and K. Wingberg, *Cohomology of number fields*, Grundlehren der Mathematischen Wissenschaften **126**, Springer-Verlag, Berlin (2000).
- [Per94] B. Perrin-Riou, *Thorie d'Iwasawa des representations  $p$ -adiques sur un corps local*, Invent. Math. **115** (1994), 81-149.
- [Per95] B. Perrin-Riou, *Fonctions  $L$   $p$ -adiques des representations  $p$ -adiques*, Astrisque. **229** (1995), 81-149.
- [Per01] B. Perrin-Riou, *Thorie d'Iwasawa des representations  $p$ -adiques semi-stables*, Mm. Soc. Math. Fr. (N.S.) **84** (2001), vi+111.