

Fundamental groups in Topology

Reza Sadoughian

University of British Columbia

January 2014

Contents

- Let X be a top. space. A **space over X** is a topological space Y together with a continuous map $p : Y \rightarrow X$.

A morphism between two spaces $p_i : Y_i \rightarrow X$ ($i=1,2$) over X is given by a continuous map $f : Y_1 \rightarrow Y_2$ making the following diagram commutes :

$$\begin{array}{ccc} Y_1 & \xrightarrow{f} & Y_2 \\ & \searrow p_1 & \downarrow p_2 \\ & & X \end{array}$$

- A **cover** of X is a space Y over X where the projection $p : Y \rightarrow X$ satisfies the following condition :
Each point of X has an open neighbourhood V for which $p^{-1}(V)$ decomposes as a disjoint union of open subsets $U_i \subseteq Y$ such that each restricted map $p|_{U_i} : U_i \rightarrow V$ induces a homeomorphism of U_i with V .

- Let X be a top. space. A **space over** X is a topological space Y together with a continuous map $p : Y \rightarrow X$.

A morphism between two spaces $p_i : Y_i \rightarrow X$ ($i=1,2$) over X is given by a continuous map $f : Y_1 \rightarrow Y_2$ making the following diagram commutes :

$$\begin{array}{ccc} Y_1 & \xrightarrow{f} & Y_2 \\ & \searrow p_1 & \downarrow p_2 \\ & & X \end{array}$$

- A **cover** of X is a space Y over X where the projection $p : Y \rightarrow X$ satisfies the following condition :

Each point of X has an open neighbourhood V for which $p^{-1}(V)$ decomposes as a disjoint union of open subsets $U_i \subseteq Y$ such that each restricted map $p|_{U_i} : U_i \rightarrow V$ induces a homeomorphism of U_i with V .

- **Example**(*Trivial cover*) : Let I be a nonempty discrete top. space. The first projection

$$X \times I \rightarrow X$$

turns $X \times I$ into a cover of X .

- **Proposition 2.1.3** : A space Y over X is a cover iff each point of X has an open neighbourhood V such that the restriction of the projection $p : Y \rightarrow X$ to $p^{-1}(V)$ is isomorphic as a space ver V to a trivial cover.
- The points of X over which the fibre of p equals I form an open subset of X .
- **Corollary 2.1.4** : If X is connected, the fibres of p are all homeomorphic to the same discrete space I .
- Let G be a group acting continuously from the left on a topological space Y . The action of G is **even** (or **properly discontinuous**) if each point $y \in Y$ has some open neighbourhood U such that $g_1 U \cap g_2 U = \emptyset$ if $g_1 \neq g_2$.

- **Example**(*Trivial cover*) : Let I be a nonempty discrete top. space. The first projection

$$X \times I \rightarrow X$$

turns $X \times I$ into a cover of X .

- **Proposition 2.1.3** : A space Y over X is a cover iff each point of X has an open neighbourhood V such that the restriction of the projection $p : Y \rightarrow X$ to $p^{-1}(V)$ is isomorphic as a space ver V to a trivial cover.
- The points of X over which the fibre of p equals I form an open subset of X .
- **Corollary 2.1.4** : If X is connected, the fibres of p are all homeomorphic to the same discrete space I .
- Let G be a group acting continuously from the left on a topological space Y . The action of G is **even** (or **properly discontinuous**) if each point $y \in Y$ has some open neighbourhood U such that $g_1 U \cap g_2 U = \emptyset$ if $g_1 \neq g_2$.

- **Example**(*Trivial cover*) : Let I be a nonempty discrete top. space. The first projection

$$X \times I \rightarrow X$$

turns $X \times I$ into a cover of X .

- **Proposition 2.1.3** : A space Y over X is a cover iff each point of X has an open neighbourhood V such that the restriction of the projection $p : Y \rightarrow X$ to $p^{-1}(V)$ is isomorphic as a space ver V to a trivial cover.
- The points of X over which the fibre of p equals I form an open subset of X .
- **Corollary 2.1.4** : If X is connected, the fibres of p are all homeomorphic to the same discrete space I .
- Let G be a group acting continuously from the left on a topological space Y . The action of G is **even** (or **properly discontinuous**) if each point $y \in Y$ has some open neighbourhood U such that $g_1 U \cap g_2 U = \emptyset$ if $g_1 \neq g_2$.

- **Example**(*Trivial cover*) : Let I be a nonempty discrete top. space. The first projection

$$X \times I \rightarrow X$$

turns $X \times I$ into a cover of X .

- **Proposition 2.1.3** : A space Y over X is a cover iff each point of X has an open neighbourhood V such that the restriction of the projection $p : Y \rightarrow X$ to $p^{-1}(V)$ is isomorphic as a space ver V to a trivial cover.
- The points of X over which the fibre of p equals I form an open subset of X .
- **Corollary 2.1.4** : If X is connected, the fibres of p are all homeomorphic to the same discrete space I .
- Let G be a group acting continuously from the left on a topological space Y . The action of G is **even** (or **properly discontinuous**) if each point $y \in Y$ has some open neighbourhood U such that $g_1 U \cap g_2 U = \emptyset$ if $g_1 \neq g_2$.

- **Example**(*Trivial cover*) : Let I be a nonempty discrete top. space. The first projection

$$X \times I \rightarrow X$$

turns $X \times I$ into a cover of X .

- **Proposition 2.1.3** : A space Y over X is a cover iff each point of X has an open neighbourhood V such that the restriction of the projection $p : Y \rightarrow X$ to $p^{-1}(V)$ is isomorphic as a space ver V to a trivial cover.
- The points of X over which the fibre of p equals I form an open subset of X .
- **Corollary 2.1.4** : If X is connected, the fibres of p are all homeomorphic to the same discrete space I .
- Let G be a group acting continuously from the left on a topological space Y . The action of G is **even** (or **properly discontinuous**) if each point $y \in Y$ has some open neighbourhood U such that $g_1 U \cap g_2 U = \emptyset$ if $g_1 \neq g_2$.

- **Example**(*Trivial cover*) : Let I be a nonempty discrete top. space. The first projection

$$X \times I \rightarrow X$$

turns $X \times I$ into a cover of X .

- **Proposition 2.1.3** : A space Y over X is a cover iff each point of X has an open neighbourhood V such that the restriction of the projection $p : Y \rightarrow X$ to $p^{-1}(V)$ is isomorphic as a space ver V to a trivial cover.
- The points of X over which the fibre of p equals I form an open subset of X .
- **Corollary 2.1.4** : If X is connected, the fibres of p are all homeomorphic to the same discrete space I .
- Let G be a group acting continuously from the left on a topological space Y . The action of G is **even** (or **properly discontinuous**) if each point $y \in Y$ has some open neighbourhood U such that $g_1 U \cap g_2 U = \emptyset$ if $g_1 \neq g_2$.

- **Lemma 2.1.7 :** If G is a group acting evenly on a connected space Y , the projection $p_G : Y \rightarrow G \backslash Y$ turns Y into a cover of $G \backslash Y$.
- **Example 2.1.8 :** 1. Let \mathbb{Z} act on \mathbb{R} by translations. We obtain a cover $\mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z} \simeq S^1$.
2. For an integer $n > 1$, we get an even action of μ_n on \mathbb{C}^* from that a cover $p_n : \mathbb{C}^* \rightarrow \mathbb{C}^*/\mu_n$. The map $z \mapsto z^n$ defines a natural homeomorphism (isomorphism) of \mathbb{C}^*/μ_n onto \mathbb{C}^* . So this induces a cover $\mathbb{C}^* \rightarrow \mathbb{C}^*$.
- For now on, assume that X is locally connected.
- $\text{Aut}(Y/X) := \{\phi : Y \rightarrow Y : p \circ \phi = p\}$.
- For each point $x \in X$, $p^{-1}(x)$ is equipped with a natural action of $\text{Aut}(Y/X)$.

- **Lemma 2.1.7 :** If G is a group acting evenly on a connected space Y , the projection $p_G : Y \rightarrow G \backslash Y$ turns Y into a cover of $G \backslash Y$.
- **Example 2.1.8 :** 1. Let \mathbb{Z} act on \mathbb{R} by translations. We obtain a cover $\mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z} \simeq S^1$.
 2. For an integer $n > 1$, we get an even action of μ_n on \mathbb{C}^* from that a cover $p_n : \mathbb{C}^* \rightarrow \mathbb{C}^*/\mu_n$. The map $z \mapsto z^n$ defines a natural homeomorphism (isomorphism) of \mathbb{C}^*/μ_n onto \mathbb{C}^* . So this induces a cover $\mathbb{C}^* \rightarrow \mathbb{C}^*$.
- For now on, assume that X is locally connected.
- $\text{Aut}(Y/X) := \{\phi : Y \rightarrow Y : p \circ \phi = p\}$.
- For each point $x \in X$, $p^{-1}(x)$ is equipped with a natural action of $\text{Aut}(Y/X)$.

- **Lemma 2.1.7 :** If G is a group acting evenly on a connected space Y , the projection $p_G : Y \rightarrow G \backslash Y$ turns Y into a cover of $G \backslash Y$.
- **Example 2.1.8 :** 1. Let \mathbb{Z} act on \mathbb{R} by translations. We obtain a cover $\mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z} \simeq S^1$.
2. For an integer $n > 1$, we get an even action of μ_n on \mathbb{C}^* from that a cover $p_n : \mathbb{C}^* \rightarrow \mathbb{C}^*/\mu_n$. The map $z \mapsto z^n$ defines a natural homeomorphism (isomorphism) of \mathbb{C}^*/μ_n onto \mathbb{C}^* . So this induces a cover $\mathbb{C}^* \rightarrow \mathbb{C}^*$.
- For now on, assume that X is locally connected.
- $\text{Aut}(Y/X) := \{\phi : Y \rightarrow Y : p \circ \phi = p\}$.
- For each point $x \in X$, $p^{-1}(x)$ is equipped with a natural action of $\text{Aut}(Y/X)$.

- **Lemma 2.1.7 :** If G is a group acting evenly on a connected space Y , the projection $p_G : Y \rightarrow G \backslash Y$ turns Y into a cover of $G \backslash Y$.
- **Example 2.1.8 :** 1. Let \mathbb{Z} act on \mathbb{R} by translations. We obtain a cover $\mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z} \simeq S^1$.
2. For an integer $n > 1$, we get an even action of μ_n on \mathbb{C}^* from that a cover $p_n : \mathbb{C}^* \rightarrow \mathbb{C}^*/\mu_n$. The map $z \mapsto z^n$ defines a natural homeomorphism (isomorphism) of \mathbb{C}^*/μ_n onto \mathbb{C}^* . So this induces a cover $\mathbb{C}^* \rightarrow \mathbb{C}^*$.
- For now on, assume that X is locally connected.
- $\text{Aut}(Y/X) := \{\phi : Y \rightarrow Y : p \circ \phi = p\}$.
- For each point $x \in X$, $p^{-1}(x)$ is equipped with a natural action of $\text{Aut}(Y/X)$.

- **Lemma 2.1.7 :** If G is a group acting evenly on a connected space Y , the projection $p_G : Y \rightarrow G \backslash Y$ turns Y into a cover of $G \backslash Y$.
- **Example 2.1.8 :** 1. Let \mathbb{Z} act on \mathbb{R} by translations. We obtain a cover $\mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z} \simeq S^1$.
2. For an integer $n > 1$, we get an even action of μ_n on \mathbb{C}^* from that a cover $p_n : \mathbb{C}^* \rightarrow \mathbb{C}^*/\mu_n$. The map $z \mapsto z^n$ defines a natural homeomorphism (isomorphism) of \mathbb{C}^*/μ_n onto \mathbb{C}^* . So this induces a cover $\mathbb{C}^* \rightarrow \mathbb{C}^*$.
- For now on, assume that X is locally connected.
- $\text{Aut}(Y/X) := \{\phi : Y \rightarrow Y : p \circ \phi = p\}$.
- For each point $x \in X$, $p^{-1}(x)$ is equipped with a natural action of $\text{Aut}(Y/X)$.

- **Lemma 2.1.7 :** If G is a group acting evenly on a connected space Y , the projection $p_G : Y \rightarrow G \backslash Y$ turns Y into a cover of $G \backslash Y$.
- **Example 2.1.8 :** 1. Let \mathbb{Z} act on \mathbb{R} by translations. We obtain a cover $\mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z} \simeq S^1$.
2. For an integer $n > 1$, we get an even action of μ_n on \mathbb{C}^* from that a cover $p_n : \mathbb{C}^* \rightarrow \mathbb{C}^*/\mu_n$. The map $z \mapsto z^n$ defines a natural homeomorphism (isomorphism) of \mathbb{C}^*/μ_n onto \mathbb{C}^* . So this induces a cover $\mathbb{C}^* \rightarrow \mathbb{C}^*$.
- For now on, assume that X is locally connected.
- $\text{Aut}(Y/X) := \{\phi : Y \rightarrow Y : p \circ \phi = p\}$.
- For each point $x \in X$, $p^{-1}(x)$ is equipped with a natural action of $\text{Aut}(Y/X)$.

- **Lemma 2.1.7 :** If G is a group acting evenly on a connected space Y , the projection $p_G : Y \rightarrow G \backslash Y$ turns Y into a cover of $G \backslash Y$.
- **Example 2.1.8 :** 1. Let \mathbb{Z} act on \mathbb{R} by translations. We obtain a cover $\mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z} \simeq S^1$.
2. For an integer $n > 1$, we get an even action of μ_n on \mathbb{C}^* from that a cover $p_n : \mathbb{C}^* \rightarrow \mathbb{C}^*/\mu_n$. The map $z \mapsto z^n$ defines a natural homeomorphism (isomorphism) of \mathbb{C}^*/μ_n onto \mathbb{C}^* . So this induces a cover $\mathbb{C}^* \rightarrow \mathbb{C}^*$.
- For now on, assume that X is locally connected.
- $\text{Aut}(Y/X) := \{\phi : Y \rightarrow Y : p \circ \phi = p\}$.
- For each point $x \in X$, $p^{-1}(x)$ is equipped with a natural action of $\text{Aut}(Y/X)$.

- **Proposition 2.2.2 :** Let $p : X \rightarrow Y$ be a cover, Z a connected topological space, $f, g : Z \rightarrow Y$ two continuous maps satisfying $p \circ f = p \circ g$. If there is a point $z \in Z$ with $f(z) = g(z)$, then $f \equiv g$.
- Apply the above proposition to $Z = Y$, $f = id$ and $g = \phi$ to get :

Lemma 2.2.1 An automorphism ϕ of a connected cover $p : Y \rightarrow X$ having a fixed point must be trivial.

- If $p : Y \rightarrow X$ is a connected cover, the action of $\text{Aut}(Y/X)$ on Y is even.
- Conversely :

Proposition 2.2.4 : If G is a group acting evenly on a connected space Y , the automorphism group of the cover $p_G : Y \rightarrow G \backslash Y$ is precisely G .

$$Y \rightarrow \text{Aut}(Y/X) \backslash Y \xrightarrow{\bar{p}} X.$$

- **Proposition 2.2.2** : Let $p : X \rightarrow Y$ be a cover, Z a connected topological space, $f, g : Z \rightarrow Y$ two continuous maps satisfying $p \circ f = p \circ g$. If there is a point $z \in Z$ with $f(z) = g(z)$, then $f \equiv g$.
- Apply the above proposition to $Z = Y$, $f = id$ and $g = \phi$ to get :

Lemma 2.2.1 An automorphism ϕ of a connected cover $p : Y \rightarrow X$ having a fixed point must be trivial.

- If $p : Y \rightarrow X$ is a connected cover, the action of $\text{Aut}(Y/X)$ on Y is even.
- Conversely :

Proposition 2.2.4 : If G is a group acting evenly on a connected space Y , the automorphism group of the cover $p_G : Y \rightarrow G \backslash Y$ is precisely G .

$$Y \rightarrow \text{Aut}(Y/X) \backslash Y \xrightarrow{\bar{p}} X.$$

- **Proposition 2.2.2** : Let $p : X \rightarrow Y$ be a cover, Z a connected topological space, $f, g : Z \rightarrow Y$ two continuous maps satisfying $p \circ f = p \circ g$. If there is a point $z \in Z$ with $f(z) = g(z)$, then $f \equiv g$.
- Apply the above proposition to $Z = Y$, $f = id$ and $g = \phi$ to get :

Lemma 2.2.1 An automorphism ϕ of a connected cover $p : Y \rightarrow X$ having a fixed point must be trivial.

- If $p : Y \rightarrow X$ is a connected cover, the action of $\text{Aut}(Y/X)$ on Y is even.
- Conversely :

Proposition 2.2.4 : If G is a group acting evenly on a connected space Y , the automorphism group of the cover $p_G : Y \rightarrow G \backslash Y$ is precisely G .

$$Y \rightarrow \text{Aut}(Y/X) \backslash Y \xrightarrow{\bar{p}} X.$$

- **Proposition 2.2.2 :** Let $p : X \rightarrow Y$ be a cover, Z a connected topological space, $f, g : Z \rightarrow Y$ two continuous maps satisfying $p \circ f = p \circ g$. If there is a point $z \in Z$ with $f(z) = g(z)$, then $f \equiv g$.
- Apply the above proposition to $Z = Y$, $f = id$ and $g = \phi$ to get :

Lemma 2.2.1 An automorphism ϕ of a connected cover $p : Y \rightarrow X$ having a fixed point must be trivial.

- If $p : Y \rightarrow X$ is a connected cover, the action of $\text{Aut}(Y/X)$ on Y is even.
- Conversely :

Proposition 2.2.4 : If G is a group acting evenly on a connected space Y , the automorphism group of the cover $p_G : Y \rightarrow G \backslash Y$ is precisely G .

$$Y \rightarrow \text{Aut}(Y/X) \backslash Y \xrightarrow{\bar{p}} X.$$

- **Proposition 2.2.2** : Let $p : X \rightarrow Y$ be a cover, Z a connected topological space, $f, g : Z \rightarrow Y$ two continuous maps satisfying $p \circ f = p \circ g$. If there is a point $z \in Z$ with $f(z) = g(z)$, then $f \equiv g$.
- Apply the above proposition to $Z = Y$, $f = id$ and $g = \phi$ to get :

Lemma 2.2.1 An automorphism ϕ of a connected cover $p : Y \rightarrow X$ having a fixed point must be trivial.

- If $p : Y \rightarrow X$ is a connected cover, the action of $\text{Aut}(Y/X)$ on Y is even.
- Conversely :

Proposition 2.2.4 : If G is a group acting evenly on a connected space Y , the automorphism group of the cover $p_G : Y \rightarrow G \backslash Y$ is precisely G .

$$Y \rightarrow \text{Aut}(Y/X) \backslash Y \xrightarrow{\bar{p}} X.$$

- **Proposition 2.2.2 :** Let $p : X \rightarrow Y$ be a cover, Z a connected topological space, $f, g : Z \rightarrow Y$ two continuous maps satisfying $p \circ f = p \circ g$. If there is a point $z \in Z$ with $f(z) = g(z)$, then $f \equiv g$.
- Apply the above proposition to $Z = Y$, $f = id$ and $g = \phi$ to get :

Lemma 2.2.1 An automorphism ϕ of a connected cover $p : Y \rightarrow X$ having a fixed point must be trivial.

- If $p : Y \rightarrow X$ is a connected cover, the action of $\text{Aut}(Y/X)$ on Y is even.
- Conversely :

Proposition 2.2.4 : If G is a group acting evenly on a connected space Y , the automorphism group of the cover $p_G : Y \rightarrow G \backslash Y$ is precisely G .

$$Y \rightarrow \text{Aut}(Y/X) \backslash Y \xrightarrow{\bar{p}} X.$$

- Given a connected cover $p : Y \rightarrow X$, form the quotient of Y by the action of $\text{Aut}(Y/X)$. The projection factors as :

$$Y \longrightarrow \text{Aut}(Y/X) \backslash Y \xrightarrow{\bar{p}} X$$

- A cover $p : Y \rightarrow X$ is said to be **Galois** if Y is connected and the induced map \bar{p} above is a homeomorphism.
- Proposition 2.2.7** A connected cover $p : Y \rightarrow X$ is Galois iff $\text{Aut}(Y/X)$ acts transitively on each fibre of p .
- We will need the following lemma :
Lemma 2.2.11 Let $q : Z \rightarrow X$ be a connected cover and $f : Y \rightarrow Z$ a continuous map. If the composite $q \circ f : Y \rightarrow X$ is a cover, then so is $f : Y \rightarrow Z$.

- Given a connected cover $p : Y \rightarrow X$, form the quotient of Y by the action of $\text{Aut}(Y/X)$. The projection factors as :

$$Y \longrightarrow \text{Aut}(Y/X) \backslash Y \xrightarrow{\bar{p}} X$$

- A cover $p : Y \rightarrow X$ is said to be **Galois** if Y is connected and the induced map \bar{p} above is a homeomorphism.
- Proposition 2.2.7** A connected cover $p : Y \rightarrow X$ is Galois iff $\text{Aut}(Y/X)$ acts transitively on each fibre of p .
- We will need the following lemma :
Lemma 2.2.11 Let $q : Z \rightarrow X$ be a connected cover and $f : Y \rightarrow Z$ a continuous map. If the composite $q \circ f : Y \rightarrow X$ is a cover, then so is $f : Y \rightarrow Z$.

- Given a connected cover $p : Y \rightarrow X$, form the quotient of Y by the action of $\text{Aut}(Y/X)$. The projection factors as :

$$Y \longrightarrow \text{Aut}(Y/X) \backslash Y \xrightarrow{\bar{p}} X$$

- A cover $p : Y \rightarrow X$ is said to be **Galois** if Y is connected and the induced map \bar{p} above is a homeomorphism.
- Proposition 2.2.7** A connected cover $p : Y \rightarrow X$ is Galois iff $\text{Aut}(Y/X)$ acts transitively on each fibre of p .
- We will need the following lemma :

Lemma 2.2.11 Let $q : Z \rightarrow X$ be a connected cover and $f : Y \rightarrow Z$ a continuous map. If the composite $q \circ f : Y \rightarrow X$ is a cover, then so is $f : Y \rightarrow Z$.

- Given a connected cover $p : Y \rightarrow X$, form the quotient of Y by the action of $\text{Aut}(Y/X)$. The projection factors as :

$$Y \longrightarrow \text{Aut}(Y/X) \backslash Y \xrightarrow{\bar{p}} X$$

- A cover $p : Y \rightarrow X$ is said to be **Galois** if Y is connected and the induced map \bar{p} above is a homeomorphism.
- Proposition 2.2.7** A connected cover $p : Y \rightarrow X$ is Galois iff $\text{Aut}(Y/X)$ acts transitively on each fibre of p .
- We will need the following lemma :

Lemma 2.2.11 Let $q : Z \rightarrow X$ be a connected cover and $f : Y \rightarrow Z$ a continuous map. If the composite $q \circ f : Y \rightarrow X$ is a cover, then so is $f : Y \rightarrow Z$.

- Given a connected cover $p : Y \rightarrow X$, form the quotient of Y by the action of $\text{Aut}(Y/X)$. The projection factors as :

$$Y \longrightarrow \text{Aut}(Y/X) \backslash Y \xrightarrow{\bar{p}} X$$

- A cover $p : Y \rightarrow X$ is said to be **Galois** if Y is connected and the induced map \bar{p} above is a homeomorphism.
- Proposition 2.2.7** A connected cover $p : Y \rightarrow X$ is Galois iff $\text{Aut}(Y/X)$ acts transitively on each fibre of p .
- We will need the following lemma :
Lemma 2.2.11 Let $q : Z \rightarrow X$ be a connected cover and $f : Y \rightarrow Z$ a continuous map. If the composite $q \circ f : Y \rightarrow X$ is a cover, then so is $f : Y \rightarrow Z$.

- Let $p : Y \rightarrow X$ be a Galois cover and $G = \text{Aut}(Y|X)$. The map $H \mapsto H \backslash Y, Z \mapsto \text{Aut}(Y|Z)$ induce a bijection between subgroups of G and intermediate covers Z as above.

$$\begin{array}{ccc}
 Y & \xrightarrow{f} & Z \\
 & \searrow p & \downarrow q \\
 & & X
 \end{array}$$

The cover $q : Z \rightarrow X$ is Galois iff H is a normal subgroup of G , in which case $\text{Aut}(Z|X) \cong G/H$.

- Let $p : Y \rightarrow X$ be a Galois cover and $G = \text{Aut}(Y|X)$. The map $H \mapsto H \backslash Y$, $Z \mapsto \text{Aut}(Y|Z)$ induce a bijection between subgroups of G and intermediate covers Z as above.

$$\begin{array}{ccc}
 Y & \xrightarrow{f} & Z \\
 & \searrow p & \downarrow q \\
 & & X
 \end{array}$$

The cover $q : Z \rightarrow X$ is Galois iff H is a normal subgroup of G , in which case $\text{Aut}(Z|X) \cong G/H$.

The Monodromy Action

- **Lemma 2.3.2** Let $p : Y \rightarrow X$ be a cover, $y \in Y$ and $x = p(y)$.
 1. Given a path $f : [0, 1] \rightarrow X$ with $f(0) = x$, there is a unique path $\tilde{f} : [0, 1] \rightarrow Y$ with $p \circ \tilde{f} = f$.
 2. Assume moreover given a second path $g : [0, 1] \rightarrow X$ homotopic to f . Then the unique $\tilde{g} : [0, 1] \rightarrow Y$ with $\tilde{g}(0) = y$ and $p \circ \tilde{g} = g$ has the same endpoint as \tilde{f} , i.e. $\tilde{f}(1) = \tilde{g}(1)$.
- *Monodromy Action* on the fibre $p^{-1}(x)$: Given $y \in p^{-1}(x)$ and $\alpha \in \pi_1(X, x)$ represented by a path $f : [0, 1] \rightarrow X$ with $f(0) = f(1) = x$. We define

$$\alpha y := \tilde{f}(1),$$

where \tilde{f} is the unique lifting \tilde{f} to Y with $\tilde{f}(0) = y$. This gives a left action of $\pi_1(X, x)$ on $p^{-1}(x)$

The Monodromy Action

- **Lemma 2.3.2** Let $p : Y \rightarrow X$ be a cover, $y \in Y$ and $x = p(y)$.
 1. Given a path $f : [0, 1] \rightarrow X$ with $f(0) = x$, there is a unique path $\tilde{f} : [0, 1] \rightarrow Y$ with $p \circ \tilde{f} = f$.
 2. Assume moreover given a second path $g : [0, 1] \rightarrow X$ homotopic to f . Then the unique $\tilde{g} : [0, 1] \rightarrow Y$ with $\tilde{g}(0) = y$ and $p \circ \tilde{g} = g$ has the same endpoint as \tilde{f} , i.e. $\tilde{f}(1) = \tilde{g}(1)$.
- *Monodromy Action* on the fibre $p^{-1}(x)$: Given $y \in p^{-1}(x)$ and $\alpha \in \pi_1(X, x)$ represented by a path $f : [0, 1] \rightarrow X$ with $f(0) = f(1) = x$. We define

$$\alpha y := \tilde{f}(1),$$

where \tilde{f} is the unique lifting \tilde{f} to Y with $\tilde{f}(0) = y$. This gives a left action of $\pi_1(X, x)$ on $p^{-1}(x)$

The Monodromy Action

- **Lemma 2.3.2** Let $p : Y \rightarrow X$ be a cover, $y \in Y$ and $x = p(y)$.
 1. Given a path $f : [0, 1] \rightarrow X$ with $f(0) = x$, there is a unique path $\tilde{f} : [0, 1] \rightarrow Y$ with $p \circ \tilde{f} = f$.
 2. Assume moreover given a second path $g : [0, 1] \rightarrow X$ homotopic to f . Then the unique $\tilde{g} : [0, 1] \rightarrow Y$ with $\tilde{g}(0) = y$ and $p \circ \tilde{g} = g$ has the same endpoint as \tilde{f} , i.e. $\tilde{f}(1) = \tilde{g}(1)$.
- *Monodromy Action on the fibre $p^{-1}(x)$* : Given $y \in p^{-1}(x)$ and $\alpha \in \pi_1(X, x)$ represented by a path $f : [0, 1] \rightarrow X$ with $f(0) = f(1) = x$. We define

$$\alpha y := \tilde{f}(1),$$

where \tilde{f} is the unique lifting \tilde{f} to Y with $\tilde{f}(0) = y$. This gives a left action of $\pi_1(X, x)$ on $p^{-1}(x)$

The Monodromy Action

- **Lemma 2.3.2** Let $p : Y \rightarrow X$ be a cover, $y \in Y$ and $x = p(y)$.
 1. Given a path $f : [0, 1] \rightarrow X$ with $f(0) = x$, there is a unique path $\tilde{f} : [0, 1] \rightarrow Y$ with $p \circ \tilde{f} = f$.
 2. Assume moreover given a second path $g : [0, 1] \rightarrow X$ homotopic to f . Then the unique $\tilde{g} : [0, 1] \rightarrow Y$ with $\tilde{g}(0) = y$ and $p \circ \tilde{g} = g$ has the same endpoint as \tilde{f} , i.e. $\tilde{f}(1) = \tilde{g}(1)$.
- *Monodromy Action on the fibre $p^{-1}(x)$:* Given $y \in p^{-1}(x)$ and $\alpha \in \pi_1(X, x)$ represented by a path $f : [0, 1] \rightarrow X$ with $f(0) = f(1) = x$. We define

$$\alpha y := \tilde{f}(1),$$

where \tilde{f} is the unique lifting \tilde{f} to Y with $\tilde{f}(0) = y$. This gives a left action of $\pi_1(X, x)$ on $p^{-1}(x)$

The Monodromy Action

- **Lemma 2.3.2** Let $p : Y \rightarrow X$ be a cover, $y \in Y$ and $x = p(y)$.
 1. Given a path $f : [0, 1] \rightarrow X$ with $f(0) = x$, there is a unique path $\tilde{f} : [0, 1] \rightarrow Y$ with $p \circ \tilde{f} = f$.
 2. Assume moreover given a second path $g : [0, 1] \rightarrow X$ homotopic to f . Then the unique $\tilde{g} : [0, 1] \rightarrow Y$ with $\tilde{g}(0) = y$ and $p \circ \tilde{g} = g$ has the same endpoint as \tilde{f} , i.e. $\tilde{f}(1) = \tilde{g}(1)$.
- *Monodromy Action* on the fibre $p^{-1}(x)$: Given $y \in p^{-1}(x)$ and $\alpha \in \pi_1(X, x)$ represented by a path $f : [0, 1] \rightarrow X$ with $f(0) = f(1) = x$. We define

$$\alpha y := \tilde{f}(1),$$

where \tilde{f} is the unique lifting \tilde{f} to Y with $\tilde{f}(0) = y$. This gives a left action of $\pi_1(X, x)$ on $p^{-1}(x)$

- Fix a space X and $x \in X$. We define a functor Fib_X from the category of covers of X to the category of sets equipped with a left $\pi_1(X, x)$ -action by sending a cover $p: Y \rightarrow X$ to the fibre $p^{-1}(x)$.
- **Theorem 2.3.4** Let X be connected and locally simply connected top. space and $x \in X$ a base point. The functor Fib_X induces an equivalence of the category of covers of X with the category of left $\pi_1(X, x)$ -sets.
Connected covers correspond to $\pi_1(X, x)$ -sets with transitive action and Galois covers to coset spaces of normal subgroups.

- Fix a space X and $x \in X$. We define a functor Fib_x from the category of covers of X to the category of sets equipped with a left $\pi_1(X, x)$ -action by sending a cover $p: Y \rightarrow X$ to the fibre $p^{-1}(x)$.
- **Theorem 2.3.4** Let X be connected and locally simply connected top. space and $x \in X$ a base point. The functor Fib_x induces an equivalence of the category of covers of X with the category of left $\pi_1(X, x)$ -sets.
Connected covers correspond to $\pi_1(X, x)$ -sets with transitive action and Galois covers to coset spaces of normal subgroups.

- Fix a space X and $x \in X$. We define a functor Fib_x from the category of covers of X to the category of sets equipped with a left $\pi_1(X, x)$ -action by sending a cover $p: Y \rightarrow X$ to the fibre $p^{-1}(x)$.
- **Theorem 2.3.4** Let X be connected and locally simply connected top. space and $x \in X$ a base point. The functor Fib_x induces an equivalence of the category of covers of X with the category of left $\pi_1(X, x)$ -sets.
Connected covers correspond to $\pi_1(X, x)$ -sets with transitive action and Galois covers to coset spaces of normal subgroups.

- The proof the Theorem 2.3.4 relies on two facts :
- For a connected and locally simply connected topological space X and a base point $x \in X$ the functor Fib_x is representable by a cover $\tilde{X}_x \rightarrow X$, i.e.
 $\text{Fib}_x(.) \cong \text{Hom}(\tilde{X}_x, .)$.
- The cover \tilde{X}_x depends on the choice of the base point x .
- It comes equipped with a canonical point in the fibre $\pi^{-1}(x)$ called the *universal element*, denotes \tilde{x} .
- For an arbitrary cover $p : Y \rightarrow X$ and element $y \in \pi^{-1}(x)$, the cover map $\pi^{-1}(y) : \tilde{X}_x \rightarrow Y$ corresponding to y via the isomorphism $\text{Fib}_x(Y) \cong \text{Hom}(\tilde{X}_x, Y)$ maps \tilde{x} to y .

- The proof the Theorem 2.3.4 relies on two facts :
- For a connected and locally simply connected topological space X and a base point $x \in X$ the functor Fib_x is representable by a cover $\tilde{X}_x \rightarrow X$, i.e.
 $\text{Fib}_x(.) \cong \text{Hom}(\tilde{X}_x, .)$.
- The cover \tilde{X}_x depends on the choice of the base point x .
- It comes equipped with a canonical point in the fibre $\pi^{-1}(x)$ called the *universal element*, denotes \tilde{x} .
- For an arbitrary cover $p : Y \rightarrow X$ and element $y \in \pi^{-1}(x)$, the cover map $\pi^{-1}(y) : \tilde{X}_x \rightarrow Y$ corresponding to y via the isomorphism $\text{Fib}_x(Y) \cong \text{Hom}(\tilde{X}_x, Y)$ maps \tilde{x} to y .

- The proof the Theorem 2.3.4 relies on two facts :
- For a connected and locally simply connected topological space X and a base point $x \in X$ the functor Fib_x is representable by a cover $\tilde{X}_x \rightarrow X$, i.e.
 $\text{Fib}_x(.) \cong \text{Hom}(\tilde{X}_x, .)$.
- The cover \tilde{X}_x depends on the choice of the base point x .
- It comes equipped with a canonical point in the fibre $\pi^{-1}(x)$ called the *universal element*, denotes \tilde{x} .
- For an arbitrary cover $p : Y \rightarrow X$ and element $y \in \pi^{-1}(x)$, the cover map $\pi^{-1}(y) : \tilde{X}_x \rightarrow Y$ corresponding to y via the isomorphism $\text{Fib}_x(Y) \cong \text{Hom}(\tilde{X}_x, Y)$ maps \tilde{x} to y .

- The proof the Theorem 2.3.4 relies on two facts :
- For a connected and locally simply connected topological space X and a base point $x \in X$ the functor Fib_x is representable by a cover $\tilde{X}_x \rightarrow X$, i.e.
 $\text{Fib}_x(.) \cong \text{Hom}(\tilde{X}_x, .)$.
- The cover \tilde{X}_x depends on the choice of the base point x .
- It comes equipped with a canonical point in the fibre $\pi^{-1}(x)$ called the *universal element*, denotes \tilde{x} .
- For an arbitrary cover $p : Y \rightarrow X$ and element $y \in \pi^{-1}(x)$, the cover map $\pi^{-1}(y) : \tilde{X}_x \rightarrow Y$ corresponding to y via the isomorphism $\text{Fib}_x(Y) \cong \text{Hom}(\tilde{X}_x, Y)$ maps \tilde{x} to y .

- The proof the Theorem 2.3.4 relies on two facts :
- For a connected and locally simply connected topological space X and a base point $x \in X$ the functor Fib_x is representable by a cover $\tilde{X}_x \rightarrow X$, i.e.
 $\text{Fib}_x(.) \cong \text{Hom}(\tilde{X}_x, .)$.
- The cover \tilde{X}_x depends on the choice of the base point x .
- It comes equipped with a canonical point in the fibre $\pi^{-1}(x)$ called the *universal element*, denotes \tilde{x} .
- For an arbitrary cover $p : Y \rightarrow X$ and element $y \in \pi^{-1}(x)$, the cover map $\pi^{-1}(y) : \tilde{X}_x \rightarrow Y$ corresponding to y via the isomorphism $\text{Fib}_x(Y) \cong \text{Hom}(\tilde{X}_x, Y)$ maps \tilde{x} to y .

- The proof the Theorem 2.3.4 relies on two facts :
- For a connected and locally simply connected topological space X and a base point $x \in X$ the functor Fib_x is representable by a cover $\tilde{X}_x \rightarrow X$, i.e.
 $\text{Fib}_x(.) \cong \text{Hom}(\tilde{X}_x, .)$.
- The cover \tilde{X}_x depends on the choice of the base point x .
- It comes equipped with a canonical point in the fibre $\pi^{-1}(x)$ called the *universal element*, denotes \tilde{x} .
- For an arbitrary cover $p : Y \rightarrow X$ and element $y \in \pi^{-1}(x)$, the cover map $\pi^{-1}(y) : \tilde{X}_x \rightarrow Y$ corresponding to y via the isomorphism $\text{Fib}_x(Y) \cong \text{Hom}(\tilde{X}_x, Y)$ maps \tilde{x} to y .

- Recovering the monodromy action : We can obtain a right action on $\text{Fib}_x(Y) \cong \text{Hom}(\tilde{X}_x, Y)$ from the left action of $\text{Aut}(\tilde{X}_x|X)$ on \tilde{X}_x .
- How is this action related to the monodromy action ?
- **Theorem 2.3.7** : The cover \tilde{X}_x is a connected Galois cover of X , with automorphism group isomorphic to $\pi_1(X, x)$, i.e. $\text{Aut}(\tilde{X}_x|X) \cong \pi_1(X, x)$.
Moreover, for each cover $Y \rightarrow X$ the left action of $\text{Aut}(\tilde{X}_x|X)^{\text{op}}$ on $\text{Fib}_x(Y)$ is exactly the monodromy action of $\pi_1(X, x)$.
- **Remark** : The cover \tilde{X}_x plays the role of a separable closure k_s . In other words, the choice of x corresponds to the choice of the separable closure.
The fundamental group is the counterpart of the absolute Galois group :
In the case of fields : $A \mapsto \text{Hom}(A, k_s)$,
in the topological case : $Y \mapsto \text{Hom}_X(\tilde{X}_x, Y)$.

- Recovering the monodromy action : We can obtain a right action on $\text{Fib}_x(Y) \cong \text{Hom}(\tilde{X}_x, Y)$ from the left action of $\text{Aut}(\tilde{X}_x|X)$ on \tilde{X}_x .
- How is this action related to the monodromy action ?
- **Theorem 2.3.7** : The cover \tilde{X}_x is a connected Galois cover of X , with automorphism group isomorphic to $\pi_1(X, x)$, i.e. $\text{Aut}(\tilde{X}_x|X) \cong \pi_1(X, x)$.
Moreover, for each cover $Y \rightarrow X$ the left action of $\text{Aut}(\tilde{X}_x|X)^{\text{op}}$ on $\text{Fib}_x(Y)$ is exactly the monodromy action of $\pi_1(X, x)$.
- **Remark** : The cover \tilde{X}_x plays the role of a separable closure k_s . In other words, the choice of x corresponds to the choice of the separable closure.
The fundamental group is the counterpart of the absolute Galois group :
In the case of fields : $A \mapsto \text{Hom}(A, k_s)$,
in the topological case : $Y \mapsto \text{Hom}_X(\tilde{X}_x, Y)$.

- Recovering the monodromy action : We can obtain a right action on $\text{Fib}_x(Y) \cong \text{Hom}(\tilde{X}_x, Y)$ from the left action of $\text{Aut}(\tilde{X}_x|X)$ on \tilde{X}_x .
- How is this action related to the monodromy action ?
- **Theorem 2.3.7** : The cover \tilde{X}_x is a connected Galois cover of X , with automorphism group isomorphic to $\pi_1(X, x)$, i.e. $\text{Aut}(\tilde{X}_x|X) \cong \pi_1(X, x)$.
Moreover, for each cover $Y \rightarrow X$ the left action of $\text{Aut}(\tilde{X}_x|X)^{\text{op}}$ on $\text{Fib}_x(Y)$ is exactly the monodromy action of $\pi_1(X, x)$.
- **Remark** : The cover \tilde{X}_x plays the role of a separable closure k_s . In other words, the choice of x corresponds to the choice of the separable closure.
The fundamental group is the counterpart of the absolute Galois group :
In the case of fields : $A \mapsto \text{Hom}(A, k_s)$,
in the topological case : $Y \mapsto \text{Hom}_X(\tilde{X}_x, Y)$.

- Recovering the monodromy action : We can obtain a right action on $\text{Fib}_x(Y) \cong \text{Hom}(\tilde{X}_x, Y)$ from the left action of $\text{Aut}(\tilde{X}_x|X)$ on \tilde{X}_x .
- How is this action related to the monodromy action ?
- **Theorem 2.3.7** : The cover \tilde{X}_x is a connected Galois cover of X , with automorphism group isomorphic to $\pi_1(X, x)$, i.e. $\text{Aut}(\tilde{X}_x|X) \cong \pi_1(X, x)$.
Moreover, for each cover $Y \rightarrow X$ the left action of $\text{Aut}(\tilde{X}_x|X)^{\text{op}}$ on $\text{Fib}_x(Y)$ is exactly the monodromy action of $\pi_1(X, x)$.
- **Remark** : The cover \tilde{X}_x plays the role of a separable closure k_s . In other words, the choice of x corresponds to the choice of the separable closure.
The fundamental group is the counterpart of the absolute Galois group :
In the case of fields : $A \mapsto \text{Hom}(A, k_s)$,
in the topological case : $Y \mapsto \text{Hom}_X(\tilde{X}_x, Y)$.

- Recovering the monodromy action : We can obtain a right action on $\text{Fib}_x(Y) \cong \text{Hom}(\tilde{X}_x, Y)$ from the left action of $\text{Aut}(\tilde{X}_x|X)$ on \tilde{X}_x .
- How is this action related to the monodromy action ?
- **Theorem 2.3.7** : The cover \tilde{X}_x is a connected Galois cover of X , with automorphism group isomorphic to $\pi_1(X, x)$, i.e. $\text{Aut}(\tilde{X}_x|X) \cong \pi_1(X, x)$.
Moreover, for each cover $Y \rightarrow X$ the left action of $\text{Aut}(\tilde{X}_x|X)^{\text{op}}$ on $\text{Fib}_x(Y)$ is exactly the monodromy action of $\pi_1(X, x)$.
- **Remark** : The cover \tilde{X}_x plays the role of a separable closure k_s . In other words, the choice of x corresponds to the choice of the separable closure.

The fundamental group is the counterpart of the absolute Galois group :

In the case of fields : $A \mapsto \text{Hom}(A, k_s)$,

in the topological case : $Y \mapsto \text{Hom}_X(\tilde{X}_x, Y)$.

- Recovering the monodromy action : We can obtain a right action on $\text{Fib}_x(Y) \cong \text{Hom}(\tilde{X}_x, Y)$ from the left action of $\text{Aut}(\tilde{X}_x|X)$ on \tilde{X}_x .
- How is this action related to the monodromy action ?
- **Theorem 2.3.7** : The cover \tilde{X}_x is a connected Galois cover of X , with automorphism group isomorphic to $\pi_1(X, x)$, i.e. $\text{Aut}(\tilde{X}_x|X) \cong \pi_1(X, x)$.
Moreover, for each cover $Y \rightarrow X$ the left action of $\text{Aut}(\tilde{X}_x|X)^{\text{op}}$ on $\text{Fib}_x(Y)$ is exactly the monodromy action of $\pi_1(X, x)$.
- **Remark** : The cover \tilde{X}_x plays the role of a separable closure k_s . In other words, the choice of x corresponds to the choice of the separable closure.
The fundamental group is the counterpart of the absolute Galois group :
In the case of fields : $A \mapsto \text{Hom}(A, k_s)$,
in the topological case : $Y \mapsto \text{Hom}_X(\tilde{X}_x, Y)$.

- A cover $Y \rightarrow X$ is finite if it has finite fibres. For connected X , these have the same cardinality, called the *degree* of X .
- **Corollary 2.3.9** : For X and x as before, the functor Fib_x induces an equivalence of the category of finite covers of X with the category of finite continuous $\widehat{\pi_1(X, x)}$ -sets. Connected covers correspond to finite $\widehat{\pi_1(X, x)}$ -sets with transitive action and Galois covers to coset spaces of open normal subgroups.
- Here $\widehat{\pi_1(X, x)}$ denotes the profinite completion of $\pi_1(X, x)$:

$$\widehat{\pi_1(X, x)} = \varprojlim_U \pi_1(X, x)/U,$$

where U runs over normal subgroups of $\pi_1(X, x)$ of finite index.

- A cover $Y \rightarrow X$ is finite if it has finite fibres. For connected X , these have the same cardinality, called the *degree* of X .
- **Corollary 2.3.9** : For X and x as before, the functor Fib_x induces an equivalence of the category of finite covers of X with the category of finite continuous $\widehat{\pi_1(X, x)}$ -sets. Connected covers correspond to finite $\widehat{\pi_1(X, x)}$ -sets with transitive action and Galois covers to coset spaces of open normal subgroups.
- Here $\widehat{\pi_1(X, x)}$ denotes the profinite completion of $\pi_1(X, x)$:

$$\widehat{\pi_1(X, x)} = \varprojlim_U \pi_1(X, x)/U,$$

where U runs over normal subgroups of $\pi_1(X, x)$ of finite index.

- A cover $Y \rightarrow X$ is finite if it has finite fibres. For connected X , these have the same cardinality, called the *degree* of X .
- **Corollary 2.3.9** : For X and x as before, the functor Fib_x induces an equivalence of the category of finite covers of X with the category of finite continuous left $\widehat{\pi_1(X, x)}$ -sets. Connected covers correspond to finite $\widehat{\pi_1(X, x)}$ -sets with transitive action and Galois covers to coset spaces of open normal subgroups.
- Here $\widehat{\pi_1(X, x)}$ denotes the profinite completion of $\pi_1(X, x)$:

$$\widehat{\pi_1(X, x)} = \varprojlim_U \pi_1(X, x)/U,$$

where U runs over normal subgroups of $\pi_1(X, x)$ of finite index.

- A cover $Y \rightarrow X$ is finite if it has finite fibres. For connected X , these have the same cardinality, called the *degree* of X .
- **Corollary 2.3.9** : For X and x as before, the functor Fib_x induces an equivalence of the category of finite covers of X with the category of finite continuous $\widehat{\pi_1(X, x)}$ -sets. Connected covers correspond to finite $\widehat{\pi_1(X, x)}$ -sets with transitive action and Galois covers to coset spaces of open normal subgroups.
- Here $\widehat{\pi_1(X, x)}$ denotes the profinite completion of $\pi_1(X, x)$:

$$\widehat{\pi_1(X, x)} = \varprojlim_U \pi_1(X, x)/U ,$$

where U runs over normal subgroups of $\pi_1(X, x)$ of finite index.