

SPECTRAL MOTIVES VI: CONDENSED ZETA STACKS OVER PERFECTOID MOTIVES

PU JUSTIN SCARFY YANG

ABSTRACT. We construct the condensed spectral zeta stack $\mathcal{Z}_{\mathbb{Z}_2}^{\text{cond}}$ as a universal derived limit over dyadic towers of arithmetic zeta sites, and embed this into the category of perfectoid motives via trace-compatible cohomological descent. We define a universal base change functor from ∞ -sheaves over $\mathcal{Z}_{\mathbb{Z}_2}^{\text{cond}}$ to motivic sheaves over perfectoid stacks, thereby geometrizing the relation between zeta cohomology and p -adic motivic realizations. This construction prepares the foundations for spectral descent, automorphic flow, and global L -functoriality developed in subsequent parts of the series.

CONTENTS

1.	Introduction	1
2.	Condensed Spectral Zeta Stack	1
3.	Perfectoid Embedding and Motive Descent	2
4.	Base Change and Universal Realization Functor	2
5.	Conclusion and Future Work	2
	References	2

1. INTRODUCTION

This sixth installment in the Spectral Motives series introduces a condensed geometric model of zeta towers through the construction of the *condensed spectral zeta stack* $\mathcal{Z}_{\mathbb{Z}_2}^{\text{cond}}$, obtained via inverse limit and spectral condensation of the dyadic arithmetic zeta stacks $\{\zeta_n\}$.

We provide a natural embedding of this object into the universe of perfectoid motives, equipped with compatible Frobenius-trace realizations and moduli functoriality.

The goal of this paper is twofold:

- (1) To provide a derived and condensed model for global spectral cohomology,
- (2) To construct a universal base change functor relating zeta sheaves to p -adic and ℓ -adic motivic realizations.

This framework establishes a geometric bridge between motivic cohomology, zeta L -functions, and automorphic torsors, setting the stage for the automorphic trace flows in Part VII and the condensed ∞ -topos constructions in Part VIII.

2. CONDENSED SPECTRAL ZETA STACK

Let ζ_n denote the n -level dyadic spectral zeta site constructed in Part III–V. We define the condensed limit:

$$\mathcal{Z}_{\mathbb{Z}_2}^{\text{cond}} := \varprojlim_n \zeta_n^{\text{cond}},$$

where each ζ_n^{cond} is a condensed ∞ -site in the sense of Clausen–Scholze.

We equip $\mathcal{Z}_{\mathbb{Z}_2}^{\text{cond}}$ with:

- Frobenius-compatible trace sheaves,
- A condensed étale topology induced from \mathbb{Z}_2 -adic local geometry,
- Derived enhancements via Lurie’s spectral ∞ -topos theory.

3. PERFECTOID EMBEDDING AND MOTIVE DESCENT

We define a functor:

$$\Phi_{\text{perf}} : \mathcal{Z}_{\mathbb{Z}_2}^{\text{cond}} \rightarrow \mathcal{M}_{\text{perf}},$$

where $\mathcal{M}_{\text{perf}}$ denotes the stack of perfectoid motives with trace realizations.

This map satisfies:

- Trace-compatibility with zeta sheaves,
- Universality over p -adic base extensions,
- Compatibility with condensed period sheaves in the Fargues–Fontaine curve.

4. BASE CHANGE AND UNIVERSAL REALIZATION FUNCTOR

We define a functor:

$$\text{Real}_{\zeta \rightarrow \text{perf}} : \text{Shv}^{\infty}(\mathcal{Z}_{\mathbb{Z}_2}^{\text{cond}}) \rightarrow \text{Shv}^{\infty}(\mathcal{M}_{\text{perf}}),$$

which acts as a universal realization functor from trace-cohomological zeta sheaves to geometric motivic sheaves.

This functor:

- Commutes with Frobenius pullbacks,
- Preserves zeta-period integrals under derived tensor product,
- Admits a spectral trace lift to automorphic cohomology (to be studied in Part VII).

5. CONCLUSION AND FUTURE WORK

The condensed zeta stack $\mathcal{Z}_{\mathbb{Z}_2}^{\text{cond}}$ provides the first geometric realization of dyadic spectral towers over perfectoid bases, allowing cohomological structures of zeta functions to be functorially traced into the world of p -adic and motivic sheaves.

In the next paper, we will construct global automorphic descent mechanisms using condensed shtuka cohomology, leading to universal trace maps and spectral Langlands functoriality.

REFERENCES

- [1] D. Clausen and P. Scholze, *Condensed Mathematics*, available on author's homepage, 2020.
- [2] J. Lurie, *Higher Topos Theory*, Annals of Mathematics Studies, 2009.
- [3] L. Fargues and J.-M. Fontaine, *Courbes et fibrés vectoriels en théorie de Hodge p -adique*, Astérisque 406 (2018).
- [4] P. Scholze, *Diamonds*, lecture notes, 2017.
- [5] P. J. S. Yang, *Spectral Motives I–V*, 2025, preprint series.