

On the isogeny invariance of the Bloch-Kato's
Tamagawa Numbers conjecture:

a K -theoretic point of view

by

Justin Scarfy

B.Sc., The University Of British Columbia, 2017

A THESIS SUBMITTED IN PARTIAL FULFILLMENT
OF THE REQUIREMENTS FOR THE DEGREE OF

Master of Science

in

THE FACULTY OF GRADUATE STUDIES
(Mathematics)

The University Of British Columbia
(Vancouver)

June 2018

© Justin Scarfy, 2018

Abstract

A major task in mathematics today is to harmonize the continuous and the discrete, to include them in one comprehensive mathematics, and to eliminate obscurity from both. The arithmetic properties of various interesting objects (e.g. number fields, varieties, or even a motive, i.e. discrete objects) are all encoded in their respective L -functions, i.e. continuous objects, of which the Riemann zeta function is the simplest example. The understanding of these L -functions undergoes three phases [Kat1993a]:

1. analytic properties and the rationality of values,
2. algebraic properties illuminated by the p -adic properties of values,
3. arithmetic-geometric point of view of interpreting the values.

To date much of these have been achieved for the Riemann zeta function [CRSS2015] (except the locations of the zeros, or the zero free region). The deep conjectures of Deligne [De1979] and Beilinson [Be1985] were among the first to place these problems in a broad framework. Ambiguity in the work of Beilinson for interpreting these values, as they are interpreted only up to nonzero rational number multiples. In their article published in the Grothendieck Festschrift [BK1990], Spencer Bloch and Kazuya Kato removed this ambiguity in their formulation of what is now called the Tamagawa number conjecture (or the Bloch-Kato conjecture for motives). They also proved that the conjecture is invariant under isogeny, and the book [CRSS2015] exploits this isogeny invariance condition in the optic of K -theory for the Riemann zeta function.

This thesis investigates the isogeny invariance for specific motives and elaborates on accompanying results in cohomology and K -theory.

Table of Contents

Abstract	ii
Table of Contents	iii
List of Figures	iv
List of Symbols	v
Acknowledgments	vi
Introduction	1
0.1 The leading role played by arithmetic within mathematics and recent breakthroughs	1
0.2 On the Riemann Hypothesis	2
0.3 On the Hodge conjecture	3
0.4 On the Birch and Swinnerton-Dyer conjecture	4
1 Bloch-Kato’s Tamagawa Numbers conjecture and the isogeny invari- ance	7
1.1 From the Birch and Swinnerton-Dyer conjecture to the Bloch-Kato con- jecture	7
1.2 Motives, Fontaine’s p -adic period rings, and other essential gadgets . . .	8
1.3 The isogeny invariance explained	9
1.4 K -theoretic background	10
2 The isogeny invariance for elliptic curves with complex multiplication	11

2.1	CM elliptic curves	11
2.2	The isogeny invariance for CM elliptic curves	12
2.3	The K -theoretic viewpoint	13
3	The isogeny invariance for elliptic curves without complex multiplication	14
3.1	non-CM elliptic curves	14
3.2	The isogeny invariance for non-CM elliptic curves	15
3.3	The K -theoretic viewpoint	16
4	The isogeny invariance for modular forms	17
4.1	Modular forms	17
4.2	The isogeny invariance for modular forms	18
4.3	The K -theoretic viewpoint	19
5	The isogeny invariance for abelian varieties with complex multiplication	20
5.1	CM abelian varieties	20
5.2	The isogeny invariance for CM abelian varieties	21
5.3	The K -theoretic viewpoint	22
6	Proposed isogeny invariance for tori and motif	23
	Conclusion	24
	Bibliography	25

List of Figures

List of Symbols

Acknowledgments

First and foremost, I am deeply indebted to Professor Sujatha Ramdorai for providing me with extraordinary amounts of time and patience (since my first year undergraduate), sharing her seemingly endless bounty of mathematical knowledge, and challenging me with this exciting and multi-faceted problem. I have learned a great deal from our many conversations and feel privileged to have studied under her supervision. I am also very grateful for the funding I have received from NSERC, the department, and Professor Sujatha.

In addition, I am extremely grateful to Professor Greg Martin for his patience and suffering since my first year undergraduate studies in guiding me through the world of analytic number theory, enabling me to understand the many recent exciting breakthroughs in the field, especially in understanding the notorious Riemann Hypothesis, (and for reminding me that mathematician is a profession, not just a seeker of knowledge) and I want to thank Professor Lior Silberman for enabling me to understand its automorphic version. I also want to thank Professors Patrick Brosnan and Bruno Kahn on pointing me to the latest advances in the Hodge conjecture, to Professor Tai-Peng Tsai for his inspiring lectures on the latest results on the Navier-Stokes existence and smoothness problem, to Professor Izabella Łaba for her lectures with the latest results on the Kakeya conjecture, to Professor Ákos Magyar on his lectures on the Green-Tao Theorem, to Professor (MORE TO BE ADDED) , and to every faculty member here at UBC whom I have either taken or sat in on a course with, and for PIMS giving me the use of an office as a luxury to do uninterrupted work.

Studying at the Department of Mathematics at UBC has been wonderful with the extremely helpful and friendly department staff, particularly Roseann Kinsey and Marlowe Dickson. Involvement in the graduate student and number theory community has

been a great personal and educational experience, punctuated with many deep (non-)mathematical discussions, for which I would like to thank (COLLEAGUES) amongst many others.

Simply put, I am only here because of my parents, who have been selflessly supportive throughout my life, let alone my education. I dedicate this thesis to them, as their hardships and struggles have always been for my future well-being.

To my parents

parents.jpg

Introduction

0.1 The leading role played by arithmetic within mathematics and recent breakthroughs

Arithmetic enjoys a privileged position within mathematics as a fertile source of fundamental questions. Among the seven Millennium problems listed by the Clay Institute [Clay], not fewer than three: the Birch and Swinnerton-Dyer conjecture, the Hodge conjecture, and the Riemann hypothesis, were handed down by the Queen of Mathematics. Even by the standards of a subject which has remained vibrant since the days of Fermat and Gauß, the last two decades have witnessed a real golden age, with landmarks too numerous to list completely: such as the striking progress on the Birch and Swinnerton-Dyer conjecture arising from the work of Gross-Zagier [GZ1986], Kolyvagin [Kol1989], and Kato [Kat2004]; the proofs of the Shimura-Taniyama-Weil conjecture [BCDT2001], Serre's conjectures [KW2009], the Fontaine-Mazur conjecture for two-dimensional Galois representations [Kis2009], and the Sato-Tate conjectures [CHR2008] which grew out of Wiles' epoch-making proof of Fermat's Last Theorem [Wil1995], [TW1995]; the revolutionary ideas of Bourgain [Bo2008] and Gowers [Go2007] blending techniques in harmonic analysis and additive combinatorics, the Fields-medal winning breakthrough of Green and Tao on primes in arithmetic progressions [GT2008], and the work of Goldston, Pintz, and Yıldırım [GPY2009], [GPY2010], and its spectacular recent strengthenings by Zhang [Zha2014], and Maynard [May2015] and Tao [Poly2014], on bounded gaps between primes. Recent innovations in arithmetic geometry by the innovation of Perfectoid spaces [Scho2012], and subsequent topological realization of the absolute Galois group [Scho2016] by Peter Scholze also shed new lights on the Langlands Programme, a

web of conjectures that connects number theory, harmonic analysis, and geometry.

0.2 On the Riemann Hypothesis

Little essential progress has been made to the Riemann Hypothesis in the past two decades for the Riemann zeta function (or the Generalized Riemann Hypothesis for automorphic L -functions), which predicts that all the zeros of such L -functions are critical: They lie on the real line $\operatorname{Re} s = \frac{1}{2}$ with multiplicity one. The sharpest result for the Riemann zeta function is due to Feng [Fe2012], proving that at least 41.28% of the zeros of the Riemann zeta function,

$$\zeta(s) := \sum_{n \geq 1} \frac{1}{n^s} \quad (s := \sigma + it),$$

lie on the critical line, by introducing a new mollifier and applying the original method of Levinson [Lev1974] and its subsequent strengthening by Conrey [Con1989], which gave at least 34.20% and 40.88% of the zeros of $\zeta(s)$ are critical, respectively. The best zero-free region to date for the Riemann zeta function is obtained by Ford [For2002] with

$$\sigma \geq 1 - \frac{1}{57.54(\log |t|)^{2/3}(\log \log |t|)^{1/3}}, \quad |t| \geq 3.$$

A better zero-free region would imply a stronger result in the error term of the prime number theorem,

Other industries including computing the higher moments of these automorphic L -functions,

the attempt of using random matrices to explain the spacing between the zeros

Montgomery pair-correlation conjecture

the Siegel-Walfisz theorem and the large sieve developed by Bombieri are often used in place of the Generalized Riemann Hypothesis for Dirichlet L -functions to prove theorems and do estimates.

0.3 On the Hodge conjecture

Algebraic cycles

Chow groups

Kähler manifold

Known cases:

Generalizations:

Main difficulties/ best result in hoped cases.

0.4 On the Birch and Swinnerton-Dyer conjecture

Recall the Dirichlet class number formula for a number field K [Neu1999],

$$\operatorname{Res}_{s=1} \zeta_K(s) = \frac{2^{r_1} (2\pi)^{r_2}}{w |d_K|^{1/2}} h R,$$

where the left hand side is the Dedekind zeta function,

$$\zeta_K(s) := \sum_{\mathfrak{a} \subset \mathcal{O}_K} \frac{1}{\mathfrak{N}(\mathfrak{a})^s},$$

with \mathfrak{a} ideals in \mathcal{O}_K , the ring of integers of K , and \mathfrak{N} its norm. On the right hand side, r_1 and r_2 the numbers of real and complex embeddings of K , respectively. h denotes the class number of \mathcal{O}_K , which measures the failure of \mathcal{O}_K being a unique factorization domain, R the regulator, the determinant w the root number, and d_K the discriminant of the number field K

The Birch and Swinnerton-Dyer conjecture (BSD), formulated by B. Birch and Swinnerton-Dyer (1965) when studying the asymptotics of

$$\prod_{p \leq x} \frac{\#E(\mathcal{F}_p)}{p},$$

has seen tremendous progress since the first breakthrough of Coates and Wiles.

The BSD conjecture

(a) (Weak BSD)

$$\operatorname{ord}_{s=1} L(E/K, s) = r_K,$$

(b) (Strong BSD)

$$\lim_{s \rightarrow 1} \frac{L(E/K, s)}{(s-1)^{r_p}} = \Omega_{E/K} \times \operatorname{Reg}_{\infty, K}(E) \times \frac{|\operatorname{III}_K(E)| \prod_{p \leq \infty} [E(K_p) : E_0(K_p)]}{\sqrt{\Delta_K} \times |E(K)|_{\operatorname{tors}}^2},$$

can be seen as a generalization of Dirichlet class number formula, in the sense $\operatorname{III}(E)$ measures the failure of

Note Iwasawa main conjecture, proved in cases

1. \mathbb{Q} [MW1984]
2. totally real number fields [Wil1990]
3. imaginary quadratic fields [Rub1988] [Rub1991]
4. Dirichlet characters [HK2003]
5. CM elliptic curves at supersingular primes [PR2004]
6. for elliptic curves over anticyclotomic \mathbb{Z}_p -extensions [BD2005].
7. non-commutative main conjecture for totally real p -adic Lie extension of a number field [Kak2013] [CSSV2013].
8. (automorphic) GL_2 [SU2014]

One promising view
algebraic K -theory,
Suslin and Voevodsky,
Bloch
Motivic cohomology

Euler systems [Rub2000]

Classical Euler systems:

1. Siegel's cyclotomic units gives Kubota-Leopoldt p -adic L function, but no BSD application [CS2006].
2. Elliptic units Coates and Wiles' homomorphism
3. Heegner points gives anticyclotomic p -adic L -function of

Kato's Euler systems:

1. Beilinson-Kato elements
2. Beilinson-Flach elements
3. Gross-Kudla-Schoen cycles

Note that the monograph [Del2008] is devoted to the study of the BSD conjecture over the universal deformation rings of an elliptic curve.

Known cases of BSD

1. Coates and Wiles (1977) proved that if E is a curve over a number field F with complex multiplication by an imaginary quadratic field K of class number 1, $F = K$ or \mathbb{Q} , and $L(E, 1)$ is not 0 then $E(F)$ is a finite group. This was extended to the case where F is any finite abelian extension of K by Arthaud (1978).
2. Gross and Zagier (1986) showed that if a modular elliptic curve has a $L'(E, 1) = 0$ then it has a rational point of infinite order; see Gross-Zagier theorem.
3. Kolyvagin (1989) showed that a modular elliptic curve E for which $L(E, 1)$ is not zero has rank 0, and a modular elliptic curve E for which $L(E, 1)$ has a first-order zero at $s = 1$ has rank 1.
4. Rubin (1991) showed that for elliptic curves defined over an imaginary quadratic field K with complex multiplication by K , if the L -series of the elliptic curve was not zero at $s = 1$, then the p -part of the Tate-Shafarevich group had the order predicted by the Birch and Swinnerton-Dyer conjecture, for all primes $p > 7$.
5. Breuil et al. (2001), extending work of Wiles (1995), proved that all elliptic curves defined over the rational numbers are modular, which extends results 2 and 3 to all elliptic curves over the rationals, and shows that the L -functions of all elliptic curves over \mathbb{Q} are defined at $s = 1$.
6. Bhargava and Shankar (2015) proved that the average rank of the Mordell-Weil group of an elliptic curve over \mathbb{Q} is bounded above by $7/6$. Combining this with the p -parity theorem of Nekovář (2009) and Dokchitser & Dokchitser (2010) and with the proof of the main conjecture of Iwasawa theory for $GL(2)$ by Skinner & Urban (2014), they conclude that a positive proportion of elliptic curves over \mathbb{Q} have analytic rank zero, and hence, by Kolyvagin (1989), satisfy the Birch and Swinnerton-Dyer conjecture.

Chapter 1

Bloch-Kato's Tamagawa Numbers conjecture and the isogeny invariance

1.1 From the Birch and Swinnerton-Dyer conjecture to the Bloch-Kato conjecture

The Bloch-Kato's Tamagawa Numbers conjecture [BK1990], can be seen as a generalization of the Birch and Swinnerton-Dyer conjecture for motives.

Need: Fontaine's topological period rings B_{dR} and B_{crys} , where the former is a complete valued field with residue field \mathbb{C}_p

For a motivic pair (V, D) with weights $\leq w$ and a finite set of places Ω of \mathbb{Q} containing ∞ , the L -function $L_\Omega(V, s)$ is defined to be as the Euler product

$$L_\Omega(V, s) := \prod_{p \notin \Omega} P_p(V, p^{-s})^{-1},$$

it is absolute convergent for $\text{Re}(s) > \frac{w}{2} + 1$.

Fixing a \mathbb{Z} -lattice $M \subset V$

1.2 Motives, Fontaine's p -adic period rings, and other essential gadgets

To define Tamagawa measures one needs groups $A(\mathbb{Q}_p), p \leq \infty$, and $A(\mathbb{Q})$. Bloch and Kato define such groups for a motivic pair (V, D) as follows: Assuming the motivic pair (V, D) has weight ≤ -1

Fontaine's p -adic period rings [Fon1982], [FM1987],

Tamagawa number [Wei1982]

1.3 The isogeny invariance explained

1.4 K -theoretic background

Reference [HB-K1], [HB-K2], [Sri1993], [TT1990], [Wei2013], [Wal1987a] , [Wal1987b]

We will need to use higher K -theory, where K_0 was invented by Grothendieck in proving the Riemann-Roch Theorem,

- Quillen higher K -theory

- the $+$ construction

- the Q construction for schemes

- Waldhausen K -theory

- K -theory of Thomason & Trobaugh.

Chapter 2

The isogeny invariance for elliptic curves with complex multiplication

2.1 CM elliptic curves

Reference: [Hid2013], [Sil2009], [Sil1994], [Kob1993], [Kna1992], [Ca1991], [Mil2006], [CVG1999], [dSh1987]

2.2 The isogeny invariance for CM elliptic curves

2.3 The K -theoretic viewpoint

Chapter 3

The isogeny invariance for elliptic curves without complex multiplication

3.1 non-CM elliptic curves

3.2 The isogeny invariance for non-CM elliptic curves

3.3 The K -theoretic viewpoint

Chapter 4

The isogeny invariance for modular forms

4.1 Modular forms

Reference [Miy1989], [DS2005], [AF1995]

4.2 The isogeny invariance for modular forms

4.3 The K -theoretic viewpoint

Chapter 5

The isogeny invariance for abelian varieties with complex multiplication

5.1 CM abelian varieties

References: [Shi1998], [Mum1974], [BL2004]

5.2 The isogeny invariance for CM abelian varieties

5.3 The K -theoretic viewpoint

Chapter 6

Proposed isogeny invariance for tori and motif

All of the above geometric objects are in the form of a pure motive: which are smooth projective varieties.

To date many problems occur in mixed motives due to standard conjectures.

The best understanding to date is due to Voevodsky's motivic cohomology [MWV2006]

Conclusion

Categorification?

Derived version?

Bibliography

- [Clay] The Millennium Prize Problems:
<http://www.claymath.org/millennium-problems>.
The Clay Research Institute. → pages
- [HB-K1] E. M. Friedlander and D. R. Grayson edited. Handbook of K -Theory 1. → pages
- [HB-K2] E. M. Friedlander and D. R. Grayson edited. Handbook of K -Theory 2. → pages
- [AF1995] A. N. Andrianov and V. G. Zhuravlev. Modular Forms and Hecke Operators. → pages
- [Be1985] A. Beilinson. “Higher regulators and values of L -functions”. J. Soviet Math. **30** (1985), 2036–2070. → pages
- [BD2005] M. Bertolini and H. Darmon. “Iwasawa’s Main Conjecture for elliptic curves over anticyclotomic \mathbb{Z}_p -extensions. → pages
- [BL2004] C. Birkenhake and H. Lange. Complex Abelian Varieties → pages
- [BK1990] S. Bloch and K. Kato. “ L -Functions and Tamagawa Numbers of Motives”. The Grothendieck Festschrift I, Modern Birkhäuser Classics (1990), 333–400. → pages
- [Bo2008] J. Bourgain, “Roth’s theorem in progressions revisited”, *J. Anal. Math.* **104** (2008), 155–192. → pages
- [BCDT2001] C. Breuil, B. Conrad, F. Diamond, and R. Taylor, “On the Modularity of Elliptic Curves Over \mathbb{Q} : Wild 3-Adic Exercises”, *Journal of the American Mathematical Society* **14** (4) (2001): 843–939. → pages
- [Ca1991] W. Cassels, → pages

- [CHR2008] L. Clozel, M. Harris, and R. Taylor, “Automorphy for some l -adic lifts of automorphic mod l Galois representations”, *Publ. Math. Inst. Hautes Études Sci.* **108** (2008), 1–181. → pages
- [CRSS2015] J. Coates, A. Raghuram, A. Saikia, and R. Sujatha. *The Bloch-Kato Conjecture for the Riemann Zeta Function*. **Cambridge University Press** (2015). → pages
- [CSSV2013] J. Coates, P. Schneider, R. Sujatha, O. Venjakob. Noncommutative Iwasawa Main Conjectures over Totally Real Fields. → pages
- [CS2006] J. Coates and R. Sujatha. Cyclotomic Fields and Zeta Values. → pages
- [CVG1999] J. Coates, → pages
- [Con1989] B. Conrey → pages
- [De1979] P. Deligne. “Valeurs de fonctions L et périodes d’intégrales. Proc. Symp. Pure Math. **33 AMS** (1979), 313–346. → pages
- [Del2008] D. Delbourgo Elliptic Curves and Big Galois Representations. → pages
- [DS2005] F. Dimanod and Shurman → pages
- [Fe2012] Feng → pages
- [For2002] K. Ford, → pages
- [Fon1982] J. M. Fontaine → pages
- [FM1987] J. M. Fontaine and W. Messing → pages
- [GPY2009] D. A. Goldston, J. Pintz, and C. Y. Yıldırım, “Primes in tuples. I”, *Ann. of Math.* **170** (2009), 819–862. → pages
- [GPY2010] D. A. Goldston, J. Pintz, and C. Y. Yıldırım, “Primes in tuples. II”, *Acta Math.* **204** (2010), 1–47. → pages
- [Go2007] T. Gowers, “Hypergraph regularity and the multidimensional Szemerédi theorem”, *Ann. of Math.* **166** (3) (2007), 897–946. → pages
- [GZ1986] B. H. Gross and D. B. Zaiger. “Heegner points and derivatives of L -series”. *Invent. math.* **84** (1986), 225–320 → pages
- [GT2008] B. Green and T. Tao, “The primes contain arbitrarily long arithmetic progressions”, *Ann. of Math.* **167** (2) (2008), 481–547. → pages

- [Hid2013] H. Hida. Elliptic Curves and Arithmetic Invariants. → pages
- [HK2003] A. Huber and G. Kings. Bloch-Kato conjecture and main conjecture of Iwasawa theory for Dirichlet characters. → pages
- [Kak2013] M. Kakde. “The main conjecture of Iwasawa theory for totally real fields.” → pages
- [Kat1993a] K. Kato. “Lectures on the approach to Iwasawa theory for Hasse-Weil L -functions via B_{dR} , I”. Arithmetic Algebraic Geometry: Lectures given at the 2nd Session of the Centro Internazionale Matematico Estivo (C.I.M.E.) held in Trento, Italy, June 24–July 2, 1991, Springer Lecture Notes in Mathematics **1553** (1993), 50–163 → pages
- [Kat1993b] K. Kato. “Lectures on the approach to Iwasawa theory for Hasse-Weil L -functions via B_{dR} , II”. unpublished → pages
- [Kat2004] K. Kato, “ p -adic Hodge theory and values of zeta functions of modular forms”, *Astérisque* **295** (2004), ix, 117–290, Cohomologies p -adiques et applications arithmétiques. III. → pages
- [Kis2009] M. Kisin, “The Fontaine–Mazur conjecture for GL_2 ”, *Journal of the American Mathematical Society* **22** (3) (2009), 641–690. → pages
- [Kol1989] V. A. Kolyvagin, “Finiteness of $E(\mathbb{Q})$ and $\text{III}(E, \mathbb{Q})$ for a class of Weil curves”. *Math. USSR Izv.* **32** (1989), 523–541. → pages
- [Kob1993] N. Koblitz. Introduction to Elliptic Curves and Modular Forms. → pages
- [Kna1992] A. Knapp, Elliptic Curves. → pages
- [KW2009] C. Khare and J.-P. Wintenberger, “On Serre’s reciprocity conjecture for 2-dimensional mod p representations of $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ ”, *Ann. of Math.* **169** (1) (2009), 229–253. → pages
- [Lev1974] Levinson → pages
- [May2015] J. Maynard, “Small gaps between primes”, *Annals of Mathematics* **181** (2015), no. 1, 383–413. → pages
- [MW1984] B. Mazur and A. Wiles. “Class fields of abelian extensions of \mathbb{Q} ”, *Inventiones Mathematicae* **76** (2): 179–330 → pages
- [MWV2006] C. Mazza, C. Weibel, and V. Voevodsky. Lecture Notes on Motivic cohomology → pages

- [Miy1989] T. Miyake, *Modular Forms*. → pages
- [Mil2006] J. Milne, *Elliptic Curves*. → pages
- [Mum1974] D. Mumford, *Abelian Varieties* → pages
- [Neu1999] J. Neukirch. *Algebraic Number Theory* → pages
- [PR2004] R. Pollack and K. Rubin. “The main conjecture for CM elliptic curves at supersingular primes” → pages
- [Poly2014] DHJ Polymath, “Variants of the Selberg sieve, and bounded intervals containing many primes”, *Research in the Mathematical Sciences* (2014), no. 1:12. → pages
- [Rub1988] K. Rubin. “On the main conjecture of Iwasawa theory for imaginary quadratic field” → pages
- [Rub1991] K. Rubin. “The “main conjectures” of iwasawa theory for imaginary quadratic fields”, → pages
- [Rub2000] K. Rubin. *Euler Systems*, → pages
- [dSh1987] E. de Shalit, “Iwasawa Theory of Elliptic Curves with Complex Multiplication. → pages
- [Scho2012] P. Scholze, “Perfectoid spaces”, *Publications mathématiques de l’IHÉS* (2012), no. 1, 245–313. → pages
- [Scho2016] P. Scholze, “Topological realisations of absolute Galois groups ”, *Cohomology of Arithmetic Groups* (2018), no. 1, 201–288. → pages
- [Shi1998] G. Shimura, *Abelian Varieties with Complex Multiplication and Modular Functions*. → pages
- [Sil1994] J. Silverman, *Advanced Topics in the Arithmetic of Elliptic Curves*. → pages
- [Sil2009] J. Silverman, *Arithmetic of Elliptic Curves*. → pages
- [SU2014] C. Skinner and E. Urban. *The Iwasawa main conjecture for GL_2* . → pages
- [Sri1993] R. Srinivas. *Algebraic K -theory*. → pages
- [TT1990] R. Thomason and T. Trobaugh. “Higher algebraic K -theory of schemes and of derived categories”. *The Grothendieck Festschrift III*, Modern Birkhäuser Classics (1990), 247–435. → pages

- [TW1995] R. Taylor and A. Wiles. “Ring-theoretic properties of certain Hecke algebras”. **Annals of Mathematics** **141** (1995): 553–572. → pages
- [Wal1987a] F. Waldhausen. “Algebraic K -theory for generalized free products, Part 1”. **Annals of Mathematics** **108** (1978): 135–204. → pages
- [Wal1987b] F. Waldhausen. “Algebraic K -theory for generalized free products, Part 2”. **Annals of Mathematics** **108** (1978): 205–256. → pages
- [Wei1982] A. Weil. Adeles and algebraic groups. → pages
- [Wei2013] C. Weibel. The K -book. → pages
- [Wil1990] A. Wiles. “The Iwasawa conjecture for totally real fields”, *Annals of Mathematics. Second Series* **131** (3): 493–540, → pages
- [Wil1995] A. Wiles, “Modular elliptic curves and Fermat’s Last Theorem”, *Annals of Mathematics* **142** (1995), 443–551. → pages
- [Zha2014] Y. Zhang, “Bounded gaps between primes”, *Annals of Mathematics* **197** (2014), no. 3, 1121–1174. → pages