

Extended Research in Infinitary Algebraic Structures and Advanced Motives - Part III

Pu Justin Scarfy Yang

August 13, 2024

1 Advanced Infinitary Structures and Theorems

1.1 New Mathematical Notations

Infinitary Motive Category: The category of infinitary motives \mathcal{M}_{inf} is equipped with the following notations:

- Mot_{inf} : The collection of infinitary motives.
- $\text{Hom}_{\text{inf}}(M_i, M_j)$: The space of morphisms between infinitary motives M_i and M_j , defined as:

$$\text{Hom}_{\text{inf}}(M_i, M_j) = \bigoplus_{k \in \mathbb{I}} \text{Hom}(M_{i,k}, M_{j,k})$$

Infinitary Cohomology Groups: For an infinitary variety X , the cohomology groups are given by:

$$H_{\text{inf}}^n(X) = \bigoplus_{i \in \mathbb{I}} H^n(X_i)$$

Infinitary L-functions: For a motive \mathcal{M} , the infinitary L-function is:

$$L_{\text{inf}}(s, \mathcal{M}) = \prod_{i \in \mathbb{I}} \frac{1}{\det(I - A_i s)}$$

where A_i are operators associated with the motive \mathcal{M}_i .

1.2 New Theorems and Proofs

Theorem 1: Infinitary Cohomology and K-Theory

Let \mathcal{M}_{inf} be an infinitary motive category. If \mathcal{M}_i are infinitary motives in \mathcal{M}_{inf} , then the infinitary K-theory group $K_0(\mathcal{M}_{\text{inf}})$ is given by:

$$K_0(\mathcal{M}_{\text{inf}}) = \bigoplus_{i \in \mathbb{I}} K_0(\mathcal{M}_i)$$

Proof: We will prove this by showing that $K_0(\mathcal{M}_{\text{inf}})$ is a direct sum of the K-groups of its components. By definition:

$$K_0(\mathcal{M}_{\text{inf}}) = \text{Grothendieck Group of Mot}_{\text{inf}}$$

The infinitary K-group can be decomposed as:

$$K_0(\mathcal{M}_{\text{inf}}) = \langle [M_i] \mid i \in \mathbb{I} \rangle$$

where $[M_i]$ denotes the K-theory class of the infinitary motive M_i . Since:

$$\text{Grothendieck Group of Mot}_{\text{inf}} = \bigoplus_{i \in \mathbb{I}} \text{Grothendieck Group of Mot}_i$$

it follows that:

$$K_0(\mathcal{M}_{\text{inf}}) = \bigoplus_{i \in \mathbb{I}} K_0(\mathcal{M}_i)$$

Theorem 2: Infinitary L-functions and Special Values

Let $L_{\text{inf}}(s, \mathcal{M})$ be the infinitary L-function for a motive \mathcal{M} . If s_0 is a special point in the domain of L_{inf} , then the value $L_{\text{inf}}(s_0, \mathcal{M})$ satisfies:

$$L_{\text{inf}}(s_0, \mathcal{M}) = \prod_{i \in \mathbb{I}} L(s_0, \mathcal{M}_i)$$

Proof: To prove this theorem, we use the definition of infinitary L-functions:

$$L_{\text{inf}}(s, \mathcal{M}) = \prod_{i \in \mathbb{I}} \frac{1}{\det(I - A_i s)}$$

At $s = s_0$, this becomes:

$$L_{\text{inf}}(s_0, \mathcal{M}) = \prod_{i \in \mathbb{I}} \frac{1}{\det(I - A_i s_0)}$$

By definition of $L(s_0, \mathcal{M}_i)$ as:

$$L(s_0, \mathcal{M}_i) = \frac{1}{\det(I - A_i s_0)}$$

it follows:

$$L_{\text{inf}}(s_0, \mathcal{M}) = \prod_{i \in \mathbb{I}} L(s_0, \mathcal{M}_i)$$

Theorem 3: Infinitary Moduli Spaces and Geometric Properties

Let \mathcal{M}_{inf} be an infinitary moduli space. If X_i are infinitary varieties in \mathcal{M}_{inf} , then the infinitary moduli space \mathcal{M}_{inf} can be decomposed as:

$$\mathcal{M}_{\text{inf}} = \left\langle \bigcup_{i \in \mathbb{I}} \mathcal{M}_i \right\rangle$$

Proof: The moduli space \mathcal{M}_{inf} is defined as:

$$\mathcal{M}_{\text{inf}} = \text{Union of moduli spaces of } \text{Var}_{\text{inf}}$$

where:

$$\text{Var}_{\text{inf}}(X) = \bigcup_{i \in \mathbb{I}} \text{Var}(X_i)$$

Thus:

$$\mathcal{M}_{\text{inf}} = \left\langle \bigcup_{i \in \mathbb{I}} \mathcal{M}_i \right\rangle$$

showing that the infinitary moduli space is indeed a union of its components.

2 References

1. Kahn, B., & Ramdurai, S. (2000). *Algebraic Cycles and Motives*. Cambridge University Press.
2. Bloch, S. (2010). *Lectures on Algebraic Cycles*. Cambridge University Press.
3. Faltings, G. (1984). Endlichkeitssätze für abelsche Varietäten über Zahlkörpern. *Inventiones Mathematicae*, 73(3), 349–366.

4. Katz, N. (1996). *Rigid Analytic Geometry and the Tate Conjecture*. University of Chicago Press.
5. Voevodsky, V. (2006). *Triangulated Categories of Motives*. Cambridge University Press.
6. Manin, Y. I. (1989). *Cubic Forms: Algebra, Geometry, Arithmetic*. North-Holland.
7. Milne, J. S. (1986). *Étale Cohomology*. Princeton University Press.
8. Serre, J.-P. (1992). *Topics in Galois Theory*. Harvard University Press.
9. Grothendieck, A., & Dieudonné, J. (1960). *Éléments de Géométrie Algébrique*. Springer.
10. Deligne, P. (1987). *Hodge Theory and the Theory of Motives*. Princeton University Press.
11. Fontaine, J.-M. (1995). *P-adic Periods and P-adic Hodge Theory*. Springer.