

lifequal: Calculating Life-table Lifespan Equality

Jonas Schöley

2016-03-10

How to install?

You can install `lifequal` by running:

```
install.packages("devtools")
devtools::install_github("jschoeley/lifequal")
```

What does it do?

`lifequal` lets you calculate three measures of lifespan equality from a life-table:

- 1) `ExDagger(x, ex, ax)` Life expectancy lost by those who die in age interval $[x, x+w)$
- 2) `EDagger(dx, exdagger, radix)` Total life expectancy lost due to death
- 3) `KeyfzEntro(edagger, e0)` Keyfitz's entropy

They are defined as follows,

Measure	Definition
Start of age interval	x
Width of age interval starting at x	w_x
Start of last age interval	ω
Average time spent in age interval $[x, x + w_x)$ when dying in that interval	a_x
Deaths in age interval $[x, x + w_x)$	d_x
Life-expectancy at age x	e_x
Life expectancy lost due to death in age interval $[x, x + w_x)$	$e_x^\dagger = \begin{cases} \frac{a_x}{w_x} e_{x+w_x} + (1 - \frac{a_x}{w_x}) e_x & x \neq \omega \\ \frac{e_\omega + 1.4}{2} & x = \omega \end{cases}$
Total life expectancy lost due to death	$e^\dagger = \sum_{x=0}^{\omega} d_x e_x^\dagger$
Keyfitz's Entropy	$\frac{e^\dagger}{e_0}$

Life expectancy versus lifespan equality for 1x1 Swedish life-tables

```
library(lifequal)
library(dplyr)
library(ggplot2)
```

The analysis starts with a demographic life-table. We want 1) age intervals ordered from low to high, 2) no gaps between subsequent age intervals, 3) an open last age interval (e.g. 110+). Something like this:

```
# Swedish 1x1 period life-tables by period and sex
sweden1x1
```

```
## Source: local data frame [58,608 x 11]
##
##      sex period      x      mx      qx      ax      lx      dx      Lx      Tx
##      (chr) (int) (int)  (dbl)  (dbl) (dbl)  (int) (int) (int)  (int)
## 1 female  1751      0 0.21223 0.18651 0.35 100000 18651 87877 3987544
## 2 female  1751      1 0.04941 0.04822 0.50  81349  3923 79388 3899667
## 3 female  1751      2 0.03225 0.03174 0.50  77427  2457 76198 3820279
## 4 female  1751      3 0.02601 0.02567 0.50  74970  1925 74007 3744080
## 5 female  1751      4 0.02370 0.02342 0.50  73045  1711 72190 3670073
## 6 female  1751      5 0.01876 0.01859 0.50  71334  1326 70671 3597884
## 7 female  1751      6 0.01296 0.01287 0.50  70008   901 69558 3527212
## 8 female  1751      7 0.00877 0.00873 0.50  69107   603 68806 3457654
## 9 female  1751      8 0.00608 0.00606 0.50  68504   415 68296 3388849
## 10 female 1751      9 0.00494 0.00493 0.50  68089   335 67921 3320553
## ..      ...      ...      ...      ...      ...      ...      ...      ...
## Variables not shown: ex (dbl)
```

First, we use `ExDagger()` on each single life-table (separate by period and sex) to calculate the life expectancy lost in each age. We then summarise each life-table into a set of 3 numbers: Life expectancy at birth, total life years lost due to death (`EDagger()`) and lifespan equality (`KeyfzEntro()`). Note that we transform Keyfitz's Entropy by taking the negative log.

```
sweden1x1 %>%
  # ...for each single life-table...
  group_by(period, sex) %>%
  #...we calculate the life years lost in age x...
  mutate(exdagger = ExDagger(x, ex)) %>%
  # ...and then summarise each life-table into a set of 3 numbers:
  # e0:      Life-expectancy at birth
  # edagger: Total life years lost due to death
  # keyfzentro: Lifespan equality
  summarise(
    e0      = ex[x == 0],
    edagger  = EDagger(dx, exdagger, radix = 100000),
    keyfzentro = -log(KeyfzEntro(edagger, e0))
  ) -> sweden1x1summary
```

The summarised life-tables look like this:


```
sweden1x1summary
```

```
## Source: local data frame [528 x 5]
## Groups: period [?]
##
##      period  sex    e0  edagger keyfzentro
##      (int) (chr) (dbl)  (dbl)  (dbl)
## 1    1751 female 39.88 26.42272 0.4116507
## 2    1751  male 36.81 26.29076 0.3365518
## 3    1752 female 36.75 28.65521 0.2488031
```

```
## 4    1752    male 33.88 28.26323 0.1812630
## 5    1753   female 41.27 27.49097 0.4062784
## 6    1753    male 38.05 27.36753 0.3295438
## 7    1754   female 39.01 27.45547 0.3512527
## 8    1754    male 35.80 27.05649 0.2800212
## 9    1755   female 37.60 27.59827 0.3092511
## 10   1755    male 35.34 27.21785 0.2611423
## ..    ...    ...    ...    ...    ...
```

For each life-table we plot the life expectancy at birth versus the lifespan equality.

```
plot_lifequal <-
  ggplot(sweden1x1summary, aes(x = e0, y = keyfzentro, color = sex)) +
  geom_point(size = 0.6, alpha = 0.5) +
  theme(aspect.ratio = 1)
plot_lifequal
```



Life expectancy versus lifespan equality for 5x5 Swedish life-tables

The same exercise as before, only now we deal with life-tables aggregated over multiple year period and age intervals.

```
# Swedish 5x5 period life-tables by period and sex
sweden5x5
```

```
## Source: local data frame [2,496 x 11]
##
##      sex    period      x      mx      qx      ax      lx      dx      Lx
##   (chr)   (chr) (int)  (dbl)  (dbl) (dbl)  (int) (int)  (int)
```

```
## 1 female 1755-1759    0 0.23517 0.20399 0.35 100000 20399 86741
## 2 female 1755-1759    1 0.04076 0.14805 1.52 79601 11785 289151
## 3 female 1755-1759    5 0.01256 0.06043 1.88 67816 4098 326313
## 4 female 1755-1759   10 0.00658 0.03239 2.53 63718 2064 313492
## 5 female 1755-1759   15 0.00622 0.03063 2.44 61654 1888 303428
## 6 female 1755-1759   20 0.00720 0.03537 2.62 59766 2114 293794
## 7 female 1755-1759   25 0.00929 0.04544 2.57 57652 2620 281902
## 8 female 1755-1759   30 0.01230 0.05971 2.53 55032 3286 267060
## 9 female 1755-1759   35 0.01108 0.05389 2.49 51746 2788 251744
## 10 female 1755-1759  40 0.01622 0.07806 2.59 48958 3822 235561
## ..      ...      ...      ...      ...      ...      ...      ...
## Variables not shown: Tx (int), ex (dbl)
```

The width of each age interval, w_x , is calculated by the `ExDagger()` function on the basis of the differences between the age interval starting points x .

```
sweden5x5 %>%
  # ...for each single life-table...
  group_by(period, sex) %>%
  #...we calculate the life years lost in age interval [x, x+wx)...
  mutate(exdagger = ExDagger(x, ex, ax)) %>%
  # ...and then summarise each life-table into a set of 3 numbers:
  # e0:           Life-expectancy at birth
  # edagger:      Total life years lost due to death
  # keyfzentro:   Lifespan equality
  summarise(
    e0      = ex[x == 0],
    edagger = EDagger(dx, exdagger, radix = 100000),
    keyfzentro = -log(KeyfzEntro(edagger, e0))
  ) -> sweden5x5summary
```

The summarised life-tables look like this:

```
sweden5x5summary

## Source: local data frame [104 x 5]
## Groups: period [?]
##
##      period      sex    e0  edagger keyfzentro
##      (chr)   (chr) (dbl)  (dbl)      (dbl)
## 1  1755-1759 female 36.80 26.33548 0.3345809
## 2  1755-1759  male 33.89 26.03847 0.2635449
## 3  1760-1764 female 36.50 26.60345 0.3162712
## 4  1760-1764  male 33.62 26.12272 0.2523156
## 5  1765-1769 female 37.81 26.52746 0.3543933
## 6  1765-1769  male 34.78 26.16651 0.2845622
## 7  1770-1774 female 31.23 25.58650 0.1993144
## 8  1770-1774  male 28.67 24.93197 0.1397002
## 9  1775-1779 female 38.08 26.55897 0.3603218
## 10 1775-1779  male 35.95 26.48334 0.3056131
## ..      ...      ...      ...      ...      ...
```

The aggregated life-tables follow the same trend as the single year life-tables.

```
plot_lifequal +  
  geom_point(shape = 22, fill = "white",  
            data = sweden5x5summary)
```

