

Important Notice to Authors

Attached is a PDF proof of your forthcoming article in *Physical Review D*. The article accession code is LK14763D.

Please note that as part of the production process, APS converts all articles, regardless of their original source, into standardized XML that in turn is used to create the PDF and online versions of the article as well as to populate third-party systems such as Portico, CrossRef, and Web of Science. We share our authors' high expectations for the fidelity of the conversion into XML and for the accuracy and appearance of the final, formatted PDF. This process works exceptionally well for the vast majority of articles; however, please check carefully all key elements of your PDF proof, particularly any equations or tables.

Figures submitted electronically as separate PostScript files containing color appear in color in the online journal. However, all figures will appear as grayscale images in the print journal unless the color figure charges have been paid in advance, in accordance with our policy for color in print (<http://journals.aps.org/authors/color-figures-print>).

No further publication processing will occur until we receive your response to this proof.

Specific Questions and Comments to Address for This Paper

The numbered items below correspond to numbers in the margin of the proof pages pinpointing the source of the question and/or comment. The numbers will be removed from the margins prior to publication.

Q: Please check that all references include complete titles.

- 1** Please note that, for the same of internal consistency, the serial comma is used throughout.
- 2** Please note that your abbreviation "A.U." was changed to "arb. units", the standard APS abbreviation for the same term.
- 3** Please note that a verb in the sentence beginning "The invariant mass of the two" was changed to effect subject verb agreement.
- 4** Please note that the sentence formerly beginning "500,000 events" was recast so that it now begins with a word, in accordance with PRD style preferences.
- 5** Please review the confirmation funding information section of the proof's cover letter and respond as appropriate. We must receive that the funding agencies have been properly identified before the article can publish.
- 6** This query was generated by an automatic reference checking system. References [2, 16, 22] could not be located in the databases used by the system. While the references may be correct, we ask that you check them so we can provide as many links to the referenced articles as possible.
- 7** NOTE: External links, which appear as blue text in the reference section, are created for any reference where a Digital Object Identifier (DOI) can be found. Please confirm that the links created in this PDF proof, which can be checked by clicking on the blue text, direct the reader to the correct references online. If there is an error, correct the information in the reference or supply the correct DOI for the reference. If no correction can be made or the correct DOI cannot be supplied, the link will be removed.
- 8** Reference [21] was updated. Please review.
- 9** In Ref. [26], please review the proceedings details that were added and supply the publisher and the publisher city.
- 10** Please review the e-print info in Ref. [27] and supply journal publication details if they have become available.

Titles in References

The editors now encourage insertion of article titles in references to journal articles and e-prints. This format is optional, but if chosen, authors should provide titles for *all* eligible references. If article titles remain missing from eligible references, the

production team will remove the existing titles at final proof stage.

Funding Information

Information about an article's funding sources is now submitted to CrossRef to help you comply with current or future funding agency mandates. Please ensure that your acknowledgments include all sources of funding for your article following any requirements of your funding sources. CrossRef's FundRef registry (<http://www.crossref.org/fundref/>) is the definitive registry of funding agencies. Please carefully check the following funder information we have already extracted from your article and ensure its accuracy and completeness:

- U.S. Department of Energy, FundRef ID <http://dx.doi.org/10.13039/1000000015> (United States/District of Columbia)

Other Items to Check

- Please note that the original manuscript has been converted to XML prior to the creation of the PDF proof, as described above. Please carefully check all key elements of the paper, particularly the equations and tabular data.
- Title: Please check; be mindful that the title may have been changed during the peer review process.
- Author list: Please make sure all authors are presented, in the appropriate order, and that all names are spelled correctly.
- Please make sure you have inserted a byline footnote containing the email address for the corresponding author, if desired. Please note that this is not inserted automatically by this journal.
- Affiliations: Please check to be sure the institution names are spelled correctly and attributed to the appropriate author(s).
- Receipt date: Please confirm accuracy.
- Acknowledgments: Please be sure to appropriately acknowledge all funding sources.
- References: Please check to ensure that titles are given as appropriate.
- Hyphenation: Please note hyphens may have been inserted in word pairs that function as adjectives when they occur before a noun, as in “x-ray diffraction,” “4-mm-long gas cell,” and “R-matrix theory.” However, hyphens are deleted from word pairs when they are not used as adjectives before nouns, as in “emission by x rays,” “was 4 mm in length,” and “the R matrix is tested.”
Note also that Physical Review follows U.S. English guidelines in that hyphens are not used after prefixes or before suffixes: superresolution, quasiequilibrium, nanoprecipitates, resonancelike, clockwise.
- Please check that your figures are accurate and sized properly. Make sure all labeling is sufficiently legible. Figure quality in this proof is representative of the quality to be used in the online journal. To achieve manageable file size for online delivery, some compression and downsampling of figures may have occurred. Fine details may have become somewhat fuzzy, especially in color figures. The print journal uses files of higher resolution and therefore details may be sharper in print. Figures to be published in color online will appear in color on these proofs if viewed on a color monitor or printed on a color printer.
- Overall, please proofread the entire *formatted* article very carefully. The redlined PDF should be used as a guide to see changes that were made during copyediting. However, note that some changes to math and/or layout may not be indicated.

Ways to Respond

- **Web:** If you accessed this proof online, follow the instructions on the web page to submit corrections.
- **Email:** Send corrections to aps-robot@luminad.com. Include the accession code LK14763D in the subject line.
- **Fax:** Return this proof with corrections to +1.855.808.3897.

If You Need to Call Us

You may leave a voicemail message at +1.855.808.3897. Please reference the accession code and the first author of your article in your voicemail message. We will respond to you via email.

Determination of the WW polarization fractions in $pp \rightarrow W^\pm W^\pm jj$ using a deep machine learning technique

Jacob Searcy,¹ Lillian Huang,¹ Marc-André Pleier,² and Junjie Zhu¹

¹*Department of Physics, University of Michigan, Ann Arbor, Michigan 48109, USA*

²*Brookhaven National Laboratory, Upton, New York 11973-5000, USA*

(Received 6 October 2015)

The unitarization of the longitudinal vector boson scattering (VBS) cross section by the Higgs boson is a fundamental prediction of the Standard Model which has not been experimentally verified. One of the most promising ways to measure VBS uses events containing two leptonically decaying same-electric-charge W bosons produced in association with two jets. However, the angular distributions of the leptons in the W boson rest frame, which are commonly used to fit polarization fractions, are not readily available in this process due to the presence of two neutrinos in the final state. In this paper we present a method to alleviate this problem by using a deep machine learning technique to recover these angular distributions from measurable event kinematics and demonstrate how the longitudinal-longitudinal scattering fraction could be studied. We show that this method doubles the expected sensitivity when compared to previous proposals.

DOI:

Studying longitudinal vector boson scattering (VBS) processes has long been a central goal of high energy colliders [1]. Without a Higgs boson, the scattering amplitude of longitudinal vector bosons ($V_L V_L \rightarrow V_L V_L$) increases with center-of-mass energy and eventually violates unitarity [2–4]. The discovery of a Higgs-like boson at the LHC [5,6] was the first step towards understanding these interactions. However, if this Higgs boson’s couplings to vector bosons deviate from the Standard Model (SM) expectation that the scattering amplitude of VBS processes can still increase with center-of-mass energy, which makes VBS a sensitive probe of anomalous Higgs couplings [7]. In addition, many new physics scenarios predict increases in VBS cross sections, through extended Higgs sectors or other new resonances [8–11]. VBS measurements hence are both a window to new physics and a constraint on fundamental properties of the Higgs boson. Measuring VBS processes at a hadron collider is experimentally challenging due to small cross sections and the difficulty of separating longitudinal states from transverse ones.

The ATLAS and CMS collaborations recently provided the first evidence for and study of a VBS process using events with two leptonically decaying same-electric-charge W bosons in association with two forward jets ($pp \rightarrow W^\pm W^\pm jj$) [12,13]. This final state has the advantage of relatively small SM background contributions compared to other VBS processes, paired with a production rate large enough to measure in early LHC data sets. While an ideal candidate for first observation of the VBS process, measuring the longitudinal fraction in these events is not straightforward since the presence of two neutrinos in the final state prevents full kinematic reconstruction of the events.

Recent studies have shown that advances in machine learning can improve the prospects for measurements at the

Large Hadron Collider [14,15]. In this paper we explore a machine learning technique that has not previously been used in the experimental high energy physics community: regression with deep neural networks. We apply this method to the difficult problem of measuring longitudinal VBS in $W^\pm W^\pm jj$.

In general, the polarization of a gauge boson can be determined from the angular distribution of its decay products. The differential cross section of a leptonically decaying W boson is related to the polarization fractions as [16]

$$\frac{d\sigma}{d\cos\theta^*} \propto \frac{3}{8}f_-(1 \mp \cos\theta^*)^2 + \frac{3}{8}f_+(1 \pm \cos\theta^*)^2 + \frac{3}{4}f_L(1 - \cos^2\theta^*), \quad \text{for } W^\pm, \quad (1)$$

where θ^* is the angle between the charged lepton in the boson rest frame and the W boson direction of motion. Fraction parameters f_- , f_+ , and f_L denote the fractions of events with the three possible polarization states of the W boson, -1 , $+1$, and 0 , respectively. They are constrained via $f_- + f_+ + f_L = 1$. In order to measure θ^* , we need to fully reconstruct the direction of motion of the W boson.

Requiring both W bosons to decay leptonically in $pp \rightarrow W^\pm W^\pm jj$ events enables the determination of the electric charge of each W boson via the charged leptons. However, since the corresponding two neutrinos in the final state are not detected, the W boson rest frames cannot be directly measured. It is thus difficult to determine polarization fractions of each boson and the fraction of longitudinal scattering events in the $W^\pm W^\pm jj$ process.

Many proposals have been made to determine the longitudinal fractions in other VBS final states, such as semileptonic W^+W^- [17] and $W^\pm Z$ [18] or fully leptonic decay modes of $W^\pm Z$ and ZZ , where the full event kinematics can be reconstructed or estimated using the W boson mass constraint. However, these channels suffer from large SM backgrounds that are not present in the $W^\pm W^\pm jj$ channel. Attempts have been made to gain sensitivity through other variables [1,18–21] than θ^* in the $W^\pm W^\pm jj$ channel. One example is the variable $R_{pT} = (p_T^{\ell_1} \times p_T^{\ell_2}) / (p_T^{j_1} \times p_T^{j_2})$ [20], where ℓ_1 and ℓ_2 denote the two leptons in no particular order and j_1 and j_2 denote the two most energetic jets in the event. This single variable does not encompass all of the sensitivity to longitudinal scattering, and better discrimination can be achieved by combining the available event information with a machine learning technique. Therefore, we develop a method to use a neural network (NN) to map measurable quantities to the true $\cos\theta^*$ values that contain the event’s polarization information. This approximation mitigates the limitation of missing kinematic information in this final state, and makes $W^\pm W^\pm jj$ a promising channel for observing the behavior of longitudinal W boson scattering.

While it has become a common practice in high energy physics to use multivariate techniques to separate signal from background, multivariate regression is not commonly used to measure underlying physics quantities of interest. Unlike classification, where the goal of the neural network is to produce discrete assignments, for example signal and background, NN regression relies on the fact that a neural network is a universal approximator [22] to instead approximate an unknown continuous function. The goal of our NN is to find the best approximation of the two truth values of $\cos\theta^*$ (one for each W boson) present in each event, using measurable quantities. Similar techniques are currently in use to address the problem of estimating parton distribution functions [23].

It has also become common practice to train these multivariate methods using variables built from basic measurable event quantities to be sensitive to a given physics process [24]. These high-level variables add some understanding of the underlying physics to the training variables and can produce better results. However, it has also recently been observed that extending a single layer neural network to a “deep” network containing many layers can regain most of the sensitivity produced by these high-level variables [14,15]. Since only a few high-level variables have been proposed for this process we choose to use deep neural networks for maximum sensitivity.

$W^\pm W^\pm jj$ events have the signature of two quarks, two same sign leptons, and two undetected neutrinos. We use all basic measurable object kinematics as input variables to the neural network: the transverse momentum (p_T), pseudorapidity (η), and azimuthal angle (ϕ) of the two leptons and two jets, and x and y components of the missing transverse

energy (E_T^x and E_T^y). The overall number of measurable quantities used hence is 14. The goal of the multivariate technique is to find the best mapping from these measurable quantities to the two truth values of $\cos\theta^*$ (one for each W boson) present in each event.

Training deep NNs has been the subject of intensive study and a good review of some of this work is presented in [25]. For completeness we list some of the properties of the NN that we utilize. We choose a multilayer neural network with a two node final output layer with linear activation to approximate the true $\cos\theta^*$ distribution of each W boson. The NN is implemented with the THEANO software packages [26,27]. The cost function is defined as

$$\mathcal{C} = \frac{1}{N} \sum_{i=1}^N [(\cos\theta_{1,i}^* - \cos\theta_{1,i}^{NN})^2 + (\cos\theta_{2,i}^* - \cos\theta_{2,i}^{NN})^2], \quad (2)$$


where N is the number of events per minibatch, $\cos\theta_{1/2,i}^*$ is the truth value of $\cos\theta^*$ for each W boson with random ordering for the i th event, and $\cos\theta_{1/2,i}^{NN}$ is the value of the two neural network outputs. A stochastic gradient descent algorithm is used to train the neural network by minimizing the cost function. Hyperparameters are tuned by hand and confirmed by a local grid search, with the best performance given by a deep network with 20 layers each with 200 hidden nodes. This network yields a 20% better cost value in a validation data set than the best single layer NN tested.

Signal $W^\pm W^\pm jj$ events are generated using the MADGRAPH event generator [28] at a proton-proton center-of-mass energy of 13 TeV. The invariant mass of the two outgoing partons was required to be greater than 150 GeV. The W bosons are decayed, assuming they are on shell and have no spin correlations, with the DECAY routine provided with MADGRAPH. We generated 500,000 events for training and another million for testing and validation.

Polarization fractions can then be obtained experimentally by fitting the two-dimensional distribution of the NN output $\cos\theta^{NN}$. In order to fit for these polarization fractions templates must be built for “pure” polarization states. These templates are created using generator level helicity information. In addition, a method was tested that used truth level reweighting and included all spin correlations and off-shell effects, and the results were found to be comparable.

Since there are two W bosons in each event, six distinct polarization states are possible. Events where both W bosons have a polarization of 0 are referred to as LL , of 1 as $++$, and of -1 as $--$. Events with differing polarizations of $(-1, 1)$ or $(1, -1)$ are referred to as $+-$, $(-1, 0)$ or $(0, -1)$ as $L-$, and likewise $(1, 0)$ or $(0, 1)$ as $L+$.

Figure 1(a) shows the comparison between the truth $\cos\theta^*$ and the NN output $\cos\theta^{NN}$ for $--$, $++$, and LL events ($+-$ / $L+$ / $L-$ are omitted for clarity, but closely




F1:1 FIG. 1. A comparison of the truth level $\cos \theta^*$ and the NN output $\cos \theta^{NN}$ distributions for --, ++, and LL events (a), R_{pT} templates
 F1:2 for the corresponding polarization states with log scaling (b), and a comparison of the $\cos \theta^{NN}$ for the signal and dominant $WZjj$
 F1:3 background (c) are shown with arbitrary units (arb. units). (a),(c) Projections of the 2D distributions onto one of the two leptons, which
 F1:4 are identical to the projections onto the other lepton due to arbitrary sorting.

resemble combinations of the templates shown). As expected, $\cos \theta^{NN}$ has less separation power for the different polarization states compared to $\cos \theta^*$ due to missing information for the two final state neutrinos. However, reasonable discrimination between each polarization state can clearly be seen from these distributions. Figure 1(b) shows the R_{pT} distribution for --, ++, and LL events. The discrimination power is seen only for large values of R_{pT} , and is only apparent with a logarithmic scale. Figure 1(c) shows the $\cos \theta^{NN}$ distribution for the signal $W^\pm W^\pm jj$ process and an important background process: $WZjj$ production. Reasonable separation power is observed, which could be utilized in a combined fit. In an actual data analysis, the $WZjj$ component would be subtracted as background from the observed data before fitting the polarization fractions.


Having established templates for each polarization state and a distribution that is sensitive to different polarization states, we fit the two-dimensional $\cos \theta_1^{NN}$ versus $\cos \theta_2^{NN}$ distribution in pseudo data to derive each polarization fraction. Five equal-size bins are used for each $\cos \theta^{NN}$ variable ranging from -1 to 1. A maximum likelihood fit is performed within the RooFit framework [29]. We combine events with both W bosons transversely polarized as “TT”

(the sum of the --, ++, and ++ combinations), events with one W boson transversely polarized and one W boson longitudinally polarized as “TL” (the sum of the $L-$ and $L+$ combinations), and events with both W bosons longitudinally polarized as “LL.” This reduces the free fitting parameters from five to two and allows for a better constraint on the LL scattering fraction of interest, under the assumption that the relative admixture of contributions within TT and TL does not change. The two-dimensional distributions for these templates are shown in Fig. 2. Statistical fit uncertainties are determined by randomly fluctuating data expectations within their Poisson uncertainties and repeating the fit, and confidence intervals are derived from these toy experiments.

Fits are performed in a range of integrated luminosities from 0.01–3 ab^{-1} . An example fit for 1 ab^{-1} is shown in Fig. 3 where the pseudo data are compared to the sum of the contributions from the TT , TL , and LL components. It is found that the LL fraction can be measured with a 68% confidence limit of $(6.7 \pm 1.4)\%$ with an ultimate luminosity of 3 ab^{-1} . When a similar fit is applied to the R_{pT} variable the precision of $(6.7^{+7.1}_{-6.7})\%$ is found to be consistent with 0. The regression technique hence greatly enhances the sensitivity to this process.



F2:1 FIG. 2. The two-dimensional distribution of $\cos \theta_1^{NN}$ versus $\cos \theta_2^{NN}$ for the two leptons in each event is shown for each of the three
 F2:2 polarization states: TT, TL, and LL, respectively.




F3:1 FIG. 3. One example fit where the pseudo data are compared to
 F3:2 the sum of contributions from the TT , TL , and LL components.
 F3:3 There are five groups with five bins inside each group. These five
 F3:4 groups represent $\cos \theta_2^{NN}$ from -1 to 1 with a step size of 0.4 ,
 F3:5 while five bins inside each group represent $\cos \theta_1^{NN}$ from -1 to 1
 F3:6 with a step size of 0.4 .

231 The above measurements are performed with parton
 232 level predictions. While they show encouraging results it is
 233 important to also check to see if this procedure will stand up
 234 to the experimental reality of finite detector resolution, and
 235 the event level selection that will be required to remove
 236 backgrounds from this analysis. To study the effects of
 237 event level cuts, we apply additional selection criteria as
 238 used by the ATLAS Collaboration [12] to obtain a tighter
 239 fiducial region which is dominated by the contribution from

240 electroweak production of $W^\pm W^\pm jj$ events: jet
 241 $p_T > 30$ GeV, lepton $p_T > 25$ GeV, missing transverse
 242 energy $E_T > 40$ GeV, dijet mass $M_{jj} > 500$ GeV, and
 243 dijet pseudorapidity difference $|\Delta\eta_{jj}| > 2.4$.

244 To emulate the response of a typical general-purpose
 245 LHC detector, these events are passed through the appli-
 246 cation of parton showering in PYTHIA [30] and then through
 247 the response simulation of the CMS detector implemented
 248 in DELPHES [31]. This detector smearing adds some degree
 249 of realism, but could neglect various effects due to the large
 250 number of overlapping interactions during the high lumi-
 251 nosity LHC runs. Since these effects are often mitigated
 252 with specific reconstruction techniques and need detailed
 253 detector modeling, we leave studies of this nature to
 254 dedicated efforts by the experiments. After detector sim-
 255 ulation, parton level quantities are then approximated by
 256 taking the leading two jets clustered with an anti- k_t
 257 algorithm [32] with jet size parameter $R = 0.5$.

258 Backgrounds to the $W^\pm W^\pm jj$ process depend largely on
 259 experimental choice, and require detailed simulation. It can
 260 be seen in Fig. 1 that background from the $WZjj$ process
 261 (where one of the leptons from the Z boson decay is not
 262 detected) has a different $\cos \theta^{NN}$ shape than the $W^\pm W^\pm jj$
 263 events, but it is likely event level cuts will still need to be
 264 applied to reduce this component. Determining the system-
 265 atics on the background shapes will require significant work
 266 from the experiments, and is not treated here. However, as
 267 already mentioned the backgrounds in the $W^\pm W^\pm jj$ channel
 268 are relatively small compared to other channels.



F4:1 FIG. 4. 68% (yellow) and 95% (green) expected confidence intervals for the measured LL fraction f_{LL} (top panels), the TL fraction
 F4:2 f_{TL} (middle panels), and the TT fraction f_{TT} (bottom panels) as a function of the integrated luminosity for, from left to right, scenarios
 F4:3 (a), (b), and (c) discussed in the text.

We determine the precision that can be achieved for fractions of TT , TL , and LL components using three different scenarios increasing in realism: (a) using all generated events at the parton level; (b) using events passing the additional selection criteria used by ATLAS at the parton level; (c) using events processed with the DELPHES detector simulation and with reconstructed objects passing additional selection criteria used by ATLAS.

The precision for the three polarization fractions as a function of the integrated luminosity is presented in Fig. 4. TT components can be measured with great precision, whereas separating pure LL scattering from LT scattering is challenging. In the most difficult and realistic scenario, (c), the cuts and object efficiencies slightly increase the mean LL fraction to 7.0% and 68% confidence limits are found $7_{-7}^{+19}\%$ ($7_{-6}^{+5}\%$) for an integrated luminosity of 0.1 (3) ab^{-1} . Equivalent limits from fits to the R_{pT} variable are found to be $7_{-7}^{+29}\%$ and $7_{-7}^{+9}\%$. The fit to the neural network almost doubles the ultimate precision to anomalous LL fractions in this scenario. Reaching the same statistical sensitivity using the R_{pT} variable would require approximately 10 ab^{-1} , more than three times the total expected luminosity of the LHC program. We have shown that large sensitivity gains can be made with NN regression. In scenario (c) our simple estimate falls short of the 5σ criteria for observation of longitudinal VBS, however, new physics beyond the SM could greatly enhance this fraction [8], making stringent limits or a future observation of this fraction very important. A comparison of scenarios

(a) and (b) to (c) illustrates potential gains in sensitivity through cut optimization and improved detector performance. In addition, the authors hope that experiments can improve on this methodology by training on fully simulated events or by improving detector performance (e.g. through upgrades) to enhance sensitivity.

In conclusion, we present a method to determine the WW polarization fractions in $W^{\pm}W^{\pm}jj$ events by using a deep machine learning technique. This method allows us to recover the charged lepton angular distributions in the W boson rest frame from measurable event kinematics. The results obtained from this method show greatly enhanced sensitivity over the example R_{pT} variable. Cuts to reject backgrounds as well as finite detector resolutions reduce the sensitivity as expected, but the method remains a powerful tool for the study of polarization fractions in VBS events, almost doubling the ultimate precision when compared to R_{pT} .

We would like to thank our colleagues Sally Dawson, Frank Paige, and Olivier Mattelaer for their help and guidance while preparing this manuscript. The contributions from J. S., L. H., and J. Z. were supported by the U.S. DOE Early Career Grant under Contract No. DE-SC0008062. The work of M.-A. P. is supported by U.S. DOE Contract No. DE-SC0012704. We would also like to acknowledge the role of the U.S. ATLAS Scholar program for fostering a collaboration between Michigan and the Brookhaven National Laboratory.

326

-
- [1] D. A. Dicus, J. F. Gunion, and R. Vega, *Phys. Lett. B* **258**, 475 (1991).
 [2] M. J. G. Veltman, *Acta Phys. Pol. B* **8**, 475 (1977).
 [3] B. W. Lee, C. Quigg, and H. B. Thacker, *Phys. Rev. Lett.* **38**, 883 (1977).
 [4] B. W. Lee, C. Quigg, and H. B. Thacker, *Phys. Rev. D* **16**, 1519 (1977).
 [5] G. Aad *et al.* (ATLAS Collaboration), *Phys. Lett. B* **716**, 1 (2012).
 [6] S. Chatrchyan *et al.* (CMS Collaboration), *Phys. Lett. B* **716**, 30 (2012).
 [7] J. M. Campbell and R. K. Ellis, *J. High Energy Phys.* **04** (2015) 030.
 [8] A. Alboteanu, W. Kilian, and J. Reuter, *J. High Energy Phys.* **11** (2008) 010.
 [9] W. Kilian, T. Ohl, J. Reuter, and M. Sekulla, *Phys. Rev. D* **91**, 096007 (2015).
 [10] S. Godfrey and K. Moats, *Phys. Rev. D* **81**, 075026 (2010).
 [11] C.-W. Chiang, S. Kanemura, and K. Yagyu, *Phys. Rev. D* **90**, 115025 (2014).
 [12] G. Aad *et al.* (ATLAS Collaboration), *Phys. Rev. Lett.* **113**, 141803 (2014).
 [13] V. Khachatryan *et al.* (CMS Collaboration), *Phys. Rev. Lett.* **114**, 051801 (2015).
 [14] P. Baldi, P. Sadowski, and D. Whiteson, *Nat. Commun.* **5**, 4308 (2014).
 [15] P. Baldi, P. Sadowski, and D. Whiteson, *Phys. Rev. Lett.* **114**, 111801 (2015).
 [16] R. K. Ellis, W. J. Stirling, and B. R. Webber, *Cambridge Monogr. Part. Phys., Nucl. Phys., Cosmol.* **8**, 1 (1996).
 [17] T. Han, D. Krohn, L.-T. Wang, and W. Zhu, *J. High Energy Phys.* **03** (2010) 082.
 [18] A. Ballestrero, D. B. Franzosi, and E. Maina, *J. High Energy Phys.* **06** (2011) 013.
 [19] A. Freitas and J. S. Gainer, *Phys. Rev. D* **88**, 017302 (2013).
 [20] K. Doroba, J. Kalinowski, J. Kuczmarski, S. Pokorski, J. Rosiek, M. Szleper, and S. Tkaczyk, *Phys. Rev. D* **86**, 036011 (2012).
 [21] M. Fabbrihesi, M. Pinamonti, A. Tonero, and A. Urbano, Vector boson scattering at the LHC: A study of the

- 369 $WW \rightarrow WW$ channels with the Warsaw cut, [Phys. Rev. D](#)
370 **8** [93](#), 015004 (2016).
371 [22] G. Cybenko, *Math Control Signals Syst.* **2**, 303 (1989).
372 [23] S. Forte, L. Garrido, J. I. Latorre, and A. Piccione, [J. High](#)
373 [Energy Phys.](#) **05** (2002) 062.
374 [24] P. C. Bhat, *Annu. Rev. Nucl. Part. Sci.* **61**, 281 (2011).
375 [25] J. Schmidhuber, *Neural Netw.* **61**, 85 (2015).
376 [26] J. Bergstra, O. Breuleux, F. Bastien, P. Lamblin, R. Pascanu,
377 G. Desjardins, J. Turian, D. Warde-Farley, and Y. Bengio, in
378 *Proceedings of the 9th Python in Science Conference, Austin,*
379 **9** *TX, 2010*, edited by S. van der Walt and J. Millman (2010), p. 3.
380 [27] F. Bastien, P. Lamblin, R. Pascanu, J. Bergstra, I. J.
381 Goodfellow, A. Bergeron, N. Bouchard, D. Warde-Farley,
382 **10** and Y. Bengio, [arXiv:1211.5590](#).
- [28] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni,
383 O. Mattelaer, H.-S. Shao, T. Stelzer, P. Torrielli, and M.
384 Zaro, [J. High Energy Phys.](#) **07** (2014) 079.
385 [29] W. Verkerke and D. P. Kirkby, *Proceedings of the 2003*
386 *Computing in High Energy and Nuclear Physics (CHEP03),*
387 *La Jolla, CA, 2003*, eConf C0303241, MOLT007
388 (2003).
389 [30] T. Sjöstrand, S. Mrenna, and P. Z. Skands, *Comput. Phys.*
390 *Commun.* **178**, 852 (2008).
391 [31] J. de Favereau, C. Delaere, P. Demin, A. Giammanco, V.
392 Lemaître, A. Mertens, and M. Selvaggi (DELPHES3
393 Collaboration), [J. High Energy Phys.](#) **02** (2014) 057.
394 [32] M. Cacciari, G. P. Salam, and G. Soyez, [J. High Energy](#)
395 [Phys.](#) **04** (2008) 063.
396
397