ESOFORM PDFLEX MACROS USER MANUAL FOR PHASE 1 PROPOSALS

European Southern Observatory

CONTENTS

- 1. Introduction
- 2. New features for Period 96
- 3. How to fill a Normal Programme template
- 4. How to fill a Large Programme template
- 5. Submission of the application

ESO DEADLINE FOR PROPOSAL SUBMISSION

The ESO deadline for Period 96 proposal submission is:

26 March 2015, 12:00 CEST

This User Manual and the whole ESOFORM Package is maintained by the Observing Programmes Office (OPO), while the background software is provided by the User Support System (USS) Department.

1 INTRODUCTION

The ESOFORM package has been designed to enable a fully electronic preparation and submission of applications for observing time on all telescopes located at the European Southern Observatory's La Silla Paranal observatory.

Getting help. Should you need assistance from ESO to prepare your proposal, please send emails to the address esoform@eso.org for questions related to the ESOFORM package as well as for more general questions about Observatory policy, etc... You should receive automatic acknowledgment from the system within 4 hours. If you do not receive any acknowledgment of your query, please email opo@eso.org directly.

For instrument specific questions, please email usd-help@eso.org.

1.1 How to Obtain the New ESOFORM Proposal Package

The ESOFORM Proposal Package may be obtained by logging into the ESO User Portal at the following address:

http://www.eso.org/UserPortal.

1.2 Description of the Content of the ESOFORM Proposal Package

The ESOFORM package consists of:

- two LATEX class files (esoform.cls and esoformlarge.cls) that, together with the style files (common2e.sty and config.sty), define all the macros required to generate the application forms for Normal and Large Programmes, respectively;
- two template proposals (template.tex for Normal Programme applications, and templatelarge.tex for Large Programme applications), which the users may edit directly in order to create a new proposal;
- this Users' Manual (usersmanual.tex), which contains all the information required to fill the templates, as well as instructions on the electronic submission of proposals (via the Web-based WASP interface in the ESO User Portal);
- a short README file.

1.3 General Features

The present manual describes the use of the ESOFORM templates, which are composed of macros that are defined in the ESOFORM class and style files. The macros allow the computer controlled typesetting of applications for observing time at ESO. If you are already familiar with TEX or LATEX, you will probably have no difficulty using the macros provided. You should follow the instructions given below and keep in mind that all your input must conform to the standard LATEX rules.

The ESOFORM package has been prepared with the following version of pdfIATEX: pdfTEX, Version 3.141592 (Web2C 7.5.5). If you encounter any serious pdfTEX or pdfIATEX problem, please send an email to the address esoform@eso.org, describing the problem and indicating which version of pdfIATEX you are using. For ease of use, we have adopted (and already included in the class files esoform.cls, and esoformlarge.cls) a number of IATEX definitions of commonly used astronomical symbols (see list in Table 1).

For every observing period, the layout of the instruments will be updated according to the anticipated availability of instruments at the La Silla Paranal Observatory. The ESO Call for Proposals for the corresponding period has all the latest information on instrument availability and any significant changes in performance. Please note that **only** proposals prepared using the **latest** version of ESOFORM will be valid and accepted by ESO.

• ESO proposal submission deadline: Please note that the ESO deadline will be strictly enforced: users should plan accordingly. It is the PI's responsibility to resolve any problems related to LaTeX, figure uploads or configuration problems and verify the LaTeX proposal form well before the deadline as ESO cannot provide support beyond 11:00 CET on the day of the deadline. The online receiver will switch off at 12:00 CET and no submissions or amendments to submitted proposals can be accepted after this time.

Table 1: Astronomical LaTeX Symbols

\ang	Å	90\deg	90°	
\halpha	$_{ m H}\alpha$	16\sqdeg	$16 \mathrm{deg}^2$	
\hbeta	$_{\mathrm{H}\beta}$	28\arcmin	28'	
\hgamma	${ m H}\gamma$	11\arcsec	11''	
\lya	$Ly\alpha$	5\fd4	$5\overset{\mathrm{d}}{\cdot}4$	
\lyb	$\text{Ly}\beta$	8\fh2	$8^{\rm h}_{\cdot}2$	
\mv	$m_{_{V}}$	2\fm56	$2^{\rm m}_{\cdot}56$	
\Mv	$M_{_{V}}$	10\fs08	$10^{\rm s}\!.08$	
\ubvr	UBVR	23\fdg12	$23^{\circ}_{\cdot}12$	
\ub	$U\!-\!B$	3\farcm6	3.6	
\bv	B - V	0\farcs27	0′′27	
\vr	$V\!-\!R$	0\fp4	0.4	
\ur	$U\!-\!R$	\onehalf	$^{1}/_{2}$	
\jhk	JHK	\onethird	$^{1}/_{3}$	
\jh	$J\!-\!H$	\twothirds	$^{2}/_{3}$	
\hk	H - K	\onequarter	$^{1}/_{4}$	
\jk	$J\!-\!K$	\threequarters	$\frac{3}{4}$	
$\inf\{C\}{4}$	C IV	$\slantfrac{{22}}{7}$	$^{22}/_{7}$	(braces unless one character)
3.6\micron	$3.6 \mu \mathrm{m}$	\$\squig\$	\sim	(math mode only)
$25\kms$	$25\mathrm{km\ s^{-1}}$	<pre>\$\lesssim\$</pre>	\lesssim	(math mode only)
\peryr	$ m yr^{-1}$	$\scriptstyle \$	\gtrsim	(math mode only)
M\subsun	${ m M}_{\odot}$	\$\la\$	\lesssim	(math mode only)
\sun	\odot	\$\ga\$	\gtrsim	(math mode only)
\earth	\oplus	\nodata	• • • •	(tables only)
\sq				

2 IMPORTANT REMINDERS AND NEW FEATURES FOR PERIOD 96

- Normal and Large Programme forms: The Large Programme template (templatelarge.tex) must be used to prepare Large Programme proposals. The total amount of time requested in a programme determines which proposal template should be used. Any programme requesting a total amount of time of 100 hours or more must use the Large Programme template. The Normal Programme template form must be used for all other programmes, including the Normal, GTO, ToO, Monitoring and Calibration proposal types.
- Large Programme Data Products: A key requirement of Large Programmes is the delivery of data products into the ESO Archive. Starting from Period 95, PIs of Large Programmes are required to provide all data products within a reasonable timeframe. Further details are available in the Large Programmes Section of the Call for Proposals document. A detailed delivery plan along with the relevant experience of proposing teams should be outlined in Sections 6 and 7 of the Large Programme proposal template (also see Section 4).
- Monitoring programme proposals: Proposers can apply for Monitoring Programmes to monitor targets over several periods. Monitoring Programme proposals should be prepared using the normal template form template.tex. The Monitoring Programme type should be specified in this form as follows: \ProgrammeType{MONITORING}, also see Table 4. More details on the definition of Monitoring Programmes and instruments available for this type of programme are given in Section 4.2 of the Call for Proposals.
- Definition of Service Mode and Visitor Mode runs: An observing programme, as described in a single proposal, may consist of one or more runs. Multiple runs should only be requested for observations

with different instruments, for different observing modes (e.g., service mode, visitor mode or pre-imaging runs). Proposers should split Visitor Mode observations at different epochs (e.g., due to different target RAs) into separate runs. Conversely, Service Mode runs should not be split according to time-critical windows, or used to group targets according to their Right Ascensions.

- Observing conditions: The definitions of the observing conditions for Phase 1 and Phase 2 can be found on the Observing Conditions webpage. Please note that the seeing constraint to be specified should be the maximum acceptable seeing value (FWHM in arcseconds) in the V-band at zenith. Please see the Call for Proposals and the Phase 2 Observing Conditions page for more details.
- Scheduling requirements in visitor mode: Any time constraints related to scheduling visitor mode runs should be flagged using the \HasTimingConstraints macro. Any time constraints specified using the \TimeCritical macro will be subject to a strict verification at proposal submission. Please take particular care to ensure that the start and end dates of suitable observation windows comply with the applicable time specification convention, and that the amount of requested time (in the \ObservingRun macro) fits within the length of the slots indicated in the \TimeCritical macro.

 Several time intervals might be acceptable for each run (e.g., transits of extrasolar planets across their host star); if this is the case the \TimeCritical macro should be repeated as many times as there are adequate alternative observation dates. Further details and examples of the correct usage are shown in Section 3.19.
- Precipitable water vapour (PWV): Users of APEX instruments in Service Mode must specify an upper limit for PWV as an observing constraint during their Phase 1 and Phase 2 preparation. Examples of the correct usage are shown in the ESOFORM package template files.
- VLTI-ATs: The baselines offered in Period 96 are listed on the VLTI baseline page.

3 HOW TO FILL A NORMAL PROGRAMME TEMPLATE

As mentioned in the Introduction, you should fill in the template file (template.tex) with your favourite editor. Instructions for Large Programmes (templatelarge.tex) are given in Sect. 4. The easiest way to write a proposal is to modify the template file template.tex by following the examples therein and the detailed instructions given in the present manual. Input in the template is allowed only within the arguments of the provided ESOFORM macros. The presence of text outside the macro arguments will lead to rejection of the proposal by the automatic proposal reception system (see Sect. 5).

Please note that it is the responsibility of the applicants to stay within the current box limits and to eliminate potential overfill/overwrite problems. A careful visual check of the generated pdf file is mandatory.

3.1 The Cycle, the Title, the Subcategory Code, the GTO, ToO, RRM, and XMM Flags: BOX 1

The first macros to check in the template.tex files are as follows:

- \Cycle contains the Period ID for this Call for Proposals, and should NOT be modified by the users.
- \Title must contain the title of the application (up to two lines).
- \SubCategoryCode must contain only one subcategory code, corresponding to the keyword (see Table 2) best summarizing the aim of your proposal. For example, a study of high-redshift clusters of galaxies will have the code A5. Please note: CALIBRATION proposals MUST have a SubCategoryCode of L0.
- \ProgrammeType should be NORMAL for Normal Programmes, CALIBRATION for Calibration, GTO for Guaranteed Time Observation (GTO), MONITORING for Monitoring Programmes, or TOO for Target of Opportunity (ToO) programmes. LARGE types are also defined for Large Programmes, but they can only be used within the corresponding specific template, templatelarge.tex (see Sect. 4).

 A CALIBRATION proposal MUST have a SubCategoryCode of LO.
 - Note that a GTO proposal should ask only for GTO time; it is compulsory to fill another, non-GTO proposal if you need more non-GTO time, even if it is for exactly the same project. If you submit a

TOO proposal, your proposal must include a completed ToO page (see Sect. 3.22). If you submit a GTO proposal, you must additionally uncomment and fill in the macro \GTOcontract, which is described below.

- \GTOcontract must be uncommented if the proposal type is GTO. This macro takes one mandatory argument: the keyword corresponding to the contract or agreement governing the allocation of Guaranteed Time under which the present proposal is submitted. The applicable keyword for each contract or agreement has been communicated by ESO to the coordinator of the respective GTO Team or to the designated contact person. Every single GTO proposal is allowed to request time within the framework of only one GTO contract or agreement. If your project involves Guaranteed Time corresponding to several different agreements, you must submit separate proposals for the time to be charged to each agreement.
- \ObservationInRRM must be uncommented if your proposal is a Target of Opportunity proposal applying for Rapid Response Mode (RRM) observations.
- \ObservationWithXMM should be left commented, as for Period 96 it is not possible to apply for time under the VLT-XMM agreement.

Your first sequence will then have the following general format:

```
\Cycle{96A}
\Title{AGN host galaxies}
\SubCategoryCode{B9}
\ProgrammeType{GT0}
\GTOcontract{<your-contract-keyword>)
```

which means that you would like to study some AGN host galaxies, with subcategory code B9, and this would be Guaranteed Time Observations within the framework of the contract or agreement referred to by keyword <your-contract-keyword>.

3.2 The Abstract/Total Time Requested: BOX 2

The ESOFORM includes a string called the "Total Time Requested" in Box 2. This field remains blank when compiling locally; this is normal and proposers should not be concerned about it. The total time requested is only computed in the version of the proposal that is reviewed by the OPC.

This macro (\Abstract) contains the abstract of the proposal, i.e., a brief summary, in up to nine lines, of your scientific aim.

```
\Abstract{ .
    .
    .
    .
    The text of your summary which will usually be several lines long. Line breaking will automatically be taken care of by LaTeX.
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
   .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
    .
```

3.3 Information about the Different Runs: BOX 3

The next macro (\ObservingRun) allows the description of the different parameters characterising your observing run(s) and is necessary for the scheduling and completion of your programme (see examples below). This macro takes ten arguments. These must be specified between the ten pairs of curly braces {}, which are related to the parameters described below.

1. RUN ID. Your programme may involve several observing runs, e.g. for complementary use of different telescopes or different instruments. Each observing run (up to 26) required by a proposal should be identified by a different letter, following the sequence A, B, C, ..., Z as needed. Provide, in the first pair of curly braces, the run identification(s). For example,

Table 2: ESO OPC categories and subcategories

A Cosmology A1 Surveys of AGNs and high-z galaxies; A2 Identification studies of extragalactic surveys; A3 Large scale structure and evolution; A4 Distance scale; A5 Groups and clusters of galaxies; A6 Gravitational lensing; A7 Intervening absorption line systems; A8 High-redshift galaxies (star formation and ISM). B1 Morphology and galactic structure; B2 Unresolved and resolved stellar populations; B4 Galaxy dynamics; B5 Peculiar/interacting galaxies; B6 Non-thermal processes in galactic nuclei (incl. QSRs, QSOs, blazars, Seyfert galaxies, BALs, radio galaxies, and LINERS); B7 Thermal processes in galactic nuclei and starburst galaxies (incl. ultraluminous IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; C2 Chemical processes in the interstellar medium; C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution; C7 Young binaries, brown dwarfs, exosolar planet searce	
A2 Identification studies of extragalactic surveys; A3 Large scale structure and evolution; A4 Distance scale; A5 Groups and clusters of galaxies; A6 Gravitational lensing; A7 Intervening absorption line systems; A8 High-redshift galaxies (star formation and ISM). B1 Morphology and galactic structure; and B2 Unresolved and resolved stellar populations; B4 Galaxy dynamics; B5 Peculiar/interacting galaxies; B6 Non-thermal processes in galactic nuclei (incl. QSRs, QSOs, blazars, Seyfert galaxies, BALs, radio galaxies, and LINERS); B7 Thermal processes in galactic nuclei and starburst galaxies (incl. ultraluminous IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
A4 Distance scale; A5 Groups and clusters of galaxies; A6 Gravitational lensing; A7 Intervening absorption line systems; A8 High-redshift galaxies (star formation and ISM). B1 Morphology and galactic structure; and B2 Unresolved and resolved stellar populations; B3 Chemical evolution; B4 Galaxy dynamics; B5 Peculiar/interacting galaxies; B6 Non-thermal processes in galactic nuclei (incl. QSRs, QSOs, blazars, Seyfert galaxies, BALs, radio galaxies, and LINERS); B7 Thermal processes in galactic nuclei and starburst galaxies (incl. ultraluminous IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
A5 Groups and clusters of galaxies; A6 Gravitational lensing; A7 Intervening absorption line systems; A8 High-redshift galaxies (star formation and ISM). B1 Galaxies B1 Morphology and galactic structure; and B2 Unresolved and resolved stellar populations; B3 Galaxy dynamics; B4 Galaxy dynamics; B5 Peculiar/interacting galaxies; B6 Non-thermal processes in galactic nuclei (incl. QSRs, QSOs, blazars, Seyfert galaxies, BALs, radio galaxies, and LINERS); B7 Thermal processes in galactic nuclei and starburst galaxies (incl. ultraluminous IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
A6 Gravitational lensing; A7 Intervening absorption line systems; A8 High-redshift galaxies (star formation and ISM). B1 Morphology and galactic structure; and B2 Unresolved and resolved stellar populations; B3 Galaxies B3 Chemical evolution; B4 Galaxy dynamics; B5 Peculiar/interacting galaxies; B6 Non-thermal processes in galactic nuclei (incl. QSRs, QSOs, blazars, Seyfert galaxies, BALs, radio galaxies, and LINERS); B7 Thermal processes in galactic nuclei and starburst galaxies (incl. ultraluminous IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
A7 Intervening absorption line systems; A8 High-redshift galaxies (star formation and ISM). B1 Morphology and galactic structure; and B2 Unresolved and resolved stellar populations; B3 Chemical evolution; B4 Galaxy dynamics; B5 Peculiar/interacting galaxies; B6 Non-thermal processes in galactic nuclei (incl. QSRs, QSOs, blazars, Seyfert galaxies, BALs, radio galaxies, and LINERS); B7 Thermal processes in galactic nuclei and starburst galaxies (incl. ultraluminous IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C ISM, C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
A8 High-redshift galaxies (star formation and ISM). B1 Morphology and galactic structure; and B2 Unresolved and resolved stellar populations; galactic nuclei B3 Chemical evolution; B4 Galaxy dynamics; B5 Peculiar/interacting galaxies; B6 Non-thermal processes in galactic nuclei (incl. QSRs, QSOs, blazars, Seyfert galaxies, BALs, radio galaxies, and LINERS); B7 Thermal processes in galactic nuclei and starburst galaxies (incl. ultraluminous IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
Galaxies and B2 Unresolved and resolved stellar populations; B3 Chemical evolution; B4 Galaxy dynamics; B5 Peculiar/interacting galaxies; B6 Non-thermal processes in galactic nuclei (incl. QSRs, QSOs, blazars, Seyfert galaxies, BALs, radio galaxies, and LINERS); B7 Thermal processes in galactic nuclei and starburst galaxies (incl. ultraluminous B8 Gentral supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; blanetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
and galactic nuclei B3	
galactic nuclei B3 Chemical evolution; B4 Galaxy dynamics; B5 Peculiar/interacting galaxies; B6 Non-thermal processes in galactic nuclei (incl. QSRs, QSOs, blazars, Seyfert galaxies, BALs, radio galaxies, and LINERS); B7 Thermal processes in galactic nuclei and starburst galaxies (incl. ultraluminous IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C ISM, Star formation and C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
B4 Galaxy dynamics; B5 Peculiar/interacting galaxies; B6 Non-thermal processes in galactic nuclei (incl. QSRs, QSOs, blazars, Seyfert galaxies, BALs, radio galaxies, and LINERS); B7 Thermal processes in galactic nuclei and starburst galaxies (incl. ultraluminous IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
B5 Peculiar/interacting galaxies; B6 Non-thermal processes in galactic nuclei (incl. QSRs, QSOs, blazars, Seyfert galaxies, BALs, radio galaxies, and LINERS); B7 Thermal processes in galactic nuclei and starburst galaxies (incl. ultraluminous IR galaxies (incl. ultraluminous IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
B6 Non-thermal processes in galactic nuclei (incl. QSRs, QSOs, blazars, Seyfert galaxies, BALs, radio galaxies, and LINERS); B7 Thermal processes in galactic nuclei and starburst galaxies (incl. ultraluminous IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
QSRs, QSOs, blazars, Seyfert galaxies, BALs, radio galaxies, and LINERS); B7 Thermal processes in galactic nuclei and starburst galaxies (incl. ultraluminous IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
radio galaxies, and LINERS); B7 Thermal processes in galactic nuclei and starburst galaxies (incl. ultraluminous IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
B7 Thermal processes in galactic nuclei and starburst galaxies (incl. ultraluminous IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
galaxies (incl. ultraluminous IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
IR galaxies, outflows, emission lines, and spectral energy distributions); B8 Central supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
spectral energy distributions); B8	
B8 Central supermassive objects; B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
B9 AGN host galaxies. C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
ISM, C1 Gas and dust, giant molecular clouds, cool and hot diffuse and translucent clouds; and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
star formation and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
and C2 Chemical processes in the interstellar medium; planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	gas,
planetary systems C3 Star forming regions, globules, protostars, HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
HII regions; C4 Pre-main-sequence stars (massive PMS stars, Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
Herbig Ae/Be stars and T Tauri stars); C5 Outflows, stellar jets, HH objects; C6 Main-sequence stars with circumstellar matter, early evolution;	
C6 Main-sequence stars with circumstellar matter, early evolution;	
C6 Main-sequence stars with circumstellar matter, early evolution;	
C7 Young hipping brown dwarfs exceller planet seems	
Ci roung biliaries, brown dwarfs, exosolar planet searc	hes;
C8 Solar system (planets, comets, small bodies).	
O Stellar D1 Main-sequence stars;	
evolution D2 Post-main-sequence stars, giants, supergiants,	
AGB stars, post-AGB stars;	
D3 Pulsating stars and stellar activity;	
D4 Mass loss and winds;	
D5 Supernovae, pulsars;	
D6 Planetary nebulae, nova remnants and	
supernova remnants;	
D7 Pre-white dwarfs and white dwarfs, neutron stars;	
D8 Evolved binaries, black-hole candidates, novae,	
X-ray binaries, CVs;	
D9 Gamma-ray and X-ray bursters;	
D10 OB associations, open and globular clusters,	
extragalactic star clusters;	
D11 Individual stars in external galaxies, resolved stellar	
D12 Distance scale – stars.	population

\ObservingRun{A}{}{}{}{}{}{}{}{}{}{}{}{}{}{}{}

A Normal Programme may have up to 26 runs. Since the space for the run description in Box 3 is limited to 10 lines, a new box containing the observing runs beyond this limit will be created at the end of the proposal form if needed.

- 2. PERIOD KEYWORD. Provide the period number in the second pair of curly braces. For normal proposals this must always be 96, while for Monitoring programmes this can span up to 4 periods.
- **3. INSTRUMENT.** Provide the keyword of the instrument required for each observing run. The complete list of keywords of all instruments offered in Period 96 for Normal Programmes is given in Table 3.

	Table 3:	Keywords	of Available	Instruments	for Normal	Programmes
--	----------	----------	--------------	-------------	------------	------------

Telescope	Instrument keywords
UT1	NACO ¹ , FORS2, KMOS
UT2	FLAMES, XSHOOTER, UVES
UT3	SPHERE ² , VIMOS, VISIR ³
UT4	HAWKI, MUSE, SINFONI
VISTA	VIRCAM
VST	$\mathrm{OMEGACAM}^4$
APEX	CHAMPP ⁵ , FLASH ⁵ , ARTEMIS, LABOCA, SHFI, SUPERCAM, SpecialAPEX
$ m NTT^6$	SOFI, EFOSC2, SOFOSC ⁷ , SpecialNTT
3.6^{6}	HARPS, Special3.6

¹ NACO availability is subject to successful re-commissioning in September 2014;

Provide, in the third pair of curly braces, the instrument(s) required for each observing run. For example,

\ObservingRun{A}{96}{FORS2}{}{}{}{}{}{}{}{}

For visitors bringing their own Visitor Instrument, please use the corresponding "Special" keyword, e.g., "Special3.6" for an instrument to be mounted on the 3.6, and "SpecialNTT" for the NTT. It is then compulsory to fill in the Visitor Instrument information page (see Sect. 3.23).

4. REQUESTED TIME. In order to allow for the automated scheduling of proposals, you must specify the amount of time that you are requesting (hours in Service Mode and nights in Visitor Mode).

For Service Mode (SM), at this stage only provide the total number of hours requested, followed by the letter h for hours. This should include also the time related to any special calibrations required in addition to the standard calibrations provided by ESO. Any more detailed information about possible particular scheduling features will be provided during the preparation of the Phase 2 Service Mode proposal.

For Visitor Mode (VM) proposals, the runs are scheduled according to the information supplied in the Phase 1 proposal form. Consequently, you are required to provide detailed information about any particular scheduling requirements for the successful completion of your programme in this form. You should code the requested time (in nights), starting with the total time requested, followed by the letter n for nights, according to the examples below. If runs are to be split they should be specified using the following convention:

² SPHERE availability is subject to successful commissioning during Period 94;

³ VISIR availability is subject to successful re-commissioning in November 2014;

⁴ OMEGACAM Normal Programmes are restricted to poor weather conditions in Period 96.

⁵ CHAMPP and FLASH are APEX PI instruments, in order to propose the use of these instruments the instrument PI must be contacted at least two weeks prior to submitting the proposal (see the Call for Proposals):

⁶Only Visitor Mode observations are allowed on telescopes at La Silla.

⁷ SOFOSC is the instrument configuration to be specified for combined EFOSC-2/SOFI runs (Section 3.3).

Table 4: Keywords of Available Instruments for Monitoring Programmes

Telescope	Instrument keywords
UT1	NACO ¹ , FORS2, KMOS
UT2	FLAMES, UVES
UT3	$SPHERE^2$, $VIMOS^3$
$UT4^4$	HAWK-I, SINFONI ⁵ , MUSE
$APEX^6$	LABOCA, SHFI

¹ NACO availability is subject to successful re-commissioning in September 2014;

4n	for 4 consecutive nights,
4n=4x1	for 4 times 1 night, with intervals in between,
6n=3x2	for 3 times 2 consecutive nights,
7n = 3x2 + 1	for 3 times 2 consecutive nights followed by 1 night.

If the programme needs run durations of a half-night, you can code the requested time (in nights and half-nights), starting with the total time requested, following the few examples below (with $\mathbf{H1} = \text{first half of the night}$, and $\mathbf{H2} = \text{second half of the night}$):

2n=4H1	for 4 consecutive first naives of a night,
2n=4x1H2	for 4 times the second half of a night, with intervals in between,
3n=3x2H1	for 3 times 2 consecutive first halves of a night,
8n = 3x2 + 4H2	for 3 times 2 consecutive nights followed by 4 consecutive second halves of a night,
8n=3x2+4x1H1	for 3 times 2 consecutive nights followed by 4 non-consecutive first halves of a night.

There is flexibility in this macro so that sub-runs can be specified for fractions of a night (not just half-nights). Where a decimal value precedes the H, H1 or H2 values, they specify the fraction of the night that is required for each observation. Only one decimal place is allowed. For example:

```
\begin{array}{ll} 0.6\mathrm{n}{=}2\mathrm{x}0.6\mathrm{H}1 & \text{for 2 consecutive 0.3n runs - starting in the first half of the night,} \\ 1.8\mathrm{n}{=}3\mathrm{x}1.2\mathrm{H}2 & \text{for 3 consecutive 0.6n runs - starting at the end of the first half of the night,} \\ 2.5\mathrm{n}{=}2\mathrm{H}1{+}2\mathrm{x}1.6\mathrm{H}2 & \text{for 2 consecutive first halves of a night followed by 2 x 0.8n runs.} \end{array}
```

Please note that designations such as 2.0n=2x2.0H1 are invalid. 2.0H1 implies a run that is one whole night and will therefore be rejected by the proposal receiver. In this case the user should specify 2n=2x1. Similarly any values greater than or equal to 2.0 preceding H, H1 or H2 will be rejected.

Provide the total amount of time that is required for the observing run in the fourth pair of curly braces, with details about possible sub-runs. For example,

² SPHERE availability is subject to successful commissioning during Period 94;

³ VIMOS Monitoring Programmes are subject to the restrictions outlined in Section 1.1 of the Call for Proposals;

⁴ Monitoring programmes for UT4 instruments must be compatible with constraints imposed by the AOF installation activities described in Section 1.3 of the Call for Proposals (taking into account that the schedule described is provisional).

⁵ Monitoring programmes using SINFONI must be compatible with the AOF installation activities described above as well as the likely changes to SINFONI in Period 96 as described in Section 1.3 of the Call for Proposals:

⁶ APEX observations for approved MPs can only be carried out in the ESO time-slots.

For all non-consecutive schedules, the details of the time intervals between the different sub-runs must be provided in Box 12 (see Sect. 3.18).

5. MONTH PREFERENCE. Provide the first three letters of the month (e.g. dec) that is your first preference for scheduling (valid months are the ones included in the current period, namely oct, nov, dec, jan, feb, mar). For Service Mode runs "any" should be entered here with any specific requirements detailed in Box 12. For example,

Please note that this month preference should be consistent with the time constraints specified in Box 12 (Sect. 3.18).

- **6. MOON REQUIREMENT.** Provide the required phase of the moon, by using only one of the following three characters (see the Call for Proposals for the exact definition), namely:
 - d for "dark time"
 - \bullet g for "grey time"
 - n for "no restriction"

For example,

7. SEEING REQUIREMENT. Provide the required maximum acceptable seeing value (FWHM in arcseconds) in the V-band at zenith (see the Call for Proposals). The value should correspond to the one given as input to the ETC. Your requirement must be one of the following values:

0.4, 0.6, 0.8, 1.0, 1.2, 1.4, n

For example,

 $\Description A = 96 = 3x2+4H2 = 3x$

8. TRANSPARENCY REQUIREMENT. Provide the transparency condition of the atmosphere required during your observations (see the Call for Proposals for the exact definition). Your requirement must be one of the following values:

CLR for clear sky, although with some rare clouds

PHO for photometric, a perfect night

THN for thin cirrus, inducing absorption up to 0.2 mag

For example,

 $\color= \color= \col$

9. OBSERVING MODE. Provide the requested observing mode: v = Visitor Mode and s = Service Mode. For example,

 $\label{lem:loss} $$ \operatorname{Nn-3x2+4H2}{\det}{dec}{d}_{0.8}{PH0}_{v}{} $$$

10. RUN TYPE. This should only be filled in if the programme is a TOO or GTO and the run is a TOO Run. The RUN TYPE field must be left blank for all Normal and Calibration Programme proposals.

For programme types that allow TOO Runs (TOO, GTO and DDT), this feature should be used only for ToO runs. Briefly, ToO Runs are defined to be runs for which the target cannot be known more than one week before the observation must be carried out. If you want to specify one of your runs as being a ToO run, then please enter "TOO" in the tenth (and final) brackets.

More details corresponding to this ToO run must be specified in the \TOORun macro (Sect. 3.22). This field should be left blank for all other (non-TOO) runs.

Alternative runs

For each requested run, you may specify one or several "alternative runs" for possible execution of the proposed observations with another instrument (in general mounted on another telescope). To this effect, add another line in Box 3, with in the first pair of curly braces, the letter identifying your primary run, followed by "/alt". For example,

```
\ObservingRun{A}{96}{FORS2}{20h}{dec}{d}{0.8}{CLR}{s}{} \ObservingRun{A/alt}{96}{VIMOS}{6n}{dec}{d}{0.8}{CLR}{v}{}
```

indicates that the observations of run A could be obtained through allocation either of 20 hours in Service Mode with FORS2 (primary choice) or of 6 nights in Visitor Mode with VIMOS (secondary choice). You may specify several alternative runs for each primary run (e.g., in the example above, an EFOSC2 run might be a suitable alternative).

Multiple runs

If more than one run is needed for execution of the programme, then fill as many lines as needed. For example,

APEX users should note that all observations for a given APEX instrument must be included in a **single run** for each period. The proposal receiver will reject any proposal with more than one run per period per APEX instrument.

Programmes using both EFOSC-2 and SOFI at La Silla

As described in the Call for Proposals, proposals involving observations with both EFOSC-2 and SOFI, for which the amount of time with either instrument is less than 3 nights can be carried out at La Silla. Note that the runs should be contiguous and the requested time summed over both the EFOSC-2 and SOFI runs is ≥ 3 nights. These types of programmes should be specified by using the pseudo-instrument SOFOSC in the Instrument field and the total amount of time over the EFOSC-2 and SOFI runs.

For instance, if you require 3 nights on EFOSC-2 and 2 nights on SOFI, the combined run would be specified as follows using the **\ObservingRun** macro:

```
\label{losservingRunA} $$ \operatorname{SOFOSC}_{5n}_{dec}_{0.8}_{PHO}_{v}$$
```

and in the justification for the requested amount of time (Box 8, macro \WhyNights), you should indicate the actual lengths of each of the two runs. SOFOSC runs (i.e. combined EFOSC-2 and SOFI runs) with a total duration of less than 3 nights are not accepted.

Please specify the instrument configuration for the longest run in the macro \INSconfig. For example, if the longest run is with the EFOSC-2 instrument:

```
\INSconfig{A}{SOFOSC}{Spectro-long-slit}{Grism\#2:510-1100}
```

Proprietary time

The default data proprietary time is 12 months. Nevertheless, you can ask to reduce it for your data by using the macro \ProprietaryTime{time}. The time is expressed in months, and only the following values can be entered: 0, 1, 2, 6, 12. For example,

```
\ProprietaryTime{6}
```

Please note that this macro does not produce any printable output at compilation, but the information that it contains will be duly stored in ESO's database when the proposal is submitted.

3.4 Past, Present, and Future of this Programme: BOX 4

In order to allow for the evaluation of the proposal within the broader context of the project of which it is part, taking into account the observations already obtained in the past and the data still to be acquired in the future, indicate in Box 4:

- \AwardedNights: the amount of time (in nights or hours) allocated to this project in previous periods, together with the programme number (e.g., 094.B-1234), and the telescope on which this time was allocated:
- \FutureNights: the amount of time (in nights or hours) still necessary, in the future, after this proposal, to complete the programme, if any, and the corresponding telescope(s).

For example,

```
\AwardedNights{UT1}{4n in 094.B-1234}
\FutureNights{UT3}{20h}
```

3.5 Special Remarks: BOX 5

Take advantage of this box to provide any special remark (up to three lines). For example,

```
\SpecialRemarks{This programme is a resubmission, in updated form, of proposal 094.B-1234, which had been granted 2n in VM with UT2+UVES and was entirely clouded out.}
```

3.6 Name and Affiliations of PI and CoI(s): BOX 6

The macro \PI must be used to identify the Principal Investigator (PI) of the proposals. Its parameters are simply the ESO User Portal username of the PI. Do not fill in the PI's full name as the receiver will convert the ESO User Portal username to the PI's full name on submission. Usage of this macro is illustrated in the following example, please note that the username is case-sensitive:

```
\PI{JSMITH99}
```

You should use the macro \CoI to specify also, for all the Co-Investigators (CoIs) of this proposal, their initial(s), last name, and institute code. Institutes and their codes are listed according to country at the following webpage:

http://www.eso.org/sci/observing/phase1/countryselect.html.

You should have one instance of the macro \CoI for each CoI of the proposal. Even if several CoIs share the same institute please use a separate macro for each CoI. The number of CoI macros you can use is in principle unlimited. However, please note that even if all CoIs do not appear in the printed version of the form, the entire CoI list is stored in the ESO database, where it can be accessed when required.

Please note: Due to the way in which the proposal receiver system parses the CoI macro, the number of pairs of curly brackets, {}, in this macro must be strictly equal to 3, i.e., the number of parameters of the macro. Accordingly, curly brackets should not be used within the parameters (e.g., to protect LaTeX signs). For instance:

```
\CoI{L.}{Ma\c con}{1150}
\CoI{R.}{Men\'endez}{1098}
are valid, while
\CoI{L.}{Ma{\c}con}{1150}
\CoI{R.}{Men{\'}endez}{1098}
```

are not. Unfortunately the receiver does not give an explicit error message when such invalid forms are used in the CoI macro, but the processing of the proposal keeps hanging indefinitely.

An example of a CoI list follows:

```
\CoI{H.}{Cerny}{1150}
\CoI{S.}{Bailer-Brown}{1088}
\CoI{K.L.}{Giorgi}{1164}
\CoI{S.}{Lichtman}{2047}
\CoI{L.}{Men\'endez}{1150}
```

When the proposal is submitted each institute code will be converted to the corresponding country and institute by the online receiver.

3.7 Description of the Proposed Programme: BOX 7

The next two pages contain the description of the proposed programme, including all text, references and figures. This is restricted to TWO pages *including* figures. The text sections are composed of two required components, which are activated by two macros.

A – Scientific rationale: this section should describe the scientific background of the project, with pertinent references; any previous work in the field plus the justification for the present proposal should be included. The content of this section should be placed between the curly braces of the macro \ScientificRationale{}.

B – Immediate objective of the proposal: this section should state what is actually going to be observed and what will be extracted from the observations, so that the feasibility becomes clear. The content of this section should be placed between the curly braces of the macro \ImmediateObjective{}.

The references should preferably use the simplified abbreviations used in Astronomy & Astrophysics.

THE RELATIVE LENGTHS OF EACH OF THESE SECTIONS ARE VARIABLE. ALL TEXT AND FIGURES MUST FIT WITHIN A TOTAL OF TWO PAGES. Any text not fitting within the allocated page can be added to the next page (with the figures). It is the responsibility of the proposers to check that their programme description does not exceed the maximum acceptable length. To this effect, proposers should carry out a careful visual inspection of a print-out of their proposal prior to submitting it. Also, when the proposal is compiled with pdfIATEX, the length of the text is checked, and a warning message is issued if it is greater than 2 pages. While this warning may easily be overlooked in the real-time terminal window from which pdfIATEX is run because of the continued scrolling resulting from other output, it is recorded in the pdfIATEX logfile.

3.8 Figures: BOX 7 (cont'd)

Up to ONE page of figures and text can be added in addition to the page detailing the Scientific Justification and the Immediate Objective. There is a size limit of 1MB for each figure to be uploaded. This material can be included using the macros \MakePicture{}{} and \MakeCaption{}. Any additional text from the sections in the previous page (e.g., Immediate Objective) can also be included using the \MakeCaption{}.

NOTE THAT POSTSCRIPT PICTURES ARE NOT ACCEPTED. Since the proposals are compiled using the pdfLATEX package, only JPEG and PDF file formats are accepted. Attachments in other formats should be converted into one of the accepted formats using appropriate tools (such as ps2pdf, convert, or gimp). In order to reduce the size of the attachments, we strongly suggest the use of the PDF format for simple plots and graphs, and JPEG for large figures (such as astronomical images).

The figure macro \MakePicture{}{} has two arguments: the name of the file of the picture, and a list of optional keywords specifying formatting parameters of the image (as defined in the graphicx package). For example:

```
\MakePicture{MyPic1.pdf}{width=15cm,height=8.0cm,angle=90}
\MakePicture{MyPic2.jpg}{width=12cm}
```

The filename should have a .jpg or .jpeg extension for JPEG files, and a .pdf extension for PDF files; other extensions are not accepted. If there is no extension in the filename the receiver will hang and submission will not be possible.

The caption macro \MakeCaption{} takes one single argument, which should contain any LATEX caption. For example:

```
\MakeCaption{Insert caption using LaTeX.}
```

These attachments will be printed on up to one page immediately following the scientific description. You must check the pdf output generated by pdfIATEX before submitting your proposal to make sure that the attachments are properly included. In particular, colour figures should still be readable if printed in black and white. Also, it is your responsibility to check that your attachments fit within the allocated 1 page. Please note that when the proposal is compiled with pdfIATEX, the space required by the attachments is checked, and a warning message is issued if it exceeds 1 page. While this warning may easily be overlooked in the real-time terminal window from which IATEX is run because of the continued scrolling resulting from other output, it is recorded in the logfile generated by LaTeX. You are strongly encouraged to check this log file.

3.9 Justification of Requested Time: BOX 8

In this box, you should provide a careful justification of the requested lunar phase and of the requested amount of time. To this effect, you should use the ESO Exposure Time Calculators whenever possible; these exist for all Paranal and La Silla instruments and are available at http://www.eso.org/observing/etc.

Links to exposure time calculators for APEX instrumentation can be found in Section 7 of the Call for Proposals.

For each telescope and instrument to be used, please specify the version of the ESO Exposure Time Calculator that you have used. Do **not** include any correction for unexpected meteorological conditions. The text should be typed as arguments of the following two macros:

\WhyLunarPhase{} \WhyNights{}

3.10 Telescope Justification: BOX 8a

This section should provide a justification for the use of the selected telescope(s) (e.g., VLT, NTT, etc...) with respect to other available alternatives. The content of this section should be placed between the curly braces of the macro \TelescopeJustification{}.

3.11 Observing Mode Justification: BOX 8b

This section should provide a justification for the observing mode requested (Visitor or Service). The content of this section should be placed between the curly braces of the macro \ModeJustification{}.

3.12 Calibration Request: BOX 8c

For Service Mode runs, the calibrations foreseen in the instrument calibration plans are absorbed by the Observatory; they do not need to be included in the amount of requested time. In Visitor Mode, up to 30 min per night is devoted to the acquisition of these calibrations by the Observatory staff. If, in order to achieve the scientific goals of your projects, calibrations not foreseen in the respective calibration plan are required, you must include the additional amount of time that is needed to obtain them in the total amount of time that you are requesting.

The macro \Calibrations must be used to specify the calibration requirements of your proposal. It takes two arguments. The first one should be set to standard if the calibrations contemplated in the calibration plan are sufficient. In this case, no input is required for the second argument:

\Calibrations{standard}{}

If, on the other hand, you need additional calibrations, the first argument must be set to special, and a brief description of non-standard calibrations that you need must be given as second argument. For example,

\Calibrations{special}{Adopt a special calibration}

Note that non-standard daytime calibrations must be specified here, but contrary to additional nighttime calibrations, the corresponding time need not be included in the total amount of requested time.

3.13 Last Use of ESO Facilities: BOX 9

The macro \LastObservationRemark must be used to provide a brief report on the use of the ESO facilities during the last 2 years. You should specify the programme identification numbers, and describe the status of the data obtained, and the scientific output generated.

3.14 ESO Archive: BOX 9a

You should use the \RequestedDataRemark macro to indicate if the data requested in the proposal are in the ESO Archive (http://archive.eso.org), and if so, to explain the need for new data.

3.15 Duplication of GTO/Public Surveys: BOX 9b

Specify whether there is any duplication of targets/regions covered by ongoing GTO and/or Public Survey programmes. If so, please explain the need for the new data here. Details on the protected target/fields in these ongoing programmes can be found at the following webpages for GTO programmes and Public surveys: http://www.eso.org/sci/observing/teles-alloc/gto.html

http://www.eso.org/sci/observing/PublicSurveys/sciencePublicSurveys.html.

3.16 Applicant's Publications: BOX 10

The macro \Publications{} should be used to provide a list of your publications related to the subject of the current proposal and published during the past two years. The A&A simplified abbreviations for references should be used. The individual references should be separated with a small amount of vertical space, to be created with the standard LATEX command \smallskip\\. For example:

```
\Publications{
Name1 A., Name2 B., 2001, ApJ, 518, 567: Title of article1
\smallskip\\
Name3 A., Name4 B., 2002, A\&A, 388, 17: Title of article2
\smallskip\\
Name5 A. et al., 2002, AJ, 118, 1567: Title of article3
}
```

3.17 List of Targets: BOX 11

Provide the complete list of targets to be observed in this programme, by using the macro \Target with the following nine parameters: run identifier (you may use the same target/field in more than one run), target field/name, Right Ascension (hh mm ss.f, or hh mm.f, or hh.f) and Declination (dd mm ss, or dd mm.f, or dd.f) for the J2000 equinox, requested time on target (in hours with overheads and calibration included), magnitude, angular diameter, additional information (see below), and reference star identifier (see below) for each target field. Please use the format {00 00 00} in case of unknown coordinates. There can be as many occurrences of the macro \Target as required to accommodate all targets of all runs of the programme. Long lists of targets will continue on the last page(s) of the proposal form.

Please note that the scheduling of your programme will take all targets given in this list into account. Only include the targets requested for Period 96. Make sure your targets are significantly observable during this period. Inclusion of targets with insufficient visibility during the current period may result in rejection of your programme by the automatic scheduler.

The additional information field (8th argument of the \Target macro) may in general be used to provide any relevant piece of information about the target that does not pertain to any other argument of the macro (e.g. the period of a variable star). However, for APEX targets, usage of this field is **mandatory** to indicate the requested Precipitable Water Vapour (PWV) and the acceptable range of Local Sidereal Time (LST) for the considered observation. The format should be similar to the one shown in the following example:

A reference source identifier must be provided for all natural guide stars (NGS), in the case of NGS observations with NACO, SINFONI and CRIRES, and all tip-tilt stars (TTS), in the case of all laser guide star (LGS) observations with SINFONI. For observations with the noAO modes of SINFONI and NACO, you do not need to provide this information. The reference source designation has to be the exact identifier of the selected star either from the Guide Star Catalog 2 (GSC2) or the 2MASS point source catalogue. Note that GSC2 stars identifiers should NOT be preceded by GSC2, but must start with either N or S. In case the reference source is not included in either catalogue, for instance because it is a supernova or a solar system object, "alt" should be entered as reference source identifier, and additional information can be provided in the \TargetNotes macro. Rules for reference star designation can be found for GSC2 at:

http://vizier.u-strasbg.fr/viz-bin/VizieR-n?-source=METAnot&catid=1271¬id=1&-out=text.

Examples of valid and invalid GSC2 identifiers are given below:

N01230121	good
S33333331	good
n01230121	bad
N012301201	bad
S01230141	bad
S3333333000001	good
S01201201234567	bad

For 2MASS, the rules for reference star designation are available at: http://www.ipac.caltech.edu/2mass/releases/allsky/doc/sec2_2a.html.

Here are some examples of correct and incorrect identifiers:

```
2MASS J01234567+7801020 good
2MASS J00000000+7801020L good
2MASS J01234567+9000000 good
2MASS J01234567+9000000W good
2MASX J01234567+7801020 bad
2MASS J97234567+7801020 bad
```

Thus the following examples illustrate the correct usage of the **\Target** macro when a reference star must be specified:

```
\Target{B}{NGC 105}{22 55 00}{-47 50 30}{9.0}{}}{$S33333331}
\Target{C}{NGC 106}{00 24 43}{-05 09 00}{2.0}{}}{$2MASS J01234567+7801020}
```

The macro \TargetsNotes{} should be used to include any comments that apply to several or all targets (or to specify reference stars that are not found in the GSC2 or 2MASS catalogues).

```
\TargetsNotes{The planned grid of pointings around the targets listed above will be defined during the first observing night.}
```

3.18 Scheduling Requirements: BOX 12

If your proposal involves any of the following:

- observations to be executed on specific dates (e.g., for simultaneity with observations at other facilities);
- observations to be executed at pre-defined time intervals (e.g., at different epochs so as to achieve phase coverage of a periodically variable target);
- Visitor Mode runs split into non-consecutive nights (for which the fourth argument specifying the number of nights in Box 3 of the proposal form includes a formula);
- Visitor Mode runs mutually linked, which need to be scheduled in a given sequence and at specified time intervals.

you must uncomment the macro \HasTimingConstraints. Please leave the brackets in \HasTimingConstraints blank. Details of time constraints can be entered in Special Remarks and using the other flags in Box 12.

Please note that the macro \HasTimingConstraints should be commented out in the following cases:

- for scheduling constraints resulting only from the genuine visibility window of the target sources (defined by their location in the sky) or from the phases of the Moon;
- for time series of observations acquired during a single night or over several consecutive nights of a contiguous Visitor Mode run;
- for Target of Opportunity observations.

In order to allow for the automated scheduling of all Visitor and Service Mode observing runs, you must provide all information related to the details of the way your programme should be scheduled. If you have any doubts about how to specify your particular requirements please email OPO at opo@eso.org well in advance of the proposal submission deadline.

1. RUN SPLITTING. For Visitor Mode runs: if the fourth argument you have provided in Box 3 indicates a simple number of consecutive nights or half-nights, e.g.:

```
\DbservingRun{A}{96}{FORS2}{4}{nov}{d}{0.8}{CLR}{v}{}
```

you do not have to do anything. If the fourth argument in Box 3 indicates a more complicated requirement, with some non-consecutive nights, e.g.:

```
\color= \col
```

you must provide some additional information. The fourth argument above, namely, 8n=3x2+4H2, means that you request a total of 8 nights, made of three sub-runs of 2 consecutive nights each, followed later by a sub-run of 4 consecutive second halves of a night.

You should use the macro \RunSplitting{}{}, and put the run identifier in the first argument. The second argument should indicate the way the run should be split into different sub-runs. If the interval between two sub-runs has to be exactly a given number of days, say 20, then this is a **strong** constraint and this number of days should be followed by the letter s. If the interval between two sub-runs has some tolerance on the number of days, say 20±5, then this is a **weak** constraint and this number of days should be followed by the letter w. Consequently, if 8n=3x2+4H2 means that you want three sub-runs of 2 consecutive nights each, the first and the second separated by 10 nights exactly, the second and the third separated by more or less 20 nights, followed exactly 15 days later by one sub-run of 4 consecutive second halves of a night, then the second argument of the macro \RunSplitting{}} should contain the following expression: 2,10s,2,20w,2,15s,4H2. Hence the following entry should appear in your proposal file:

```
\mathbb{A}_{2,10s,2,20w,2,15s,4H2}
```

Note that it is necessary to specify explicitly the time intervals among all non consecutive sub-runs.

The macro \RunSplitting is meaningless for Service Mode runs, for which constraints of the considered type should be fully specified at Phase 2 (but the macro \HasTimingConstraints should be uncommented for such runs, so that they are duly flagged). However please note that in Service Mode, monitoring programmes are executed on a best effort basis only. In particular, a monitoring sequence can be interrupted by unsuitable weather conditions or by runs scheduled in Visitor Mode.

2. LINK FOR COORDINATED OBSERVATIONS. If you have requested three different runs in Box 3, e.g.:

```
\label{losservingRun} $$ \operatorname{S}_{96}_{FORS2}_{2n}_{nov}_{d}_{0.8}_{v}_{v}_{0bservingRun}_{96}_{FORS2}_{3n}_{nov}_{d}_{0.8}_{PH0}_{v}_{0bservingRun}_{96}_{UVES}_{20h}_{nov}_{d}_{0.8}_{CLR}_{v}_{}$
```

and would like some of them to be simultaneous and some later than others, independently of the exact period of scheduling, then use simultaneous, after and the macro \Link{}{}{} in the following way:

```
\Link{B}{after}{A}{10}
\Link{B}{after}{A}{}
\Link{B}{simultaneous}{C}{}
```

3. UNSUITABLE PERIOD(S) OF TIME. If you have requested two nights in Box 3 and would like them to be scheduled to avoid some unsuitable periods of time, for some reason, then use the macro \UnsuitableTimes{}{}{} in the following way:

```
\UnsuitableTimes{A}{15-jan-16}{18-jan-16}{International Conference} \UnsuitableTimes{B}{15-jan-16}{18-jan-16}{International Conference} \UnsuitableTimes{C}{15-jan-16}{18-jan-16}{International Conference} \UnsuitableTimes{C}{1-jan-16}{3-jan-16}{Committee Meeting}
```

Dates correspond to 12:00 noon Local Time at the Observatory location (i.e., in Chile). In other words the first date refers to the start of the first night of the unsuitable period; the second date refers to the end of the last night. As with the TimeCritical macro, only one run can be specified in each UnsuitableTimes macro. Further explanation can also be entered in \SpecialRemarks{} (Box 5).

3.19 Scheduling Requirements contd..: BOX 12

4. SPECIFIC DATE(S) FOR TIME CRITICAL OBSERVATIONS. If you have requested 2 nights in Box 3, e.g.:

and if for some reason (e.g., specific phase of a variable object or parallel observations with already scheduled HST observations, etc.) you need these two nights scheduled between some specific dates, then use the macro \TimeCritical{}{}} in the following way:

Please note the following points:

- As of P87, the constraints set by this macro are checked against the time specified by the **\ObservingRun** in more detail. Some examples of correct and incorrect uses of this pair of macros are shown below.
- The indicated dates correspond to 12:00 noon LOCAL Time at the Observatory location (i.e., in Chile). In other words, the first date refers to the start of the first night of the acceptable interval, and the second to the end of the last night. Please convert event times from Universal Time to Local Time.
- Only one run can be specified in each TimeCritical field. However, as an observing run can have multiple Time critical dates, users can specify multiple separate \TimeCritical macros for a single observing run (as shown above for Observing Run A).
- ESO can still access the information for this macro (and all other macros in Box 12) so do not worry if the constraints overfill the available space.

Examples of correct usages of the \TimeCritical macro:

If the user requires an observing run to be split into three observations, each to taken in the second half of the night over a specific 8-day period, this could be specified using the following macros:

Conversely if the ObservingRun was specified as above but the TimeCritical macro was configured as follows it would be rejected as the configuration 1.5n=3H2 implies there must be a minimum time separation of three nights.

```
\TimeCritical{A}{12-oct-15}{14-oct-15}{reason here}
```

3.20 Instrument configuration: BOX 13

The two template proposals (template.tex for normal applications, and templatelarge.tex for Large Programme applications) contain the full list of configurations for all available instruments at all available ESO telescopes (Paranal, La Silla and Chajnantor). In order to provide general information about the setup of the ESO instrument(s) you plan to use, please uncomment only the lines related to the instrument modes and configurations needed for the acquisition of your desired observations. For some lines related, e.g., to special filters or central wavelength, please add the required information where appropriate (between the already existing curly braces).

Note that you **must** put the run ID within the first pair of curly braces of the relevant lines. **Do not** specify any instrument configuration for alternative runs or visitor instruments (see Box 3). Note that all parameters are **mandatory** for the \INSconfig macro (do not use empty fields).

3.21 Interferometry page

For VLTI runs, a separate run must be specified for each requested baseline configuration. If your proposal includes VLTI runs, you MUST uncomment and fill in the arguments of the macro \VLTITarget with run ID, target name, visual magnitude, magnitude at wavelength of observation, wavelength of observation (in microns), size at wavelength of observation (in mas), baseline (see the following website for available configurations: http://www.eso.org/paranal/insnews/vlti_overview.html), range of visibilities for the specified configuration (at preferred hour angle or at hour angle 0), correlated magnitude, and time on target (ToT) in hours. For example,

```
\VLTITarget{E}{NGC 106}{-0.7}{-3.5}{10.6}{40}{UT1-UT2-UT3}{0.84}{-2.5}{6}
\VLTITarget{F}{NGC 107}{-0.7}{-3.5}{2.1}{40}{UT1-UT2-UT3}{0.84/1.0/0.1}{1./0.5/2.}{6}
\VLTITarget{G}{NGC 108}{-0.7}{1.5}{2.1}{40}{D0-K0-G2-J3}{0.90/0.80/0.70}{1.6/1.7/1.9}{6}
```

For UT observations, please specify one of the available VLTI baselines.

Please specify one of the available AT quadruplets listed in the template proposal files. At Phase 2, the time can then be split among the AMBER triplets of the chosen quadruplet.

For observations with AT baselines, please use one typical baseline of the quadruplet that you have specified in order to compute representative visibility and correlated magnitude values.

For VLTI observations using UT baselines, you should specify the visibilities corresponding to the various baselines as values separated by a slash (/); up to two of the three values can be replaced by a star (*). Similarly, the correlated magnitudes for the various baselines are also specified as three values separated by slash.

You can use the macro \VLTITargetNotes to insert comments about some or all of your VLTI targets. You should take advantage of this macro to indicate suitable alternative baselines for your observations.

3.22 ToO page

If your programme has Target of Opportunity runs (ToO) runs, you should specify these in the \ObservingRun macro using "TOO" in the corresponding field. The ToO information must be filled in for the run using the corresponding \TOORun macro. The arguments in this macro are, in order: the run identifier, the nature of the observation, the number of targets per run, and the number of triggers per targets. There must be one occurrence of the macro \TOORun for each of the ToO runs specified in Run Type field in the \ObservingRun macro (Box 3). The second argument (nature of the observation) may be one of the following keywords:

- RRM, for observations to be triggered via the automated Rapid Response Mode system;
- ToO-hard, for observations to be triggered manually that need to be carried out within 48 hours of receipt of the trigger by the Observatory (and in most cases, as soon as possible), or that involve a strict time constraint (i.e., that must be executed during a specific night);
- ToO-soft, for manually triggered observations for which the Observatory can receive notification more than 48 hours before execution, and which can be scheduled for execution with a flexibility of at least ± 1 day.

Only one keyword can be specified for each run. If observations pertaining to different categories are needed, several runs must be defined. The number of triggers must be indicated for RRM, ToO-hard and ToO-soft observations only. An occurrence of the macro \TOORun looks like the following example:

Table 5: Keywords of Available Instruments (Large Programmes)

Telescope	Instrument keywords
UT1	NACO ¹ , FORS2, KMOS
UT2	FLAMES, UVES
UT3	$SPHERE^2$, $VIMOS^3$
$UT4^4$	HAWK-I, SINFONI ⁵ , MUSE
APEX	LABOCA, SHFI
NTT	SOFI, EFOSC2, SpecialNTT
3.6	HARPS, Special 3.6

¹ NACO availability is subject to successful re-commissioning in September 2014;

$\texttt{T00Run}\{A\}\{\texttt{To0-hard}\}\{2\}\{3\}$

You have the opportunity to add some notes to the ToO page by using the macro \TOONotes. As a rule, ToO proposals should involve at least one trigger of one of either the RRM or ToO-hard or ToO-soft types.

3.23 The Visitor Instrument Page

The following commands are only needed for proposals involving a Visitor Instrument, in which case they are also **mandatory**. You should uncomment them and provide the required information between the different pairs of curly braces. Please do not enter anything in Box 13 (i.e. \INSconfig) for Visitor Instrument runs.

If applying for the VLTI visitor instrument, SpecialVLTI, please remember to uncomment and fill in the \VLTITarget macro and this Visitor Instrument page.

4 HOW TO FILL A LARGE PROGRAMME TEMPLATE

The ESOFORM package includes a specific template (templatelarge.tex) that must be used to generate a Large Programme proposal. This template is only slightly different from the template for Normal Programmes. Hereafter is the list of the differences between the two templates. Any feature not appearing in this list is identical in both templates though the box number may differ.

• Box 2: the abstract for Large Programme proposals can be slightly longer (up to 13 lines) than for Normal Programme proposals (up to 9 lines).

² SPHERE availability is subject to successful commissioning during Period 94;

³ VIMOS LPs must comply with the RA restrictions outlined in the Call for Proposals;

⁴ Large Programmes for UT4 instruments must be compatible with constraints imposed by the AOF installation activities described in Sections 1.1 and 1.3 of Call for Proposals (taking into account that the schedule described is provisional);

⁵ Large programmes using SINFONI must be compatible with the AOF installation activities described above as well as the likely changes to SINFONI in Period 96 as described in Section 1.2 of the Call for Proposals.

- Box 3: proposers must specify all the runs across all the Periods covered by the Large Programme. For Paranal instruments this means a maximum of 4 periods, while for La Silla instruments up to Period 97 is allowed (see the Call for Proposals for more information). A Large Programme can have up to 26 runs. Since the space for the runs description in Box 3 is limited to 12 lines, a new box containing the observing runs beyond this limit will be created at the end of the proposal form if needed.
- Box 3: the list of keywords of all instruments offered in Period 96 for Large Programmes is given in Table 5.
- Box 5: while two pages (including figures) are allowed for the scientific description of Normal Programmes, the description of Large Programmes may take up to three pages followed by two additional pages for figures.

The scientific description is comprised of the following four subsections: \ScientificRationale{}, \ImmediateObjective{}, \TelescopeJustification{}, and

\ModeJustification{}. The \TelescopeJustification{} and \ModeJustification{} macros are the same as those in Boxes 8b and 8c in the Normal Programme template.

As noted above there are also there are two additional pages for attachments (e.g., figures, etc).

- Boxes 6 and 7: specific to the Large Programme proposals, these two boxes should contain the required information describing the experience of the applicants with telescopes, instrumentation, data reduction and delivery of data products to the ESO Archive (\Experience{}). The strategy for data reduction, analysis and a delivery plan for data products into the ESO Archive should be presented along with a brief description of the resources available to the observing team, such as: computing facilities and research assistants (\Resources{}).
- Box 8: Additional remarks or comments can be provided using the macro \SpecialRemarks{}.

5 SUBMISSION OF THE APPLICATION

Proposals must be prepared as pdfLATEX source files, making use of the latest ESOFORM package, corresponding to the ESO Period for which they are submitted. Proposals received in any other format, or with modified ESOFORM macros, will be automatically rejected by the automated proposal handling system.

When the LATEX source file of your application is complete, **please process it with pdfLATEX** so as to identify any possible LATEX format errors. In particular, we **strongly** recommend that you

- review the log file generated by LATEX so as to check for the presence of warning messages issued by the ESOFORM macros. Such messages report, among others, instances in which a text field is too long, so that your input is truncated in the pdf file that is generated, and part of the information that you submit will not be communicated to the OPC;
- carefully inspect a printed copy of the output to make sure that all parts of the application are completed, and that their formatting is appropriate.

Please note that while several checks are performed by the ESOFORM package when running pdfIATEX, successful compilation does not guarantee that a proposal is fully compliant. Indeed, many key checks can only be performed by the proposal reception system after the proposal has been uploaded within the ESO User Portal.

You should verify that your proposal complies with ESO requirements using a "skeleton" version of your proposal that only contains the technical details of your programme. This should be done well before the Phase 1 Deadline. Please upload the "skeleton" proposal as if you were submitting it and follow the instructions online. If the proposal passes all the technical checks you will get the message: "Your proposal is verified and passed all the checks (but not submitted yet)!". You can then either log out of the User Portal or return to the submission page by selecting the option to "Go back to the beginning". More details on the full proposal submission procedure are given below.

Proposals are submitted by logging into the ESO User Portal. In order to do this, proposers must be registered and have activated their User Portal accounts. To submit please go to:

http://www.eso.org/UserPortal

Once you have logged in, Once you have logged in, you should select the item called "Download ESOFORM packages" under the Phase 1 menu items. In order to submit, select "Submit a proposal" from the same menu. You will then be taken to the WASP page (Web Application for Submitting Proposals). On choosing the relevant cycle, you will be asked to upload the LATEX file of your proposal and should follow the subsequent instructions.

A number of checks are executed at the various steps of the submission process; please follow the instructions online. If a problem is detected it will be clearly reported by the system: fix it in your proposal and resubmit it. Once the proposal passes through all the checks, you will be requested to finalise the submission by clicking on the corresponding button. It is essential that you execute this final step: your proposal will not be submitted until this is done, even though you have uploaded all the necessary files!

Upon submission of a correctly completed proposal, the ESO proposal validation software will return an identifier assigned to the valid proposal. This identifier, and the acknowledgment page in which it appears, represent the official confirmation that the proposal successfully entered the proposal handling system. We recommend that you take note of the identifier; you may also want to print the acknowledgment page for your records. In addition, an email confirmation is sent to the submitter and to the PI of the proposal.

Some common problems are described below.

- BibTeX formats are not permitted within the ESOFORM package.
- Figures without the corresponding .jpg or .pdf extensions will make the receiver hang.
- The proposal reception system checks for **the presence of text outside the argument fields of the ESOFORM macros** in the LATEX source of the proposal, and rejects proposals in which such text is found. If there is text outside the macros it appears as text or extra space above the ESO logo on top of the first page. This input **must be commented out or relocated within the relevant macro** before the proposal is submitted.

Submission Problems

The proposal submission acknowledgment page normally appears within seconds of completion of a a submission. However, during the last few hours before the Phase 1 deadline, the system may be slowed down by the high load, and the acknowledgment process may take several minutes. Please **be patient**: even though it may look like "nothing is happening", the system most likely is actually busy processing a queue of proposals. Please **do not** abort your submission or make a new attempt at submitting the same proposal as this would only increase the load on the system and make it slower.

If you have not succeeded in completing your submission within 1 hour, please contact ESO via email at esoform@eso.org. Do not under any circumstance attach your application (in any format) to this email. Do not try to resubmit your application before receiving further instructions from ESO. Do not panic! Once you have uploaded the LATEX source of your proposal, your attempt, and the time at which you initiated it, are recorded in the ESO system, so anomalous delays due to the proposal reception system will be duly identified. Be aware that if you experience difficulties due to the proposal reception system, you are most likely not the only user in this situation. As problem reports must be handled sequentially it may take some time before you receive feedback from ESO.

As mentioned above, the acknowledgment Web page providing the identifier of your proposal is the official confirmation of its successful submission. The subsequent email notification is only sent to you as a secondary confirmation, and delay in its delivery should not represent a concern. However, if you have not received it within 24 hours of your submission, please report this anomaly to esoform@eso.org.

A safe way to avoid submission problems (often related to heavy system load during the last few hours before the deadline) is to submit your proposal early. We strongly encourage you to send in your applications and all attachments several days before the deadline.

All proposals and their attachments must reach the ESO servers via the WASP interface BEFORE 12:00 noon (Central European Summer Time) on the date of the deadline. Please note that ESO cannot provide support beyond 11:00 on that day. Responsibility for verifying that ESO has correctly received, processed, and acknowledged your proposal before the proposal

submission deadline rests entirely with you. Revisions, corrections, and/or modifications will not be accepted after the deadline.

IMPORTANT NOTICE

Electronic proposal submission does not allow applicants to sign their proposals. Therefore ESO assumes that PI's take full responsibility for the contents of the proposal, in particular in regard to the names of co-investigators and the agreement to act according to the instructions for visiting astronomers, should observing time be granted.