

# **Representation Theory in Braid Groups**

by

Jaxon Green

None

A dissertation submitted in partial satisfaction of the  
requirements for the degree of  
Master of Science

in

Mathematics

in the

GRADUATE DIVISION

of the

UNIVERSITY OF CALIFORNIA, SAN LUIS OBISPO

Committee in charge:  
Professor Ben Richert  
Professor of Mathematics, Chair  
Professor Sean Gasiorek

June 2024

# Representation Theory in Braid Groups

Copyright 2024

by

Jaxon Green

# **Abstract**

i

Representation Theory in Braid Groups

by

Jaxon Green

Master of Science in Mathematics

University of California, San Luis Obispo

Professor Ben Richert

Professor of Mathematics, Chair

Write an abstract here.

## Acknowledgments

ii

Any acknowledgements?

## Contents

<b>List of Tables</b>	<b>iv</b>
<b>List of Figures</b>	<b>v</b>
<b>1 Sample Chapter 1</b>	<b>1</b>
1.1 First Section of Introduction . . . . .	1
<b>2 Sample Chapter 2</b>	<b>2</b>
2.1 MY FIRST SECTION . . . . .	2
2.1.1 Interesting subsection title . . . . .	2
2.1.2 Another interesting subsection title . . . . .	2
<b>3 Representation Theory</b>	<b>4</b>
3.1 Introduction to Representations . . . . .	4
<b>Bibliography</b>	<b>8</b>
<b>APPENDICES</b>	
A Appendix A Title . . . . .	9

## List of Tables

## List of Figures

2.1	This is a sample figure. It is not interesting. Note that figures will “float” to wherever $\text{\LaTeX}$ wants to put them. . . . .	3
-----	---	---

## Chapter 1

## SAMPLE CHAPTER 1

## 1.1 First Section of Introduction

This is an equation:

$$c^2 = a^2 + b^2. \tag{1.1}$$



## Chapter 2

## SAMPLE CHAPTER 2

## 2.1 MY FIRST SECTION

There are lots of great resources on the internet to help you learn L<sup>A</sup>T<sub>E</sub>X. Perhaps start with examples like the ones at

[http://en.wikibooks.org/wiki/LaTeX/Sample\\_LaTeX\\_documents](http://en.wikibooks.org/wiki/LaTeX/Sample_LaTeX_documents).

It is important to cite references. [1, 2, 5]

Organize the paper into sections and subsections.

## 2.1.1 Interesting subsection title

You get the idea. Hey, this one has some displayed math,

$$\frac{2}{x} = \sin(\epsilon),$$

not that it makes any sense whatsoever. And here is how you do a numbered equation,

$$\int_0^{y^2} f(x) dx = \sqrt{z+y}. \quad (2.1)$$

Don't forget to punctuate your equations as part of the sentence. You can do inline math, too, as in  $f(x) = \lim_{n \rightarrow \infty} n f(x)/n$ , which is trivial.

## 2.1.2 Another interesting subsection title

Okay, not really.



Figure 2.1: This is a sample figure. It is not interesting. Note that figures will “float” to wherever  $\text{\LaTeX}$  wants to put them.

## Chapter 3

## REPRESENTATION THEORY

## 3.1 Introduction to Representations

Over the course of this chapter, we will develop the theory and utility of representations. At a glance, representations give us the ability to dial back the complexity of a mysterious group by viewing its elements as matrices. Thanks to the rigorous development and study of linear algebra, groups of matrices are well-understood structures. Representations allow us to unravel the mystery of an unknown group structure and reveal a group's fundamental properties as results of linear algebra techniques.

**Definition 3.1.** (*First Definition*) A **representation** of degree  $n$  is a group homomorphism that maps a group into  $GL_n(\mathbb{C})$

$$\phi : G \rightarrow GL_n(\mathbb{C})$$

We say that  $\phi$  is a representation of  $G$ . If  $\phi$  is an injective homomorphism, we say that the representation is **faithful**. Otherwise, the representation is called **degenerate**.

To illustrate the concept of representations, we will consider the group of all roots of unity,  $G$ , for the following examples. We can construct multiple homomorphisms from  $G$  to showcase different kinds of representations.

Note: Since  $GL_1(\mathbb{C})$  as  $\mathbb{C}$  are isomorphic, we identify each  $1 \times 1$  matrix with its corresponding entry with its element in  $\mathbb{C}$ .

**Example 3.2.** (*Trivial Representation*)

$$\text{Let } \begin{array}{l} \phi : G \rightarrow GL_1(\mathbb{C}) \\ g \mapsto 1 \end{array}$$

This map is the trivial homomorphism from  $G$  to  $GL_1(\mathbb{C})$  and therefore it easily satisfies the requirement of a degree 1 representation of  $G$ . We say that  $\phi$  is the **trivial representation** of  $G$ .

**Example 3.3.** (*Nontrivial Degree 1 Representation*)

By construction of  $G$ , if  $g \in G$ , then  $g = e^{\frac{2\pi im}{n}}$  where  $m, n \in \mathbb{Z}$

$$\begin{aligned} \text{Let } \phi : G &\rightarrow GL_1(\mathbb{C}) \\ g &\mapsto g \end{aligned}$$

where we view  $G$  as a multiplicative subgroup of  $\mathbb{C}$ . This observation trivializes the argument that  $\phi$  is a homomorphism. Therefore,  $\phi$  is a degree 1 representation of  $G$ .

**Example 3.4.** (Degree 2 Representation)

$$\begin{aligned} \phi : G &\rightarrow GL_2(\mathbb{C}) \\ \text{Let } e^{2\pi i \frac{m}{n}} &\mapsto \begin{bmatrix} \cos(\frac{2\pi m}{n}) & \sin(\frac{2\pi m}{n}) \\ -\sin(\frac{2\pi m}{n}) & \cos(\frac{2\pi m}{n}) \end{bmatrix} \end{aligned}$$

To show this map is a homomorphism, we will take two elements of  $G$ , say  $e^{2\pi i \frac{x}{y}}$  and  $e^{2\pi i \frac{a}{b}}$  and track the image of their product under  $\phi$ .

$$\begin{aligned} \phi(e^{2\pi i \frac{x}{y}} * e^{2\pi i \frac{a}{b}}) &= \phi(e^{2\pi i (\frac{x}{y} + \frac{a}{b})}) \\ &= \begin{bmatrix} \cos(2\pi(\frac{x}{y} + \frac{a}{b})) & \sin(2\pi(\frac{x}{y} + \frac{a}{b})) \\ -\sin(2\pi(\frac{x}{y} + \frac{a}{b})) & \cos(2\pi(\frac{x}{y} + \frac{a}{b})) \end{bmatrix} \\ &= \begin{bmatrix} \cos(2\pi \frac{x}{y}) \cos(2\pi \frac{a}{b}) - \sin(2\pi \frac{x}{y}) \sin(2\pi \frac{a}{b}) & \sin(2\pi \frac{x}{y}) \cos(2\pi \frac{a}{b}) + \cos(2\pi \frac{x}{y}) \sin(2\pi \frac{a}{b}) \\ -\sin(2\pi \frac{x}{y}) \cos(2\pi \frac{a}{b}) - \cos(2\pi \frac{x}{y}) \sin(2\pi \frac{a}{b}) & \cos(2\pi \frac{x}{y}) \cos(2\pi \frac{a}{b}) - \sin(2\pi \frac{x}{y}) \sin(2\pi \frac{a}{b}) \end{bmatrix} \\ &= \begin{bmatrix} \cos(2\pi \frac{x}{y}) & \sin(2\pi \frac{x}{y}) \\ -\sin(2\pi \frac{x}{y}) & \cos(2\pi \frac{x}{y}) \end{bmatrix} \begin{bmatrix} \cos(2\pi \frac{a}{b}) & \sin(2\pi \frac{a}{b}) \\ -\sin(2\pi \frac{a}{b}) & \cos(2\pi \frac{a}{b}) \end{bmatrix} \\ &= \phi(e^{2\pi i \frac{x}{y}}) * \phi(e^{2\pi i \frac{a}{b}}) \end{aligned} \tag{3.1}$$

Since,  $\phi$  has been shown to be a homomorphism, we can conclude that  $\phi$  is also a degree 2 representation of  $G$ .

Is  $\phi$  faithful or degenerate? A faithful representation would have a trivial kernel. Suppose  $\phi(e^{2\pi i \frac{x}{y}}) = I_2$  ( $I_n$  is the Identity Matrix of dimension  $n \times n$ ).

$$\begin{bmatrix} \cos(2\pi \frac{x}{y}) & \sin(2\pi \frac{x}{y}) \\ -\sin(2\pi \frac{x}{y}) & \cos(2\pi \frac{x}{y}) \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \tag{3.2}$$

Comparing entrywise, we see that  $\cos(2\pi\frac{x}{y}) = 1$  and  $\pm\sin(2\pi\frac{x}{y}) = 0$ . Using any of these<sup>6</sup> equations, we see that  $\frac{x}{y} = n$  for some  $n \in \mathbb{Z}$ . Therefore,  $\ker(\phi) = \mathbb{Z}$  and this representation is degenerate.

Alternatively, we can formulate the definition of a representation in a different context, illuminating a useful interpretation that will be used extensively throughout this paper.

**Definition 3.5.** (Second Definition) Let  $G$  be a group, let  $V$  be a linear vector space, and let  $\mathcal{L}(V)$  be the group of linear operators on  $V$  together with the operation of composition. A **representation** of  $G$  is a group homomorphism that maps  $G$  into  $\mathcal{L}(V)$ .

$$\phi : G \rightarrow \mathcal{L}(V)$$

The degree of the representation is the dimension of  $V$ .

**Remark 3.6.** In the case where we have a finite dimensional vector space, we can make an interesting observation. Suppose  $V$  is finite dimensional and  $G$  is a group. It is easy to identify both definitions of representations with one another. Let  $\{e_i\}_{i=1}^n$  be a basis for  $V$ . Let  $\phi : G \rightarrow \mathcal{L}(V)$  be a representation of  $G$ . Then  $\forall g \in G$ ,  $U_g := \phi(g)$  is a linear operator on  $V$ .  $U_g$  has a corresponding matrix,  $M(U_g)$ , with coefficients defined by the image of our basis vectors of  $V$ .

$$M(U_g) = \begin{array}{cccc|c} U_g(e_1) & U_g(e_2) & \dots & U_g(e_n) & \\ \left[ \begin{array}{cccc} m_{11} & m_{12} & \dots & m_{1n} \\ m_{21} & m_{22} & \dots & m_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ m_{n1} & m_{n2} & \dots & m_{nn} \end{array} \right] & e_1 \\ & e_2 \\ & \vdots \\ & e_n \end{array}$$

$$U_g(e_j) = \sum_{i=1}^n m_{ij} e_i$$

Does the map  $\begin{array}{l} \psi : G \rightarrow GL_n(\mathbb{C}) \\ g \mapsto M(U_g) \end{array}$  satisfy the criteria to be considered a representation (by the first definition)? If  $g, h \in G$ , then

$$\psi(gh) = M(U_{gh}) = M(U_g \circ U_h) = M(U_g) * M(U_h) = \psi(g)\psi(h)$$

Where the homomorphism property of  $\phi$  is used in succession with link between composition and multiplication of operators and their corresponding matrices. Therefore, we have established a very special link between th

## Bibliography

- [1] A. F. Blumberg and G. L. Mellor. A description of a three-dimensional coastal circulation model. In N. Heaps, editor, *Three Dimensional Coastal Ocean Models*, pages 1–16. Amer. Geophys. Union, 1987.
- [2] A. Gill. *Atmosphere - Ocean Dynamics*. Academic Press, 1982.
- [3] P. F. Choboter, R. M. Samelson, and J. S. Allen. A New Solution of a Nonlinear Model of Upwelling. *J. Phys Oceanogr.*, 35:532–544, 2005.
- [4] S. J. Lentz and D. C. Chapman. The importance of non-linear cross-shelf momentum flux during wind-driven coastal upwelling. *J. Phys. Oceanogr.*, 34:2444–2457, 2004.
- [5] J. Pedlosky. A Nonlinear Model of the Onset of Upwelling. *J. Phys Oceanogr.*, 8:178–187, 1978.

## APPENDICES

## Appendix A

## APPENDIX A TITLE

Blah blah