Elementary Analysis

Chapter 2: Sections 7–10

All 45 exercises, with every subpart
Detailed worked solutions

Mathjock

Text and numbering. Kenneth A. Ross, *Elementary Analysis: The Theory of Calculus*, second edition (Springer, 2013), Chapter 2. Exercise numbers follow the copy already linked on Mathjock. These are independently written worked solutions, not the publisher's solutions manual. Each exercise is identified by a short mathematical description.

Reading philosophy. Every argument states its goal, records the assumptions being used, and explains the intermediate algebra or logical inference. Definitions, sign conditions, and both directions of an "if and only if" are made explicit. A cited rule is named at the point where it is used.

Conventions. $\mathbb{N} = \{1, 2, 3, \ldots\}$, $\mathbb{Z} = \{\ldots, -2, -1, 0, 1, 2, \ldots\}$, and $\mathbb{Q} = \{p/q : p, q \in \mathbb{Z},\ q \neq 0\}$. Unless a different domain is stated, sequence indices run through $\mathbb{N}$ and all terms and finite limits are real. The empty sum is 0, and $0! = 1$.

Coverage. Section 7: Exercises 7.1–7.5. Section 8: Exercises 8.1–8.10. Section 9: Exercises 9.1–9.18. Section 10: Exercises 10.1–10.12.

Contents

7 Limits of sequences **4**

7.1 Exercise 7.1: The first five terms of four sequences 4

7.2 Exercise 7.2: Convergence of the four preceding sequences 6

7.3 Exercise 7.3: A catalogue of twenty sequences and their limits 7

7.4 Exercise 7.4: Rational and irrational terms can have different kinds of limits 17

7.5 Exercise 7.5: Three differences of large square roots 19

8 A discussion about proofs **22**

8.1 Exercise 8.1: Four limits proved directly from the definition 22

8.2 Exercise 8.2: Determine five limits and justify each one formally 25

8.3 Exercise 8.3: Taking square roots when the limit is zero 28

8.4 Exercise 8.4: A sequence tending to zero times a bounded sequence 29

8.5 Exercise 8.5: The squeeze lemma and an absolute-value comparison 30
8.6 Exercise 8.6: Absolute values at zero and an alternating counterexample 31
8.7 Exercise 8.7: Three divergent sequences 33
8.8 Exercise 8.8: Three radical limits with explicit error bounds 34
8.9 Exercise 8.9: Eventual bounds pass to a convergent limit 37
8.10 Exercise 8.10: A strict bound for the limit becomes an eventual strict bound 39

9 Limit theorems for sequences 41
9.1 Exercise 9.1: Three quotients evaluated by the finite limit laws 41
9.2 Exercise 9.2: Limits obtained from two given convergent sequences 43
9.3 Exercise 9.3: A polynomial expression in convergent sequences 43
9.4 Exercise 9.4: A nested-square-root recursion 44
9.5 Exercise 9.5: A positive rational recursion 46
9.6 Exercise 9.6: A necessary finite-limit equation and an actual infinite limit 47
9.7 Exercise 9.7: Finish the estimate for the root of the index 48
9.8 Exercise 9.8: Finite and infinite possibilities for five limits 49
9.9 Exercise 9.9: Order preserved by existing extended limits 51
9.10 Exercise 9.10: Scaling and reversing an infinite limit 52
9.11 Exercise 9.11: Adding a sequence that is bounded below 53
9.12 Exercise 9.12: The ratio test for the size of a sequence 54
9.13 Exercise 9.13: All real bases in a geometric sequence 56
9.14 Exercise 9.14: Geometric growth compared with a real power 57
9.15 Exercise 9.15: Every fixed geometric sequence is dominated by a factorial 58
9.16 Exercise 9.16: Three applications of the infinite-limit results 59
9.17 Exercise 9.17: A definition-only proof for the sequence of squares 60
9.18 Exercise 9.18: Finite geometric sums and their limits 61

10 Monotone sequences and Cauchy sequences 63
10.1 Exercise 10.1: Classifying six sequences 63
10.2 Exercise 10.2: Convergence of a bounded decreasing sequence 66
10.3 Exercise 10.3: A finite bound for decimal approximations 67
10.4 Exercise 10.4: Why the monotone and Cauchy theorems require real completeness . 68
10.5 Exercise 10.5: An unbounded decreasing sequence tends to negative infinity 70
10.6 Exercise 10.6: Small successive differences: one sufficient rate and one insufficient rate 70

10.7 Exercise 10.7: Approaching an unattained supremum by points of the set 72
10.8 Exercise 10.8: Averages of an increasing positive sequence 73
10.9 Exercise 10.9: A decreasing recurrence with a squared term 74
10.10 Exercise 10.10: A linear recurrence approaching one half 76
10.11 Exercise 10.11: An infinite-product recurrence 77
10.12 Exercise 10.12: A product recurrence with an exact formula 80

7 Limits of sequences

A sequence assigns a real number s_n to each index n. Unless the exercise states otherwise, the indices here are $n = 1, 2, 3, \ldots$. The order of the terms matters, even when their values repeat.

We use Ross's Definition 7.1: $s_n \to s$ means that, for every $\varepsilon > 0$, there is a number N such that

$$n > N \quad \Longrightarrow \quad |s_n - s| < \varepsilon \qquad (n \in \mathbb{N}).$$

The choice of N may depend on ε but cannot depend on the later index n. A real threshold N is permitted; the Archimedean property ensures that there are integer indices past it. Convergence here means convergence to a finite real number. Divergence means that no such real limit exists.

Although several exercises in this section ask only for answers, we include the algebra and estimates that explain those answers. The proofs use elementary inequalities and the limit definition, rather than the limit laws introduced in Section 9.

7.1 Exercise 7.1: The first five terms of four sequences

What we must show. Compute the terms with indices $n = 1, 2, 3, 4, 5$ for

$$\text{(a) } s_n = \frac{1}{3n+1}, \qquad \text{(b) } b_n = \frac{3n+1}{4n-1}, \qquad \text{(c) } c_n = \frac{n}{3^n}, \qquad \text{(d) } d_n = \sin\left(\frac{n\pi}{4}\right).$$

Plan. Substitute the index into every occurrence of n, evaluate the numerator and denominator separately, and simplify where possible. Indices start at 1. In part (a), the denominator $3n+1$ means 3 times n, followed by addition of 1.

1. Part (a): A reciprocal with a linear denominator

For $n \in \mathbb{N}$, we have $3n + 1 \geq 4 > 0$, so all the indicated fractions are defined. The five substitutions give

$$\begin{aligned}
s_1 &= \frac{1}{3 \cdot 1 + 1} = \frac{1}{3+1} = \frac{1}{4}, \\
s_2 &= \frac{1}{3 \cdot 2 + 1} = \frac{1}{6+1} = \frac{1}{7}, \\
s_3 &= \frac{1}{3 \cdot 3 + 1} = \frac{1}{9+1} = \frac{1}{10}, \\
s_4 &= \frac{1}{3 \cdot 4 + 1} = \frac{1}{12+1} = \frac{1}{13}, \\
s_5 &= \frac{1}{3 \cdot 5 + 1} = \frac{1}{15+1} = \frac{1}{16}.
\end{aligned}$$

2. Part (b): A quotient of two linear expressions

Here $4n - 1 \geq 3 > 0$ for every $n \geq 1$, so there is again no zero denominator. Compute both parts of each fraction:

$$b_1 = \frac{3 \cdot 1 + 1}{4 \cdot 1 - 1} = \frac{3+1}{4-1} = \frac{4}{3},$$

$$b_2 = \frac{3\cdot 2+1}{4\cdot 2-1} = \frac{6+1}{8-1} = \frac{7}{7} = 1,$$
$$b_3 = \frac{3\cdot 3+1}{4\cdot 3-1} = \frac{9+1}{12-1} = \frac{10}{11},$$
$$b_4 = \frac{3\cdot 4+1}{4\cdot 4-1} = \frac{12+1}{16-1} = \frac{13}{15},$$
$$b_5 = \frac{3\cdot 5+1}{4\cdot 5-1} = \frac{15+1}{20-1} = \frac{16}{19}.$$

3. Part (c): A linear numerator and an exponential denominator

The denominator is the positive integer power 3^n. In particular, $3^1 = 3$, $3^2 = 9$, $3^3 = 27$, $3^4 = 81$, and $3^5 = 243$. Substitution yields

$$c_1 = \frac{1}{3^1} = \frac{1}{3},$$
$$c_2 = \frac{2}{3^2} = \frac{2}{9},$$
$$c_3 = \frac{3}{3^3} = \frac{3}{27} = \frac{1}{9},$$
$$c_4 = \frac{4}{3^4} = \frac{4}{81},$$
$$c_5 = \frac{5}{3^5} = \frac{5}{243}.$$

The cancellation in the third term divides numerator and denominator by 3; it does not change the value of that term.

4. Part (d): Sine at multiples of an eighth-turn angle

On the unit circle at angle $\pi/4$, the two coordinates are equal and positive. If each is t, then $t^2+t^2 = 1$, so $t^2 = 1/2$ and $t = \sqrt{2}/2$. Thus $\sin(\pi/4) = \sqrt{2}/2$. The unit-circle values at $\pi/2$ and π are 1 and 0. Using these values and the identities $\sin(\pi-\theta) = \sin\theta$ and $\sin(\pi+\theta) = -\sin\theta$, we obtain

$$d_1 = \sin\left(\frac{1\pi}{4}\right) = \sin\left(\frac{\pi}{4}\right) = \frac{\sqrt{2}}{2},$$
$$d_2 = \sin\left(\frac{2\pi}{4}\right) = \sin\left(\frac{\pi}{2}\right) = 1,$$
$$d_3 = \sin\left(\frac{3\pi}{4}\right) = \sin\left(\pi - \frac{\pi}{4}\right) = \frac{\sqrt{2}}{2},$$
$$d_4 = \sin\left(\frac{4\pi}{4}\right) = \sin\pi = 0,$$
$$d_5 = \sin\left(\frac{5\pi}{4}\right) = \sin\left(\pi + \frac{\pi}{4}\right) = -\frac{\sqrt{2}}{2}.$$

Conclusion. The requested five-term lists are

Part	1	2	3	4	5
(a)	$\frac{1}{4}$	$\frac{1}{7}$	$\frac{1}{10}$	$\frac{1}{13}$	$\frac{1}{16}$
(b)	$\frac{4}{3}$	1	$\frac{10}{11}$	$\frac{13}{15}$	$\frac{16}{19}$
(c)	$\frac{1}{3}$	$\frac{2}{9}$	$\frac{1}{9}$	$\frac{4}{81}$	$\frac{5}{243}$
(d)	$\frac{\sqrt{2}}{2}$	1	$\frac{\sqrt{2}}{2}$	0	$-\frac{\sqrt{2}}{2}$

7.2 Exercise 7.2: Convergence of the four preceding sequences

What we must show. Decide which sequences in Exercise 7.1 converge and give each finite limit when it exists.

Plan. Use exact error estimates for the first three sequences. For the sine sequence, find values 1 and -1 at arbitrarily large indices, then compare them with any proposed common limit. The arguments use the limit definition directly: $u_n \to L$ means that for every $\varepsilon > 0$ there is a threshold N such that $n > N$ implies $|u_n - L| < \varepsilon$.

1. Part (a): The limit of $1/(3n+1)$

For every positive integer n,

$$\left|\frac{1}{3n+1} - 0\right| = \frac{1}{3n+1} < \frac{1}{3n}.$$

The reciprocal inequality follows because $3n+1 > 3n > 0$. Let $\varepsilon > 0$. By the Archimedean property, choose an integer $N > 1/(3\varepsilon)$. If $n > N$, then $3n > 1/\varepsilon$ and

$$\left|\frac{1}{3n+1} - 0\right| < \frac{1}{3n} < \varepsilon.$$

This verifies the definition and proves $s_n \to 0$.

2. Part (b): The limit of $(3n+1)/(4n-1)$

The ratio of the leading coefficients suggests $3/4$. Compute the error over a common denominator to verify it:

$$\begin{aligned} b_n - \frac{3}{4} &= \frac{3n+1}{4n-1} - \frac{3}{4} \\ &= \frac{4(3n+1) - 3(4n-1)}{4(4n-1)} \\ &= \frac{12n + 4 - 12n + 3}{4(4n-1)} \\ &= \frac{7}{4(4n-1)}. \end{aligned}$$

The denominator is positive for $n \geq 1$, so this is also the absolute error. Moreover, $4n - 1 - 3n = n - 1 \geq 0$, hence $4n - 1 \geq 3n$. Therefore

$$\left|b_n - \frac{3}{4}\right| = \frac{7}{4(4n-1)} \leq \frac{7}{12n}.$$

Given $\varepsilon > 0$, choose an integer $N > \max\{1, 7/(12\varepsilon)\}$. For $n > N$ we get

$$\left|b_n - \frac{3}{4}\right| \leq \frac{7}{12n} < \varepsilon.$$

Thus $b_n \to 3/4$. No quotient limit law is needed for this conclusion.

3. Part (c): Bound the exponential denominator by a square

We first prove $3^n \geq n^2$ for every integer $n \geq 2$. For $n = 2$, this reads $9 \geq 4$. Suppose it holds for an integer $n \geq 2$. Then

$$3^{n+1} = 3 \cdot 3^n \geq 3n^2.$$

Expanding the next square gives

$$3n^2 - (n+1)^2 = 3n^2 - (n^2 + 2n + 1) = 2n(n-1) - 1 \geq 3 > 0,$$

where $n \geq 2$ gives $2n(n-1) \geq 4$. Hence $3n^2 \geq (n+1)^2$, so $3^{n+1} \geq (n+1)^2$. Mathematical induction proves the claimed bound.

Because all denominators are positive, the bound implies, for $n \geq 2$,

$$0 < \frac{n}{3^n} \leq \frac{n}{n^2} = \frac{1}{n}.$$

Let $\varepsilon > 0$ and choose an integer $N > \max\{2, 1/\varepsilon\}$. Then $n > N$ gives

$$\left| \frac{n}{3^n} - 0 \right| = \frac{n}{3^n} \leq \frac{1}{n} < \varepsilon.$$

It follows directly that $c_n \to 0$.

4. Part (d): Values separated by 2 persist arbitrarily far out

For every nonnegative integer k, periodicity of sine gives

$$d_{8k+2} = \sin\left(\frac{(8k+2)\pi}{4}\right) = \sin\left(2k\pi + \frac{\pi}{2}\right) = 1,$$
$$d_{8k+6} = \sin\left(\frac{(8k+6)\pi}{4}\right) = \sin\left(2k\pi + \frac{3\pi}{2}\right) = -1.$$

Both families of indices can be made larger than any given real threshold N: choose an integer $k > \max\{0, N/8\}$. Then $8k + 2 > N$ and $8k + 6 > N$.

Suppose, for a contradiction, that $d_n \to L \in \mathbb{R}$. Use $\varepsilon = 1/2$ in the definition. There is a threshold N such that $n > N$ implies $|d_n - L| < 1/2$. Choose k so that both indices above exceed this N. Then

$$|1 - L| < \frac{1}{2}, \qquad |-1 - L| < \frac{1}{2}.$$

The triangle inequality, together with $|L + 1| = |-1 - L|$, would imply

$$2 = |1 - (-1)| = |(1 - L) + (L + 1)| \leq |1 - L| + |L + 1| < \frac{1}{2} + \frac{1}{2} = 1.$$

This contradicts $2 > 1$. Therefore the sine sequence has no real limit.

Conclusion. Parts (a), (b), and (c) converge to 0, 3/4, and 0, respectively. Part (d) diverges.

7.3 Exercise 7.3: A catalogue of twenty sequences and their limits

What we must show. Determine convergence or divergence for every sequence in parts (a)–(t), and give the real limit whenever one exists. Although the exercise does not require proofs, we include a justification of every answer. The arguments use the definition of convergence directly, rather than the limit laws developed later in Section 9.

Plan. For a convergent sequence, subtract the proposed limit and bound the resulting error by a reciprocal that becomes arbitrarily small. For a divergent sequence, find either arbitrarily late values that are arbitrarily large, or two arbitrarily late values separated by a fixed positive distance.

Elementary facts used in the individual parts

Fact 1: Reciprocal error estimates imply convergence.

The definition says that $u_n \to L$ if, for every $\varepsilon > 0$, there is a threshold N such that

$$n > N \quad \Longrightarrow \quad |u_n - L| < \varepsilon.$$

Suppose we prove $|u_n - L| \le C/n$ for every $n \ge n_0$, where $C > 0$ is fixed. Given $\varepsilon > 0$, choose a positive integer $N > \max\{n_0, C/\varepsilon\}$, using the Archimedean property. Then $n > N$ implies both $n \ge n_0$ and $n > C/\varepsilon$. Multiplying the latter inequality by $\varepsilon > 0$ and dividing by $n > 0$ gives

$$|u_n - L| \le \frac{C}{n} < \varepsilon.$$

This proves convergence from the definition. In particular, $1/n \to 0$. If an error is identically 0, it is less than every $\varepsilon > 0$ without any additional estimate.

We will also use a shifted version. If, for all sufficiently large n, $|u_n - L| \le C/(n - r)$, with fixed real r and $n > r$, choose N larger than the starting index and $r + C/\varepsilon$. Then $n > N$ gives $n - r > C/\varepsilon$, and the same calculation gives $|u_n - L| < \varepsilon$. This proves the shifted version without any new limit theorem.

Only the tail is used in either proof. Thus a finite number of unusual initial values does not affect the conclusion: increase N to exceed all their indices, and every $n > N$ still satisfies the required estimate. This observation will matter when an initial denominator is negative but never zero.

Fact 2: Powers of two dominate a simple linear expression.

For every integer $m \ge 0$,

$$2^m \ge m + 1.$$

At $m = 0$, both sides equal 1. If the inequality holds at m, then multiplication by $2 > 0$ gives

$$2^{m+1} = 2 \cdot 2^m \ge 2(m + 1) = 2m + 2 \ge m + 2,$$

where the last difference is $(2m + 2) - (m + 2) = m \ge 0$. Induction proves the assertion. Taking reciprocals of positive quantities consequently gives

$$0 < 2^{-m} \le \frac{1}{m + 1}.$$

Fact 3: Separated arbitrarily late values rule out a real limit.

Suppose some fixed $D > 0$ has this property: for every threshold N, there are indices $p, q > N$ with $|u_p - u_q| \ge D$. Assume, to obtain a contradiction, that $u_n \to L \in \mathbb{R}$. Apply the definition with $\varepsilon = D/3$. It supplies N such that both late terms obey $|u_p - L| < D/3$ and $|u_q - L| < D/3$. The triangle inequality would then give

$$|u_p - u_q| = |(u_p - L) + (L - u_q)| \le |u_p - L| + |L - u_q| < \frac{D}{3} + \frac{D}{3} = \frac{2D}{3} < D,$$

contradicting the stipulated separation. Thus such a sequence does not converge to a real number. This argument does not assume any theorem about subsequences.

Fact 4: Arbitrarily large late values also rule out a real limit.

Suppose that for every $B \in \mathbb{R}$ and every threshold N, some $n > N$ satisfies $u_n > B$. If $u_n \to L \in \mathbb{R}$, the definition with $\varepsilon = 1$ would supply N for which $|u_n - L| < 1$ whenever $n > N$. Choose a late value with $u_n > L + 1$. It satisfies $u_n - L > 1$, and therefore $|u_n - L| = u_n - L > 1$, a contradiction. This is a direct nonconvergence argument; it does not invoke a later boundedness theorem for convergent sequences.

Solutions of parts (a)–(t)

(a) The sequence $a_n = n/(n+1)$.

For every $n \in \mathbb{N}$, $n + 1 \geq 2 > 0$, so every term is defined. To test the candidate 1, subtract it using the common denominator $n + 1$:

$$a_n - 1 = \frac{n}{n+1} - \frac{n+1}{n+1} = \frac{n - (n+1)}{n+1} = -\frac{1}{n+1}.$$

It follows that

$$|a_n - 1| = \frac{1}{n+1} \leq \frac{1}{n}.$$

The inequality uses $n + 1 \geq n > 0$ and the reversal of order for positive reciprocals. Fact 1 with $C = 1$ now proves the limit.

Conclusion. Part (a) converges, and $\lim_{n\to\infty} \dfrac{n}{n+1} = 1$.

(b) The sequence $b_n = (n^2 + 3)/(n^2 - 3)$.

The denominator is nonzero at every positive integer. At $n = 1$ it is $1 - 3 = -2$, while $n \geq 2$ implies $n^2 \geq 4$ and $n^2 - 3 \geq 1 > 0$. Thus there is no undefined term, including in the initial part of the sequence.

Subtracting the candidate 1 gives

$$b_n - 1 = \frac{n^2 + 3 - (n^2 - 3)}{n^2 - 3} = \frac{n^2 + 3 - n^2 + 3}{n^2 - 3} = \frac{6}{n^2 - 3}.$$

For $n \geq 3$, we have $n^2 \geq 9 \geq 6$, so $3 \leq n^2/2$. Subtracting 3 from n^2 therefore gives

$$n^2 - 3 \geq n^2 - \frac{n^2}{2} = \frac{n^2}{2} > 0.$$

Consequently

$$|b_n - 1| = \frac{6}{n^2 - 3} \leq \frac{6}{n^2/2} = \frac{12}{n^2} \leq \frac{12}{n}.$$

The last inequality follows from $n^2 \geq n > 0$ for $n \geq 1$. Fact 1 with $C = 12$ and starting index 3 proves convergence. The negative first denominator does not enter the tail estimate.

Conclusion. Part (b) converges to 1.

(c) The sequence $c_n = 2^{-n}$.

The expression means the positive reciprocal $1/2^n$. Fact 2 gives

$$|c_n - 0| = \frac{1}{2^n} \leq \frac{1}{n+1} \leq \frac{1}{n} \qquad (n \geq 1).$$

The error is thus bounded by $1/n$. Applying Fact 1 proves the limit directly.

Conclusion. Part (c) converges to 0.

(d) The sequence $t_n = 1 + 2/n$.

Since $n > 0$, every term is defined, and subtracting 1 gives

$$|t_n - 1| = \left|1 + \frac{2}{n} - 1\right| = \left|\frac{2}{n}\right| = \frac{2}{n}.$$

For each $\varepsilon > 0$, choosing $N > 2/\varepsilon$ makes this error smaller than ε whenever $n > N$. This is Fact 1 with $C = 2$.

Conclusion. Part (d) converges to 1.

(e) The sequence $x_n = 73 + (-1)^n$.

If n is even, $(-1)^n = 1$ and $x_n = 73 + 1 = 74$. If n is odd, $(-1)^n = -1$ and $x_n = 73 - 1 = 72$. These values do not merely occur at the start: for any threshold N, choose a positive integer $m > N$. Then $2m > N$ and $2m + 1 > N$, and

$$x_{2m} = 74, \qquad x_{2m+1} = 72, \qquad |x_{2m} - x_{2m+1}| = |74 - 72| = 2.$$

Thus Fact 3 applies with $D = 2$. No real number can be approached by every term, because every tail still contains these two values at distance 2.

Conclusion. Part (e) does not converge.

(f) The sequence $s_n = 2^{1/n}$.

Write $r_n = 2^{1/n}$ for the positive real nth root of 2; thus $r_n > 0$ and $r_n^n = 2$. We prove the concrete estimate

$$0 \le r_n - 1 \le \frac{1}{n}.$$

It will imply convergence by Fact 1 without any theorem about limits of roots or logarithms.

First prove the auxiliary Bernoulli inequality. For every $x \ge 0$ and every positive integer m,

$$(1 + x)^m \ge 1 + mx.$$

Fix $x \ge 0$. At $m = 1$ the two sides are equal. If the inequality holds at m, multiplying by the positive number $1 + x$ gives

$$\begin{aligned}(1 + x)^{m+1} &\ge (1 + mx)(1 + x)\\ &= 1 + x + mx + mx^2\\ &= 1 + (m + 1)x + mx^2\\ &\ge 1 + (m + 1)x,\end{aligned}$$

because $mx^2 \ge 0$. This is the required step at $m + 1$, so induction proves the inequality for the fixed x, and hence for every $x \ge 0$.

Apply it with $x = 1/n$ and $m = n$. We obtain

$$\left(1 + \frac{1}{n}\right)^n \ge 1 + n\frac{1}{n} = 2.$$

For positive real numbers, raising to a positive integer power preserves strict order. For example, if $u > v > 0$, then $u/v > 1$, so the product of n factors u/v is greater than 1; multiplication by $v^n > 0$ gives $u^n > v^n$. If $r_n > 1 + 1/n$, this order property would give

$$2 = r_n^n > \left(1 + \frac{1}{n}\right)^n \geq 2,$$

a contradiction. Hence $r_n \leq 1 + 1/n$. Also $r_n \geq 1$: if $0 < r_n < 1$, multiplying n factors each less than 1 would give $r_n^n < 1$, contradicting $r_n^n = 2$.

Thus $1 \leq r_n \leq 1 + 1/n$, and subtraction of 1 yields

$$|r_n - 1| = r_n - 1 \leq \frac{1}{n}.$$

Fact 1 proves the asserted limit.

Conclusion. Part (f) converges to 1.

(g) The sequence $y_n = n!$.

For $n \geq 1$, the factorial can be written

$$n! = n\,(1 \cdot 2 \cdots (n-1)).$$

At $n = 1$, the parenthesized product is empty and equals 1. For $n \geq 2$, every factor in that product is at least 1, so its product is at least 1. In all cases, therefore,

$$n! \geq n.$$

Now let a real number B and a threshold N be arbitrary. Choose $n \in \mathbb{N}$ with $n > \max\{N, B, 1\}$. Then $n > N$ and $y_n = n! \geq n > B$. Thus arbitrarily late values exceed every prescribed real size. Fact 4 rules out convergence to any real number. This argument establishes the needed nonconvergence without citing a boundedness theorem.

Conclusion. Part (g) does not converge to a real limit; its values grow beyond every real bound.

(h) The sequence $d_n = (-1)^n n$.

At an even index $2m$ the value is $2m$, while at an odd index $2m + 1$ the value is $-(2m + 1)$. Given any threshold N, choose a positive integer $m > N$. Both indices $2m$ and $2m + 1$ are then greater than N, and

$$\begin{aligned} |d_{2m} - d_{2m+1}| &= |2m - (-(2m+1))| \\ &= |4m + 1| \\ &= 4m + 1 \geq 5 > 1. \end{aligned}$$

Here $m \geq 1$, so $4m + 1$ is positive, justifying the removal of the absolute value. Every tail has two values separated by at least 1. Fact 3 with $D = 1$ rules out a real limit.

Conclusion. Part (h) does not converge. The signs alternate while the magnitudes increase.

(i) The sequence $u_n = (-1)^n/n$.

Although its sign alternates, its distance from 0 is

$$|u_n - 0| = \frac{|(-1)^n|}{|n|} = \frac{1}{n},$$

because $|(-1)^n| = 1$ and $n > 0$. Fact 1 applies with $C = 1$. In contrast with (h), the alternating values now have small magnitude for every sufficiently large index.

Conclusion. Part (i) converges to 0.

(j) The sequence $u_n = (7n^3 + 8n)/(2n^3 - 3)$.

At $n = 1$, the denominator is $2 - 3 = -1$, so it is nonzero. For $n \geq 2$, $n^3 \geq 8$, giving $2n^3 - 3 \geq 16 - 3 = 13 > 0$. Thus the sequence is defined at every positive integer.

We test the candidate $7/2$ by exact subtraction:

$$\begin{aligned} u_n - \frac{7}{2} &= \frac{2(7n^3 + 8n) - 7(2n^3 - 3)}{2(2n^3 - 3)} \\ &= \frac{14n^3 + 16n - 14n^3 + 21}{2(2n^3 - 3)} \\ &= \frac{16n + 21}{2(2n^3 - 3)}. \end{aligned}$$

For $n \geq 2$, we have $n^3 \geq 8 > 3$, so

$$2n^3 - 3 \geq n^3 > 0.$$

Also $n \geq 1$ implies $21 \leq 21n$ and hence $16n + 21 \leq 37n$. Using these positive denominator bounds gives

$$\left| u_n - \frac{7}{2} \right| = \frac{16n + 21}{2(2n^3 - 3)} \leq \frac{37n}{2n^3} = \frac{37}{2n^2} \leq \frac{37}{2n}.$$

The last step again uses $n^2 \geq n > 0$. Fact 1, with $C = 37/2$ and starting index 2, proves convergence.

Conclusion. Part (j) converges to $7/2$.

(k) The sequence $u_n = (9n^2 - 18)/(6n + 18)$.

The denominator $6n + 18$ is positive for every $n \geq 1$. We establish a lower estimate that becomes arbitrarily large. For $n \geq 3$, $n^2 \geq 9$, so $3n^2 \geq 27 > 18$ and

$$9n^2 - 18 \geq 9n^2 - 3n^2 = 6n^2 > 0.$$

Also $n \geq 3$ implies $18 \leq 6n$, giving

$$0 < 6n + 18 \leq 12n.$$

First lower the numerator while keeping the positive denominator fixed, then increase that denominator:

$$u_n = \frac{9n^2 - 18}{6n + 18} \geq \frac{6n^2}{6n + 18} \geq \frac{6n^2}{12n} = \frac{n}{2}.$$

The second comparison follows from order reversal for positive reciprocals, followed by multiplication by $6n^2 > 0$.

Let $B \in \mathbb{R}$ and a threshold N be arbitrary. Choose a positive integer $n > \max\{N, 3, 2B\}$. The estimate applies and gives $u_n \geq n/2 > B$, with $n > N$. Fact 4 therefore rules out convergence to every real candidate.

Conclusion. Part (k) does not converge to a real limit; its values grow beyond every real bound.

(l) The sequence $u_n = \sin(n\pi/2)$.

Every positive integer has the form $n = 4q + r$, with $q \geq 0$ an integer and $r \in \{1, 2, 3, 4\}$. Since $n\pi/2 = 2q\pi + r\pi/2$, the period 2π of sine gives the repeating values

r	1	2	3	4
$\sin(r\pi/2)$	1	0	-1	0

The repetition alone suggests nonconvergence; we now check the arbitrarily late separation required by Fact 3. Given a threshold N, choose $m \in \mathbb{N}$ with $m > N$. Then $p = 4m+1 > N$ and $q = 4m+3 > N$, and periodicity gives

$$u_p = \sin(2m\pi + \pi/2) = 1, \qquad u_q = \sin(2m\pi + 3\pi/2) = -1.$$

Their distance is $|1 - (-1)| = 2$. Fact 3 applies with $D = 2$ and proves nonconvergence.

Conclusion. Part (l) does not converge.

(m) The sequence $u_n = \sin(n\pi)$.

For every integer n, the angle $n\pi$ lies at an integer multiple of a half-turn, so its sine is 0. Explicitly, if $n = 2m$ is even, then $\sin(n\pi) = \sin(2m\pi) = 0$ by periodicity. If $n = 2m + 1$ is odd, then

$$\sin(n\pi) = \sin(2m\pi + \pi) = \sin \pi = 0.$$

These two cases cover every positive integer. Thus $u_n = 0$ for every index, and for every $\varepsilon > 0$,

$$|u_n - 0| = 0 < \varepsilon$$

holds at every index. This verifies convergence directly from the definition.

Conclusion. Part (m) converges to 0.

(n) The sequence $u_n = \sin(2n\pi/3)$.

Write $n = 3q + r$, with $q \geq 0$ and $r \in \{1, 2, 3\}$. Then $2n\pi/3 = 2q\pi + 2r\pi/3$, giving the period table

r	1	2	3
$\sin(2r\pi/3)$	$\sqrt{3}/2$	$-\sqrt{3}/2$	0

For any threshold N, choose a positive integer $m > N$. The indices $p = 3m + 1$ and $q = 3m + 2$ both exceed N, and

$$u_p = \frac{\sqrt{3}}{2}, \qquad u_q = -\frac{\sqrt{3}}{2}.$$

Their distance is

$$|u_p - u_q| = \left| \frac{\sqrt{3}}{2} + \frac{\sqrt{3}}{2} \right| = \sqrt{3} > 0.$$

Consequently Fact 3 with $D = \sqrt{3}$ excludes every real limit.

Conclusion. Part (n) does not converge.

(o) The sequence $u_n = (\sin n)/n$.

Here the argument is n radians, with no factor of π. We do not need to classify the behavior of $\sin n$ itself. The elementary trigonometric inequality $|\sin x| \leq 1$ for every real x gives

$$|u_n - 0| = \left| \frac{\sin n}{n} \right| = \frac{|\sin n|}{n} \leq \frac{1}{n},$$

where division by $n > 0$ preserves the inequality. Fact 1 with $C = 1$ proves the limit.

Conclusion. Part (o) converges to 0.

(p) The sequence $u_n = (2^{n+1} + 5)/(2^n - 7)$.

The denominator is $2^n - 7$. At $n = 1$ it is $2 - 7 = -5$, and at $n = 2$ it is $4 - 7 = -3$. For $n \geq 3$, $2^n \geq 8$ and therefore $2^n - 7 \geq 1 > 0$. These cases show that every term is defined. For example the first three terms are

$$u_1 = \frac{4+5}{2-7} = -\frac{9}{5}, \qquad u_2 = \frac{8+5}{4-7} = -\frac{13}{3}, \qquad u_3 = \frac{16+5}{8-7} = 21.$$

They do not by themselves determine the limiting behavior.

Subtract the candidate 2 exactly:

$$\begin{aligned} u_n - 2 &= \frac{2^{n+1} + 5 - 2(2^n - 7)}{2^n - 7} \\ &= \frac{2 \cdot 2^n + 5 - 2 \cdot 2^n + 14}{2^n - 7} \\ &= \frac{19}{2^n - 7}. \end{aligned}$$

For $n \geq 4$, $2^n \geq 16 \geq 14$, so $7 \leq 2^n/2$. It follows that

$$2^n - 7 \geq 2^n - \frac{2^n}{2} = \frac{2^n}{2} > 0.$$

Using this bound and Fact 2 yields

$$\begin{aligned} |u_n - 2| &= \frac{19}{2^n - 7} \\ &\leq \frac{19}{2^n/2} = \frac{38}{2^n} \leq \frac{38}{n+1} \leq \frac{38}{n}. \end{aligned}$$

Fact 1 with $C = 38$ and starting index 4 proves convergence. It also explains why the negative initial terms and the relatively large value $u_3 = 21$ do not change the tail conclusion.

Conclusion. Part (p) converges to 2.

(q) The sequence $a_n = 3^n/n!$.

Both 3^n and $n!$ are positive for every $n \in \mathbb{N}$, so $a_n > 0$ and every ratio below is defined. Compare successive terms using $(n+1)! = (n+1)n!$:

$$\begin{aligned} \frac{a_{n+1}}{a_n} &= \frac{3^{n+1}}{(n+1)!} \cdot \frac{n!}{3^n} \\ &= \frac{3 \cdot 3^n \cdot n!}{(n+1)n!\, 3^n} \\ &= \frac{3}{n+1}. \end{aligned}$$

For $n \geq 5$, $n + 1 \geq 6$, giving

$$0 < \frac{3}{n+1} \leq \frac{3}{6} = \frac{1}{2}, \qquad a_{n+1} \leq \frac{1}{2} a_n.$$

We will turn this ratio estimate into an explicit reciprocal bound; no theorem about ratios of sequences is being invoked.

The value at the starting index is

$$C = a_5 = \frac{3^5}{5!} = \frac{243}{120} = \frac{81}{40} > 0.$$

We claim, for every integer $m \geq 0$, that

$$a_{5+m} \leq C\,2^{-m}.$$

At $m = 0$, this is equality $a_5 = C$. If the estimate holds at m, the ratio bound applies at $n = 5+m \geq 5$, and gives

$$a_{5+(m+1)} = a_{6+m} \leq \frac{1}{2}a_{5+m} \leq \frac{1}{2}C\,2^{-m} = C\,2^{-(m+1)}.$$

This is the next estimate, so induction proves the claim. Putting $m = n - 5$ now yields, for every $n \geq 5$,

$$0 < a_n \leq C\,2^{-(n-5)}.$$

Fact 2, applied to $m = n - 5 \geq 0$, supplies

$$2^{n-5} \geq (n-5) + 1 = n - 4 > 0.$$

Taking reciprocals and multiplying by $C > 0$ therefore gives

$$|a_n - 0| = a_n \leq C\,2^{-(n-5)} \leq \frac{C}{n-4} = \frac{81/40}{n-4}.$$

For any $\varepsilon > 0$, choose an integer $N > \max\{5, 4 + C/\varepsilon\}$. If $n > N$, then $n - 4 > C/\varepsilon$, so the displayed error is less than ε. This is the shifted estimate of Fact 1, proving the limit directly.

Conclusion. Part (q) converges to 0.

(r) The sequence $u_n = (1 + 1/n)^2$.

Expand by multiplying its two factors, not by assuming a rule about limits of squares:

$$\begin{aligned} u_n &= \left(1 + \frac{1}{n}\right)\left(1 + \frac{1}{n}\right) \\ &= 1 + \frac{1}{n} + \frac{1}{n} + \frac{1}{n^2} \\ &= 1 + \frac{2}{n} + \frac{1}{n^2}. \end{aligned}$$

All terms after the leading 1 are positive. Since $n^2 \geq n > 0$, we have $1/n^2 \leq 1/n$, giving

$$|u_n - 1| = \frac{2}{n} + \frac{1}{n^2} \leq \frac{2}{n} + \frac{1}{n} = \frac{3}{n}.$$

Fact 1 with $C = 3$ proves convergence.

Conclusion. Part (r) converges to 1.

(s) The sequence $u_n = (4n^2 + 3)/(3n^2 - 2)$.

For all $n \geq 1$, $n^2 \geq 1$, so $3n^2 - 2 \geq 3 - 2 = 1 > 0$. Thus every term is defined and the denominator has a fixed positive sign. Subtracting $4/3$ with a common denominator gives

$$\begin{aligned} u_n - \frac{4}{3} &= \frac{3(4n^2+3) - 4(3n^2-2)}{3(3n^2-2)} \\ &= \frac{12n^2 + 9 - 12n^2 + 8}{3(3n^2-2)} \\ &= \frac{17}{3(3n^2-2)}. \end{aligned}$$

The estimate $n^2 \geq 1$ also yields

$$(3n^2 - 2) - n^2 = 2n^2 - 2 = 2(n^2 - 1) \geq 0,$$

so $3n^2 - 2 \geq n^2 > 0$. Therefore

$$\left| u_n - \frac{4}{3} \right| = \frac{17}{3(3n^2-2)} \leq \frac{17}{3n^2} \leq \frac{17}{3n}.$$

Fact 1 with $C = 17/3$ verifies the limit from the definition.

Conclusion. Part (s) converges to $4/3$.

(t) The sequence $u_n = (6n + 4)/(9n^2 + 7)$.

For $n \geq 1$, both $6n + 4$ and $9n^2 + 7$ are positive, so the absolute value of u_n is u_n itself. We have

$$9n^2 + 7 \geq 9n^2 > 0, \qquad 6n + 4 \leq 6n + 4n = 10n,$$

where $4 \leq 4n$ uses $n \geq 1$. Taking reciprocals in the denominator comparison and then applying the numerator comparison,

$$|u_n - 0| = \frac{6n+4}{9n^2+7} \leq \frac{6n+4}{9n^2} \leq \frac{10n}{9n^2} = \frac{10}{9n}.$$

Fact 1 with $C = 10/9$ proves convergence to the proposed value 0.

Conclusion. Part (t) converges to 0.

Complete answer list

In this list, "does not converge" means that there is no finite real limit. Every entry is justified in its corresponding part above. The growth cases (g) and (k), the persistent oscillations (e), (h), (l), and (n), and the shrinking oscillations (i) and (o) have been distinguished by direct estimates and the definition.

Part	Answer	Part	Answer
(a)	Converges to 1	(k)	Does not converge
(b)	Converges to 1	(l)	Does not converge
(c)	Converges to 0	(m)	Converges to 0
(d)	Converges to 1	(n)	Does not converge
(e)	Does not converge	(o)	Converges to 0
(f)	Converges to 1	(p)	Converges to 2
(g)	Does not converge	(q)	Converges to 0
(h)	Does not converge	(r)	Converges to 1
(i)	Converges to 0	(s)	Converges to 4/3
(j)	Converges to 7/2	(t)	Converges to 0

7.4 Exercise 7.4: Rational and irrational terms can have different kinds of limits

What we must show. Give (a) a sequence of irrational numbers converging to a rational number, and (b) a sequence of rational numbers converging to an irrational number.

Plan. For part (a), divide $\sqrt{2}$ by larger positive integers. For part (b), use the lower decimal truncations of $\sqrt{2}$. We verify both the nature of every term and convergence.

1. Recall why $\sqrt{2}$ is irrational

The irrationality of $\sqrt{2}$ was established in Section 2. Here is the short integer argument as well. If $\sqrt{2} = p/q$ in lowest terms, with integers p, q and $q > 0$, then $p^2 = 2q^2$, so p^2 is even. The square of an odd integer $2j + 1$ is $4j(j + 1) + 1$, which is odd. Thus p must be even; write $p = 2j$. Substitution gives

$$4j^2 = 2q^2, \qquad q^2 = 2j^2.$$

Now q^2 is even, so the same odd-square argument makes q even. The numerator and denominator would have the common factor 2, contrary to lowest terms. Hence $\sqrt{2} \notin \mathbb{Q}$.

2. Part (a): Irrational terms with rational limit

Define, for every $n \in \mathbb{N}$,

$$x_n = \frac{\sqrt{2}}{n}.$$

Since $n \geq 1$, division is legitimate. If any term x_n were rational, multiplication by the rational integer n would make $nx_n = \sqrt{2}$ rational. This contradicts Step 1. Thus every term x_n is irrational.

To prove convergence, let $\varepsilon > 0$ and choose an integer $N > \sqrt{2}/\varepsilon$ by the Archimedean property. For $n > N$,

$$|x_n - 0| = \frac{\sqrt{2}}{n} < \varepsilon.$$

Therefore $x_n \to 0$. The limit is rational because $0 = 0/1$.

3. Part (b): Define each decimal truncation using a finite maximum

For a real $t \geq 0$, the notation $\lfloor t \rfloor$ denotes the integer m satisfying

$$m \leq t < m + 1.$$

To justify existence, choose a positive integer $M > t$ using the Archimedean property. The finite set

$$F = \{j \in \mathbb{Z} : 0 \le j \le M \text{ and } j \le t\}$$

is nonempty because it contains 0. Let $m = \max F$, which exists for a nonempty finite set of integers. Then $m \le t$. If $m + 1 \le t$, the inequalities $0 \le m + 1 \le t < M$ would give $m + 1 \in F$, contradicting the maximality of m. Hence $t < m + 1$, as required.

This integer is unique. If two integers $m < j$ satisfied their respective inequalities, then $m+1 \le j \le t$ would contradict $t < m + 1$. Interchanging them handles the other possible order. Thus the floor notation is well-defined here.

Apply this construction to $t = 10^n\sqrt{2}$ and put

$$m_n = \left\lfloor 10^n\sqrt{2} \right\rfloor, \qquad r_n = \frac{m_n}{10^n} \quad (n \in \mathbb{N}).$$

Every m_n is an integer and 10^n is a positive integer. Therefore every r_n is rational by the definition of $\mathbb{Q}$. The defining floor inequalities give

$$m_n \le 10^n\sqrt{2} < m_n + 1.$$

Dividing by $10^n > 0$ preserves order and yields

$$r_n \le \sqrt{2} < r_n + \frac{1}{10^n}.$$

After subtracting r_n, this becomes the precise truncation error estimate

$$0 \le \sqrt{2} - r_n < \frac{1}{10^n}.$$

In fact the left inequality is strict, because a rational r_n cannot equal the irrational number $\sqrt{2}$.

4. Prove convergence of the rational truncations

We may compare the error with $1/n$. The inequality $10^n \ge n$ holds for all positive integers n: the base case is $10 \ge 1$; if it holds for n, then

$$10^{n+1} = 10 \cdot 10^n \ge 10n \ge n + 1,$$

because $10n - (n+1) = 9n - 1 \ge 8 > 0$ for $n \ge 1$. Induction proves the inequality. Taking reciprocals of positive quantities gives $1/10^n \le 1/n$.

Let $\varepsilon > 0$ and choose an integer $N > 1/\varepsilon$. For $n > N$, we obtain

$$\left|r_n - \sqrt{2}\right| = \sqrt{2} - r_n < \frac{1}{10^n} \le \frac{1}{n} < \varepsilon.$$

Thus $r_n \to \sqrt{2}$, and its limit is irrational by Step 1.

Conclusion. For (a), take $x_n = \sqrt{2}/n \to 0$, with every term irrational and the limit rational. For (b), take $r_n = \lfloor 10^n\sqrt{2} \rfloor / 10^n \to \sqrt{2}$, with every term rational and the limit irrational.

7.5 Exercise 7.5: Three differences of large square roots

What we must show. Find the limits of

$$\text{(a) } \sqrt{n^2+1}-n, \qquad \text{(b) } \sqrt{n^2+n}-n, \qquad \text{(c) } \sqrt{4n^2+n}-2n.$$

Plan. Multiply each difference by its conjugate divided by that same conjugate. The difference-of-squares identity removes the subtraction in the numerator. We then bound the error from the proposed limit directly, so no later limit laws are needed.

1. Part (a): Expand the product with the conjugate

Let $a_n = \sqrt{n^2+1}-n$. For $n \geq 1$, the square root is nonnegative and $n > 0$, so the conjugate $\sqrt{n^2+1}+n$ is positive. Dividing by it is valid. Using $(u-v)(u+v) = u^2 - v^2$ gives

$$\begin{aligned} a_n &= \frac{(\sqrt{n^2+1}-n)(\sqrt{n^2+1}+n)}{\sqrt{n^2+1}+n} \\ &= \frac{(\sqrt{n^2+1})^2 - n^2}{\sqrt{n^2+1}+n} \\ &= \frac{n^2+1-n^2}{\sqrt{n^2+1}+n} \\ &= \frac{1}{\sqrt{n^2+1}+n}. \end{aligned}$$

Since $n^2+1 > n^2$ and $n \geq 0$, the nonnegative square root $\sqrt{n^2+1}$ is greater than n. Indeed, the contrary inequality would give $n^2+1 \leq n^2$ upon squaring nonnegative quantities. Therefore

$$0 < a_n = \frac{1}{\sqrt{n^2+1}+n} \leq \frac{1}{2n}.$$

Given $\varepsilon > 0$, choose an integer $N > 1/(2\varepsilon)$. For $n > N$, this estimate gives

$$|a_n - 0| = a_n \leq \frac{1}{2n} < \varepsilon.$$

Thus the limit in part (a) is 0.

2. Part (b): Rationalize and divide by the positive index

Write $b_n = \sqrt{n^2+n}-n$. Its conjugate is positive for $n \geq 1$, and the same multiplication gives

$$\begin{aligned} b_n &= \frac{(\sqrt{n^2+n}-n)(\sqrt{n^2+n}+n)}{\sqrt{n^2+n}+n} \\ &= \frac{(n^2+n)-n^2}{\sqrt{n^2+n}+n} \\ &= \frac{n}{\sqrt{n^2+n}+n}. \end{aligned}$$

The radicand factors as $n^2+n = n^2(1+1/n)$. Because $n > 0$,

$$\sqrt{n^2+n} = n\sqrt{1+\frac{1}{n}}.$$

To check this identity, the right-hand side is nonnegative, and its square is $n^2(1+1/n) = n^2+n$; uniqueness of the nonnegative square root then gives the equality. Define $q_n = \sqrt{1+1/n}$. Substitution and cancellation of $n \neq 0$ yield

$$b_n = \frac{n}{nq_n + n} = \frac{n}{n(q_n+1)} = \frac{1}{q_n+1}.$$

Here $q_n > 1$, since $q_n \geq 0$ and $q_n^2 = 1 + 1/n > 1$; a nonnegative $q_n \leq 1$ would have square at most 1. Hence $q_n + 1 > 2$, so $0 < b_n < 1/2$.

3. Part (b): Compute the exact error from $1/2$

All the denominators below are positive. Since $b_n < 1/2$,

$$\begin{aligned}
\left| b_n - \frac{1}{2} \right| &= \frac{1}{2} - \frac{1}{q_n+1} \\
&= \frac{q_n + 1 - 2}{2(q_n+1)} \\
&= \frac{q_n - 1}{2(q_n+1)} \\
&= \frac{(q_n-1)(q_n+1)}{2(q_n+1)^2} \\
&= \frac{q_n^2 - 1}{2(q_n+1)^2} \\
&= \frac{(1+1/n) - 1}{2(q_n+1)^2} \\
&= \frac{1}{2n(q_n+1)^2} \leq \frac{1}{8n}.
\end{aligned}$$

The last inequality uses $(q_n+1)^2 \geq 2^2 = 4$ and $n > 0$. For any $\varepsilon > 0$, choose an integer $N > 1/(8\varepsilon)$. Then $n > N$ implies $|b_n - 1/2| \leq 1/(8n) < \varepsilon$. Therefore the limit in part (b) is $1/2$.

4. Part (c): Rationalize and factor the square root

Put $c_n = \sqrt{4n^2+n} - 2n$. Since $n \geq 1$, its conjugate $\sqrt{4n^2+n} + 2n$ is positive. Consequently

$$\begin{aligned}
c_n &= \frac{(\sqrt{4n^2+n} - 2n)(\sqrt{4n^2+n} + 2n)}{\sqrt{4n^2+n} + 2n} \\
&= \frac{(4n^2+n) - (2n)^2}{\sqrt{4n^2+n} + 2n} \\
&= \frac{4n^2 + n - 4n^2}{\sqrt{4n^2+n} + 2n} \\
&= \frac{n}{\sqrt{4n^2+n} + 2n}.
\end{aligned}$$

Now $4n^2 + n = n^2(4 + 1/n)$. The positive-index square-root identity gives

$$\sqrt{4n^2+n} = n\sqrt{4 + \frac{1}{n}},$$

because both sides are nonnegative and the square of the right-hand side is $n^2(4 + 1/n) = 4n^2 + n$. Let $p_n = \sqrt{4 + 1/n}$. Canceling the nonzero factor n gives

$$c_n = \frac{n}{np_n + 2n} = \frac{n}{n(p_n + 2)} = \frac{1}{p_n + 2}.$$

Since $p_n \geq 0$ and $p_n^2 = 4 + 1/n > 4$, we have $p_n > 2$. Thus $p_n + 2 > 4$ and $0 < c_n < 1/4$.

5. Part (c): Compute the exact error from $1/4$

Using $c_n < 1/4$ and then multiplying numerator and denominator by the positive number $p_n + 2$, we obtain

$$\begin{aligned}
\left| c_n - \frac{1}{4} \right| &= \frac{1}{4} - \frac{1}{p_n + 2} \\
&= \frac{p_n + 2 - 4}{4(p_n + 2)} \\
&= \frac{p_n - 2}{4(p_n + 2)} \\
&= \frac{(p_n - 2)(p_n + 2)}{4(p_n + 2)^2} \\
&= \frac{p_n^2 - 4}{4(p_n + 2)^2} \\
&= \frac{(4 + 1/n) - 4}{4(p_n + 2)^2} \\
&= \frac{1}{4n(p_n + 2)^2} \leq \frac{1}{64n}.
\end{aligned}$$

Here $(p_n + 2)^2 \geq 4^2 = 16$, so the denominator is at least $4n \cdot 16 = 64n$. For any $\varepsilon > 0$, choose an integer $N > 1/(64\varepsilon)$. Then $n > N$ gives $|c_n - 1/4| \leq 1/(64n) < \varepsilon$. This proves that the limit in part (c) is $1/4$.

Conclusion. The limits are (a) 0, (b) 1/2, and (c) 1/4. The displayed conjugate identities and error estimates justify all three values directly.

8 A discussion about proofs

Every limit proof in this section starts with an arbitrary $\varepsilon > 0$, chooses an appropriate threshold, and verifies the required inequality for every integer index beyond that threshold. An exploratory rearrangement helps discover N; the formal argument then runs forward from $n > N$. Divergence proofs instead show that the definition cannot hold for any proposed real limit. No limit law from Section 9 is assumed.

A proof template for the first exercises

Definition 7.1 says that $x_n \to L$ means

for every real $\varepsilon > 0$ there exists a number N such that every integer $n > N$ satisfies $|x_n - L| < \varepsilon$.

The order of these choices matters: the reader supplies an arbitrary positive ε; we choose N in terms of that ε; then we verify the estimate for *every* sequence index $n > N$. Our choice of N cannot depend on which later index is tested.

Ross permits a real threshold N. If an integer threshold is preferred, the Archimedean property supplies a natural number larger than our chosen N, and the same proof applies to all indices above that larger threshold. The first finitely many terms need not obey the error estimate: the definition requires it only after the threshold.

All the limit assertions below are proved from this definition. Algebraic simplifications, sign rules, and absolute-value identities are used as rules about individual real numbers, without invoking the limit theorems of Section 9.

8.1 Exercise 8.1: Four limits proved directly from the definition

What we must show. Prove each of the four stated limits with a complete ε–N argument.

Part (a): the alternating reciprocal sequence.

We must prove

$$\lim_{n\to\infty} \frac{(-1)^n}{n} = 0.$$

For an integer n, $(-1)^n$ is either 1 or -1, according to the parity of n. Its absolute value is therefore always 1. Since $n \in \mathbb{N}$ implies $n > 0$, the exact error is

$$\begin{aligned}\left|\frac{(-1)^n}{n} - 0\right| &= \frac{|(-1)^n|}{n}, \\ &= \frac{1}{n}.\end{aligned}$$

We need this error to be strictly less than ε.

Let $\varepsilon > 0$ be arbitrary and choose

$$N = \max\left\{1, \frac{1}{\varepsilon}\right\}.$$

Take any integer $n > N$. Then $n > 1/\varepsilon$. Multiplication by the positive number ε gives $\varepsilon n > 1$; division by the positive number n gives $\varepsilon > 1/n$. Consequently

$$\left|\frac{(-1)^n}{n} - 0\right| = \frac{1}{n} < \varepsilon.$$

The estimate holds for every integer $n > N$, as the definition requires.

Conclusion. For every $\varepsilon > 0$ the displayed threshold works, so $\boxed{(-1)^n/n \longrightarrow 0}$. Alternation does not interfere with the estimate because both possible signs have absolute value 1.

Part (b), Step 1: compute the error and the needed threshold.

We must prove $1/n^{1/3} \to 0$, where $n^{1/3}$ is the positive real cube root of n. Since $n^{1/3} > 0$,

$$\left|\frac{1}{n^{1/3}} - 0\right| = \frac{1}{n^{1/3}}.$$

To make this less than ε, it suffices to have $n^{1/3} > 1/\varepsilon$, or $n > 1/\varepsilon^3$. These implications concern ordinary positive real numbers. For example, strict comparison of their cubes follows from

$$u^3 - v^3 = (u - v)(u^2 + uv + v^2) \qquad (u, v > 0).$$

If $u > v > 0$, both factors on the right are positive, so $u^3 > v^3$. Conversely, if $u^3 > v^3$ but $u \le v$, cubing the nonnegative numbers would give $u^3 \le v^3$, a contradiction. Thus comparing positive cubes is equivalent to comparing their positive cube roots.

Part (b), Step 2: give the formal proof.

Let $\varepsilon > 0$ be arbitrary and choose

$$N = \max\left\{1, \frac{1}{\varepsilon^3}\right\}.$$

For an arbitrary integer $n > N$, we have

$$n > \frac{1}{\varepsilon^3} = \left(\frac{1}{\varepsilon}\right)^3.$$

Taking positive cube roots gives $n^{1/3} > 1/\varepsilon$. Multiplying this inequality by $\varepsilon > 0$ and then dividing by $n^{1/3} > 0$ gives

$$\varepsilon n^{1/3} > 1 \quad \Longrightarrow \quad \frac{1}{n^{1/3}} < \varepsilon.$$

Hence the exact error satisfies $\left|1/n^{1/3} - 0\right| < \varepsilon$ for every integer $n > N$.

Conclusion. Definition 7.1 therefore gives $\boxed{1/n^{1/3} \longrightarrow 0}$. The threshold depends on ε^{-3} because the denominator is a cube root.

Part (c), Step 1: simplify the exact difference from the proposed limit.

The sequence in the printed exercise is

$$x_n = \frac{2n - 1}{3n + 2}, \qquad \text{and the proposed limit is } \frac{2}{3}.$$

For every $n \in \mathbb{N}$, $3n+2 > 0$, so the denominator is nonzero. Putting the two fractions over the common denominator $3(3n+2)$ gives

$$\begin{aligned} x_n - \frac{2}{3} &= \frac{3(2n-1) - 2(3n+2)}{3(3n+2)}, \\ &= \frac{6n - 3 - 6n - 4}{9n+6}, \\ &= \frac{-7}{9n+6}. \end{aligned}$$

Because $9n+6 > 0$, taking absolute values yields

$$\left| x_n - \frac{2}{3} \right| = \frac{7}{9n+6}.$$

Also $9n+6 \geq 9n > 0$, so increasing the positive denominator decreases the quotient:

$$\frac{7}{9n+6} \leq \frac{7}{9n}.$$

Part (c), Step 2: choose a threshold and check all later indices.

Let $\varepsilon > 0$ and set

$$N = \max\left\{1, \frac{7}{9\varepsilon}\right\}.$$

If an integer $n > N$, then $n > 7/(9\varepsilon)$. Multiplication by $9\varepsilon > 0$ gives $9\varepsilon n > 7$; division by $9n > 0$ gives $\varepsilon > 7/(9n)$. Thus

$$\left| x_n - \frac{2}{3} \right| = \frac{7}{9n+6} \leq \frac{7}{9n} < \varepsilon.$$

Conclusion. The estimate holds for every integer $n > N$, proving

$$\boxed{\lim_{n\to\infty} \frac{2n-1}{3n+2} = \frac{2}{3}}.$$

Part (d), Step 1: isolate a positive tail of the denominator.

We must prove

$$x_n = \frac{n+6}{n^2-6} \longrightarrow 0.$$

The denominator is nonzero for every positive integer n: if $n \leq 2$, then $n^2 \leq 4 < 6$; if $n \geq 3$, then $n^2 \geq 9 > 6$. For the limit estimate, we will use only indices $n \geq 4$. For such n, we have $n^2/2 \geq 8 \geq 6$, and therefore

$$n^2 - 6 \geq n^2 - \frac{n^2}{2} = \frac{n^2}{2} > 0.$$

This is a lower bound on the denominator, with its sign recorded.

Part (d), Step 2: bound the numerator and then the entire error.

For $n \geq 1$, the inequality $6 \leq 6n$ gives

$$n + 6 \leq n + 6n = 7n.$$

If $n \geq 4$, both the numerator $n+6$ and denominator n^2-6 are positive. Thus

$$\begin{aligned} |x_n - 0| &= \frac{n+6}{n^2-6}, \\ &\leq \frac{7n}{n^2-6} && \text{by the upper bound on the numerator,} \\ &\leq \frac{7n}{n^2/2} && \text{by the positive lower bound on the denominator,} \\ &= \frac{14n}{n^2} = \frac{14}{n}. \end{aligned}$$

The cancellation in the last line is valid because $n > 0$.

Part (d), Step 3: give the formal threshold.

Let $\varepsilon > 0$ and choose

$$N = \max\left\{4, \frac{14}{\varepsilon}\right\}.$$

An arbitrary integer $n > N$ satisfies $n > 4$, so every preceding tail estimate applies. It also satisfies $n > 14/\varepsilon$; multiplying by $\varepsilon > 0$ and dividing by $n > 0$ gives $14/n < \varepsilon$. Hence

$$\left|\frac{n+6}{n^2-6} - 0\right| \leq \frac{14}{n} < \varepsilon \qquad \text{for every integer } n > N.$$

Conclusion. The threshold supplies the estimate required by Definition 7.1, so

$$\boxed{\frac{n+6}{n^2-6} \longrightarrow 0}.$$

We did not require the first few terms to have a positive denominator.

8.2 Exercise 8.2: Determine five limits and justify each one formally

What we must show. Find the limits of the five stated sequences and prove every answer directly from Definition 7.1. The candidate for a quotient of linear expressions can be suggested by its leading coefficients, but the error calculations below are what establish the limits.

Part (a), Step 1: estimate the difference from zero.

For $a_n = n/(n^2+1)$, the numerator and denominator are positive for every $n \in \mathbb{N}$. Since $n^2 + 1 \geq n^2 > 0$,

$$\begin{aligned} |a_n - 0| &= \frac{n}{n^2+1}, \\ &\leq \frac{n}{n^2}, \\ &= \frac{1}{n}. \end{aligned}$$

We will prove that 0 is the limit.

Part (a), Step 2: supply the quantified proof.

Let $\varepsilon > 0$ and set

$$N = \max\left\{1, \frac{1}{\varepsilon}\right\}.$$

For every integer $n > N$, we have $n > 1/\varepsilon$. Because ε and n are positive, this implies $1/n < \varepsilon$. The error bound from Step 1 then gives

$$|a_n - 0| \leq \frac{1}{n} < \varepsilon.$$

Conclusion. The limit is $\boxed{\lim_{n\to\infty} \frac{n}{n^2+1} = 0}$, as verified for every positive error tolerance.

Part (b), Step 1: compute the exact difference from $7/3$.

Let $b_n = (7n-19)/(3n+7)$. Since $n \geq 1$, its denominator satisfies $3n+7 \geq 10 > 0$. Compute

$$\begin{aligned} b_n - \frac{7}{3} &= \frac{3(7n-19) - 7(3n+7)}{3(3n+7)}, \\ &= \frac{21n - 57 - 21n - 49}{9n+21}, \\ &= \frac{-106}{9n+21}. \end{aligned}$$

Thus

$$\left|b_n - \frac{7}{3}\right| = \frac{106}{9n+21} \leq \frac{106}{9n},$$

because $9n + 21 \geq 9n > 0$.

Part (b), Step 2: choose and verify a threshold.

Given any $\varepsilon > 0$, choose

$$N = \max\left\{1, \frac{106}{9\varepsilon}\right\}.$$

Take any integer $n > N$. Then $n > 106/(9\varepsilon)$, so $9\varepsilon n > 106$ and hence $106/(9n) < \varepsilon$. It follows that

$$\left|b_n - \frac{7}{3}\right| = \frac{106}{9n+21} \leq \frac{106}{9n} < \varepsilon.$$

Conclusion. Since this works for every $\varepsilon > 0$,

$$\boxed{\lim_{n\to\infty} \frac{7n-19}{3n+7} = \frac{7}{3}}.$$

Part (c), Step 1: compute the exact difference from $4/7$.

Let $c_n = (4n+3)/(7n-5)$. Its denominator is positive for all natural indices, since $7n-5 \geq 7-5 = 2 > 0$. Cross-multiplication gives

$$\begin{aligned} c_n - \frac{4}{7} &= \frac{7(4n+3) - 4(7n-5)}{7(7n-5)}, \\ &= \frac{28n + 21 - 28n + 20}{49n-35}, \end{aligned}$$

$$= \frac{41}{49n - 35}.$$

The error is already positive, so its absolute value has the same expression.

Part (c), Step 2: bound the denominator below by a multiple of n.

For $n \geq 1$,

$$(7n - 5) - 2n = 5n - 5 = 5(n - 1) \geq 0.$$

Consequently $7n - 5 \geq 2n > 0$, and multiplying by $7 > 0$ gives $49n - 35 \geq 14n > 0$. Hence

$$\left| c_n - \frac{4}{7} \right| = \frac{41}{49n - 35} \leq \frac{41}{14n}.$$

Part (c), Step 3: prove the definition with an explicit threshold.

Let $\varepsilon > 0$ and put

$$N = \max\left\{1, \frac{41}{14\varepsilon}\right\}.$$

For an arbitrary integer $n > N$, the inequality $n > 41/(14\varepsilon)$ gives $14\varepsilon n > 41$ and thus $41/(14n) < \varepsilon$. Therefore

$$\left| c_n - \frac{4}{7} \right| \leq \frac{41}{14n} < \varepsilon.$$

The estimates used here hold for every such index, not merely for a selected collection of large indices.

Conclusion. The limit is

$$\boxed{\lim_{n\to\infty} \frac{4n + 3}{7n - 5} = \frac{4}{7}}.$$

Part (d), Step 1: simplify the difference from $2/5$.

For $d_n = (2n + 4)/(5n + 2)$, the denominator satisfies $5n + 2 \geq 7 > 0$. The exact calculation is

$$\begin{aligned} d_n - \frac{2}{5} &= \frac{5(2n + 4) - 2(5n + 2)}{5(5n + 2)}, \\ &= \frac{10n + 20 - 10n - 4}{25n + 10}, \\ &= \frac{16}{25n + 10}. \end{aligned}$$

Thus

$$\left| d_n - \frac{2}{5} \right| = \frac{16}{25n + 10} \leq \frac{16}{25n},$$

where the comparison uses $25n + 10 \geq 25n > 0$.

Part (d), Step 2: choose a threshold and verify the required error.

For an arbitrary $\varepsilon > 0$, set

$$N = \max\left\{1, \frac{16}{25\varepsilon}\right\}.$$

If an integer $n > N$, then $n > 16/(25\varepsilon)$. Multiplication by $25\varepsilon > 0$ gives $25\varepsilon n > 16$; division by $25n > 0$ gives $16/(25n) < \varepsilon$. Hence

$$\left| d_n - \frac{2}{5} \right| \leq \frac{16}{25n} < \varepsilon.$$

Conclusion. Definition 7.1 gives

$$\boxed{\lim_{n\to\infty} \frac{2n+4}{5n+2} = \frac{2}{5}}.$$

Part (e), Step 1: bound the error using the elementary sine bound.

The sequence is $s_n = (\sin n)/n$. For every real x, the elementary trigonometric inequality $-1 \leq \sin x \leq 1$ gives $|\sin x| \leq 1$. Applying it at $x = n$ and using $n > 0$ yields

$$\begin{aligned} |s_n - 0| &= \left|\frac{\sin n}{n}\right|, \\ &= \frac{|\sin n|}{n}, \\ &\leq \frac{1}{n}. \end{aligned}$$

The bound holds for every natural n regardless of the sign of $\sin n$.

Part (e), Step 2: carry out the formal proof.

Let $\varepsilon > 0$ and choose

$$N = \max\left\{1, \frac{1}{\varepsilon}\right\}.$$

For every integer $n > N$, multiplication of $n > 1/\varepsilon$ by $\varepsilon > 0$ and division by $n > 0$ give $1/n < \varepsilon$. Consequently

$$|s_n - 0| \leq \frac{1}{n} < \varepsilon.$$

Conclusion. Thus $\boxed{(\sin n)/n \longrightarrow 0}$. The proof uses a uniform bound on sine, not a claim that the sequence $(\sin n)$ has a limit.

8.3 Exercise 8.3: Taking square roots when the limit is zero

What we must show. Suppose $s_n \geq 0$ for every $n \in \mathbb{N}$ and $s_n \to 0$. Prove that $\sqrt{s_n} \to 0$ directly from Definition 7.1.

Plan. To make the nonnegative square root smaller than ε, it suffices to make the original number smaller than ε^2. The input error tolerance for the given convergence will therefore be $\delta = \varepsilon^2$.

Step 1: choose the input tolerance and use the hypothesis.

Let $\varepsilon > 0$ be arbitrary. Then

$$\delta = \varepsilon^2 > 0.$$

The assertion $s_n \to 0$ means that the definition applies to every positive tolerance, including this particular δ. Hence there is an N such that every integer $n > N$ satisfies

$$|s_n - 0| < \delta = \varepsilon^2.$$

Because $s_n \geq 0$, its absolute value is itself, and the last estimate is exactly

$$0 \leq s_n < \varepsilon^2.$$

Step 2: justify the strict comparison of square roots.

Fix any integer $n > N$ and put $u = \sqrt{s_n}$. The nonnegative square root exists because $s_n \geq 0$, and it satisfies $u \geq 0$ and $u^2 = s_n$. If $u \geq \varepsilon$, then

$$u^2 - \varepsilon^2 = (u - \varepsilon)(u + \varepsilon) \geq 0,$$

since $u - \varepsilon \geq 0$ and $u + \varepsilon > 0$. This would imply $s_n = u^2 \geq \varepsilon^2$, contradicting the estimate from Step 1. Thus $u < \varepsilon$.

Step 3: verify the target absolute-value estimate.

Since $u \geq 0$,

$$|\sqrt{s_n} - 0| = \sqrt{s_n} = u < \varepsilon \qquad \text{for every integer } n > N.$$

The threshold N was supplied by the given convergence for the tolerance ε^2; it does not depend on the later index n.

Conclusion. For every target tolerance $\varepsilon > 0$ we obtained the required threshold, proving $\boxed{\sqrt{s_n} \longrightarrow 0}$. No division by $\sqrt{s_n}$ was used, so the argument includes terms equal to zero.

8.4 Exercise 8.4: A sequence tending to zero times a bounded sequence

What we must show. Assume (t_n) is bounded, so there is a real number M with $|t_n| \leq M$ for every $n \in \mathbb{N}$, and assume $s_n \to 0$. Prove $s_n t_n \to 0$ directly from the definition.

Step 1: choose a positive bound that can be used as a divisor.

The given bound satisfies $M \geq |t_1| \geq 0$, so $M \geq 0$. It need not be strictly positive. Define instead

$$B = \max\{1, M\}.$$

Then $B \geq 1 > 0$, and for every n,

$$|t_n| \leq M \leq B.$$

The constant B is fixed for the whole sequence; it does not depend on the index n or on our later choice of error tolerance.

Step 2: ask the sequence tending to zero for a sufficiently small error.

Let $\varepsilon > 0$ be arbitrary and put

$$\delta = \frac{\varepsilon}{B} > 0.$$

Since $s_n \to 0$, there is an N such that, for every integer $n > N$,

$$|s_n - 0| < \delta = \frac{\varepsilon}{B}.$$

Step 3: verify the product estimate term by term.

Take an arbitrary integer $n > N$. Multiplication of $|t_n| \leq B$ by the nonnegative number $|s_n|$ gives $|s_n|\,|t_n| \leq B\,|s_n|$. Multiplication of $|s_n| < \varepsilon/B$ by the strictly positive number B preserves the strict inequality. Therefore

$$|s_n t_n - 0| = |s_n|\,|t_n|\,,$$

$$\begin{aligned} &\leq B\,|s_n|\,, \\ &< B\frac{\varepsilon}{B}, \\ &= \varepsilon. \end{aligned}$$

This bound holds for every integer $n > N$.

Step 4: check the possible zero-bound case.

If the original M was zero, then $0 \leq |t_n| \leq 0$ forces $t_n = 0$ for every n. The products are then all zero. Our proof already covers this case because it used $B = \max\{1, 0\} = 1$ rather than dividing by $M = 0$.

Conclusion. The error estimate establishes $\boxed{s_n t_n \longrightarrow 0}$ by Definition 7.1. The sequence (t_n) need only be bounded; its own convergence was not assumed or used.

8.5 Exercise 8.5: The squeeze lemma and an absolute-value comparison

What we must show. Prove (a) that $a_n \leq s_n \leq b_n$ for all n, together with $a_n \to s$ and $b_n \to s$, implies $s_n \to s$; and (b) that $|s_n| \leq t_n$ for all n, together with $t_n \to 0$, implies $s_n \to 0$.

Part (a), Step 1: obtain two thresholds for the same tolerance.

Let $\varepsilon > 0$ be arbitrary. Because $a_n \to s$, there is an N_1 such that

$$n > N_1 \quad \Longrightarrow \quad |a_n - s| < \varepsilon.$$

Because $b_n \to s$, there is an N_2 such that

$$n > N_2 \quad \Longrightarrow \quad |b_n - s| < \varepsilon.$$

Each implication is required for every integer index exceeding the corresponding threshold.

Part (a), Step 2: make both estimates valid at once.

Choose

$$N = \max\{N_1, N_2\}.$$

For an arbitrary integer $n > N$, we have both $n > N_1$ and $n > N_2$. Thus both error estimates apply at this same index. The elementary absolute-value equivalence from Exercise 3.7(a) gives

$$-\varepsilon < a_n - s < \varepsilon, \qquad -\varepsilon < b_n - s < \varepsilon.$$

Part (a), Step 3: insert the middle term between these errors.

The order assumption holds for every n, so in particular for our chosen $n > N$ we have $a_n \leq s_n \leq b_n$. Add $-s$ throughout:

$$a_n - s \leq s_n - s \leq b_n - s.$$

Combine the strict lower bound from a_n with the strict upper bound from b_n:

$$-\varepsilon < a_n - s \leq s_n - s \leq b_n - s < \varepsilon.$$

The middle inequalities are permitted to be weak. The outer strict inequalities still give $-\varepsilon < s_n - s < \varepsilon$ by transitivity. Applying Exercise 3.7(a) once more gives $|s_n - s| < \varepsilon$.

Part (a), Step 4: record the quantifiers of the conclusion.

The number $\varepsilon > 0$ was arbitrary. We chose a single N for that tolerance and verified $|s_n - s| < \varepsilon$ for every integer $n > N$. This is exactly the definition of convergence to s.

Conclusion. The squeeze lemma is

$$\boxed{a_n \leq s_n \leq b_n, \quad a_n \to s, \quad b_n \to s \quad \Longrightarrow \quad s_n \to s}.$$

The two enclosing sequences may approach s from either side; only their common limit and the displayed order comparisons were used.

Part (b), Step 1: extract the sign information contained in the hypothesis.

For each n, $|s_n| \geq 0$. The assumed inequality therefore gives

$$0 \leq |s_n| \leq t_n.$$

In particular, $t_n \geq 0$ for every index. This is a consequence of the hypothesis, not an extra assumption.

Part (b), Step 2: apply the definition to the given sequence.

Let $\varepsilon > 0$ be arbitrary. Since $t_n \to 0$, there is an N such that every integer $n > N$ satisfies

$$|t_n - 0| < \varepsilon.$$

As $t_n \geq 0$, this is the same as $t_n < \varepsilon$.

Part (b), Step 3: finish the required error estimate.

For every integer $n > N$, combine Step 1 with Step 2:

$$0 \leq |s_n - 0| = |s_n| \leq t_n < \varepsilon.$$

Thus $|s_n - 0| < \varepsilon$ for every such n.

Conclusion. For every $\varepsilon > 0$ the preceding threshold works, so $\boxed{s_n \longrightarrow 0}$. This direct proof records why the sign of t_n causes no difficulty.

8.6 Exercise 8.6: Absolute values at zero and an alternating counterexample

What we must show. Prove (a) $s_n \to 0$ if and only if $|s_n| \to 0$, and (b) the sequence $s_n = (-1)^n$ has convergent absolute values but is itself divergent.

Throughout these exercises, convergence is used in the precise form of Definition 7.1: $x_n \to L$ means that for every $\varepsilon > 0$ there is a real number N such that

$$n \in \mathbb{N}, \quad n > N \quad \Longrightarrow \quad |x_n - L| < \varepsilon.$$

A threshold N need not be an integer. Whenever an index above a specified threshold is needed, the Archimedean property supplies a positive integer above it.

Part (a), Step 1: compare the two errors exactly.

For every $n \in \mathbb{N}$, the number $|s_n|$ is nonnegative. Therefore

$$||s_n| - 0| = ||s_n|| = |s_n| = |s_n - 0|.$$

The errors in the two proposed zero limits are the same real number, not merely bounded by each other.

Part (a), Step 2: prove the forward implication.

Assume $s_n \to 0$, and let $\varepsilon > 0$ be arbitrary. The definition supplies N such that

$$n > N \quad \Longrightarrow \quad |s_n - 0| < \varepsilon.$$

For each such index, the identity in Step 1 gives

$$||s_n| - 0| = |s_n - 0| < \varepsilon.$$

Thus the same threshold N verifies convergence of $|s_n|$ to 0.

Part (a), Step 3: prove the reverse implication.

Assume $|s_n| \to 0$, and let $\varepsilon > 0$ be arbitrary. Choose N from this convergence so that

$$n > N \quad \Longrightarrow \quad ||s_n| - 0| < \varepsilon.$$

Using Step 1 again yields

$$|s_n - 0| = ||s_n| - 0| < \varepsilon \qquad (n > N).$$

This is the definition of $s_n \to 0$.

Conclusion. Both implications have been proved with the same error and the same threshold:

$$\boxed{s_n \to 0 \quad \Longleftrightarrow \quad |s_n| \to 0.}$$

Part (b), Step 1: prove convergence of the absolute values.

If $s_n = (-1)^n$, then s_n equals either 1 or -1. Hence $|s_n| = 1$ for every n. For any $\varepsilon > 0$, choose $N = 0$. Every positive integer $n > N$ satisfies

$$||s_n| - 1| = |1 - 1| = 0 < \varepsilon.$$

Therefore $|s_n| \to 1$.

Part (b), Step 2: rule out every possible limit of the original sequence.

Suppose, for a contradiction, that $s_n \to L$ for some real L. Apply the definition with the fixed positive tolerance $\varepsilon = 1$. There is a real N such that

$$n > N \quad \Longrightarrow \quad |s_n - L| < 1.$$

Choose an integer $k > \max\{N, 1\}$. Then both $2k$ and $2k + 1$ are positive integers larger than N. Their terms are

$$s_{2k} = 1, \qquad s_{2k+1} = -1.$$

The triangle inequality gives

$$\begin{aligned} 2 &= |1 - (-1)| \\ &= |(1 - L) + (L - (-1))| \end{aligned}$$

$$\begin{aligned}
&\leq |1 - L| + |L - (-1)| \\
&= |s_{2k} - L| + |s_{2k+1} - L| \\
&< 1 + 1 = 2.
\end{aligned}$$

The equality of the absolute values in the fourth line uses $|-x| = |x|$. The resulting strict inequality $2 < 2$ is impossible.

Conclusion. The absolute-value sequence converges to 1, but $((-1)^n)$ has no real limit. Thus existence of a limit for $|s_n|$ alone does not guarantee existence of a limit for s_n.

8.7 Exercise 8.7: Three divergent sequences

What we must show. Show that none of the following sequences converges:

$$\cos\left(\frac{n\pi}{3}\right), \qquad (-1)^n n, \qquad \sin\left(\frac{n\pi}{3}\right).$$

Plan. We will use only the definition of convergence and the triangle inequality. If a sequence converged to L, then for any chosen $\varepsilon > 0$, all sufficiently late terms would be within ε of L. Any two such terms would satisfy

$$|s_p - s_q| = |(s_p - L) + (L - s_q)| \leq |s_p - L| + |s_q - L| < 2\varepsilon.$$

For each proposed sequence, we will produce two arbitrarily late terms whose distance contradicts this inequality. This displayed observation has just been proved directly, so no theorem about subsequences or bounded convergent sequences is needed.

Part (a), Step 1: select a tolerance under an assumed limit.

Suppose

$$s_n = \cos\left(\frac{n\pi}{3}\right) \to L.$$

With $\varepsilon = 1$, choose N such that every positive integer $n > N$ satisfies $|s_n - L| < 1$.

Part (a), Step 2: find late terms equal to 1 and -1.

Choose an integer $k > \max\{N, 1\}$, and put

$$p = 6k, \qquad q = 6k + 3.$$

Both indices exceed N. The period 2π of cosine gives

$$s_p = \cos(2k\pi) = 1, \qquad s_q = \cos(2k\pi + \pi) = -1.$$

Their distance is therefore $|s_p - s_q| = 2$. The two-late-terms estimate gives

$$2 = |s_p - s_q| \leq |s_p - L| + |s_q - L| < 1 + 1 = 2,$$

a contradiction.

Conclusion. The sequence $(\cos(n\pi/3))$ does not converge.

Part (b), Step 1: use the definition with a fixed tolerance.

Suppose $s_n = (-1)^n n \to L$. Choose N from the definition with $\varepsilon = 1$.

Part (b), Step 2: compute the distance between late even and odd terms.

Choose an integer $k > \max\{N, 1\}$. Then $p = 2k$ and $q = 2k + 1$ are larger than N and

$$s_p = 2k, \qquad s_q = -(2k + 1).$$

Their difference is

$$s_p - s_q = 2k + (2k + 1) = 4k + 1 > 2.$$

Since this difference is positive, its absolute value is also $4k + 1$. But convergence would give

$$4k + 1 = |s_p - s_q| \leq |s_p - L| + |s_q - L| < 1 + 1 = 2.$$

This contradicts $4k + 1 > 2$.

Conclusion. The sequence $((-1)^n n)$ does not converge. The proof uses two explicit terms in every sufficiently late tail, rather than assuming a boundedness theorem from a later section.

Part (c), Step 1: choose a tolerance suited to the sine values.

Suppose

$$s_n = \sin\left(\frac{n\pi}{3}\right) \to L.$$

Choose the positive tolerance $\varepsilon = \sqrt{3}/2$. The definition gives N such that $n > N$ implies $|s_n - L| < \sqrt{3}/2$.

Part (c), Step 2: use two positions in the six-term period.

Choose an integer $k > \max\{N, 1\}$ and let

$$p = 6k + 1, \qquad q = 6k + 4.$$

Both are larger than N. By the period 2π of sine,

$$s_p = \sin\left(2k\pi + \frac{\pi}{3}\right) = \frac{\sqrt{3}}{2}, \qquad s_q = \sin\left(2k\pi + \frac{4\pi}{3}\right) = -\frac{\sqrt{3}}{2}.$$

In particular, $|s_p - s_q| = \sqrt{3}$. The convergence estimates would imply

$$\sqrt{3} = |s_p - s_q| \leq |s_p - L| + |s_q - L| < \frac{\sqrt{3}}{2} + \frac{\sqrt{3}}{2} = \sqrt{3}.$$

This is impossible.

Conclusion. The sequence $(\sin(n\pi/3))$ does not converge. Every sequence in the exercise has now been ruled out by the definition of convergence itself.

8.8 Exercise 8.8: Three radical limits with explicit error bounds

What we must show. Prove the three stated limits:

$$\sqrt{n^2 + 1} - n \longrightarrow 0,$$

$$\sqrt{n^2+n}-n \longrightarrow \frac{1}{2},$$
$$\sqrt{4n^2+n}-2n \longrightarrow \frac{1}{4}.$$

The index n is always a positive integer. In particular, division by n is legitimate, and $\sqrt{n^2} = n$ rather than $-n$.

Plan. Multiplication by a conjugate removes the subtraction of large radicals. We will then bound each *error from its claimed limit* by a positive multiple of $1/n$. The estimates below are ordinary algebraic identities and inequalities, not applications of limit laws.

Part (a), Step 1: rationalize the expression and check its sign.

Let $u_n = \sqrt{n^2+1}-n$. Because $n \geq 1$, both n and $\sqrt{n^2+1}$ are positive. Moreover, $\sqrt{n^2+1} > n$: otherwise squaring a supposed inequality $\sqrt{n^2+1} \leq n$ between nonnegative numbers would give $n^2+1 \leq n^2$, a contradiction. Thus $u_n > 0$. Its conjugate $\sqrt{n^2+1}+n$ is positive, so

$$\begin{aligned} u_n &= \frac{(\sqrt{n^2+1}-n)(\sqrt{n^2+1}+n)}{\sqrt{n^2+1}+n} \\ &= \frac{(n^2+1)-n^2}{\sqrt{n^2+1}+n} \\ &= \frac{1}{\sqrt{n^2+1}+n}. \end{aligned}$$

Part (a), Step 2: obtain an error estimate valid for every index.

The denominator satisfies $\sqrt{n^2+1}+n \geq 2n > 0$. Taking reciprocals of positive quantities reverses the comparison, so

$$|u_n - 0| = u_n \leq \frac{1}{2n}.$$

Part (a), Step 3: choose and verify a threshold.

Let $\varepsilon > 0$ be arbitrary, and choose

$$N = \frac{1}{2\varepsilon}.$$

For every positive integer $n > N$, multiplying by $2\varepsilon > 0$ gives $2\varepsilon n > 1$. Division by $2n > 0$ gives $1/(2n) < \varepsilon$. Hence

$$n > N \quad \Longrightarrow \quad |u_n - 0| \leq \frac{1}{2n} < \varepsilon.$$

This is precisely $u_n \to 0$.

Conclusion. $\boxed{\lim_{n\to\infty}(\sqrt{n^2+1}-n) = 0}$. An explicit error bound is $1/(2n)$.

Part (b), Step 1: rationalize and factor out the positive index.

Let $v_n = \sqrt{n^2+n}-n$. Multiplying by its positive conjugate gives

$$v_n = \frac{(n^2+n)-n^2}{\sqrt{n^2+n}+n} = \frac{n}{\sqrt{n^2+n}+n}.$$

Set

$$r_n = \sqrt{1 + \frac{1}{n}}.$$

The quantity nr_n is positive, and its square is

$$(nr_n)^2 = n^2\left(1 + \frac{1}{n}\right) = n^2 + n.$$

By uniqueness of the nonnegative square root, $\sqrt{n^2+n} = nr_n$. Substitution gives the exact identity

$$v_n = \frac{n}{nr_n + n} = \frac{1}{r_n + 1}.$$

The number r_n is greater than 1, because $r_n^2 = 1 + 1/n > 1$. Thus $r_n + 1 > 2$, and $0 < v_n < 1/2$.

Part (b), Step 2: rationalize the error from $1/2$.

The sign information in Step 1 gives

$$\begin{aligned}\left|v_n - \tfrac{1}{2}\right| &= \frac{1}{2} - \frac{1}{r_n + 1} \\ &= \frac{(r_n + 1) - 2}{2(r_n + 1)} \\ &= \frac{r_n - 1}{2(r_n + 1)}.\end{aligned}$$

A second conjugate calculation yields

$$r_n - 1 = \frac{(r_n - 1)(r_n + 1)}{r_n + 1} = \frac{r_n^2 - 1}{r_n + 1} = \frac{1/n}{r_n + 1}.$$

Therefore

$$\left|v_n - \tfrac{1}{2}\right| = \frac{1}{2n(r_n + 1)^2}.$$

Since $r_n \geq 1$, we have $(r_n + 1)^2 \geq 4$, so

$$\left|v_n - \tfrac{1}{2}\right| \leq \frac{1}{8n}.$$

Part (b), Step 3: verify the definition of convergence.

Let $\varepsilon > 0$ be arbitrary and choose $N = 1/(8\varepsilon)$. For every positive integer $n > N$, multiplication and division by positive quantities give $1/(8n) < \varepsilon$. Consequently

$$\left|v_n - \tfrac{1}{2}\right| \leq \frac{1}{8n} < \varepsilon \qquad (n > N).$$

The definition proves the desired limit.

Conclusion. $\boxed{\lim_{n\to\infty}(\sqrt{n^2+n} - n) = \tfrac{1}{2}}$. An explicit error bound is $1/(8n)$.

Part (c), Step 1: rationalize and identify a simpler denominator.

Let $w_n = \sqrt{4n^2+n} - 2n$. Its conjugate is positive, and multiplication gives

$$w_n = \frac{(4n^2+n) - 4n^2}{\sqrt{4n^2+n} + 2n} = \frac{n}{\sqrt{4n^2+n} + 2n}.$$

Define $R_n = \sqrt{4 + 1/n}$. Since $n > 0$, the quantity nR_n is the nonnegative square root of $4n^2 + n$:

$$(nR_n)^2 = n^2\left(4 + \frac{1}{n}\right) = 4n^2 + n.$$

Hence

$$w_n = \frac{n}{nR_n + 2n} = \frac{1}{R_n + 2}.$$

Because $R_n^2 = 4 + 1/n > 4$ and $R_n \geq 0$, we have $R_n > 2$. Thus $R_n + 2 > 4$ and $0 < w_n < 1/4$.

Part (c), Step 2: compute and bound the error from $1/4$.

Using the sign of the error, put the fractions over one denominator:

$$\begin{aligned}\left|w_n - \tfrac{1}{4}\right| &= \frac{1}{4} - \frac{1}{R_n + 2} \\ &= \frac{(R_n + 2) - 4}{4(R_n + 2)} \\ &= \frac{R_n - 2}{4(R_n + 2)}.\end{aligned}$$

The second rationalization gives

$$R_n - 2 = \frac{R_n^2 - 4}{R_n + 2} = \frac{1/n}{R_n + 2}.$$

It follows that

$$\left|w_n - \tfrac{1}{4}\right| = \frac{1}{4n(R_n + 2)^2} \leq \frac{1}{64n},$$

because $R_n \geq 2$ implies $(R_n + 2)^2 \geq 16$.

Part (c), Step 3: give the final tolerance-dependent threshold.

Let $\varepsilon > 0$ be arbitrary and choose $N = 1/(64\varepsilon)$. Then $n > N$ implies $1/(64n) < \varepsilon$, so

$$\left|w_n - \tfrac{1}{4}\right| \leq \frac{1}{64n} < \varepsilon \qquad (n \in \mathbb{N},\ n > N).$$

This verifies convergence directly.

Conclusion. $\boxed{\lim_{n\to\infty}(\sqrt{4n^2+n} - 2n) = \tfrac{1}{4}}$. All three limits were proved by explicit inequalities, without using algebraic limit theorems from Section 9.

8.9 Exercise 8.9: Eventual bounds pass to a convergent limit

What we must show. Let $s_n \to s$. Prove:

(a) If $s_n \geq a$ for all but finitely many indices, then $s \geq a$.

(b) If $s_n \le b$ for all but finitely many indices, then $s \le b$.

(c) If $s_n \in [a, b]$ for all but finitely many indices, then $s \in [a, b]$.

A preliminary detail: replace finitely many exceptions by one threshold.

If a property fails only on a finite set $E \subseteq \mathbb{N}$ of indices, define $K = 0$ if E is empty and $K = \max E$ otherwise. Every positive integer $n > K$ is outside E, so every such index satisfies the property. This explains each use of an eventual-bound threshold below.

Part (a), Step 1: suppose the limit violates the eventual lower bound.

Choose K so that $n > K$ implies $s_n \ge a$. Suppose, for a contradiction, that $s < a$. Then the gap $a - s$ is positive. Set

$$\varepsilon = \frac{a - s}{2} > 0.$$

Convergence supplies N_0 such that $n > N_0$ implies $|s_n - s| < \varepsilon$.

Part (a), Step 2: combine the two tail conditions.

Choose a positive integer $n > \max\{K, N_0\}$. The condition $n > K$ gives $s_n \ge a$. The condition $n > N_0$ gives

$$s_n - s < \varepsilon, \qquad s_n < s + \varepsilon = \frac{s + a}{2}.$$

But

$$a - \frac{s + a}{2} = \frac{a - s}{2} > 0.$$

Thus $s_n < (s + a)/2 < a$, contradicting $s_n \ge a$.

Conclusion. The inequality $s < a$ is impossible. Therefore $\boxed{s \ge a}$. Equality $s = a$ is permitted; for example, the constant sequence $s_n = a$ has limit a and satisfies every lower bound in the hypothesis.

Part (b), Step 1: suppose the limit violates the eventual upper bound.

Choose K so that $n > K$ implies $s_n \le b$. Suppose $s > b$, and choose the positive tolerance

$$\varepsilon = \frac{s - b}{2}.$$

Convergence gives N_0 such that $n > N_0$ implies $|s_n - s| < \varepsilon$.

Part (b), Step 2: choose one index after both thresholds.

Take a positive integer $n > \max\{K, N_0\}$. Then $s_n \le b$, whereas the convergence inequality gives

$$-\varepsilon < s_n - s, \qquad s_n > s - \varepsilon = \frac{s + b}{2}.$$

The midpoint exceeds b because

$$\frac{s + b}{2} - b = \frac{s - b}{2} > 0.$$

Hence $s_n > b$, contradicting $s_n \le b$.

Conclusion. The inequality $s > b$ is impossible. Therefore $\boxed{s \le b}$. The endpoint $s = b$ is allowed.

Part (c), Step 1: unpack membership of a closed interval.

For $a \le b$,

$$x \in [a, b] \quad \Longleftrightarrow \quad a \le x \le b.$$

If all but finitely many s_n belong to this interval, then all but finitely many satisfy $s_n \ge a$, and all but finitely many satisfy $s_n \le b$. The finite-exception threshold may be chosen from the indices where interval membership fails, so both inequalities hold beyond the same threshold.

Part (c), Step 2: apply both proved statements to the same limit.

Part (a) gives $s \ge a$, and part (b) gives $s \le b$. Together these say $a \le s \le b$, which is exactly $s \in [a, b]$. If $a = b$, the two inequalities force $s = a = b$, so the singleton case is included as well.

Conclusion. For a convergent sequence,

$$\boxed{s_n \in [a, b] \text{ for all but finitely many } n \quad \Longrightarrow \quad \lim s_n \in [a, b].}$$

The finitely many exceptional early terms do not affect any of these conclusions because an index can be chosen after every exception and after the convergence threshold.

8.10 Exercise 8.10: A strict bound for the limit becomes an eventual strict bound

What we must show. If $s_n \to s$ and $s > a$, prove that there is a real N such that

$$n \in \mathbb{N}, \quad n > N \quad \Longrightarrow \quad s_n > a.$$

Plan. The positive gap $s - a$ allows us to place a whole ε-neighborhood of the limit above a.

Step 1: choose a tolerance smaller than the gap.

Because $s > a$, the real number $s - a$ is positive. Define

$$\varepsilon = \frac{s - a}{2} > 0.$$

The tolerance is fixed from the limit and the bound; it does not depend on the index n.

Step 2: obtain a threshold from convergence.

The definition of $s_n \to s$ gives a real number N such that

$$n \in \mathbb{N}, \quad n > N \quad \Longrightarrow \quad |s_n - s| < \varepsilon.$$

For any index in this tail, the absolute-value inequality implies

$$-\varepsilon < s_n - s < \varepsilon.$$

The left inequality gives $s_n > s - \varepsilon$.

Step 3: check that the lower edge is above the desired bound.

Substitution and simplification give

$$s - \varepsilon = s - \frac{s - a}{2} = \frac{2s - s + a}{2} = \frac{s + a}{2}.$$

Also

$$\frac{s+a}{2} - a = \frac{s-a}{2} > 0,$$

so $(s+a)/2 > a$. Combining these strict comparisons yields

$$s_n > s - \varepsilon = \frac{s+a}{2} > a \qquad (n \in \mathbb{N},\ n > N).$$

Conclusion. The threshold supplied for $\varepsilon = (s-a)/2$ satisfies the required implication:

$$\boxed{n > N \quad \Longrightarrow \quad s_n > a.}$$

The strict hypothesis $s > a$ matters: if $s = a$ and $s_n = a$ for all n, the sequence converges to a but none of its terms is greater than a.

9 Limit theorems for sequences

This section proves and applies rules for limits. Each application checks the hypotheses: the sequences must have the required finite limits, and a quotient needs a nonzero limiting denominator.

Definition 9.8 extends the phrase *a limit exists* to include divergence to $+\infty$ or $-\infty$. These are not finite real limits. We use the extended convention where the exercise permits it, while checking separately whether a finite-limit theorem applies. A recurrence's fixed-point equation gives a necessary condition for a finite limit; it does not prove convergence.

9.1 Exercise 9.1: Three quotients evaluated by the finite limit laws

What we must show. Use Theorems 9.2–9.7, with their hypotheses checked, to prove

$$\text{(a)} \quad \lim_{n\to\infty} \frac{n+1}{n} = 1,$$
$$\text{(b)} \quad \lim_{n\to\infty} \frac{3n+7}{6n-5} = \frac{1}{2},$$
$$\text{(c)} \quad \lim_{n\to\infty} \frac{17n^5+73n^4-18n^2+3}{23n^5+13n^3} = \frac{17}{23}.$$

Plan. Divide each polynomial expression by the appropriate positive power of n. The resulting expressions involve constants and reciprocal powers, whose limits are known. For each use of the quotient theorem, verify that the denominator terms are nonzero and that their finite limit is nonzero.

1. Record the finite limit rules being used

A constant sequence with value $c \in \mathbb{R}$ converges to c, because $|c-c| = 0 < \varepsilon$ for every $\varepsilon > 0$. Theorem 9.7(a) gives

$$\frac{1}{n^p} \longrightarrow 0 \qquad (p > 0).$$

The scalar rule is Theorem 9.2, the sum rule is Theorem 9.3, and the product rule is Theorem 9.4. Subtraction follows by combining the scalar rule with scalar -1 and the sum rule. The quotient rule, Theorem 9.6, requires a convergent numerator, a convergent denominator, a nonzero denominator at every index, and a nonzero denominator limit. All limits used in these rules are finite real numbers.

2. Part (a): Separate the constant term

Since $n \in \mathbb{N}$, the denominator n is positive. The identity

$$\frac{n+1}{n} = \frac{n}{n} + \frac{1}{n} = 1 + \frac{1}{n}$$

therefore holds for every index. The constant sequence 1 converges to 1, and Theorem 9.7(a), with $p = 1$, gives $1/n \to 0$. Applying Theorem 9.3 yields

$$\frac{n+1}{n} = 1 + \frac{1}{n} \longrightarrow 1 + 0 = 1.$$

Here the sum rule applies after rewriting; there is no need to treat the unbounded sequence n as a finite convergent sequence.

3. Part (b): Normalize numerator and denominator

For every $n \geq 1$, the original denominator satisfies $6n - 5 \geq 1 > 0$. Dividing numerator and denominator by $n > 0$ gives

$$\frac{3n+7}{6n-5} = \frac{(3n+7)/n}{(6n-5)/n} = \frac{3+7/n}{6-5/n}.$$

Let $u_n = 3+7/n$ and $v_n = 6-5/n$. By Theorem 9.7(a), $1/n \to 0$. The scalar rule gives $7/n \to 7 \cdot 0 = 0$ and $-5/n \to -5 \cdot 0 = 0$. Adding the constant sequences by Theorem 9.3 gives

$$u_n \longrightarrow 3 + 0 = 3, \qquad v_n \longrightarrow 6 + 0 = 6.$$

For each n, we also have

$$v_n = \frac{6n-5}{n} > 0.$$

Thus every denominator term is nonzero, and its limit 6 is nonzero. All hypotheses of Theorem 9.6 have now been verified, so

$$\frac{3n+7}{6n-5} = \frac{u_n}{v_n} \longrightarrow \frac{3}{6} = \frac{1}{2}.$$

4. Part (c): Divide by the largest power in the denominator

The original denominator is positive for every index:

$$23n^5 + 13n^3 = n^3(23n^2 + 13) > 0,$$

because $n \geq 1$. Divide both numerator and denominator by $n^5 > 0$. Each term then simplifies as follows:

$$\frac{n^4}{n^5} = \frac{1}{n}, \qquad \frac{n^2}{n^5} = \frac{1}{n^3}, \qquad \frac{n^3}{n^5} = \frac{1}{n^2}.$$

Consequently

$$\frac{17n^5 + 73n^4 - 18n^2 + 3}{23n^5 + 13n^3} = \frac{17 + 73/n - 18/n^3 + 3/n^5}{23 + 13/n^2}.$$

Write the normalized numerator and denominator as U_n and V_n. Theorem 9.7(a), applied with $p = 1, 2, 3, 5$, gives

$$\frac{1}{n} \to 0, \qquad \frac{1}{n^2} \to 0, \qquad \frac{1}{n^3} \to 0, \qquad \frac{1}{n^5} \to 0.$$

Multiplication by the indicated constants, using Theorem 9.2, gives

$$\frac{73}{n} \to 0, \qquad -\frac{18}{n^3} \to 0, \qquad \frac{3}{n^5} \to 0, \qquad \frac{13}{n^2} \to 0.$$

Repeated applications of the sum rule, together with the constant limits, therefore yield

$$U_n \longrightarrow 17 + 0 + 0 + 0 = 17, \qquad V_n \longrightarrow 23 + 0 = 23.$$

Moreover, $V_n = 23 + 13/n^2 > 23 > 0$ for every n, and its limit 23 is nonzero. The quotient rule applies and gives

$$\frac{U_n}{V_n} \longrightarrow \frac{17}{23}.$$

Conclusion. The three limits are (a) 1, (b) 1/2, and (c) 17/23. Every use of the quotient theorem has a denominator that is nonzero at every index and has a nonzero finite limit.

9.2 Exercise 9.2: Limits obtained from two given convergent sequences

What we must show. Suppose $x_n \to 3$, $y_n \to 7$, and $y_n \neq 0$ for every n. Find

$$\text{(a)} \quad \lim_{n\to\infty}(x_n + y_n), \qquad \text{(b)} \quad \lim_{n\to\infty} \frac{3y_n - x_n}{y_n^2}.$$

Plan. Use the sum rule in part (a). In part (b), build the numerator from scalar multiples and a sum, build the denominator with the product rule, and then check the quotient rule.

1. Part (a): Add the two limits

The hypotheses give two finite convergent sequences. Theorem 9.3 therefore applies directly:

$$x_n + y_n \longrightarrow 3 + 7 = 10.$$

2. Part (b): Evaluate the numerator

Theorem 9.2 gives

$$3y_n \longrightarrow 3 \cdot 7 = 21, \qquad -x_n \longrightarrow (-1) \cdot 3 = -3.$$

Theorem 9.3 then gives

$$3y_n - x_n = 3y_n + (-x_n) \longrightarrow 21 + (-3) = 18.$$

The numerator here is $3y_n - x_n$; only the denominator contains a square.

3. Part (b): Evaluate the denominator and take the quotient

Since $y_n \to 7$, Theorem 9.4 applied to the two factors y_n gives

$$y_n^2 = y_n\, y_n \longrightarrow 7 \cdot 7 = 49.$$

The assumption $y_n \neq 0$ makes $y_n^2 > 0$ at every index. Thus every denominator is nonzero, and its limit 49 is also nonzero. Theorem 9.6 now gives

$$\frac{3y_n - x_n}{y_n^2} \longrightarrow \frac{18}{49}.$$

Conclusion. The limits are (a) 10 and (b) 18/49.

9.3 Exercise 9.3: A polynomial expression in convergent sequences

What we must show. Assume $a_n \to a \in \mathbb{R}$ and $b_n \to b \in \mathbb{R}$. For

$$s_n = \frac{a_n^3 + 4a_n}{b_n^2 + 1},$$

prove carefully that

$$s_n \longrightarrow \frac{a^3 + 4a}{b^2 + 1}.$$

Plan. Form the powers one multiplication at a time. Then use the scalar and sum rules. The added 1 makes the denominator and its limit positive even when $b_n = 0$ or $b = 0$.

1. Construct the square and cube of the numerator sequence

The two factors in $a_n\,a_n$ both converge to a. The product rule, Theorem 9.4, gives

$$a_n^2 = a_n\,a_n \longrightarrow a \cdot a = a^2.$$

We may now apply the product rule again, this time to a_n^2 and a_n, since both have finite limits:

$$a_n^3 = (a_n^2)a_n \longrightarrow a^2 \cdot a = a^3.$$

Separately, Theorem 9.2 gives $4a_n \to 4a$. The sum rule therefore yields

$$a_n^3 + 4a_n \longrightarrow a^3 + 4a.$$

2. Construct the denominator limit

Apply Theorem 9.4 to $b_n\,b_n$ to obtain

$$b_n^2 \longrightarrow b \cdot b = b^2.$$

The constant sequence 1 converges to 1, so Theorem 9.3 gives

$$b_n^2 + 1 \longrightarrow b^2 + 1.$$

For every real b_n, its square is nonnegative, and thus $b_n^2 + 1 \geq 1 > 0$. The denominator terms are consequently nonzero. Likewise $b^2 + 1 \geq 1 > 0$, so their limit is nonzero.

3. Apply the quotient theorem with all hypotheses established

Step 1 proves convergence of the numerator, Step 2 proves convergence of the denominator, and the positivity checks verify both nonzero requirements. Theorem 9.6 therefore gives the desired limit.

Conclusion. $\lim\limits_{n\to\infty} s_n = \dfrac{a^3 + 4a}{b^2 + 1}$ for all finite real limits a and b, including $b = 0$.

9.4 Exercise 9.4: A nested-square-root recursion

What we must show. For $s_1 = 1$ and $s_{n+1} = \sqrt{s_n + 1}$, list the first four terms and, assuming convergence, prove that the limit is $(1 + \sqrt{5})/2$.

Plan. First check that every radical is defined and every term is at least 1. Under the stated convergence assumption, square the recursion and apply the product and sum limit laws. The lower bound will select the appropriate quadratic root.

1. Verify that the recursion stays defined and positive

The initial value is $s_1 = 1 \geq 1$. If $s_n \geq 1$, then $s_n + 1 \geq 2 > 0$, so its nonnegative real square root exists. Moreover, $\sqrt{s_n + 1} \geq 1$: a nonnegative number less than 1 would have square less than 1, whereas this square is at least 2. Thus $s_{n+1} \geq 1$. Induction shows that all terms are defined and satisfy $s_n \geq 1$.

2. Part (a): Substitute successively four times

The successive terms are

$$\begin{aligned} s_1 &= 1, \\ s_2 &= \sqrt{s_1 + 1} = \sqrt{1+1} = \sqrt{2}, \\ s_3 &= \sqrt{s_2 + 1} = \sqrt{\sqrt{2}+1} = \sqrt{1+\sqrt{2}}, \\ s_4 &= \sqrt{s_3 + 1} = \sqrt{\sqrt{1+\sqrt{2}}+1} = \sqrt{1+\sqrt{1+\sqrt{2}}}. \end{aligned}$$

3. A shift of a convergent sequence has the same limit

We will use this elementary observation for this and the next two recursions. If $u_n \to u \in \mathbb{R}$, then $u_{n+1} \to u$. Indeed, given $\varepsilon > 0$, choose N so that $m > N$ implies $|u_m - u| < \varepsilon$. If $n > N$, then also $n + 1 > N$, so $|u_{n+1} - u| < \varepsilon$. This is exactly the limit definition for the shifted sequence.

4. Part (b): Record the restriction on a finite limit

Assume, as requested, that $s_n \to L \in \mathbb{R}$. Since $s_n \geq 1$ for every n, we have $L \geq 1$. To see the bound directly, if $L < 1$, set $\varepsilon = (1 - L)/2 > 0$. Eventually $|s_n - L| < \varepsilon$, giving

$$s_n < L + \varepsilon = \frac{L+1}{2} < 1,$$

contrary to the bound on the terms. This is also the eventual-bound principle of Exercise 8.9(a).

5. Pass to limits in a squared identity

The defining radical equality implies

$$s_{n+1}^2 = s_n + 1$$

for every n. By Step 3, $s_{n+1} \to L$. Theorem 9.4 applied to $s_{n+1}\, s_{n+1}$ gives $s_{n+1}^2 \to L^2$. Theorem 9.3 gives $s_n + 1 \to L + 1$. These two expressions are equal term by term, so uniqueness of finite limits, proved at the end of Section 7, gives

$$L^2 = L + 1.$$

Only the product and sum laws were used; a square-root limit rule is not needed.

6. Solve the quadratic equation and select its permitted root

Rearrange and complete the square:

$$L^2 - L - 1 = 0, \qquad 4L^2 - 4L = 4, \qquad (2L-1)^2 = 5.$$

Consequently $2L - 1 = \sqrt{5}$ or $2L - 1 = -\sqrt{5}$, which gives

$$L = \frac{1+\sqrt{5}}{2} \quad \text{or} \quad L = \frac{1-\sqrt{5}}{2}.$$

We have $\sqrt{5} > 1$, since the opposite nonnegative inequality would imply $5 \leq 1$ after squaring. Thus $(1 - \sqrt{5})/2 < 0$, contradicting $L \geq 1$. The only remaining root is $(1 + \sqrt{5})/2$.

Conclusion. The first four terms are 1, $\sqrt{2}$, $\sqrt{1+\sqrt{2}}$, and $\sqrt{1+\sqrt{1+\sqrt{2}}}$. Under the given convergence assumption, $\lim s_n = (1 + \sqrt{5})/2$.

9.5 Exercise 9.5: A positive rational recursion

What we must show. Let $t_1 = 1$ and

$$t_{n+1} = \frac{t_n^2 + 2}{2t_n} \qquad (n \geq 1).$$

Assuming that (t_n) converges to a finite real number, find its limit.

Plan. Prove positivity so that every term is defined, and strengthen it to the uniform lower bound $t_n \geq 1$. The lower bound makes the assumed limit nonzero before the quotient theorem is applied.

1. Check that the recursion is defined at every index

We start with $t_1 = 1 > 0$. If $t_n > 0$, then its denominator $2t_n$ is positive and its numerator $t_n^2 + 2$ is positive. The quotient is therefore a well-defined positive real number. Induction proves $t_n > 0$ for all n, and in particular no denominator in the recursion vanishes.

2. Obtain a positive uniform lower bound

For every n, use the positive denominator to compute

$$\begin{aligned} t_{n+1} - 1 &= \frac{t_n^2 + 2}{2t_n} - \frac{2t_n}{2t_n} \\ &= \frac{t_n^2 - 2t_n + 2}{2t_n} \\ &= \frac{(t_n - 1)^2 + 1}{2t_n} > 0. \end{aligned}$$

The last numerator is positive because $(t_n - 1)^2 \geq 0$. Thus every term after t_1 is greater than 1, and $t_1 = 1$. It follows that $t_n \geq 1$ for every n.

3. The convergence assumption gives a nonzero limit

Suppose $t_n \to L \in \mathbb{R}$. The bound $t_n \geq 1$ passes to the limit by Exercise 8.9(a), so $L \geq 1 > 0$. Equivalently, if $L < 1$, the positive tolerance $(1 - L)/2$ would force eventual terms to be less than $L + (1 - L)/2 = (L + 1)/2 < 1$, a contradiction. In particular, $2L \neq 0$.

4. Apply the finite limit laws to the recursion

The product rule, Theorem 9.4, gives

$$t_n^2 = t_n \, t_n \longrightarrow L^2.$$

The constant sequence 2 converges to 2, and hence the sum rule gives $t_n^2 + 2 \to L^2 + 2$. The scalar rule gives $2t_n \to 2L$. The terms $2t_n$ are nonzero by Step 1, and their limit $2L$ is nonzero by Step 3. Thus Theorem 9.6 applies:

$$\frac{t_n^2 + 2}{2t_n} \longrightarrow \frac{L^2 + 2}{2L}.$$

The shift observation in Exercise 9.4 also gives $t_{n+1} \to L$. The recursion identifies these two sequences, so uniqueness of finite limits gives

$$L = \frac{L^2 + 2}{2L}.$$

5. Solve and select the positive root

Since $2L \neq 0$, multiplying the last equality by $2L$ is valid:

$$2L^2 = L^2 + 2, \qquad L^2 = 2.$$

Thus $L = \sqrt{2}$ or $L = -\sqrt{2}$. The root $-\sqrt{2}$ is negative and violates $L \geq 1$. Consequently the assumed finite limit is $\sqrt{2}$.

Conclusion. Under the convergence assumption, $\lim_{n\to\infty} t_n = \sqrt{2}$. The nonzero denominator limit was established before the quotient theorem was used.

9.6 Exercise 9.6: A necessary finite-limit equation and an actual infinite limit

What we must show. For $x_1 = 1$ and $x_{n+1} = 3x_n^2$, address all three questions: (a) determine the possibilities for a finite real limit; (b) decide whether the actual sequence has a limit; (c) explain why the conclusions are consistent.

Plan. A hypothetical finite limit must satisfy the recursion's fixed-point equation. Separately, use an induction estimate for the actual terms. Distinguish finite convergence from the extended infinite-limit convention in Definition 9.8.

1. Part (a): Assume a finite limit before applying the product rule

Suppose $x_n \to a \in \mathbb{R}$. The shift observation proved in Exercise 9.4 gives $x_{n+1} \to a$. Theorem 9.4 gives $x_n^2 \to a^2$, and Theorem 9.2 then gives $3x_n^2 \to 3a^2$. Since $x_{n+1} = 3x_n^2$ for every n, uniqueness of finite limits implies

$$a = 3a^2.$$

Subtracting a and factoring gives

$$0 = 3a^2 - a = a(3a - 1).$$

The zero-product property in $\mathbb{R}$ says that one of the two factors must be zero. Thus $a = 0$ or $3a - 1 = 0$; the second equation gives $a = 1/3$ after division by $3 \neq 0$.

This proves only a necessary condition: *if the sequence has a finite real limit, that limit must be* 0 *or* $1/3$. The computation does not prove that such a finite limit exists.

2. Part (b): Bound the actual terms below by their indices

We prove $x_n \geq n$ for every $n \in \mathbb{N}$. For $n = 1$, the equality $x_1 = 1$ supplies the base case. Suppose $x_n \geq n$ for some $n \geq 1$. In particular $x_n \geq 1$, so $x_n^2 - x_n = x_n(x_n - 1) \geq 0$. It follows that

$$x_{n+1} = 3x_n^2 \geq 3x_n \geq 3n \geq n + 1.$$

The final inequality holds because $3n - (n + 1) = 2n - 1 \geq 1 > 0$ for $n \geq 1$. This completes the induction.

Now let $M > 0$ and take the threshold $N = M$. For every integer $n > N$, the bound gives

$$x_n \geq n > M.$$

This is exactly Definition 9.8 for $x_n \to +\infty$. Thus the actual sequence has the extended limit $+\infty$.

It cannot have a finite real limit. For a direct check, if $x_n \to a \in \mathbb{R}$, then eventually $|x_n - a| < 1$, which implies $x_n < a + 1$. Choose an integer n larger than both that eventual threshold and $a + 1$, using the Archimedean property. The bound $x_n \geq n$ would instead give $x_n > a + 1$, a contradiction. This is also consistent with Theorem 9.1, which says every finitely convergent sequence is bounded.

3. Part (c): Explain the two meanings of the limit statement

Part (a) began with the hypothesis of a *finite real* limit a. Its use of the finite product and scalar limit laws is valid under that hypothesis. Part (b) shows that the hypothesis fails for this initial value: the terms grow without bound. Consequently the necessary alternatives in part (a) are not attained by this sequence.

Under the convention introduced after Definition 9.8, saying that "the limit exists" also allows divergence to $+\infty$ or $-\infty$. In that stated convention the answer to part (b) is yes, with limit $+\infty$; the sequence does not converge to a finite real number. Theorems 9.2–9.6 were proved for finite real limits, so their product calculation in part (a) cannot be applied with $a = +\infty$. The extended infinite-limit conclusion therefore supplies no contradiction to the finite necessary condition.

Conclusion. (a) Any hypothetical finite real limit is 0 or 1/3. (b) The actual sequence satisfies $\lim x_n = +\infty$ in the sense of Definition 9.8, and has no finite real limit. (c) A necessary condition for finite convergence does not assert its existence, and finite product laws cannot be applied at an infinite limit.

9.7 Exercise 9.7: Finish the estimate for the root of the index

What we must show. Complete the argument that

$$s_n = n^{1/n} - 1 \longrightarrow 0,$$

and hence finish the proof of Theorem 9.7(c), $n^{1/n} \to 1$.

Plan. Use the nonnegative binomial terms in $(1 + s_n)^n$ to obtain an explicit bound on s_n. Then choose a threshold directly in terms of the prescribed positive tolerance.

1. Verify the sign and the defining power identity

Let $u_n = n^{1/n}$ denote the positive real nth root of n. Then $u_n^n = n \geq 1$. If $u_n < 1$, the positivity $0 < u_n < 1$ would give $u_n^n < 1$, a contradiction. Hence $u_n \geq 1$ and

$$s_n = u_n - 1 \geq 0, \qquad (1 + s_n)^n = u_n^n = n.$$

For $n = 1$, this gives $s_1 = 1 - 1 = 0$.

2. Derive the binomial bound for $n \geq 2$

For an integer $n \geq 2$, the binomial theorem gives

$$n = (1 + s_n)^n = 1 + ns_n + \frac{n(n-1)}{2} s_n^2 + \sum_{j=3}^{n} \binom{n}{j} s_n^j.$$

When $n = 2$, the final sum is empty and equals 0. All terms of that sum are nonnegative, since $s_n \geq 0$ and all the binomial coefficients are nonnegative. Also $ns_n \geq 0$, and the constant term 1 is strictly positive. Therefore

$$n > \frac{n(n-1)}{2} s_n^2.$$

Dividing by the positive number n gives $1 > (n-1)s_n^2/2$. Multiplying by $2/(n-1) > 0$ yields

$$s_n^2 < \frac{2}{n-1}.$$

Both s_n and the square root of the right-hand side are nonnegative, so the order of their squares gives

$$0 \le s_n < \sqrt{\frac{2}{n-1}} \qquad (n \ge 2).$$

3. Specify the threshold and verify the limit definition

Let $\varepsilon > 0$ be arbitrary and set

$$N = \max\left\{2,\ 1 + \frac{2}{\varepsilon^2}\right\}.$$

A real threshold is permitted by the definition of convergence; an integer larger than it may also be chosen by the Archimedean property. If $n > N$, then $n \ge 2$ and

$$n - 1 > \frac{2}{\varepsilon^2} > 0.$$

Taking reciprocals of positive quantities and multiplying by 2 gives $2/(n-1) < \varepsilon^2$. As $\varepsilon > 0$, this implies

$$\sqrt{\frac{2}{n-1}} < \varepsilon.$$

Using the bound from Step 2, we conclude

$$|s_n - 0| = s_n < \sqrt{\frac{2}{n-1}} < \varepsilon \qquad (n > N).$$

This is the required proof that $s_n \to 0$.

4. Return to the root sequence

The constant sequence 1 converges to 1, and $s_n \to 0$ was proved without assuming a limit for $n^{1/n}$. The sum rule, Theorem 9.3, therefore gives

$$n^{1/n} = 1 + s_n \longrightarrow 1 + 0 = 1.$$

Conclusion. The explicit estimate $0 \le n^{1/n} - 1 < \sqrt{2/(n-1)}$ for $n \ge 2$, together with $N = \max\{2, 1 + 2/\varepsilon^2\}$, proves $n^{1/n} - 1 \to 0$ and hence $n^{1/n} \to 1$.

9.8 Exercise 9.8: Finite and infinite possibilities for five limits

What we must show. Determine each of the five requested limits, using the convention introduced in Definition 9.8: an existing limit may be a real number, $+\infty$, or $-\infty$. The answer "NOT EXIST" therefore means that none of these three possibilities works.

Plan. Use the defining threshold tests. A sequence has limit $+\infty$ if, for every $M > 0$, all sufficiently late terms exceed M. It has limit $-\infty$ if, for every $M < 0$, all sufficiently late terms are below M.

Neither definition describes finite convergence, and the finite-limit arithmetic laws cannot simply be applied to expressions involving infinity.

(a) The sequence n^3.

Let $M > 0$ be arbitrary. Set $N = \max\{1, M\}$. If $n > N$, then $n \geq 1$ and

$$n^3 = n \cdot n^2 \geq n > M.$$

Thus every prescribed positive threshold is exceeded by all sufficiently late terms, exactly as Definition 9.8 requires.

Conclusion. In part (a), $\lim_{n\to\infty} n^3 = +\infty$.

(b) The sequence $-n^3$.

Let $M < 0$. The number $-M$ is positive. Choose $N = \max\{1, -M\}$. If $n > N$, the estimate used in (a) gives $n^3 \geq n > -M$. Multiplication by -1 reverses the inequality, giving $-n^3 < M$. This is the negative-infinity definition.

Conclusion. In part (b), $\lim_{n\to\infty}(-n^3) = -\infty$.

(c) The sequence $(-n)^n$.

At an even index $n = 2m$ the value is $(2m)^{2m} > 0$; at an odd index $n = 2m+1$ it is $-(2m+1)^{2m+1} < 0$. There are even and odd indices beyond every threshold: choose a positive integer $m > N$ to make both $2m$ and $2m + 1$ exceed a prescribed N.

The limit cannot be $+\infty$, because the test with $M = 1$ would require all sufficiently late terms to be greater than 1, whereas arbitrarily late odd terms are negative. It cannot be $-\infty$, because the test with $M = -1$ would require all sufficiently late terms to be less than -1, whereas arbitrarily late even terms are positive.

It also cannot be a finite real number L. If it converged to L, the definition with $\varepsilon = 1$ would give a threshold N_1 such that

$$n > N_1 \quad \Longrightarrow \quad |(-n)^n - L| < 1.$$

Choose an even positive integer $n > N_1$ with $n > \max\{1, L + 1\}$. Such an index exists by taking $n = 2m$ with m sufficiently large. Since $n \geq 1$, $n^n \geq n$, so at this even index

$$(-n)^n = n^n \geq n > L + 1.$$

Consequently $|(-n)^n - L| = n^n - L > 1$, contradicting the required error bound. All three possible types of limit have now been excluded.

Conclusion. In part (c), the limit does NOT EXIST, even in the extended sense of Definition 9.8.

(d) The sequence $(1.01)^n$.

Write $1.01 = 1 + 1/100$. The binomial theorem, or Bernoulli's inequality, gives

$$\left(1 + \frac{1}{100}\right)^n \geq 1 + \frac{n}{100}.$$

For completeness, the inequality $(1 + x)^n \geq 1 + nx$ for $x \geq 0$ follows by induction: the case $n = 1$ is equality, and from the inequality at n one obtains

$$(1 + x)^{n+1} \geq (1 + nx)(1 + x) = 1 + (n + 1)x + nx^2 \geq 1 + (n + 1)x.$$

Now let $M > 0$ and set $N = 100M$. If $n > N$, then $n/100 > M$, and hence

$$(1.01)^n \geq 1 + \frac{n}{100} > M.$$

This proves divergence to positive infinity by the definition.

Conclusion. In part (d), $\lim_{n\to\infty}(1.01)^n = +\infty$.

(e) The sequence n^n.

For $n \geq 1$, the product of n factors all equal to n satisfies $n^n \geq n$. Let $M > 0$ and set $N = \max\{1, M\}$. If $n > N$, then $n^n \geq n > M$. The definition therefore applies.

Conclusion. In part (e), $\lim_{n\to\infty} n^n = +\infty$.

9.9 Exercise 9.9: Order preserved by existing extended limits

What we must show. Assume there is a threshold N_0 such that $s_n \leq t_n$ whenever $n > N_0$. Prove the three assertions, including every possible finite or infinite case in part (c).

Plan. Combine the threshold at which the comparison becomes valid with the threshold supplied by the relevant limit. In the finite case, two limits in the wrong order would force the late terms to have the wrong order as well.

(a) From $s_n \to +\infty$ to $t_n \to +\infty$.

Fix $M > 0$. Since $s_n \to +\infty$, some N_1 satisfies $s_n > M$ for all $n > N_1$. Set $N = \max\{N_0, N_1\}$. For every $n > N$, both the comparison and this threshold test hold, so

$$t_n \geq s_n > M.$$

This holds for the arbitrary $M > 0$, proving $t_n \to +\infty$.

(b) From $t_n \to -\infty$ to $s_n \to -\infty$.

Fix $M < 0$. Since $t_n \to -\infty$, some N_1 satisfies $t_n < M$ for all $n > N_1$. With $N = \max\{N_0, N_1\}$, every $n > N$ gives

$$s_n \leq t_n < M.$$

This is the definition of $s_n \to -\infty$.

(c) Account for the infinite limit cases.

Write $S = \lim s_n$ and $T = \lim t_n$, which exist by the assumption in this part. Their possible values lie in $\mathbb{R} \cup \{-\infty, +\infty\}$, with the usual extended order. The finite and infinite alternatives cannot overlap. Indeed, convergence to a real c puts the late terms between $c - 1$ and $c + 1$. A simultaneous limit $+\infty$ would put them above $\max\{1, c + 2\}$, contradicting the upper restriction; a simultaneous limit $-\infty$ would put them below $\min\{-1, c - 2\}$, contradicting the lower restriction. Limits of both signs of infinity are incompatible as well, since they would eventually require a term to be both greater than 1 and less than -1. The following cases exhaust all possibilities involving infinity:

1. If $S = -\infty$, then $S \leq T$ for every possible T.

2. If $T = +\infty$, then $S \leq T$ for every possible S.

3. If $S = +\infty$, part (a) forces $T = +\infty$, so $S = T$ and in particular $S \leq T$.

4. If $T = -\infty$, part (b) forces $S = -\infty$, so again $S = T$.

After these cases, both S and T must be finite real numbers.

(c) Prove the finite case by contradiction.

Suppose, contrary to the desired inequality, that $S > T$. Set $\delta = (S-T)/3 > 0$. By finite convergence there are thresholds N_1, N_2 such that

$$n > N_1 \Longrightarrow |s_n - S| < \delta, \qquad n > N_2 \Longrightarrow |t_n - T| < \delta.$$

Thus $s_n > S - \delta$ and $t_n < T + \delta$ at the corresponding late indices. But

$$(S-\delta) - (T+\delta) = (S-T) - 2\delta = (S-T) - \frac{2(S-T)}{3} = \frac{S-T}{3} > 0.$$

For $n > \max\{N_0, N_1, N_2\}$ this gives

$$s_n > S - \delta > T + \delta > t_n,$$

contradicting $s_n \leq t_n$. Hence $S \leq T$ in the finite case as well.

Conclusion. The eventual order implies both infinite-limit comparisons in (a) and (b), and $\lim s_n \leq \lim t_n$ whenever both limits exist in the extended sense.

9.10 Exercise 9.10: Scaling and reversing an infinite limit

What we must show. Prove the effects of multiplying a sequence with limit $+\infty$ by a positive constant or a negative constant, and prove both directions of the sign-reversal equivalence.

Plan. Choose a threshold for s_n by dividing the target threshold by the multiplying constant. A negative constant reverses the inequality, so the signs must be recorded.

(a) Multiplication by $k > 0$.

Assume $s_n \to +\infty$ and fix $k > 0$. To prove $ks_n \to +\infty$, let $M > 0$. Then $M/k > 0$, so the assumed limit supplies N with

$$n > N \quad \Longrightarrow \quad s_n > \frac{M}{k}.$$

Multiplying by $k > 0$ preserves the direction and gives $ks_n > M$. This verifies the target definition.

In fact this implication is reversible. If $ks_n \to +\infty$ and $M > 0$, use the positive threshold kM for ks_n. Eventually $ks_n > kM$, and division by $k > 0$ gives $s_n > M$. Thus positive multiplication preserves the $+\infty$ limit in both directions.

(b) Prove $s_n \to +\infty \Longrightarrow -s_n \to -\infty$.

Let $M < 0$ be an arbitrary threshold for $-s_n$. Because $-M > 0$, the assumption $s_n \to +\infty$ supplies N such that $n > N$ implies $s_n > -M$. Multiplication by -1 reverses the inequality, giving $-s_n < M$. Thus $-s_n \to -\infty$.

(b) Prove the reverse implication.

Assume $-s_n \to -\infty$ and let $M > 0$. The number $-M$ is negative, so some N satisfies $n > N \implies -s_n < -M$. Multiplication by -1 reverses this inequality and gives $s_n > M$. Consequently $s_n \to +\infty$, completing the equivalence.

(c) Multiplication by $k < 0$.

Assume $s_n \to +\infty$ and $k < 0$. Let $M < 0$ be an arbitrary target threshold for ks_n. The quotient M/k is positive, since both numerator and denominator are negative. Hence the assumed limit supplies N such that $n > N$ implies $s_n > M/k$. Multiplying by $k < 0$ reverses the comparison:

$$n > N \quad \implies \quad ks_n < M.$$

This proves $ks_n \to -\infty$.

The corresponding reverse implication also holds. If $ks_n \to -\infty$, fix $M > 0$ and use the negative threshold kM. Eventually $ks_n < kM$, and division by $k < 0$ reverses the inequality, yielding $s_n > M$. Therefore $s_n \to +\infty$.

Conclusion. Positive scaling preserves a positive-infinity limit; negative scaling changes it to negative infinity. In particular, $s_n \to +\infty$ if and only if $-s_n \to -\infty$, with each direction justified by its own threshold test.

9.11 Exercise 9.11: Adding a sequence that is bounded below

What we must show. Prove the sum has limit $+\infty$ in each of the three stated situations. Distinguish a lower bound valid for all terms from a lower bound valid only after an initial segment.

Plan. Once $t_n \geq m$ holds in the tail, it suffices to make $s_n > M - m$. If this number is not positive, replace it by a larger positive threshold before invoking Definition 9.8.

(a) Use the stated global lower bound.

Assume $s_n \to +\infty$ and $\inf\{t_n : n \in \mathbb{N}\} > -\infty$. The set is nonempty and consists of real numbers; its infimum is at most the finite member t_1. Thus this hypothesis provides a real lower bound m, for example the real infimum itself, such that $t_n \geq m$ for every n.

Let $M > 0$ be arbitrary, and define

$$B = \max\{1, M - m\} > 0.$$

Since $s_n \to +\infty$, there is N such that $n > N$ implies $s_n > B$. For those indices,

$$s_n + t_n \geq s_n + m > B + m \geq (M - m) + m = M.$$

The first inequality uses the global lower bound; the last uses $B \geq M - m$. This proves the positive-infinity limit.

(b) Derive an eventual lower bound in every allowed limit case.

Assume $s_n \to +\infty$ and $T = \lim t_n > -\infty$. The existing limit T is either a real number or $+\infty$.

- If $T \in \mathbb{R}$, apply finite convergence with $\varepsilon = 1$. For some N_1, every $n > N_1$ has $|t_n - T| < 1$, implying $t_n > T - 1$. Thus we may take $m = T - 1$ as an eventual lower bound.

- If $T = +\infty$, use the positive threshold 1 in Definition 9.8. For some N_1, every $n > N_1$ has $t_n > 1$. Thus we may take $m = 1$.

In either case there are a real m and a threshold N_1 such that $t_n \geq m$ whenever $n > N_1$. This is a tail bound; no assertion about the earlier terms is needed.

Now fix $M > 0$. Set $B = \max\{1, M - m\} > 0$, and use $s_n \to +\infty$ to choose N_2 such that $s_n > B$ for $n > N_2$. For $n > \max\{N_1, N_2\}$, both estimates hold and

$$s_n + t_n \geq s_n + m > B + m \geq M.$$

This proves the desired limit using the eventual bound directly, without applying a global hypothesis to an unexamined prefix.

(c) A bounded sequence is bounded below.

If (t_n) is bounded, there is a real $C \geq 0$ with $|t_n| \leq C$ for every n. Therefore $t_n \geq -C$ for every n, a global lower bound, and part (a) applies. One can also see the threshold explicitly: for $M > 0$, choose N such that $s_n > M + C$ whenever $n > N$; this is a valid positive threshold because $M + C > 0$. Then

$$s_n + t_n \geq s_n - C > M + C - C = M.$$

Conclusion. In (a), (b), and (c), the sum $s_n + t_n$ tends to $+\infty$. An eventual lower bound on the added sequence already suffices, as the proof of (b) shows.

9.12 Exercise 9.12: The ratio test for the size of a sequence

What we must show. Suppose $s_n \neq 0$ for every n, and the limit

$$L = \lim_{n\to\infty} r_n, \qquad r_n = \left|\frac{s_{n+1}}{s_n}\right|,$$

exists. Prove $s_n \to 0$ if $L < 1$, and $|s_n| \to +\infty$ if $L > 1$, including the case $L = +\infty$.

Plan. A ratio eventually less than a fixed $q < 1$ gives a geometric bound. For ratios tending to a value larger than 1, apply that first result to the reciprocal magnitudes.

First locate the possible values of L.

Every r_n is positive, since its numerator and denominator are nonzero in absolute value. Therefore its existing limit can only be a nonnegative real number or $+\infty$. For example, if a finite limit were $L < 0$, convergence with $\varepsilon = -L/2 > 0$ would eventually give $r_n < L + (-L/2) = L/2 < 0$, a contradiction. A limit $-\infty$ would eventually give $r_n < -1$ by its definition, also a contradiction. This verifies the claimed restriction on L.

(a) Choose a single ratio bound valid at every sufficiently late index.

Suppose $L < 1$. Then $0 \leq L < 1$, so

$$q = \frac{L+1}{2} \quad \text{satisfies} \quad 0 < q < 1 \quad \text{and} \quad L < q.$$

By convergence of r_n to the finite number L, some threshold N_0 gives $|r_n - L| < q - L$ for all $n > N_0$. In particular $r_n < q$ at all those indices. Choose an integer $N \in \mathbb{N}$ with $N > N_0$. Then every integer $n \geq N$ has $n > N_0$, so

$$|s_{n+1}| = r_n \, |s_n| < q \, |s_n| \qquad (n \geq N).$$

The integer choice makes the starting index and the validity of every subsequent ratio inequality explicit.

(a) Prove the geometric estimate by induction.

For $n = N + 1$, the preceding inequality gives

$$|s_{N+1}| < q \, |s_N| = q^{(N+1)-N} \, |s_N| \, .$$

Assume for some integer $n > N$ that $|s_n| < q^{n-N} \, |s_N|$. Since the ratio estimate also applies at this n, multiplication by $q > 0$ gives

$$|s_{n+1}| < q \, |s_n| < q \, q^{n-N} \, |s_N| = q^{(n+1)-N} \, |s_N| \, .$$

Induction proves

$$|s_n| < q^{n-N} \, |s_N| \qquad (n > N).$$

Since $0 < q < 1$, Theorem 9.7(b) gives $q^m \to 0$ as $m \to \infty$. To spell out the use of this fact, let $\varepsilon > 0$. The number $C = |s_N|$ is positive, because $s_N \neq 0$. Choose a threshold K such that $m > K$ implies $q^m < \varepsilon / C$. If $n > N + \max\{K, 0\}$, then $m = n - N > K$ and $n > N$, so

$$|s_n - 0| < q^{n-N} C < \frac{\varepsilon}{C} C = \varepsilon.$$

This is precisely finite convergence to 0.

(b) Form the reciprocal sequence and compute its ratio.

Suppose $L > 1$, and define

$$t_n = \frac{1}{|s_n|} > 0.$$

It is defined and nonzero for every n. Its absolute ratio is

$$\begin{aligned} \left| \frac{t_{n+1}}{t_n} \right| &= \frac{1/\,|s_{n+1}|}{1/\,|s_n|} \\ &= \frac{|s_n|}{|s_{n+1}|} = \frac{1}{r_n}. \end{aligned}$$

There are two distinct cases for the ratio limit:

- If $1 < L < +\infty$, the finite reciprocal law applies to $r_n \to L$, since every $r_n \neq 0$ and $L \neq 0$. It gives $1/r_n \to 1/L$, with $0 < 1/L < 1$.
- If $L = +\infty$, every r_n is positive, so Theorem 9.10 gives $1/r_n \to 0$. This conclusion uses the infinite-limit reciprocal theorem, not the finite quotient law with an infinite denominator.

In either case, the absolute ratio for t_n has a finite limit strictly less than 1. Part (a) therefore proves $t_n \to 0$. Finally apply Theorem 9.10 to the positive sequence $|s_n|$: its reciprocal is $t_n \to 0$, so

$$|s_n| \longrightarrow +\infty.$$

Conclusion. If $L < 1$, the sequence itself converges to 0. If $L > 1$, including $L = +\infty$, its magnitudes tend to $+\infty$. No conclusion for the boundary value $L = 1$ is asserted by either part.

9.13 Exercise 9.13: All real bases in a geometric sequence

What we must show. Prove the complete classification of a^n as the fixed real base a varies, including $a = 0$, $a = 1$, $a = -1$, and both intervals of negative bases.

Plan. Use Exercise 9.12 where the terms are nonzero. For a negative base with magnitude at least 1, check that no finite limit and neither infinite limit can accommodate the alternating signs.

Case 1: $a = 0$, and then $0 < |a| < 1$.

If $a = 0$, every term a^n is 0 because $n \geq 1$. Thus the sequence converges to 0 directly. If $0 < |a| < 1$, every term is nonzero and

$$\left|\frac{a^{n+1}}{a^n}\right| = |a| < 1.$$

This constant ratio has limit $|a|$, so Exercise 9.12(a) proves $a^n \to 0$. The two cases together cover all $|a| < 1$, including negative a of magnitude less than 1.

Case 2: $a = 1$.

Every term equals $1^n = 1$. Thus for every $\varepsilon > 0$, $|a^n - 1| = 0 < \varepsilon$ at every index, giving limit 1.

Case 3: $a > 1$.

The terms a^n are positive and nonzero. Their absolute ratio is the constant $a > 1$. Exercise 9.12(b) gives $|a^n| \to +\infty$. Since $a^n > 0$, $|a^n| = a^n$, so the sequence itself tends to $+\infty$.

Case 4: $a = -1$.

The terms are 1 at even indices and -1 at odd indices. If there were a finite limit L, convergence with $\varepsilon = 1/2$ would put every sufficiently late term within $1/2$ of L. Choose a late even index p and a late odd index q. The triangle inequality would give

$$2 = |1 - (-1)| \leq |a^p - L| + |L - a^q| < \frac{1}{2} + \frac{1}{2} = 1,$$

a contradiction. A limit $+\infty$ fails the threshold $M = 1$ because late odd terms equal -1. A limit $-\infty$ fails the threshold $M = -1$ because late even terms equal 1. Thus no extended limit exists.

Case 5: $a < -1$.

Set $b = |a| = -a > 1$. By Case 3, $b^n \to +\infty$. The terms satisfy

$$a^n = (-1)^n b^n, \qquad |a^n| = b^n.$$

Suppose first that a^n had a finite limit L. Convergence with $\varepsilon = 1$ would give, eventually,

$$|a^n| \leq |a^n - L| + |L| < 1 + |L|\,.$$

But $b^n \to +\infty$ means that eventually $b^n > 1 + |L|$, contradicting $|a^n| = b^n$. Thus there is no finite limit.

There is also no limit $+\infty$, since for every threshold index there is a later odd index with $a^n < 0$, defeating the positive threshold $M = 1$. There is no limit $-\infty$, since there is always a later even index with $a^n > 0$, defeating the negative threshold $M = -1$. The required even and odd indices can be taken as $2m$ and $2m + 1$, where m is any positive integer larger than the proposed threshold index.

Conclusion. For every real a,

$$\lim_{n\to\infty} a^n = \begin{cases} 0, & |a| < 1, \\ 1, & a = 1, \\ +\infty, & a > 1, \\ \text{does not exist}, & a \le -1. \end{cases}$$

The last line excludes finite limits and both infinite limits.

9.14 Exercise 9.14: Geometric growth compared with a real power

What we must show. Fix $p > 0$ and determine the limit of a^n/n^p for every real a, using Exercise 9.12 where its hypotheses apply. In particular, include both boundary bases $a = 1$ and $a = -1$.

Plan. The absolute ratio is $|a|\ (n/(n+1))^p$. Prove its second factor tends to 1, then distinguish ratios below 1, above 1, and the boundary cases where the ratio test does not decide the answer.

1. State the real-power facts and justify the needed limit.

As in Theorem 9.7(a), positive bases with real exponents use the usual real-power algebra and order: products and quotients obey the exponent rules; and $r^v \le r^u$ whenever $0 < r < 1$ and $0 < u < v$. These are the elementary real-power properties already implicit in the statement of Theorem 9.7(a). We do not assume a continuity theorem for real powers from a later section.

Set $r_n = n/(n+1)$, so $0 < r_n < 1$. Choose an integer $m > p$, possible by the Archimedean property. Then

$$r_n^m \le r_n^p \le 1, \qquad 0 \le 1 - r_n^p \le 1 - r_n^m.$$

For the integer exponent m, finite multiplication gives the factorization

$$1 - r_n^m = (1 - r_n)(1 + r_n + \cdots + r_n^{m-1}).$$

Every one of the m terms in the parenthesized sum is at most 1, and $1 - r_n = 1/(n+1)$. Therefore

$$0 \le 1 - r_n^p \le 1 - r_n^m \le \frac{m}{n+1}.$$

The final expression tends to 0: for $\varepsilon > 0$, $n > m/\varepsilon$ makes $m/(n+1) < \varepsilon$. This direct error estimate proves

$$\left(\frac{n}{n+1}\right)^p \longrightarrow 1$$

for the possibly nonintegral exponent $p > 0$.

2. Compute the ratio when $a \neq 0$.

Let $u_n = a^n/n^p$. Since $n > 0$, its denominator is positive. If $a \neq 0$, all u_n are nonzero, and the real-power quotient rule yields

$$\begin{aligned} \left|\frac{u_{n+1}}{u_n}\right| &= \left|\frac{a^{n+1}}{(n+1)^p} \cdot \frac{n^p}{a^n}\right| \\ &= |a|\ \frac{n^p}{(n+1)^p} \end{aligned}$$

$$= |a| \left(\frac{n}{n+1} \right)^p.$$

Step 1 and the finite scalar-multiple limit theorem show that this ratio has limit $|a|$.

3. Treat $a = 0$ and the cases $0 < |a| < 1$.

For $a = 0$, every numerator is 0 and every denominator is positive, so $u_n = 0$ and the limit is 0. For $0 < |a| < 1$, Step 2 gives an absolute ratio limit less than 1, so Exercise 9.12(a) applies and gives $u_n \to 0$.

4. Treat the two boundary cases $|a| = 1$.

If $a = 1$ or $a = -1$, then $|a^n| = 1$, and hence

$$|u_n - 0| = \frac{1}{n^p}.$$

Theorem 9.7(a), whose hypothesis is precisely $p > 0$, says $1/n^p \to 0$. Thus $u_n \to 0$ in both boundary cases. The ratio limit here is 1; the conclusion comes from the direct error bound rather than from an inapplicable strict case of Exercise 9.12.

5. Treat $a > 1$.

The absolute ratio limit is $a > 1$, so Exercise 9.12(b) gives $|u_n| \to +\infty$. Since $a^n > 0$ and $n^p > 0$, the terms themselves are positive: $u_n = |u_n|$. Consequently $u_n \to +\infty$.

6. Treat $a < -1$ and exclude all three types of limit.

Exercise 9.12(b) gives $|u_n| \to +\infty$, since the absolute ratio limit is $|a| > 1$. If u_n converged to a finite L, then eventually

$$|u_n| \leq |u_n - L| + |L| < 1 + |L|$$

by the error bound $|u_n - L| < 1$. This contradicts $|u_n| \to +\infty$. At even indices $u_n > 0$, while at odd indices $u_n < 0$. Every tail has an odd index, so the positive-infinity test with $M = 1$ fails. Every tail also has an even index, so the negative-infinity test with $M = -1$ fails. Thus no extended limit exists in this last case.

Conclusion. For $p > 0$ and real a,

$$\lim_{n\to\infty} \frac{a^n}{n^p} = \begin{cases} 0, & |a| \leq 1, \\ +\infty, & a > 1, \\ \text{does not exist}, & a < -1. \end{cases}$$

In particular, the base $a = -1$ now gives limit 0, unlike the unscaled sequence in Exercise 9.13.

9.15 Exercise 9.15: Every fixed geometric sequence is dominated by a factorial

What we must show. Prove $a^n/n! \to 0$ for every $a \in \mathbb{R}$, including zero and negative values.

Plan. After the zero-base case is separated, the absolute ratio is $|a|/(n+1)$, which tends to 0. This gives the strict ratio condition from Exercise 9.12(a).

1. Handle $a = 0$ without dividing by a zero term.

When $a = 0$ and $n \geq 1$, $a^n = 0$. Since $n! > 0$, every term is 0, and the limit is 0.

2. Compute the absolute ratio for $a \neq 0$.

Fix a nonzero real a and write $s_n = a^n/n!$. Then every s_n is nonzero. By the factorial identity $(n+1)! = (n+1)n!$,

$$\begin{aligned}\left|\frac{s_{n+1}}{s_n}\right| &= \left|\frac{a^{n+1}}{(n+1)!} \cdot \frac{n!}{a^n}\right| \\ &= \left|\frac{a\, n!}{(n+1)n!}\right| \\ &= \frac{|a|}{n+1}.\end{aligned}$$

For $\varepsilon > 0$, choosing $N > |a|/\varepsilon$ makes $n > N$ imply $|a|/(n+1) < \varepsilon$. Thus the ratio limit is $L = 0 < 1$. Exercise 9.12(a) now proves $s_n \to 0$. Its hypothesis is about absolute ratios, so this argument applies to negative a as well as positive a.

Conclusion. For every real a, $\lim_{n\to\infty} a^n/n! = 0$.

9.16 Exercise 9.16: Three applications of the infinite-limit results

What we must show. Establish positive-infinity limits for all three given expressions, using the proved infinite-limit theorems or Exercises 9.9–9.15 with their hypotheses checked.

Plan. In (a), isolate a factor with limit $+\infty$ and another with a positive finite limit. In (b) and (c), use Exercise 9.14 for the growing term and then control the added term from below.

(a) The expression $(n^4 + 8n)/(n^2 + 9)$.

Every denominator $n^2 + 9$ is positive. Factor the expression exactly as

$$\frac{n^4 + 8n}{n^2 + 9} = n^2\,\frac{1 + 8/n^3}{1 + 9/n^2}.$$

Indeed, multiplying the numerator and denominator of the right-hand fraction by n^2 gives

$$n^2\,\frac{n^2 + 8/n}{n^2 + 9} = \frac{n^4 + 8n}{n^2 + 9}.$$

Theorem 9.7(a) gives $1/n^3 \to 0$ and $1/n^2 \to 0$. The finite scalar and sum laws therefore give $1 + 8/n^3 \to 1$ and $1 + 9/n^2 \to 1$. All the latter denominators are positive, and the limit denominator is $1 \neq 0$, so the finite quotient theorem applies to this fraction and gives

$$v_n = \frac{1 + 8/n^3}{1 + 9/n^2} \longrightarrow 1 > 0.$$

Also $n^2 \to +\infty$: for each $M > 0$, every $n > \sqrt{M}$ has $n^2 > M$. Thus Theorem 9.9 applies to $s_n = n^2$ and $t_n = v_n$, since its first limit is $+\infty$ and its second limit is the positive number 1. It gives the desired product limit.

Conclusion. In part (a), $\lim_{n\to\infty} \dfrac{n^4 + 8n}{n^2 + 9} = +\infty$.

(b) The expression $2^n/n^2 + (-1)^n$.

Exercise 9.14 with $a = 2 > 1$ and $p = 2 > 0$ gives

$$u_n = \frac{2^n}{n^2} \longrightarrow +\infty.$$

The second sequence is bounded: $-1 \leq (-1)^n \leq 1$ for every n. Exercise 9.11(c) therefore applies to the sum. In explicit threshold form, given $M > 0$, choose N so that $n > N$ implies $u_n > M + 1$. Then

$$\frac{2^n}{n^2} + (-1)^n \geq u_n - 1 > M.$$

This verifies the limit without requiring the oscillating sequence $(-1)^n$ itself to have a limit.

Conclusion. In part (b), $\lim_{n\to\infty} \left[\frac{2^n}{n^2} + (-1)^n\right] = +\infty$.

(c) The expression $3^n/n^3 - 3^n/n!$.

Exercise 9.14 with $a = 3$, $p = 3$ and Exercise 9.15 with $a = 3$ give, respectively,

$$u_n = \frac{3^n}{n^3} \longrightarrow +\infty, \qquad v_n = \frac{3^n}{n!} \longrightarrow 0.$$

Thus $-v_n \to 0$ by the finite scalar-multiple theorem. Exercise 9.11(b) applies to $u_n + (-v_n)$, since the second limit 0 is greater than $-\infty$. To verify the threshold directly, fix $M > 0$. Choose N_1 such that $n > N_1$ implies $u_n > M + 1$. Choose N_2 such that $n > N_2$ implies $|v_n| < 1$, using its convergence to 0. For $n > \max\{N_1, N_2\}$ we have $v_n < 1$ and hence

$$u_n - v_n > u_n - 1 > M.$$

This is a growing term minus a term tending to 0; the proof does not subtract two infinite limits.

Conclusion. In part (c), $\lim_{n\to\infty} \left[\frac{3^n}{n^3} - \frac{3^n}{n!}\right] = +\infty$.

9.17 Exercise 9.17: A definition-only proof for the sequence of squares

What we must show. Use only Definition 9.8 to prove $n^2 \to +\infty$.

Plan. Given the positive height M, solve $n^2 > M$ by requiring $n > \sqrt{M}$. Both sides are positive, so squaring this comparison preserves its direction.

Formal proof with the quantifiers explicit.

Let $M > 0$ be arbitrary, and set $N = \sqrt{M}$. If $n \in \mathbb{N}$ and $n > N$, then $n > \sqrt{M} > 0$. Multiplying positive inequalities, or using the strictly increasing square function on the positive numbers, gives

$$n^2 > (\sqrt{M})^2 = M.$$

More explicitly, $n > \sqrt{M}$ gives $n^2 > n\sqrt{M} > (\sqrt{M})^2$ after multiplication first by $n > 0$ and then by $\sqrt{M} > 0$. We have therefore proved that for every $M > 0$ there is $N = \sqrt{M}$ such that every positive integer $n > N$ satisfies $n^2 > M$. This is exactly Definition 9.8.

Conclusion. $\lim_{n\to\infty} n^2 = +\infty$, with the explicit threshold $N = \sqrt{M}$ for each $M > 0$.

9.18 Exercise 9.18: Finite geometric sums and their limits

What we must show. Prove the finite-sum identity in (a), then determine the limits in (b), (c), and (d), including the base $a = 1$ where the identity's quotient is not defined.

Plan. Subtract a times the finite sum from the original sum. Every interior power cancels. After the identity is proved, use the established geometric-sequence limit for $|a| < 1$, and a direct lower comparison for $a \geq 1$.

(a) Prove the identity for every allowed base and every index.

Fix a real $a \neq 1$ and a positive integer n, and define

$$S_n = 1 + a + a^2 + \cdots + a^n = \sum_{k=0}^{n} a^k.$$

Here a^0 denotes the constant empty product 1, including when $a = 0$. Multiplication and reindexing a finite sum give

$$aS_n = \sum_{k=0}^{n} a^{k+1} = \sum_{j=1}^{n+1} a^j = a + a^2 + \cdots + a^n + a^{n+1}.$$

The reindexing sets $j = k + 1$, so the lower bound $k = 0$ becomes $j = 1$ and the upper bound $k = n$ becomes $j = n + 1$. Subtracting the two finite sums cancels exactly the terms $a, a^2, \ldots, a^n$:

$$\begin{aligned} S_n - aS_n &= (1 + a + \cdots + a^n) - (a + \cdots + a^n + a^{n+1}) \\ &= 1 - a^{n+1}. \end{aligned}$$

The left side factors as $(1 - a)S_n$. Since $a \neq 1$, $1 - a \neq 0$, and division by this nonzero number yields

$$S_n = \frac{1 - a^{n+1}}{1 - a}.$$

The base and index were arbitrary, so this proves the identity for every real $a \neq 1$ and every positive integer n. The cancellation also works for $n = 0$, if that index is included, since then the identity is $1 = (1 - a)/(1 - a)$.

(b) Pass to the limit for $|a| < 1$.

Now fix a with $|a| < 1$. Then $a \neq 1$, so (a) applies. Exercise 9.13 gives $a^n \to 0$. Since $a^{n+1} = a\,a^n$, multiplication by the fixed real number a gives $a^{n+1} \to a \cdot 0 = 0$ by the finite scalar-multiple theorem. The denominator $1 - a$ is a fixed nonzero real number. Consequently the finite sum and quotient laws applied to the identity give

$$\lim_{n\to\infty} S_n = \frac{1 - 0}{1 - a} = \frac{1}{1 - a}.$$

One can read the same conclusion directly as an error estimate, avoiding any ambiguity about division:

$$\begin{aligned} \left| S_n - \frac{1}{1 - a} \right| &= \left| \frac{1 - a^{n+1} - 1}{1 - a} \right| \\ &= \frac{|a|^{n+1}}{|1 - a|} \longrightarrow 0. \end{aligned}$$

The denominator in this expression is positive and constant; the numerator tends to 0. This includes $a = 0$, when every sum is exactly 1.

Conclusion. For part (b), if $|a| < 1$, then $\lim_{n\to\infty}(1 + a + \cdots + a^n) = 1/(1 - a)$.

(c) Substitute $a = 1/3$ and simplify the fraction.

The displayed sequence of sums is

$$1 + \frac{1}{3} + \frac{1}{9} + \frac{1}{27} + \cdots + \frac{1}{3^n} = \sum_{k=0}^{n} \left(\frac{1}{3}\right)^k.$$

Since $0 < 1/3 < 1$, part (b) applies. Its limit is

$$\frac{1}{1 - 1/3} = \frac{1}{(3 - 1)/3} = \frac{1}{2/3} = \frac{3}{2}.$$

Conclusion. The limit in part (c) is $3/2$.

(d) Handle $a = 1$ and $a > 1$ explicitly.

If $a = 1$, the sum contains $n + 1$ terms, each equal to 1, so $S_n = n + 1$. For an arbitrary $M > 0$, $n > M$ implies $S_n = n + 1 > M$. Therefore $S_n \to +\infty$ in this case. The quotient in (a) was not used at $a = 1$, where its denominator would be zero.

If $a > 1$, every term a^k for $k = 0, 1, \ldots, n$ is at least 1: the exponent 0 gives 1, and a positive exponent gives a product of factors greater than 1. Adding the $n + 1$ inequalities gives

$$S_n = \sum_{k=0}^{n} a^k \geq \sum_{k=0}^{n} 1 = n + 1.$$

Since $n + 1 \to +\infty$, Exercise 9.9(a) proves $S_n \to +\infty$. Equivalently, the same threshold $N = M$ works directly: $n > N$ implies $S_n \geq n + 1 > M$.

Conclusion. For part (d), every base $a \geq 1$ gives $\lim_{n\to\infty}(1 + a + \cdots + a^n) = +\infty$.

10 Monotone sequences and Cauchy sequences

An increasing sequence satisfies $s_n \le s_{n+1}$; a decreasing sequence satisfies $s_n \ge s_{n+1}$. Both definitions permit equality. A sequence is bounded if all its terms lie between two fixed real numbers.

The Cauchy condition says that, for every $\varepsilon > 0$, there is one threshold N such that

$$m > N,\ n > N \quad \Longrightarrow \quad |s_m - s_n| < \varepsilon.$$

It controls every pair of late terms. Merely making successive differences small is not enough, as Exercise 10.6 explains. The solutions first justify convergence and then identify a limit.

Conventions for monotonicity and Cauchy estimates

In Definition 10.1, "increasing" means $s_n \le s_{n+1}$ and "decreasing" means $s_n \ge s_{n+1}$. Equality is permitted in both definitions. A constant sequence therefore has both properties. Repeated application of an increasing inequality gives $s_n \le s_m$ whenever $n < m$; for a decreasing sequence it gives $s_n \ge s_m$.

Definition 10.8 says that (s_n) is Cauchy if

> for every $\varepsilon > 0$ there is an N such that every pair of integers $m, n > N$ satisfies $|s_m - s_n| < \varepsilon$.

Both indices are arbitrary after the same threshold. An estimate for consecutive terms alone is not yet the estimate for an arbitrary pair.

10.1 Exercise 10.1: Classifying six sequences

What we must show. Decide whether each sequence is increasing, decreasing, and bounded, with the weak monotonicity conventions just stated.

Part (a): the sequence $s_n = 1/n$.

For every $n \in \mathbb{N}$, the denominators n and $n+1$ are positive, and

$$s_{n+1} - s_n = \frac{1}{n+1} - \frac{1}{n} = \frac{n-(n+1)}{n(n+1)} = -\frac{1}{n(n+1)} < 0.$$

Thus $s_{n+1} < s_n$ for every n, so the sequence is decreasing. It is not increasing: for example, $s_2 = 1/2 < 1 = s_1$. Because $n \ge 1$, we also have $0 < 1/n \le 1$, so 0 and 1 are a lower and an upper bound.

Conclusion. The sequence $1/n$ is **decreasing and bounded**, and is not increasing.

Part (b), Step 1: test monotonicity with exact terms.

Here $s_n = (-1)^n/n^2$. Its first three terms are

$$s_1 = -1, \qquad s_2 = \frac{1}{4}, \qquad s_3 = -\frac{1}{9}.$$

The inequality $s_1 < s_2$ rules out a decreasing sequence, which would require $s_1 \ge s_2$. The inequality $s_2 > s_3$ rules out an increasing sequence, which would require $s_2 \le s_3$. Hence it is neither.

Part (b), Step 2: give bounds valid at every index.

If n is odd, then $s_n = -1/n^2$, so $-1 \le s_n < 0$. If n is even, then $n \ge 2$ and $s_n = 1/n^2$, so $0 < s_n \le 1/4$. Both cases give

$$-1 \le s_n \le \frac{1}{4} \qquad (n \in \mathbb{N}).$$

These bounds are attained at $n = 1$ and $n = 2$, respectively.

Conclusion. The sequence $(-1)^n/n^2$ is **bounded but neither increasing nor decreasing**.

Part (c), Step 1: check the growth of $s_n = n^5$.

For every $n \ge 1$, the binomial expansion gives

$$\begin{aligned} s_{n+1} - s_n &= (n+1)^5 - n^5, \\ &= n^5 + 5n^4 + 10n^3 + 10n^2 + 5n + 1 - n^5, \\ &= 5n^4 + 10n^3 + 10n^2 + 5n + 1 > 0. \end{aligned}$$

All five terms in the final sum are positive. Thus $s_{n+1} > s_n$ for every index, giving an increasing sequence and ruling out a decreasing one.

Part (c), Step 2: prove unboundedness above.

Since $n \ge 1$, we have $n^4 \ge 1$, so $n^5 \ge n \ge 1$. The number 1 is a lower bound. To show there is no upper bound, let $U \in \mathbb{R}$ be any proposed upper bound. By the Archimedean property, choose $n \in \mathbb{N}$ with $n > \max\{1, U\}$. Then

$$s_n = n^5 \ge n > U.$$

Thus U fails to bound the sequence above. This works for every real candidate U.

Conclusion. The sequence n^5 is **increasing and unbounded above**. It is not decreasing or bounded, although it is bounded below by 1.

Part (d), Step 1: give a rise and a fall for the sine sequence.

Let $s_n = \sin(n\pi/7)$. Put $c = \sin(\pi/7)$. Since $0 < \pi/7 < \pi$, the elementary sign property of sine gives $c > 0$. The sine values at three indices are

$$s_1 = c > 0, \qquad s_7 = \sin \pi = 0, \qquad s_{15} = \sin(2\pi + \pi/7) = c > 0.$$

An increasing sequence would satisfy $s_1 \le s_7$, contradicting $s_1 > 0 = s_7$. A decreasing sequence would satisfy $s_7 \ge s_{15}$, contradicting $s_7 = 0 < s_{15}$. Comparisons at nonadjacent indices are legitimate because monotonicity implies the corresponding comparison at any pair of ordered indices.

Part (d), Step 2: give global bounds.

The elementary trigonometric inequality $-1 \le \sin x \le 1$ is valid for every real x. Applying it at $x = n\pi/7$ gives $-1 \le s_n \le 1$ for every n.

Conclusion. The sequence $\sin(n\pi/7)$ is **bounded but neither increasing nor decreasing**.

Part (e), Step 1: rule out both monotonicity directions.

For $s_n = (-2)^n$, the first three terms are

$$s_1 = -2, \qquad s_2 = 4, \qquad s_3 = -8.$$

The rise from -2 to 4 rules out a decreasing sequence. The fall from 4 to -8 rules out an increasing sequence.

Part (e), Step 2: record an elementary power estimate.

We will use

$$2^j \geq j+1 \qquad (j \in \mathbb{N}).$$

For $j = 1$, this is $2 \geq 2$. If it holds for $j \geq 1$, then

$$2^{j+1} = 2 \cdot 2^j \geq 2(j+1) = 2j+2 \geq j+2,$$

since $(2j+2) - (j+2) = j \geq 0$. Induction proves the estimate.

Part (e), Step 3: show that both signs have arbitrarily large magnitude.

Let $R > 0$ be arbitrary. Choose $k \in \mathbb{N}$ with $k > R$ by the Archimedean property. At an even index,

$$s_{2k} = 2^{2k} \geq 2k+1 > R.$$

At the following odd index,

$$s_{2k+1} = -2^{2k+1} \leq -(2k+2) < -R.$$

Thus there are terms above R and below $-R$ for every positive R. In particular, no real number is an upper bound and no real number is a lower bound.

Conclusion. The sequence $(-2)^n$ is **neither increasing nor decreasing, and is unbounded both above and below**.

Part (f), Step 1: compare consecutive positive terms.

Here $s_n = n/3^n > 0$. Since we are dividing positive quantities, we can compute the consecutive-term ratio:

$$\begin{aligned} \frac{s_{n+1}}{s_n} &= \frac{n+1}{3^{n+1}} \frac{3^n}{n}, \\ &= \frac{n+1}{3n}. \end{aligned}$$

For $n \geq 1$, the denominator is positive and $3n-(n+1) = 2n-1 \geq 1 > 0$. Hence $0 < (n+1)/(3n) < 1$. Multiplying this strict upper inequality by $s_n > 0$ gives $s_{n+1} < s_n$.

Part (f), Step 2: bound the whole sequence.

Every term is positive. Repeated application of the decreasing inequality gives $s_n \leq s_1 = 1/3$. Therefore

$$0 < s_n \leq \frac{1}{3} \qquad (n \in \mathbb{N}).$$

The strict decrease also rules out an increasing sequence.

Conclusion. The sequence $n/3^n$ is **decreasing and bounded**, and is not increasing.

Summary of all six classifications.

Sequence	Increasing?	Decreasing?	Bounded?
$1/n$	No	Yes	Yes
$(-1)^n/n^2$	No	No	Yes
n^5	Yes	No	No
$\sin(n\pi/7)$	No	No	Yes
$(-2)^n$	No	No	No
$n/3^n$	No	Yes	Yes

10.2 Exercise 10.2: Convergence of a bounded decreasing sequence

What we must show. Prove the decreasing case of Theorem 10.2: every bounded decreasing real sequence converges.

Step 1: define the candidate limit and justify its existence.

Let (s_n) be bounded and decreasing, and let

$$S = \{s_n : n \in \mathbb{N}\}.$$

The set is nonempty because $s_1 \in S$, and it is bounded because the sequence is bounded. In particular, it is bounded below. The lower-bound form of completeness gives a real number

$$L = \inf S.$$

Being a lower bound means $L \leq s_n$ for every n.

Step 2: obtain one term close to the infimum.

Let $\varepsilon > 0$. The number $L + \varepsilon$ is strictly larger than the greatest lower bound L, so it cannot itself be a lower bound of S. Therefore there is an element of S below $L + \varepsilon$. By the definition of S, that element is s_N for some index $N \in \mathbb{N}$. Thus

$$L \leq s_N < L + \varepsilon.$$

Step 3: use decrease to control every subsequent term.

Take any integer $n > N$. Since the sequence is decreasing, $s_n \leq s_N$. Since L is a lower bound, $L \leq s_n$. Combining these facts with Step 2 gives

$$L \leq s_n \leq s_N < L + \varepsilon.$$

Subtract L throughout to get $0 \leq s_n - L < \varepsilon$. The difference $s_n - L$ is nonnegative, so

$$|s_n - L| = s_n - L < \varepsilon \qquad \text{for every integer } n > N.$$

Conclusion. For every $\varepsilon > 0$ we obtained the threshold required by Definition 7.1. Hence $\boxed{s_n \longrightarrow \inf\{s_n : n \in \mathbb{N}\}}$. The proof uses the greatest-lower-bound property, without assuming the infimum is one of the terms.

10.3 Exercise 10.3: A finite bound for decimal approximations

What we must show. Let K be a nonnegative integer and let each digit d_j belong to $\{0, 1, \ldots, 9\}$. For

$$s_n = K + \sum_{j=1}^{n} \frac{d_j}{10^j},$$

prove $s_n < K + 1$ for every $n \in \mathbb{N}$.

Step 1: bound each digit contribution.

For every $j \geq 1$, $0 \leq d_j \leq 9$ and $10^j > 0$. Division by the positive denominator gives

$$0 \leq \frac{d_j}{10^j} \leq \frac{9}{10^j}.$$

Add these finitely many inequalities and then add K:

$$K \leq s_n \leq K + \sum_{j=1}^{n} \frac{9}{10^j}.$$

Step 2: evaluate the finite upper-bound sum by cancellation.

For an individual index j,

$$\frac{9}{10^j} = \frac{10-1}{10^j} = \frac{1}{10^{j-1}} - \frac{1}{10^j}.$$

Consequently

$$\begin{aligned} \sum_{j=1}^{n} \frac{9}{10^j} &= \left(1 - \frac{1}{10}\right) + \left(\frac{1}{10} - \frac{1}{10^2}\right) + \cdots + \left(\frac{1}{10^{n-1}} - \frac{1}{10^n}\right), \\ &= 1 - \frac{1}{10^n}. \end{aligned}$$

Every intermediate negative term cancels the identical positive term in the next parentheses. This is a finite algebraic identity; no limit or infinite sum has been used to derive it.

Step 3: preserve the strict final inequality.

Substitute the identity from Step 2 into Step 1:

$$K \leq s_n \leq K + 1 - \frac{1}{10^n}.$$

Since $10^n > 0$, its reciprocal is strictly positive. Subtracting that positive number makes $K+1-1/10^n$ strictly less than $K + 1$. Thus

$$s_n \leq K + 1 - \frac{1}{10^n} < K + 1.$$

Conclusion. Every finite decimal approximation satisfies $\boxed{s_n < K + 1}$. Moreover $s_{n+1} - s_n = d_{n+1}/10^{n+1} \geq 0$, so the approximations are increasing and bounded. Their convergence follows from Theorem 10.2 *after* the finite bound has been established.

10.4 Exercise 10.4: Why the monotone and Cauchy theorems require real completeness

What we must show. Explain why Theorems 10.2 and 10.11 would be false if the permitted terms and limits were restricted to $\mathbb{Q}$.

Plan. We construct a sequence of rational decimal truncations to $\sqrt{2}$. It is increasing, bounded by rational numbers, and Cauchy, but its only real limit is the irrational number $\sqrt{2}$. The construction itself uses finite sets and rational comparisons, rather than assuming that $\mathbb{Q}$ has the completeness property being tested.

Step 1: construct each truncation using integers only.

For each $n \in \mathbb{N}$, consider the finite set

$$E_n = \{k \in \mathbb{Z} : 0 \le k \le 2 \cdot 10^n,\ k^2 < 2 \cdot 10^{2n}\}.$$

It is nonempty because $0 \in E_n$. A finite nonempty set of integers has a largest member, so define

$$k_n = \max E_n, \qquad q_n = \frac{k_n}{10^n}.$$

The denominator is a positive integer, and the numerator is an integer. Hence every q_n is rational. The operations and comparisons used to define E_n involve only integers. In particular, this construction does not presuppose a supremum in $\mathbb{Q}$ for an infinite set.

Step 2: prove rational lower and upper bounds.

The integer 10^n belongs to E_n, since it is between 0 and $2 \cdot 10^n$ and its square 10^{2n} is strictly less than $2 \cdot 10^{2n}$. Thus $k_n \ge 10^n$. The integer $2 \cdot 10^n$ does not belong to E_n, since its square is $4 \cdot 10^{2n} > 2 \cdot 10^{2n}$. Therefore $k_n < 2 \cdot 10^n$. Dividing by $10^n > 0$ gives

$$1 \le q_n < 2.$$

Both bounds are rational, so the sequence is bounded even when all bounding numbers are required to lie in $\mathbb{Q}$.

Step 3: prove that the sequence is increasing.

Since $k_n^2 < 2 \cdot 10^{2n}$, multiplication by 100 gives

$$(10k_n)^2 < 2 \cdot 10^{2(n+1)}.$$

Also $0 \le 10k_n \le 2 \cdot 10^{n+1}$. Thus $10k_n \in E_{n+1}$. The maximality of k_{n+1} implies $10k_n \le k_{n+1}$. Dividing by $10^{n+1} > 0$ yields

$$q_n = \frac{10k_n}{10^{n+1}} \le \frac{k_{n+1}}{10^{n+1}} = q_{n+1}.$$

Equality may occur at some indices, which is permitted by the definition of increasing.

Step 4: give a rational upper approximation to the same square.

The integer $k_n + 1$ is still at most $2 \cdot 10^n$, because $k_n < 2 \cdot 10^n$ and both numbers are integers. If $(k_n + 1)^2 < 2 \cdot 10^{2n}$, then $k_n + 1$ would belong to E_n, contradicting maximality. Therefore

$$k_n^2 < 2 \cdot 10^{2n} \le (k_n + 1)^2.$$

Divide by $10^{2n} > 0$. The result is

$$q_n^2 < 2 \leq \left(q_n + \frac{1}{10^n}\right)^2.$$

Every number in this display is rational, so the comparisons remain meaningful entirely within $\mathbb{Q}$.

Step 5: verify the Cauchy condition through rational estimates.

For integers $m > n$, Step 3 gives $q_n \leq q_m$. The positive numbers q_m and $q_n + 10^{-n}$ satisfy

$$q_m^2 < 2 \leq (q_n + 10^{-n})^2.$$

Hence $q_m < q_n + 10^{-n}$. Indeed, the contrary inequality $q_m \geq q_n + 10^{-n} \geq 0$ would imply the contrary comparison between their squares. We conclude

$$0 \leq q_m - q_n < 10^{-n} \qquad (m > n).$$

We may use $10^j \geq j$ for every $j \in \mathbb{N}$: the case $j = 1$ is $10 \geq 1$; if $10^j \geq j$, then $10^{j+1} \geq 10j \geq j + 1$ because $j \geq 1$. This proves the estimate by induction, and taking reciprocals gives $10^{-j} \leq 1/j$.

Now let $\varepsilon > 0$ and choose $N = \max\{1, 1/\varepsilon\}$. Take any two integer indices $m, n > N$. If $m = n$, then $|q_m - q_n| = 0 < \varepsilon$. If $m > n$, the preceding estimate gives

$$|q_m - q_n| = q_m - q_n < 10^{-n} \leq \frac{1}{n} < \varepsilon.$$

If $n > m$, interchange their roles and use the symmetry of absolute value to obtain $|q_m - q_n| < 10^{-m} \leq 1/m < \varepsilon$. Thus the Cauchy estimate holds for every pair of indices above N. In particular, it holds for every positive rational tolerance, as a Cauchy condition formulated within $\mathbb{Q}$ requires.

Step 6: identify the only possible real limit.

In the ambient real numbers, let $\alpha = \sqrt{2}$. It is positive, $\alpha^2 = 2$, and it is irrational by Section 2. Step 4 and comparison of squares of positive numbers give

$$q_n < \alpha \leq q_n + 10^{-n}.$$

The upper equality is impossible because $q_n + 10^{-n}$ is rational and α is irrational. Therefore

$$0 < \alpha - q_n < 10^{-n} \leq \frac{1}{n}.$$

For any real $\varepsilon > 0$, the choice $N = \max\{1, 1/\varepsilon\}$ makes the right side strictly smaller than ε whenever $n > N$. Hence Definition 7.1 gives $q_n \to \alpha$ as a sequence in $\mathbb{R}$.

Step 7: rule out every rational limit.

A limit in $\mathbb{Q}$ would have to be some $r \in \mathbb{Q}$. Restricting the definition to positive rational tolerances makes no difference: given any real tolerance $\varepsilon > 0$, the Archimedean property gives a positive rational $1/j < \varepsilon$, and the estimate for that smaller rational tolerance implies the estimate for ε. Thus a limit in $\mathbb{Q}$ would also be the same limit in $\mathbb{R}$.

Because r is rational and α is irrational, they are distinct. Put $d = |\alpha - r| > 0$. If q_n converged to both numbers, there would be thresholds N_1, N_2 such that, for $n > \max\{N_1, N_2\}$,

$$|q_n - \alpha| < \frac{d}{3}, \qquad |q_n - r| < \frac{d}{3}.$$

The triangle inequality would then give

$$d = |\alpha - r| \leq |\alpha - q_n| + |q_n - r| < \frac{2d}{3} < d,$$

a contradiction. Thus the rational sequence has no rational limit.

Conclusion. The same rational sequence is bounded and increasing, so it contradicts the version of Theorem 10.2 with limits restricted to $\mathbb{Q}$. It is also Cauchy, so it contradicts the corresponding version of Theorem 10.11. Both failures arise because the required limit $\sqrt{2}$ is missing from $\mathbb{Q}$, not because the sequence is unbounded or fails its Cauchy estimates.

10.5 Exercise 10.5: An unbounded decreasing sequence tends to negative infinity

What we must show. Prove Theorem 10.4(ii): if a real sequence (s_n) is decreasing and unbounded, then $s_n \to -\infty$.

Step 1: determine which direction must be unbounded.

Since the sequence is decreasing, every term satisfies $s_n \leq s_1$. Thus s_1 is an upper bound of its set of terms. If the sequence also had a real lower bound, it would be bounded above and below, contrary to the hypothesis that it is unbounded. Therefore the sequence is unbounded below.

Step 2: find a term below an arbitrary target.

Let $M < 0$ be any real target. Since the sequence is unbounded below, M cannot be a lower bound for all its terms. Hence some index $N \in \mathbb{N}$ satisfies

$$s_N < M.$$

Step 3: show that every later term remains below that target.

For every integer $n > N$, decrease gives $s_n \leq s_N$. Thus

$$s_n \leq s_N < M \qquad (n > N).$$

We have proved that for each $M < 0$ there exists a threshold N such that every later term is less than M.

Conclusion. This is the definition of divergence to negative infinity, so $\boxed{s_n \longrightarrow -\infty}$. The choice of s_N would in fact work for any real target M, not just negative ones.

10.6 Exercise 10.6: Small successive differences: one sufficient rate and one insufficient rate

What we must show. Prove (a) that $|s_{n+1} - s_n| < 2^{-n}$ for every natural n implies (s_n) is Cauchy and convergent; and (b) decide whether the same conclusion follows from the weaker estimate $|s_{n+1} - s_n| < 1/n$.

Part (a), Step 1: compare arbitrary terms by telescoping.

Fix integers $m > n$. Then

$$s_m - s_n = \sum_{k=n}^{m-1} (s_{k+1} - s_k).$$

To see the cancellation, expand the right side as $(s_{n+1} - s_n) + (s_{n+2} - s_{n+1}) + \cdots + (s_m - s_{m-1})$; every intermediate term occurs once with each sign. Applying the finite triangle inequality gives

$$|s_m - s_n| \leq \sum_{k=n}^{m-1} |s_{k+1} - s_k| < \sum_{k=n}^{m-1} 2^{-k}.$$

The final inequality is strict because the finite sum is nonempty and every one of its summands is strictly less than the corresponding 2^{-k}.

Part (a), Step 2: bound this finite geometric sum.

For an individual k, $2^{-k} = 2^{1-k} - 2^{-k}$. Substituting that identity into the sum makes the consecutive terms telescope:

$$\begin{aligned}\sum_{k=n}^{m-1} 2^{-k} &= (2^{1-n} - 2^{-n}) + (2^{-n} - 2^{-n-1}) + \cdots + (2^{2-m} - 2^{1-m}),\\ &= 2^{1-n} - 2^{1-m},\\ &< 2^{1-n}.\end{aligned}$$

The last inequality holds because $2^{1-m} > 0$. We have therefore shown

$$|s_m - s_n| < 2^{1-n} \qquad (m > n).$$

Part (a), Step 3: choose one threshold for all pairs.

Let $\varepsilon > 0$. By the Archimedean property choose $N \in \mathbb{N}$ such that

$$N > \max\left\{1, \frac{2}{\varepsilon}\right\}.$$

The elementary estimate $2^N \geq N + 1$ was proved in Exercise 10.1(e). It gives

$$2^{1-N} = \frac{2}{2^N} \leq \frac{2}{N+1} < \frac{2}{N} < \varepsilon.$$

Now let $m, n > N$ be arbitrary integer indices. If $m = n$, their terms have distance $0 < \varepsilon$. If $m > n$, Steps 1–2 give

$$|s_m - s_n| < 2^{1-n} \leq 2^{1-N} < \varepsilon.$$

If $n > m$, interchange the indices and use $|s_m - s_n| = |s_n - s_m|$ to get the same conclusion with 2^{1-m} in place of 2^{1-n}.

Conclusion. For every $\varepsilon > 0$ all pairs of terms after the chosen threshold have distance less than ε. Thus (s_n) is Cauchy by Definition 10.8, and Theorem 10.11 for real sequences implies that it converges to a real number.

Part (b), Step 1: construct a sequence satisfying the strict hypothesis.

Define the harmonic partial sums

$$H_n = \sum_{k=1}^{n} \frac{1}{k} = 1 + \frac{1}{2} + \cdots + \frac{1}{n}.$$

The consecutive difference is exactly

$$H_{n+1} - H_n = \frac{1}{n+1}.$$

Both sides are positive, and $n + 1 > n > 0$, so

$$|H_{n+1} - H_n| = \frac{1}{n+1} < \frac{1}{n} \qquad (n \in \mathbb{N}).$$

The required inequality is strict at every index, including $n = 1$.

Part (b), Step 2: show directly that the sequence is not Cauchy.

Fix the positive tolerance $\varepsilon_0 = 1/2$. Given any proposed threshold N, choose an integer $n > \max\{N, 1\}$ and put $m = 2n$. Both indices are greater than N. The difference of the terms is

$$H_{2n} - H_n = \sum_{k=n+1}^{2n} \frac{1}{k}.$$

There are exactly n summands. For each index in this sum, $k \leq 2n$, so $1/k \geq 1/(2n)$. Adding the n inequalities gives

$$|H_{2n} - H_n| = H_{2n} - H_n \geq n\frac{1}{2n} = \frac{1}{2} = \varepsilon_0.$$

Thus, no matter how N is chosen, some pair of indices $m, n > N$ violates the required strict Cauchy estimate.

Conclusion. The answer to part (b) is **no**. The sequence (H_n) satisfies every strict bound $|H_{n+1} - H_n| < 1/n$, but it is not Cauchy. Since every convergent sequence is Cauchy by Lemma 10.9, it does not converge.

10.7 Exercise 10.7: Approaching an unattained supremum by points of the set

What we must show. Let $S \subseteq \mathbb{R}$ be bounded and nonempty, and suppose $\sup S \notin S$. Construct a sequence (s_n) with $s_n \in S$ for every n and $s_n \to \sup S$.

Step 1: use the least-upper-bound property at each precision.

Completeness gives the real number $M = \sup S$. For each $n \in \mathbb{N}$, we have $1/n > 0$, and consequently $M - 1/n < M$. The smaller number $M - 1/n$ cannot be an upper bound of S, because M is its least upper bound. Therefore there is a point $s_n \in S$ satisfying $M - 1/n < s_n$. Since M is an upper bound, also $s_n \leq M$. Equality is ruled out by $M \notin S$. Thus, choosing such a point for each n, we obtain

$$M - \frac{1}{n} < s_n < M \qquad (n \in \mathbb{N}).$$

Step 2: translate the order estimate into an error estimate.

Subtracting s_n and rearranging the left inequality gives $M - s_n < 1/n$. The right inequality gives $M - s_n > 0$, so

$$|s_n - M| = M - s_n < \frac{1}{n}.$$

This bound holds at every index; the actual choices of the points s_n need not produce a monotone sequence.

Step 3: verify convergence with an explicit threshold.

Let $\varepsilon > 0$ and choose

$$N = \max\left\{1, \frac{1}{\varepsilon}\right\}.$$

For every integer $n > N$, $n > 1/\varepsilon$ gives $1/n < \varepsilon$, and therefore

$$|s_n - M| < \frac{1}{n} < \varepsilon.$$

Conclusion. The constructed sequence consists entirely of points of S and satisfies $\boxed{s_n \longrightarrow \sup S}$. The supremum itself is excluded from the set, so each approximation is strictly below its target.

10.8 Exercise 10.8: Averages of an increasing positive sequence

What we must show. Let (s_n) be increasing with $s_n > 0$ for every n, and let

$$\sigma_n = \frac{1}{n}\sum_{k=1}^{n} s_k.$$

Prove that the average sequence (σ_n) is increasing in the sense of Definition 10.1.

Step 1: compare each average with the last term it averages.

Fix $n \in \mathbb{N}$. Since (s_n) is increasing, every k with $1 \le k \le n$ satisfies $s_k \le s_n$. Add these n inequalities:

$$s_1 + s_2 + \cdots + s_n \le \underbrace{s_n + s_n + \cdots + s_n}_{n \text{ terms}} = ns_n.$$

Divide by the positive integer n to obtain $\sigma_n \le s_n$. Since $s_n \le s_{n+1}$, transitivity gives

$$\sigma_n \le s_n \le s_{n+1}.$$

In particular, $s_{n+1} - \sigma_n \ge 0$.

Step 2: express the next average in terms of the previous one.

The definition of σ_n gives $s_1 + \cdots + s_n = n\sigma_n$. Therefore

$$\begin{aligned}\sigma_{n+1} &= \frac{s_1 + \cdots + s_n + s_{n+1}}{n+1},\\ &= \frac{n\sigma_n + s_{n+1}}{n+1}.\end{aligned}$$

Step 3: calculate the exact difference between consecutive averages.

Put the terms over the same positive denominator $n+1$:

$$\begin{aligned}\sigma_{n+1}-\sigma_n &= \frac{n\sigma_n+s_{n+1}}{n+1}-\frac{(n+1)\sigma_n}{n+1},\\ &= \frac{n\sigma_n+s_{n+1}-(n+1)\sigma_n}{n+1},\\ &= \frac{n\sigma_n+s_{n+1}-n\sigma_n-\sigma_n}{n+1},\\ &= \frac{s_{n+1}-\sigma_n}{n+1}.\end{aligned}$$

The numerator is nonnegative by Step 1, and the denominator $n+1$ is positive. Hence $\sigma_{n+1}-\sigma_n \geq 0$, or $\sigma_n \leq \sigma_{n+1}$.

Conclusion. This comparison holds for every $n \in \mathbb{N}$, proving $\boxed{(\sigma_n) \text{ is increasing}}$. Equality can occur: for a constant positive sequence, all the averages are equal. Thus the weak meaning of "increasing" is the one required by the hypotheses and conclusion.

10.9 Exercise 10.9: A decreasing recurrence with a squared term

What we must show. For

$$s_1 = 1, \qquad s_{n+1} = \frac{n}{n+1}s_n^2 \quad (n \in \mathbb{N}),$$

compute s_2, s_3, s_4, prove that a finite limit exists, and prove that its value is 0.

Part (a): substitute the indices one at a time.

For $n = 1$, the recurrence gives

$$s_2 = \frac{1}{1+1}s_1^2 = \frac{1}{2}\cdot 1^2 = \frac{1}{2}.$$

For $n = 2$,

$$s_3 = \frac{2}{2+1}s_2^2 = \frac{2}{3}\left(\frac{1}{2}\right)^2 = \frac{2}{3}\cdot\frac{1}{4} = \frac{1}{6}.$$

For $n = 3$,

$$s_4 = \frac{3}{3+1}s_3^2 = \frac{3}{4}\left(\frac{1}{6}\right)^2 = \frac{3}{4}\cdot\frac{1}{36} = \frac{3}{144} = \frac{1}{48}.$$

Conclusion. $\boxed{s_2 = \frac{1}{2}, \quad s_3 = \frac{1}{6}, \quad s_4 = \frac{1}{48}}$.

Part (b), Step 1: prove positivity and an upper bound by induction.

We claim $0 < s_n \leq 1$ for every $n \in \mathbb{N}$.

The initial case is $s_1 = 1$, so $0 < s_1 \leq 1$ holds. For the induction step, assume $0 < s_n \leq 1$. Since $n \geq 1$, the coefficient satisfies

$$0 < \frac{n}{n+1} < 1.$$

Also $0 < s_n^2 \leq 1$. Multiplying these positive quantities gives

$$0 < s_{n+1} = \frac{n}{n+1}s_n^2 < 1.$$

This proves the claim by induction.

Part (b), Step 2: compare consecutive terms.

The inequality $s_n \le 1$, multiplied by the positive number s_n, gives $s_n^2 \le s_n$. The coefficient $n/(n+1)$ is strictly below 1, and $s_n^2 > 0$, so

$$s_{n+1} = \frac{n}{n+1} s_n^2 < s_n^2 \le s_n.$$

Hence the sequence is strictly decreasing. It is bounded below by 0 and above by 1.

Part (b), Step 3: apply monotone convergence before manipulating a limit.

Theorem 10.2 for bounded monotone sequences now applies. There is a real number L with $s_n \to L$. The inequalities $s_n > 0$ imply $L \ge 0$ by Exercise 8.9(a). Since the sequence is decreasing and $s_2 = 1/2$,

$$0 < s_n \le \frac{1}{2} \qquad (n \ge 2).$$

Exercise 8.9 also passes this eventual upper bound to the limit. Therefore

$$0 \le L \le \frac{1}{2}.$$

Conclusion. A finite limit exists, and it belongs to $[0, 1/2]$. Its existence has been proved independently of any equation for its value.

Part (c), Step 1: justify the shifted limit and the coefficient limit.

If $s_n \to L$, then $s_{n+1} \to L$ as well. Indeed, for any $\varepsilon > 0$ choose a threshold N with $m > N \Rightarrow |s_m - L| < \varepsilon$. For $n > N$, we have $n + 1 > N$, so the same estimate holds with $m = n + 1$.

Next put $c_n = n/(n+1)$. Its error from 1 is

$$|c_n - 1| = \left| \frac{n - (n+1)}{n+1} \right| = \frac{1}{n+1}.$$

For any $\varepsilon > 0$, if $n > 1/\varepsilon$, then $n + 1 > 1/\varepsilon$, so $1/(n+1) < \varepsilon$. Thus $c_n \to 1$.

Part (c), Step 2: apply the product limit law to the established limits.

Because $s_n \to L$ and $c_n \to 1$, the product laws from Section 9 give

$$\lim_{n\to\infty} c_n s_n^2 = 1 \cdot L \cdot L = L^2.$$

The recurrence says that this is the same sequence as s_{n+1}, whose limit is L. Uniqueness of a finite limit gives

$$L = L^2.$$

Rearranging and factoring gives

$$L^2 - L = 0, \qquad L(L-1) = 0.$$

The zero-product property therefore gives $L = 0$ or $L = 1$.

Part (c), Step 3: use the tail bound to remove the extraneous candidate.

Part (b) proved $L \le 1/2$, so $L = 1$ is impossible. The only remaining candidate is $L = 0$.

Conclusion. $\boxed{\lim_{n\to\infty} s_n = 0}$.

An additional direct error bound.

The same recurrence supplies a quantitative check. Since $s_n^2 \le s_n$,

$$s_{n+1} \le \frac{n}{n+1} s_n.$$

Starting with $s_1 = 1 = 1/1$, induction gives $s_n \le 1/n$: if $s_n \le 1/n$, then

$$s_{n+1} \le \frac{n}{n+1} \cdot \frac{1}{n} = \frac{1}{n+1}.$$

Consequently

$$|s_n - 0| = s_n \le \frac{1}{n} < \varepsilon \qquad \left(n > \frac{1}{\varepsilon}\right).$$

This also verifies the zero limit directly from Definition 7.1.

10.10 Exercise 10.10: A linear recurrence approaching one half

What we must show. For

$$s_1 = 1, \qquad s_{n+1} = \frac{s_n + 1}{3} \quad (n \in \mathbb{N}),$$

compute s_2, s_3, s_4, prove $s_n > 1/2$ by induction, prove that the sequence is decreasing, and identify its limit.

Part (a): calculate the first three new terms.

Using the recurrence successively gives

$$s_2 = \frac{1+1}{3} = \frac{2}{3},$$

$$s_3 = \frac{\frac{2}{3}+1}{3} = \frac{\frac{2}{3}+\frac{3}{3}}{3} = \frac{5/3}{3} = \frac{5}{9},$$

and

$$s_4 = \frac{\frac{5}{9}+1}{3} = \frac{\frac{5}{9}+\frac{9}{9}}{3} = \frac{14/9}{3} = \frac{14}{27}.$$

Conclusion. $\boxed{s_2 = \frac{2}{3}, \quad s_3 = \frac{5}{9}, \quad s_4 = \frac{14}{27}}$.

Part (b), Step 1: verify the initial case of the proposed lower bound.

The proposition for induction is P_n: $s_n > 1/2$. For $n = 1$, it says $1 > 1/2$, which is true.

Part (b), Step 2: prove the induction implication.

Assume $s_n > 1/2$. Adding 1 to both sides gives $s_n + 1 > 3/2$. Dividing by the positive number 3 preserves the strict inequality:

$$s_{n+1} = \frac{s_n + 1}{3} > \frac{3/2}{3} = \frac{1}{2}.$$

Thus P_n implies P_{n+1}. Induction proves the bound for every positive integer n.

Conclusion. $\boxed{s_n > \frac{1}{2} \text{ for every } n \in \mathbb{N}}$.

Part (c): subtract consecutive terms and determine the sign.

Use a common denominator:

$$s_{n+1} - s_n = \frac{s_n + 1}{3} - s_n = \frac{s_n + 1 - 3s_n}{3} = \frac{1 - 2s_n}{3}.$$

Part (b) gives $2s_n > 1$, so the numerator is negative. The denominator 3 is positive. Hence $s_{n+1} - s_n < 0$, which means $s_{n+1} < s_n$ for every n.

Conclusion. The sequence is strictly decreasing.

Part (d), Step 1: establish convergence from bounds and monotonicity.

The terms satisfy $1/2 < s_n \leq s_1 = 1$, and part (c) proves that the sequence is decreasing. Theorem 10.2 therefore gives a finite limit L. Exercise 8.9 implies

$$\frac{1}{2} \leq L \leq 1.$$

Part (d), Step 2: solve the limit equation after convergence is known.

The shifted sequence s_{n+1} has the same limit L, by the threshold argument given in Exercise 10.9(c). The addition and scalar-multiplication laws from Section 9 give

$$\lim_{n\to\infty} \frac{s_n + 1}{3} = \frac{L + 1}{3}.$$

The recurrence identifies these two sequences, so their limits agree:

$$L = \frac{L + 1}{3}.$$

Multiplying by 3 and collecting terms gives

$$3L = L + 1, \qquad 2L = 1, \qquad L = \frac{1}{2}.$$

Conclusion. $\boxed{\lim_{n\to\infty} s_n = \frac{1}{2}}$. Every term is strictly greater than $1/2$, but the limit can equal that lower bound.

10.11 Exercise 10.11: An infinite-product recurrence

What we must show. For

$$t_1 = 1, \qquad t_{n+1} = \left(1 - \frac{1}{4n^2}\right) t_n \quad (n \in \mathbb{N}),$$

(a) prove that a finite limit exists and (b) investigate its value. Part (b) asks what the limit appears to be. We will distinguish a numerical guess from a proof; an optional elementary-calculus argument will then verify the exact value.

Part (a), Step 1: bound the coefficient and prove positivity.

For every $n \geq 1$, we have $4n^2 \geq 4$, so

$$0 < \frac{1}{4n^2} \leq \frac{1}{4}.$$

Subtracting from 1 gives

$$\frac{3}{4} \leq 1 - \frac{1}{4n^2} < 1.$$

In particular, every recurrence coefficient is strictly positive. The initial value is $t_1 = 1 > 0$. If $t_n > 0$, the recurrence is a product of two positive numbers and gives $t_{n+1} > 0$. Induction shows $t_n > 0$ for all n.

Part (a), Step 2: prove monotonicity and apply the theorem.

Since $1 - 1/(4n^2) < 1$ and $t_n > 0$,

$$t_{n+1} = \left(1 - \frac{1}{4n^2}\right) t_n < t_n.$$

Thus (t_n) is strictly decreasing, bounded below by 0 and above by its first term 1. Theorem 10.2 gives a finite real limit L. The bound on the terms implies $0 \leq L \leq 1$ by Exercise 8.9.

Conclusion. The limit of (t_n) exists as a real number. This conclusion uses only positivity, monotonicity, and boundedness.

Part (b), Step 1: compute initial values and a numerical table.

The first terms are

$$t_2 = \left(1 - \frac{1}{4}\right) 1 = \frac{3}{4},$$

$$t_3 = \left(1 - \frac{1}{16}\right) \frac{3}{4} = \frac{15}{16} \cdot \frac{3}{4} = \frac{45}{64},$$

and

$$t_4 = \left(1 - \frac{1}{36}\right) \frac{45}{64} = \frac{35}{36} \cdot \frac{45}{64} = \frac{1575}{2304} = \frac{175}{256}.$$

Continuing the recurrence produces the following rounded values:

n	t_n
1	1.000000000000
2	0.750000000000
3	0.703125000000
4	0.683593750000
10	0.653587392299
100	0.638221323951
1000	0.636779026837

The numbers suggest a limit near 0.63662. The familiar constant $2/\pi$ has the decimal value $0.636619772368\ldots$, which suggests the guess

$$L = \frac{2}{\pi}.$$

The numerical agreement motivates this guess; finitely many numerical calculations by themselves do not prove the exact identification.

Part (b), Step 2: explain why a direct limit equation gives no value.

The coefficient $1 - 1/(4n^2)$ tends to 1. Taking limits in the recurrence, after convergence has been proved, only yields

$$L = 1 \cdot L.$$

Every real L satisfies this identity, so it does not determine the limit. Further information is needed to justify the guess $2/\pi$.

Optional verification of the guess: the elementary-calculus input.

The remaining argument uses integration by parts and the order property of definite integrals. Those are additional calculus tools; they were not needed for the existence proof in part (a). We derive the required product formula here rather than assume Wallis's product.

For an integer $m \geq 0$, define

$$I_m = \int_0^{\pi/2} \sin^m x \, dx,$$

where $\sin^0 x$ means the constant function 1. The integrands are continuous and are positive in the interior when $m \geq 1$, so all I_m are positive. Direct integration gives

$$I_0 = \frac{\pi}{2}, \qquad I_1 = 1.$$

Optional verification, Step 1: derive the integral recurrence.

For $m \geq 2$, write $I_m = \int_0^{\pi/2} \sin^{m-1} x \sin x \, dx$. In integration by parts take

$$u = \sin^{m-1} x, \quad dv = \sin x \, dx, \quad du = (m-1)\sin^{m-2} x \cos x \, dx, \quad v = -\cos x.$$

The boundary term $[-\sin^{m-1} x \cos x]_0^{\pi/2}$ is zero: $\cos(\pi/2) = 0$, and at 0 the factor $\sin^{m-1} 0$ is zero because $m - 1 \geq 1$. Consequently

$$\begin{aligned} I_m &= (m-1) \int_0^{\pi/2} \sin^{m-2} x \cos^2 x \, dx \\ &= (m-1) \int_0^{\pi/2} \sin^{m-2} x (1 - \sin^2 x) \, dx \\ &= (m-1)(I_{m-2} - I_m). \end{aligned}$$

Adding $(m-1)I_m$ to both sides and dividing by $m > 0$ gives

$$I_m = \frac{m-1}{m} I_{m-2}.$$

Optional verification, Step 2: express the even and odd integrals.

Repeated use of the recurrence, ending at I_0 or I_1, gives for every integer $N \geq 1$,

$$I_{2N} = \frac{\pi}{2} \prod_{k=1}^{N} \frac{2k-1}{2k}, \qquad I_{2N+1} = \prod_{k=1}^{N} \frac{2k}{2k+1}.$$

For example, the even recurrence successively contributes the factors $(2N-1)/(2N)$, $(2N-3)/(2N-2)$, down to $1/2$. The odd recurrence contributes $(2N)/(2N+1)$ down to $2/3$. These are exactly the displayed finite products.

Unrolling the recurrence for t_n gives

$$t_{N+1} = \prod_{k=1}^{N} \left(1 - \frac{1}{4k^2}\right) = \prod_{k=1}^{N} \frac{(2k-1)(2k+1)}{(2k)^2}.$$

The initial empty product for t_1 is 1, and each new recurrence step multiplies by the next factor, which justifies this finite-product identity. Dividing the formulas for the positive integrals gives

$$\frac{I_{2N}}{I_{2N+1}} = \frac{\pi}{2} \prod_{k=1}^{N} \frac{(2k-1)(2k+1)}{(2k)^2} = \frac{\pi}{2}\, t_{N+1}.$$

Optional verification, Step 3: bound the integral ratio.

On $[0, \pi/2]$, the quantity $\sin x$ lies in $[0, 1]$. Multiplication by a number in $[0, 1]$ cannot increase a nonnegative power, so

$$\sin^{2N+1} x \le \sin^{2N} x \le \sin^{2N-1} x.$$

Integrating preserves these inequalities:

$$I_{2N+1} \le I_{2N} \le I_{2N-1}.$$

Divide by $I_{2N+1} > 0$. The integral recurrence with $m = 2N + 1$ gives $I_{2N+1} = (2N)/(2N + 1)\, I_{2N-1}$, so

$$1 \le \frac{I_{2N}}{I_{2N+1}} \le \frac{I_{2N-1}}{I_{2N+1}} = \frac{2N+1}{2N} = 1 + \frac{1}{2N}.$$

Substituting $I_{2N}/I_{2N+1} = (\pi/2)t_{N+1}$ and multiplying by $2/\pi > 0$ yields

$$0 \le t_{N+1} - \frac{2}{\pi} \le \frac{1}{\pi N}.$$

For any $\varepsilon > 0$, a positive integer index $n > 1 + 1/(\pi\varepsilon)$ has $N = n - 1 > 1/(\pi\varepsilon)$. Therefore

$$\left| t_n - \frac{2}{\pi} \right| \le \frac{1}{\pi(n-1)} < \varepsilon.$$

This proves the conjectured limit by Definition 7.1.

Conclusion. Part (a) proves existence using Section 10 alone. For part (b), the computed terms suggest $2/\pi$. The explicitly marked additional calculus argument verifies that

$$\boxed{\lim_{n\to\infty} t_n = \frac{2}{\pi}.}$$

No product formula was taken on trust; the finite-product identity and its limiting estimate were derived.

10.12 Exercise 10.12: A product recurrence with an exact formula

What we must show. For

$$t_1 = 1, \qquad t_{n+1} = \left(1 - \frac{1}{(n+1)^2}\right) t_n \quad (n \in \mathbb{N}),$$

(a) establish existence of the limit, (b) guess its value, (c) prove $t_n = (n+1)/(2n)$ by induction, and (d) use the formula to verify the limit.

Part (a), Step 1: prove positivity and monotonicity.

For every $n \geq 1$, we have $n+1 \geq 2$, so

$$0 < \frac{1}{(n+1)^2} \leq \frac{1}{4}, \qquad \frac{3}{4} \leq 1 - \frac{1}{(n+1)^2} < 1.$$

The initial term is positive. If $t_n > 0$, multiplying it by this positive coefficient gives $t_{n+1} > 0$. Induction proves positivity for every term.

Because the coefficient is strictly less than 1 and $t_n > 0$, we also have $t_{n+1} < t_n$ for every n. Thus the sequence is strictly decreasing, bounded below by 0 and above by $t_1 = 1$.

Part (a), Step 2: apply bounded monotone convergence.

Theorem 10.2 now gives a finite limit L. Exercise 8.9 implies $0 \leq L \leq 1$.

Conclusion. The limit exists as a real number. Its value has not been assumed in establishing existence.

Part (b): calculate early terms to formulate a guess.

Successive substitutions give

$$t_2 = \left(1 - \frac{1}{4}\right) 1 = \frac{3}{4},$$

$$t_3 = \left(1 - \frac{1}{9}\right) \frac{3}{4} = \frac{8}{9} \cdot \frac{3}{4} = \frac{2}{3},$$

$$t_4 = \left(1 - \frac{1}{16}\right) \frac{2}{3} = \frac{15}{16} \cdot \frac{2}{3} = \frac{5}{8},$$

and

$$t_5 = \left(1 - \frac{1}{25}\right) \frac{5}{8} = \frac{24}{25} \cdot \frac{5}{8} = \frac{3}{5}.$$

The values

$$1, \quad \frac{3}{4}, \quad \frac{2}{3}, \quad \frac{5}{8}, \quad \frac{3}{5}$$

fit the proposed pattern $t_n = (n+1)/(2n)$. Since this expression equals $1/2 + 1/(2n)$, it suggests the limit $1/2$. At this stage, the pattern and the limit are guesses to be proved.

Part (c), Step 1: state the induction proposition and verify its base.

Let P_n be the assertion

$$t_n = \frac{n+1}{2n}.$$

For $n = 1$, the right side is $(1+1)/(2 \cdot 1) = 1$, which equals t_1. Thus P_1 holds.

Part (c), Step 2: substitute the induction hypothesis.

Assume P_n for an arbitrary positive integer n. The recurrence gives

$$t_{n+1} = \left(1 - \frac{1}{(n+1)^2}\right) \frac{n+1}{2n}$$

$$
\begin{aligned}
&= \frac{(n+1)^2 - 1}{(n+1)^2} \frac{n+1}{2n} \\
&= \frac{n^2 + 2n}{(n+1)^2} \frac{n+1}{2n} \\
&= \frac{n(n+2)}{(n+1)^2} \frac{n+1}{2n} \\
&= \frac{n+2}{2(n+1)} \\
&= \frac{(n+1)+1}{2(n+1)}.
\end{aligned}
$$

The cancellation in the penultimate line is legitimate because $n > 0$ and $n+1 > 0$, so neither cancelled factor is zero. The final expression is precisely the formula in P_{n+1}.

Conclusion. Mathematical induction proves

$$\boxed{t_n = \frac{n+1}{2n} \quad \text{for every } n \in \mathbb{N}.}$$

Part (d), Step 1: separate the constant term from the error.

The proved formula gives the identity

$$t_n = \frac{n+1}{2n} = \frac{n}{2n} + \frac{1}{2n} = \frac{1}{2} + \frac{1}{2n}.$$

Since $1/n \to 0$, the limit laws from Section 9 give $t_n \to 1/2$.

Part (d), Step 2: verify the conclusion directly with a threshold.

For any $\varepsilon > 0$, choose $N = 1/(2\varepsilon)$. Every positive integer $n > N$ satisfies $1/(2n) < \varepsilon$, and

$$\left| t_n - \frac{1}{2} \right| = \left| \frac{1}{2n} \right| = \frac{1}{2n} < \varepsilon.$$

This is the definition of convergence to $1/2$.

Conclusion. The guess in part (b) is now proved:

$$\boxed{\lim_{n \to \infty} t_n = \frac{1}{2}.}$$

The exact formula also gives the error $t_n - 1/2 = 1/(2n)$ for every positive integer index.