

Trig Identities

$$\begin{aligned}\sin(x+y) &= \sin x \cos y + \cos x \sin y \\ \cos(x+y) &= \cos x \cos y - \sin x \sin y \\ \tan(x+y) &= \frac{\tan x + \tan y}{1 - \tan x \tan y}\end{aligned}$$

$$\begin{aligned}\sin(2x) &= 2 \sin x \cos x \\ \cos(2x) &= \cos^2 x - \sin^2 x \\ &= 2 \cos^2 x - 1 \\ &= 1 - 2 \sin^2 x \\ \tan(2x) &= \frac{2 \tan x}{1 - \tan^2 x}\end{aligned}$$

Chapter 10

Parametric Eq, Polar

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}, \text{ if } \frac{dx}{dt} \neq 0$$

$$\text{Area} = \int_a^b g(t) f'(t) dt \quad \begin{matrix} x = f(t) \\ y = g(t) \\ \alpha \leq t \leq \beta \end{matrix}$$

$$\text{Arc Length} = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = \int_a^b |v(t)| dt$$

$$\text{Surface Area (around x-axis)} = 2\pi \int_a^b y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$\begin{aligned}x &= r \cos \theta & y &= r \sin \theta \\ r^2 &= x^2 + y^2 & \tan \theta &= \frac{y}{x}\end{aligned}$$

$$\text{Tangent of Polar: } \frac{dy}{dx} = \frac{\frac{dr}{d\theta} \sin \theta + r \cos \theta}{\frac{dr}{d\theta} \cos \theta - r \sin \theta}$$

$$\begin{aligned}\text{Cardioid/Limacon: } r &= a \pm b \cos \theta, r = a \pm b \sin \theta \\ \text{Roses: } r &= a \cos(n\theta) \quad r = a \sin(n\theta) \\ \text{odd: petals} &= n \quad \text{even: petals} = 2n\end{aligned}$$

$$\text{Area Polar} = \frac{1}{2} \int_a^b r^2 d\theta$$

$$\text{Arc Length} = \int_a^b \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$$

Chapter 12

Vectors, Space

$$a \cdot b = |a||b| \cos \theta \quad 0 \leq \theta \leq \pi$$

$$\text{comp}_a b = \frac{a \cdot b}{|a|} \quad \text{proj}_a b = \frac{a \cdot b}{|a|^2} \vec{a}$$

$$|a \times b| = |a||b| \sin \theta \quad 0 \leq \theta \leq \pi$$

$$\text{Vol of parallelepiped} = |\vec{a} \cdot (\vec{b} \times \vec{c})|$$

$$\text{Dist from point to plane} = \frac{|Ax_1 + By_1 + Cz_1 + D|}{\sqrt{A^2 + B^2 + C^2}}$$

$$\text{Dist from point to line: } h = \frac{|\vec{AB} \times \vec{AP}|}{|\vec{AP}|}$$

Intersections

- Check if direction vector parallel
 - Set $x_1 = x_2, y_1 = y_2, z_1 = z_2$ in terms of t and s
- Find distance between lines using cross product and comp. b

$$\text{Ellipsoid: } \frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$$

$$\text{Elliptic Paraboloid: } z = \frac{x^2}{a^2} + \frac{y^2}{b^2}$$

$$\text{Hyperbolic Paraboloid: } z = \frac{x^2}{a^2} - \frac{y^2}{b^2}$$

$$\text{Cone: } z^2 = \frac{x^2}{a^2} + \frac{y^2}{b^2}$$

$$\text{Hyperboloid One Sheet: } \frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$$

$$\text{Hyperboloid Two Sheets: } -\frac{x^2}{a^2} - \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$$

$$\text{Parallel } a, b \\ a \times b = 0$$

$$1. a \times b = -b \times a$$

$$2. (a \times b) \cdot c = a \cdot (b \times c)$$

$$3. a \cdot (a \times b) = 0$$

$$4. (a \times b) \times c = (a \cdot c)b - (b \cdot c)a$$

$$5. a \cdot (a \times b) = 0$$

Chapter 13

Vector Func

$$\text{Arc length vector} = \int_a^b |\vec{r}'(t)| dt$$

$$\text{Parametrize in terms of } s \quad \vec{r}(t) \Rightarrow \vec{r}(s)$$

- Solve for s
- Use eq to solve for t in terms of s
- Plug + equation into $\vec{r}(t)$

$$\text{Unit Tangent Vector: } \vec{T}(t) = \frac{\vec{r}'(t)}{|\vec{r}'(t)|}$$

$$\vec{N}(t) = \frac{\vec{T}'(t)}{|\vec{T}'(t)|} \quad \vec{B}(t) = \vec{T} \times \vec{N}$$

$$\text{Curvature: } K = \frac{|\vec{T}'(t)|}{|\vec{r}'(t)|} = \frac{|\vec{r}'(t) \times \vec{r}''(t)|}{|\vec{r}'(t)|^3}$$

$$\text{Radius of osculating plane: } r = \frac{1}{K}$$

$$a = \frac{v^2}{r} \quad \text{where } v = |\vec{v}|$$

$$a_T = \frac{r'(t) \cdot r''(t)}{|r'(t)|} \quad a_N = \frac{|r'(t) \times r''(t)|}{|r'(t)|^2}$$

Chapter 14

Partials

$$\lim_{(x,y) \rightarrow (a,b)} f(x,y) = L \quad \begin{matrix} \text{Approach from diff lines} \\ \text{convert to polar coord} \end{matrix}$$

$$\text{Tangent Plane: } z - z_0 = f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0)$$

$$\text{Linear approx: } L(x,y) = f_x(x-x_0) + f_y(y-y_0) + z_0$$

$$\text{Case 1: } z = f(x,y) \quad x = g(t) \quad y = h(t)$$

$$\frac{dz}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}$$

$$\text{Case 2: } z = f(x,y) \quad x = g(s,t) \quad y = h(s,t)$$

$$\frac{\partial z}{\partial s} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial s} \quad \frac{\partial z}{\partial t} = \dots$$

$$\frac{\partial y}{\partial x} = -\frac{f_x}{f_y} \quad \frac{\partial z}{\partial x} = -\frac{f_x}{f_z} \quad \frac{\partial z}{\partial y} = -\frac{f_y}{f_z}$$

$$D_u f(x,y) = \nabla f \cdot \vec{u} \quad \text{in direction unit vector } \vec{u}$$

$$\nabla f(x,y) = (f_x, f_y) \quad \text{max } Df = |\nabla f|$$

$$\text{or orth} = \vec{AP} - \text{proj}_{\vec{AB}} \vec{AP} \quad \vec{v} \text{ is } \vec{n} \text{ of tangent plane}$$

Max/Min

$$\text{Critical point } f_x = f_y = 0 \text{ or one DNE}$$

$$D = f_{xx}(a,b) f_{yy}(a,b) - [f_{xy}(a,b)]^2$$

$$\text{① } D > 0, f_{xx} > 0 \Rightarrow \text{local min}$$

$$\text{② } D > 0, f_{xx} < 0 \Rightarrow \text{local max}$$

$$\text{③ } D < 0 \Rightarrow \text{not min or max (saddle point)}$$

$$\text{④ } D = 0 \Rightarrow \text{no info, any}$$

$$\text{Lagrange: max/min: } f \text{ constraint: } g = k$$

$$\nabla f(x,y,z) = \lambda \nabla g(x,y,z)$$

Chapter 15

Mult Integrals

$$V = \iint_R f(x,y) dA$$

$$\text{Avg value} = \frac{\iint_R f(x,y) dA}{\text{area of } R} = \frac{\text{Volume}}{\text{Area}}$$

$$\text{Polar: } \iint_R f(r \cos \theta, r \sin \theta) r dr d\theta$$

$$\text{Applications}$$

$$m = \iint_D \rho(x,y) dA \quad \rho: \text{lamina}$$

$$= \iiint_D \rho(x,y,z) dV$$

$$\text{Moment: } M_x = \iint_D y \rho(x,y) dA$$

$$M_y = \iint_D x \rho(x,y) dA$$

$$\text{Center of mass}$$

$$\bar{x} = \frac{M_y}{m} = \frac{1}{m} \iint_D x \rho(x,y) dA$$

$$\bar{y} = \frac{M_x}{m} = \frac{1}{m} \iint_D y \rho(x,y) dA$$

$$\text{Moment of Inertia:}$$

$$I_x = \iint_D y^2 \rho(x,y) dA$$

$$I_y = \iint_D x^2 \rho(x,y) dA$$

$$I_z(\text{origin}) = \iint_D (x^2 + y^2) \rho(x,y) dA$$

$$\text{Probability: } f(x,y) \geq 0 \quad P = \iint_R f(x,y) dA = 1$$

$$\text{Triple Integrals}$$

$$\text{Cylindrical}$$

$$\int_{\alpha}^{\beta} \int_{h(\theta)}^{u_2} \int_u^{v_2} f(x,y,z) r dz dr d\theta$$

$$x = r \cos \theta \quad y = r \sin \theta \quad z = z$$

$$\text{Spherical}$$

$$\int_{\alpha}^{\beta} \int_{\phi_1}^{\phi_2} \int_{\rho_1}^{\rho_2} f(x,y,z) \rho^2 \sin \phi d\rho d\phi d\theta$$

$$x = \rho \sin \phi \cos \theta$$

$$y = \rho \sin \phi \sin \theta$$

$$z = \rho \cos \phi \quad x^2 + y^2 + z^2 = \rho^2$$

$$\iint_R F(x,y) dA = \iint_S F(x(u,v), y(u,v)) \left| \frac{\partial(x,y)}{\partial(u,v)} \right| du dv$$

Jacobian: $x = g(u,v)$ $y = h(u,v)$

$$\frac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u}$$

- ① Find u, v suitable
- ② Find x, y
- ③ Find Jacobian plug in above val
- ④ Plug into equation, solve

Chapter 16 Vector Calc

Conservative if some f that $F = \nabla f$

16.2 Line Integrals

$$\int_C F \cdot dr = \int_a^b F(r(t)) \cdot r'(t) dt$$

$$\int_C f(x,y) ds = \int_a^b f(x(t), y(t)) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

16.3 Fundamental Theorem for Line Integrals

Cond

- ① C smooth curve
- ② f differentiable
- ③ ∇f continuous on C

Then

$$\int_C \nabla f \cdot d\vec{r} = f(r(b)) - f(r(a))$$

Thm

Cond

- ① $\int_C F \cdot dr = 0$ for every closed path C in D

Then

$\int_C F \cdot dr$ is independent of path in D

- ① open (no boundary points)
- ② connected (2 pts joined by path)
- ③ independent of path

There is f where $\nabla f = F$

- ① Open, simply connected (no holes)
- ② P and Q have continuous first partials
- ③ $\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}$ throughout D

\vec{F} is conservative

16.4 Green's Theorem

- ① C is positively oriented
- ② piecewise-smooth
- ③ simple closed curve
- ④ P and Q have continuous partial derivatives on open region

let D be region bound by C

$$\int_C P dx + Q dy =$$

$$\iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA$$

16.5 Curl and Divergence

$$\text{Curl } F = \nabla \times F$$

$$\text{div } F = \nabla \cdot F$$

Func 3 vars continuous partials $\Rightarrow \text{Curl}(\nabla f) = 0$

\vec{F} conservative $\Rightarrow \text{Curl}(F) = 0$

$\text{Curl}(F) \neq 0 \Rightarrow \vec{F}$ not conservative

$\text{Curl}(F) = 0$ & simply connected & continuous partials $\Rightarrow \vec{F}$ conservative

has second partials $\Rightarrow \text{div}(\text{Curl}(F)) = 0$

$\text{div}(F) \neq 0 \Rightarrow F$ not a curl

16.6 Parametric Surfaces and Areas

Parametric Surface: $r(u,v) = \langle x(u,v), y(u,v), z(u,v) \rangle$

Surface of Revolution: $x = x$ $y = f(x)\cos\theta$ $z = f(x)\sin\theta$
(about x -axis)

Normal vector to tangent plane is $r_u \times r_v$

$$\text{Surface Area: } A(s) = \iint_D |r_u \times r_v| dA = \iint_D \sqrt{1 + \left(\frac{dz}{dx}\right)^2 + \left(\frac{dz}{dy}\right)^2} dA$$

16.7 Surface Integrals

Surface Integral / Flux:

$$\iint_S F \cdot dS = \iint_S F \cdot n dS = \iint_S F \cdot (r_u \times r_v) dA = \iint_D (-P \frac{\partial z}{\partial x} - Q \frac{\partial z}{\partial y} + R) dA$$

$$= \iint_D f(r(u,v)) |r_u \times r_v| dA = \iint_D f(x,y,g(x,y)) \sqrt{\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2 + 1} dA$$

16.8 Stokes' Theorem

- ① S is piecewise smooth surface
- ② Bounded by simple, closed piecewise-smooth boundary curve C w/ positive orientation
- ③ Continuous partial derivatives

$$\int_C F \cdot dr =$$

$$\iint_S \text{Curl } \vec{F} \cdot dS$$

16.9 Divergence Theorem

- ① E is simple solid region
- ② S is boundary surface of E w/ positive outward orientation
- ③ continuous partials on open region contains E

$$\iint_S F \cdot dS =$$

$$\iiint_E \text{div } F dV$$

$$\text{Vol(Sphere)} = \frac{4}{3} \pi r^3$$