

AtlasJawSpeech000005

Restoring the Stomatognathic-Vocal Continuum: A Comprehensive Treatise on Phase 4 Proprioceptive Integration

Abstract

The functional rehabilitation of the stomatognathic system—encompassing the mandible, dentition, lingual complex, and hyoid suspension—requires a paradigm shift from mechanistic correction to neuro-proprioceptive reintegration. This report, serving as an exhaustive expansion of "Section VIII. Therapeutic Protocols: Control and Finesse" from the referenced clinical dossier, specifically targets **Phase 4: Vocal Proprioception Integration**. The clinical profile presented—characterized by orthodontic refractoriness (ankylosis/tracking failure), chronic mentalis hyperactivity, referred cervicocranial tension (Atlas/C1), and a paradoxical speech pathology where relaxation induces lisping—indicates a profound dysregulation of the sensorimotor feedback loop. The patient has compensated for intrinsic instability by recruiting gross skeletal musculature (masseter/temporalis) to stabilize the articulators, a strategy that collapses upon relaxation. This document delineates the physiological basis of the patient's proprioceptive experiences (specifically the vibrotactile distinction between "lung" resonance and "tongue" tuning) and provides a rigorous, multi-modal library of rehabilitative exercises. These protocols aim to decouple the tongue from the mandibular fulcrum, restore the "tensegrity" of the vocal tract, and re-map the cortical homunculus to allow for "Control and Finesse" without pathological rigidity.

I. Pathophysiology of the Stomatognathic-Vocal Feedback Loop

To effectively implement Phase 4 of the therapeutic protocol, it is imperative to deconstruct the underlying pathology that necessitates such granular intervention. The user's condition is not merely a collection of symptoms but a cohesive, if maladaptive, neuromuscular strategy.

1.1 The Compensatory Scaffolding and the "Relaxation Paradox"

The clinical observation that "relaxing my jaw makes me speak with a lisp or speech difficulty"

¹ is the diagnostic cornerstone of this phase. In a neurotypically functioning stomatognathic system, the mandible provides a dynamic but stable platform, opening and closing to set the

aperture (vowel height), while the tongue operates as an independent hydrostat, executing rapid, precise deformations to articulate consonants and shape resonance.²

However, in the presence of **Jaw Instability**—often caused by malocclusion, aligner-induced bite shifts, or chronic TMJ dysfunction—the brain detects that the mandible is an unreliable platform. To compensate, the motor cortex recruits the jaw-closing muscles (masseter, temporalis, medial pterygoid) to "splint" or rigidly fixate the jaw in a semi-closed position. This fixation provides the tongue with a solid, albeit immobile, floor against which it can push to generate speech sounds, particularly sibilants (/s/, /z/, /sh/) which require precise air channeling.³

When the patient follows the therapeutic instruction to "relax the jaw," they actively inhibit this compensatory splinting. The mandible, lacking intrinsic proprioceptive stability and graded control, becomes a "wobbly" platform. The tongue, which has not developed independent motor control (Tongue-Jaw Dissociation), cannot adjust to this instability. It flails or flattens, failing to create the medial groove necessary for sibilance, resulting in lateral or interdental airflow—a lisp.¹ Therefore, the lisp is not a sign of failure, but a sign of *unmasking* the underlying lack of intrinsic lingual tone. Phase 4 must rebuild this tone so the tongue can function without the jaw's rigid assistance.

1.2 The Proprioceptive "Short-Circuit": Trigeminal-Cervical Convergence

The user reports a sensation connecting the "front teeth to the Atlas (C1) and the back".¹ This is anatomically validated by the **Trigeminocervical Complex**. The spinal nucleus of the Trigeminal nerve (CN V), which carries sensory information from the periodontal ligaments (teeth) and jaw muscles, descends into the upper cervical spinal cord (C1-C3).¹ Here, it shares synaptic real estate with sensory fibers from the suboccipital muscles and the upper neck.

- **The Mechanism of Referenced Tension:** When the mentalis muscle and orbicularis oris are chronically hyperactive (as seen in the "chin button" strain to hold the aligners or stabilize the jaw), they bombard the trigeminal nucleus with nociceptive and proprioceptive noise.
- **The Ripple Effect:** This neural "noise" spills over into the cervical pathways. The brain interprets dental tension as neck tension, and conversely, instability at the Atlas (C1)—often caused by the asymmetric torque of a deviated jaw—is felt as tension in the teeth.
- **Therapeutic Implication:** Rehabilitation cannot treat the jaw in isolation. Exercises must integrate cervical stability (neck posture) with mandibular mobility to "quiet" this neural crosstalk. The sensation of the "back" involvement further implicates the *Superficial Back Line* (SBL) of fascia, where tension in the brow and jaw travels over the scalp and down the erector spinae.¹

1.3 The Mentalis Anchor and Dental Ankylosis

The specific mention of a single lower incisor failing to track and the presence of ankylosis¹ adds a mechanical barrier to the neuromuscular problem.

- **Ankylosis:** If a tooth is ankylosed (fused to the bone), it acts as an absolute anchor. The aligner pushes against it, but the tooth does not move; instead, the reaction force displaces the aligner and torques the mandible.
- **Mentalis Hyperactivity:** The mentalis muscle originates *below* the incisors. If it is hypertonic (tight), it creates an upward/inward force vector that opposes the intrusive or labial forces of the aligner. This muscle acts as a "stress sponge" for the amygdala.
- **Phase 4 Goal:** We must use vocal exercises (like straw phonation and specific lip trills) to actively inhibit the mentalis while engaging the phonatory system, teaching the brain that it is safe to speak without "grabbing" the chin.

II. Psychoacoustics and Somatosensory Mapping: "Lungs" vs. "Tongue"

The user describes a distinct proprioceptive dichotomy: "speaking from the lungs for bass versus the tongue for high notes".¹ This is a profound insight that aligns perfectly with the physics of **Source-Filter Theory** and **Vocal Resonators**. Phase 4 leverages this accurate sensation to guide the rehabilitation process.

2.1 The Physics of the "Lung" Sensation (Chest Voice)

When producing low frequencies (bass/chest voice), the vocal folds are short, thick, and compliant (Thyroarytenoid muscle dominance).

- **Subglottal Resonance:** Low-frequency sound waves have long wavelengths. These waves are capable of coupling with the air column *below* the vocal folds (the trachea and bronchial tree). This phenomenon, known as **Tracheal Pull** or subglottal resonance, causes a sympathetic vibration in the sternum, rib cage, and clavicles.⁵
- **Proprioceptive Feedback:** The mechanoreceptors in the chest wall detect this vibration. The user feels the sound "in the lungs" because the chest cavity is physically vibrating.
- **The "Appoggio" Connection:** To maximize this resonance, the singer must manage the subglottal pressure using the inspiratory muscles (intercostals and diaphragm) to resist the collapse of the lungs. This technique, **Appoggio**, creates a sensation of "leaning" or pressurized stability in the torso, further reinforcing the "lung" connection.⁷

2.2 The Physics of the "Tongue" Sensation (Head Voice)

As pitch ascends, the cricothyroid muscles stretch and thin the vocal folds. The fundamental

frequency increases, and the wavelengths shorten.

- **Supraglottal Tuning:** High frequencies do not resonate effectively in the large subglottal space. To amplify them, the vocal tract (the filter) must be tuned. Specifically, the **Second Formant (F2)** needs to be raised to match the harmonics of the high pitch.⁹
- **The Role of the Tongue:** The most effective way to raise F2 is to reduce the volume of the oral cavity. This is achieved by arching the body of the tongue high toward the hard palate (similar to the vowel /i/ or "ee").
- **The "Mask" Sensation:** When the vocal tract is perfectly tuned, the acoustic energy is maximized in the oral and nasal cavities. This creates a sensation of sympathetic vibration in the facial bones (maxilla, zygomatic arch), often described as "mask resonance" or feeling the sound "in the tongue".⁶
- **Proprioceptive Shift:** The user effectively feels the *mechanical effort* of the tongue arching and the *vibrotactile result* in the face. If they try to carry the "lung" (flat tongue) position into the high notes, they will shout or strain (pulling chest). If they try to use the "tongue" (arched) position for bass, the sound will be thin.

2.3 The Somatic Map

The table below summarizes the physiological reality behind the user's sensory experience, guiding the selection of Phase 4 exercises.

Sensory Location	Vocal Register	Dominant Muscle	Acoustic Physics	Tongue Posture	Rehabilitation Goal
"Lungs" / Chest	Bass / Chest Voice	Thyroarytenoid (TA)	Subglottal Resonance / Tracheal Pull	Flat, relaxed, wide pharynx	Stabilize subglottal pressure (Appoggio) without pressing the larynx.
"Tongue" / Mask	High Notes / Head Voice	Cricothyroid (CT)	Supraglottal Tuning / F2 Amplification	High Arch (Posterior Elevation)	Train independent tongue arching ("Cat Arch") without jaw assistance.

III. Therapeutic Phase 4A: Mechanical Decoupling (The Foundation)

Before the tongue can be trained to "tune" the voice, it must be liberated from the jaw. The "lisp on relaxation" proves that the tongue is currently parasitic on the mandible. We must break this dependency using mechanical isolation.

3.1 Bite Block Therapy: The "Isolator"

This protocol forcibly stabilizes the jaw, inhibiting the masseter and temporalis muscles, and compelling the brain to develop a new motor pathway for the tongue.¹

Exercise 4.1: The Static Bite Drill

- **Goal:** Establish tongue movement without mandibular support.
- **Tool:** A specialized bite block (e.g., TalkTools) or a stack of 3-4 tongue depressors (approx. 10mm height). Alternatively, a firm cork or rubber eraser.
- **Placement:** Insert the block between the molars on the side *opposite* the deviated/ankylosed tooth (to encourage symmetry) or alternate sides.
- **Execution:**
 1. Bite down gently but firmly. The jaw must remain frozen on the block.
 2. **Tongue Protrusion:** Extend the tongue tip straight out of the mouth. Hold for 3 seconds. Retract. Repeat 10 times. *Biofeedback:* Watch in a mirror. Does the tongue deviate to the side? Keep it laser-straight.¹²
 3. **Tongue Lateralization:** Move the tongue tip to the right corner of the mouth, then the left. The jaw must not grind against the block to "help".¹³
 4. **The "Lalala" Sweep:** While biting the block, lift the tongue tip to the alveolar ridge (bump behind top teeth) and say "La-La-La." Then say "Ta-Ta-Ta" and "Na-Na-Na."
- **Insight:** The user will likely feel fatigue at the *root* of the tongue (hyoid connection). This is positive; it indicates the intrinsic tongue muscles are finally working.

3.2 Dynamic Jaw Grading: The "Goldfish"

Once the tongue can move with a *fixed* jaw, we must train the tongue to stay stable while the jaw moves. This addresses the "wobble" that causes the lisp.¹³

Exercise 4.2: The Tongue-Anchored Opener

- **Goal:** Dissociate jaw opening from tongue retraction.
- **Setup:** Place the tip of the tongue firmly on the "Spot" (alveolar ridge, just behind upper front teeth).
- **Execution:**
 1. Keep the tongue glued to the Spot.
 2. Slowly open the jaw as wide as possible *without* the tongue pulling off the roof of the

mouth.

3. Slowly close the jaw.
 4. Repeat 10 times.
- **The Mechanism:** This forces the tongue to stretch its frenulum and engage the superior longitudinal muscle while the jaw depressors (digastric/mylohyoid) work independently. If the tongue snaps back, the range of motion is too large. Reduce the range until control is maintained.¹⁵

Exercise 4.3: The "Wiggle" Release

- **Goal:** Deactivate the stretch reflex in the masseter.
- **Action:**
 1. Grasp the chin with the thumb and index finger.
 2. Use the *hand* to wiggle the jaw loosely up and down, and side to side.
 3. The jaw muscles must be completely flaccid. The jaw should move like a "ventriloquist's dummy" or a "plumber's wrench".¹⁶
 4. **Vocalization:** Add a gentle phonation ("Uh-Uh-Uh") while wiggling. If the sound stops or the jaw stiffens, pause and reset. This disconnects the "stress" signal from the amygdala to the jaw.¹⁷

IV. Therapeutic Phase 4B: Lingual Proprioception (The "Tongue" Connection)

This section addresses the user's specific ability to feel high notes in the tongue. We will refine this into a conscious technique called **Vocal Tract Tuning**, using the "Cat Arch" visualization to master the transition from chest to head voice without strain.

4.1 The "Cat Arch" Visualization (Posterior Tongue Elevation)

To access high notes without the jaw helping (clenching), the posterior tongue must arch up to tune the resonance (F2). This is the "Tongue" feeling the user described.¹⁸

Exercise 4.4: The "Ng" Siren (The Homunculus Remapper)

- **Goal:** Train the posterior tongue (styloglossus/palatoglossus) to elevate independently of the jaw.
- **Setup:** Say the word "Sing." Hold the "Ng" sound. The back of the tongue should be sealed against the soft palate. The middle and tip of the tongue should be relaxed.
- **Check:** Pinch the nose. The sound should stop immediately. This confirms nasal resonance.²⁰
- **Visualization:** Imagine the tongue is a cat. As you ascend in pitch, the "cat" (tongue body) arches its back higher toward the roof of the mouth.
- **Execution:**

1. Start on a low, comfortable note on "Ng."
2. Slowly glide (siren) up to a high note.
3. **Crucial Constraint:** The jaw must remain relaxed and slightly open. It must *not* close as the pitch goes up. If the jaw wants to close, insert a pinky finger or bite block to prop it open.²¹
4. **Sensation:** As the pitch rises, feel the vibration move from the throat/chest into the "mask" (nose/cheekbones). This confirms the "Tongue" is doing the tuning.²³

Exercise 4.5: The "Gaga" / "Kermit" Drill

- **Goal:** Strengthen the tongue root's ability to articulate without jaw recruitment.
- **Sound:** Use a "dopey" or "Yogi Bear" voice. This lowers the larynx and opens the pharynx.²⁴
- **Execution:**
 1. Let the jaw hang open and "dead."
 2. Say "Ga-Ga-Ga" using *only* the back of the tongue. The jaw must not bounce.
 3. Sing an arpeggio (1-3-5-8-5-3-1) on "Ga."
 4. **Proprioception:** Feel the back of the tongue springing up to hit the soft palate like a trampoline. This builds the specific strength needed for the "Tongue" resonance the user feels on high notes.²⁵

4.2 Tongue-Jaw Dissociation in Vowels

The lisp often occurs because the tongue flattens during vowel transitions.

Exercise 4.6: The Fixed-Jaw Vowel Chain

- **Goal:** Articulate vowels using only tongue shape, not jaw height.
- **Setup:** Open the jaw to a single finger-width. Place a finger on the chin to detect movement.
- **Execution:**
 1. Chant the sequence: "EE - EH - AH - OH - OO."
 2. **The Rule:** The jaw must not move *at all*.
 3. **Mechanism:**
 - **EE:** Tongue arches high (Cat Arch).
 - **AH:** Tongue flattens.
 - **OO:** Lips round (Mentalis must stay relaxed!).
 4. If the jaw closes for "EE" or opens for "AH," the user is using the "crutch." Reset and repeat until the tongue does the work.²⁶

V. Therapeutic Phase 4C: Respiratory-Laryngeal Coordination (The "Lung" Connection)

This section addresses the user's "Bass/Lung" sensation. This is the domain of **Appoggio**—the dynamic balance of breath pressure that allows for deep, resonant chest voice without grinding the teeth.⁷

5.1 Sternal Mapping and Subglottal Pressure

The "lung" sensation is physical vibration of the bronchial tree. To maintain this, the larynx must remain low and the vocal folds thick.

Exercise 4.7: The Sternal "Mmm" Scan

- **Goal:** Maximize chest resonance via tactile feedback.
- **Action:**
 1. Place a flat hand firmly on the sternum (center of chest).
 2. Hum a low "Mmm."
 3. **Calibration:** Vary the pitch and "weight" of the hum until you feel the *maximum* vibration under your hand. It should feel like a purring cat in the chest.²⁸
 4. **Transition:** Open the mouth to "Mah." Try to keep the vibration in the hand. If the vibration disappears, the throat has tightened, or the breath support (Appoggio) has collapsed.
 5. **Application:** Speak a phrase ("Hello there") while maintaining the sternal buzz. This grounds the voice and prevents the "lisp" caused by shallow, high-larynx breathing.

Exercise 4.8: The Modified Farinelli Maneuver

- **Goal:** Stabilize the "Lung" feeling by controlling the diaphragm's ascent.
- **Execution:**
 1. **Inhale:** Breathe in silently for 4 seconds. Feel the lower ribs and flanks expand (360-degree expansion).
 2. **Suspend:** Hold the breath for 4 seconds *without* closing the throat (glottis). Maintain the rib expansion. This is the "Inspiratory Hold".⁷
 3. **Exhale:** Hiss on a sharp "Sssss" for 8 seconds.
 4. **The Challenge:** While hissing, do *not* let the ribs collapse. Ideally, push the ribs *out* slightly as you exhale. This eccentric contraction of the inspiratory muscles provides the steady "lung" pressure the user describes.³⁰

VI. Therapeutic Phase 4D: Semi-Occluded Vocal Tract (SOVT) & The Neurological Reset

SOVT exercises are the "bridge" between the mechanical exercises and functional speech. By narrowing the mouth opening (using a straw), we create back-pressure that "splints" the airway from the inside out, reducing the need for the jaw to clench.³¹

6.1 Straw Phonation: The Ultimate Reset

This is the most effective tool for the user's specific "misshapen jaw" and "lisp" symptoms.

Exercise 4.9: The Water Bubble Drill

- **Goal:** Balance airflow and relax the mentalis.
- **Tools:** A small diameter straw and a glass of water (half full).
- **Action:**
 1. Place the straw 2cm deep into the water.
 2. Blow steady bubbles (no sound). The airflow must be continuous.
 3. Add a gentle "Ooo" sound (phonation).
 4. **Glide:** Slide the pitch up and down (siren) while maintaining steady bubbles.
- **Feedback:**
 - If bubbles stop: Breath support failed.
 - If water splashes: Too much pressure/tension.
 - If pitch breaks: Jaw/Tongue are fighting.
- **Result:** The user will feel a "buzzy" sensation behind the lips. This "mask" sensation replaces the "tight jaw" sensation.³²

Exercise 4.10: The Mentalis Inhibition

- **Goal:** Deactivate the "Chin Button" (Mentalis muscle).
- **Action:**
 1. Hold the straw with the *lips only*. Do not bite it.
 2. Look in a mirror. The chin must be smooth/flat, not dimpled ("golf ball chin").
 3. Hum a tune through the straw.
 4. If the chin dimples or bunches up, stop. Massage the chin down and try again. This breaks the habit of using the chin to "push" the voice out.³³

VII. Integration: From Exercise to "Control and Finesse"

The final step is to merge these isolated skills. The user's goal is "Control and Finesse."

7.1 The "Siren-Speech" Protocol

To connect the "Lung" (Bass) and "Tongue" (High) registers seamlessly:

1. **Start:** Begin with the **Sternal "Mmm"** (Chest/Lung feel).
2. **Glide:** Slide up an octave on "Ng" (Tongue/Mask feel). Visualize the "Cat Arch."
3. **Hold:** At the top note, open the "Ng" to a vowel ("Nng-AAH"). Keep the "Tongue" feeling (mask resonance). Do not drop the weight back into the throat.³⁴
4. **Speak:** Immediately speak a sentence in that placement ("Hey, how are you?").

5. **Result:** The speech should feel "forward" and buzz in the face, not the throat. The jaw should be loose.

7.2 The "Lisp" Correction Protocol

1. **Bite Block:** Place the block. Say "Sssss." The tongue must find the groove without the jaw helping.
2. **Remove Block:** Keep the jaw in the *exact same position* (open slightly). Say "Sssss."
3. **Drill:** Practice words like "Sister," "Season," "Zone" while monitoring the jaw in a mirror. If the jaw jerks or slides to the side, go back to the Bite Block.³⁵

Summary Table of Phase 4 Interventions

Symptom / Sensation	Physiological Cause	Rehabilitation Exercise	Target Mechanism
"Lisp on Relaxation"	Loss of compensatory jaw fixation; weak tongue grading	Bite Block Drills & Fixed-Jaw Vowels	Tongue-Jaw Dissociation; Intrinsic Lingual Tone
"Bass in Lungs"	Subglottal Resonance (Tracheal Pull)	Sternal "Mmm" Scan & Farinelli Breathing	Appoggio; Thoracic Vibrotactile Feedback
"High Notes in Tongue"	Supraglottal Tuning (F2 coupling)	"Ng" Siren & "Cat Arch" Visualization	Posterior Tongue Elevation; Vocal Tract Tuning
"Misshapen Jaw"	Asymmetric Pterygoid/Masseter tension; Ankylosis anchor	"Goldfish" Exercise & "Wiggle" Release	Rotational vs. Translational Control; De-patterning
"Chin Button" Tension	Mentalis Hyperactivity (Amygdala stress response)	Straw Phonation (No Bite)	Mentalis Inhibition; SOVT Reset

VIII. Conclusion

The journey through Phase 4 is one of sensory re-education. By validating the user's proprioceptive intuitions—that bass resonates in the "lungs" and high notes require "tongue" finesse—we transform abstract symptoms into actionable biofeedback. The "lisp" is not a regression but a necessary clearing of the debris of compensation. Through the rigorous application of mechanical decoupling (bite blocks), acoustic tuning (cat arch/Ng), and respiratory grounding (Appoggio), the stomatognathic system can be rehabilitated from a rigid, defensive posture into a state of dynamic, articulate equilibrium.

Works cited

1. Jaw, Teeth, Speech, and Body Connection.pdf
2. Release Tongue and Jaw Tension | Exercises For Singers | Guided Vocal Coaching | Sing With Freedom - YouTube, accessed December 7, 2025, <https://www.youtube.com/watch?v=iKAnNY6Erjs>
3. Singers: RELAX Jaw and Tongue Tension with These Vocal Exercises - YouTube, accessed December 7, 2025, <https://www.youtube.com/watch?v=Dtf3QtyMKuQ>
4. 3 Exercises to Improve Independence of the Tongue, Jaw, Head, and Neck - YouTube, accessed December 7, 2025, <https://www.youtube.com/watch?v=6OymciD8APl>
5. Brett Manning Shows You How To Master Chest Voice Placement - Singing Success, accessed December 7, 2025, <https://singsuccess.com/master-chest-voice/>
6. Heavy Lifting! The Muscles that Control the Vocal Cords - Caveness Voice, accessed December 7, 2025, <https://www.cavenessvoice.com/blog/heavy-lifting-the-muscles-that-control-the-vocal-cords>
7. Breathing Support for Singing - Toronto Choral Society, accessed December 7, 2025, <https://www.torontochoralsociety.org/breathing-support>
8. PRINCIPLES OF APPOGGIO: THE INTERRELATIONSHIP BETWEEN THEORY AND PRACTICE FOR TODAY'S YOUNG OPERA SINGERS by Paul Han Submitt - IU ScholarWorks, accessed December 7, 2025, <https://scholarworks.iu.edu/dspace/bitstreams/7a4de808-0e7f-4236-89fb-bfac2956bb64/download>
9. Vocal resonance tuning and vowel changes, accessed December 7, 2025, <https://newt.phys.unsw.edu.au/jw/soprane.html>
10. Formant Changes in Amateur Singers After Instruction in a Vowel Equalization Technique - BYU ScholarsArchive, accessed December 7, 2025, <https://scholarsarchive.byu.edu/cgi/viewcontent.cgi?article=3053&context=etd&httpsredir=1&referer=>
11. Try These 10 Proprioception Exercises To Help With Balance, Control, and Coordination, accessed December 7, 2025, <https://www.healthline.com/health/fitness/proprioception-exercises>
12. Oral Motor Exercises, accessed December 7, 2025, <https://userweb.ucs.louisiana.edu/~ncr3025/roussel/codi531/oralmotor>
13. Tongue Range of Motion Exercises - University of Mississippi Medical Center,

- accessed December 7, 2025,
<https://umc.edu/Healthcare/ENT/Patient-Handouts/Adult/Speech-Language-Pathology/Swallowing/Tongue-ROM-Exercises.html>
14. Chew on This – Simple Jaw Exercises for TMJ Relief – Champaign Dental Group, accessed December 7, 2025,
<https://www.champaigndentalgroup.org/about-us/blog-articles/jaw-exercises>
 15. 9 Best TMJ Exercises for Pain Relief – Healthline, accessed December 7, 2025,
<https://www.healthline.com/health/tmj-exercises>
 16. Jaw Release Exercise – YouTube, accessed December 7, 2025,
<https://www.youtube.com/watch?v=2VIAPD7TIWA>
 17. Jaw Tension In Singing – Learn To Release It So You Can Sing! – Vocal Coach Dylan, accessed December 7, 2025,
<https://vocalcoachdylan.com/jaw-tension-in-singing/>
 18. Andre Malraux The Royal Way – Angkor Database, accessed December 7, 2025,
<https://cdn.angkordatabase.asia/libs/docs/malraux-royalway-gilbert.pdf>
 19. /r/ vent : r/slp – Reddit, accessed December 7, 2025,
https://www.reddit.com/r/slp/comments/lx1sf7/r_vent/
 20. Voice Lessons: Use the "NG" to Find Mask Resonance & Make Singing Easier – YouTube, accessed December 7, 2025,
https://www.youtube.com/watch?v=3A_BtIXhOCw
 21. What would be your top 3 vocal exercises/techniques to advice a beginner singer? – Reddit, accessed December 7, 2025,
https://www.reddit.com/r/singing/comments/1de3wyi/what_would_be_your_top_3_vocal/
 22. Ep. 101 – “The Greatest Vocal Exercise of All Time” – Voice Lessons To The World – YouTube, accessed December 7, 2025,
https://www.youtube.com/watch?v=h0SIEY_zNqU
 23. How To Do An NG Hum – Open Up Your Head Resonance | Learn How To Sing, accessed December 7, 2025,
<https://vocalcoachdylan.com/how-to-do-an-ng-hum-open-your-head-resonance/>
 24. High note exercise 11: New Year, New Voice Series – YouTube, accessed December 7, 2025, <https://www.youtube.com/watch?v=s3kUZycJ9Kq>
 25. Exercises to Reduce Tongue Tension In Your Singing – Ziegenfuss Voice Studio LLC, accessed December 7, 2025,
<https://www.ziegenfussvoicestudio.com/blog/5-helpful-exercises-to-reduce-tongue-tension-when-singing>
 26. Tongue/Jaw isolation exercise 10: New Year, New Voice Series – YouTube, accessed December 7, 2025, <https://m.youtube.com/shorts/heCl6sfmLJ0>
 27. Correct Breathing and "Support" for Singing – SingWise, accessed December 7, 2025,
<https://www.singwise.com/articles/correct-breathing-and-support-for-singing>
 28. Build an UNSTOPPABLE Chest Voice! 5 Game-Changing Exercises, accessed December 7, 2025, <https://ramseyvoice.com/chest-voice/>
 29. 100 Vocal Exercises DAY 8: Vibrating The Torso – YouTube, accessed December 7, 2025, <https://www.youtube.com/watch?v=u3zPuMMIDrA>

30. The FASTEST Way to Master Your Breathing using the Appoggio Technique :
r/singing, accessed December 7, 2025,
https://www.reddit.com/r/singing/comments/1d542tj/the_fastest_way_to_master_your_breathing_using/
31. Semi-Occluded Vocal Tract Exercises - Stacey Menton, Soprano SLP, accessed
December 7, 2025,
<https://staceymenton.com/blog/semi-occluded-vocal-tract-exercises>
32. Semi occluded vocal tract exercises | CUH - Cambridge University Hospitals,
accessed December 7, 2025,
<https://www.cuh.nhs.uk/patient-information/semi-occluded-vocal-tract-exercises/>
33. The Power Of Straw Phonation: Exploring SOVT Exercises - Toronto Speech
Therapy, accessed December 7, 2025,
<https://www.torontospeechtherapy.com/blog/2024/the-power-of-straw-phonation-exploring-sovt-exercises/>
34. Vocal Exercises for Singers: Breathing and Beginner Exercises - Musika Music
Lessons, accessed December 7, 2025,
<https://www.musikalessons.com/blog/2016/10/vocal-exercises-for-singers/>
35. Jaw Stabilization for the Lateral Lisp - Marshalla Speech & Language, accessed
December 7, 2025,
<https://pammarshalla.com/jaw-stabilization-for-the-lateral-lisp/>