

CSMHuman00000000015

\The Resonant Human: An Exhaustive Synthesis of Frequency-Dependent Biodynamics and Radiofrequency Dosimetry

1. Introduction: The Spectral Continuum of Biological Interaction

The human organism, a complex amalgamation of viscoelastic tissues, fluid-filled cavities, and electrochemical networks, exists within a ceaseless ocean of oscillatory energy. This energy, spanning the spectrum from the infrasonic judder of heavy machinery to the hyper-frequency pulsations of modern radar and telecommunications, does not merely pass through the body; it interacts with it. The nature of this interaction—whether the energy is absorbed, transmitted, or amplified—is dictated by the principles of resonance and impedance matching. This report presents a comprehensive, high-resolution synthesis of the physical mechanisms governing these interactions, anchored in the seminal research of the U.S. Air Force School of Aerospace Medicine (USAF SAM) and the definitive biodynamic investigations of the mid-20th century.

The scope of this analysis is bifurcated yet deeply interconnected. On one end of the spectrum lies the domain of **Mechanical Biodynamics** (0 Hz to 100 Hz), where the body acts as a system of coupled masses and springs, susceptible to structural resonance and inertial deformation. On the other end lies **Radiofrequency (RF) Dosimetry** (100 kHz to 100 GHz), where the body functions as a lossy dielectric antenna, absorbing electromagnetic energy and converting it into thermal or acoustic phenomena.

The convergence of these fields is found in the concept of the "critical frequency"—specific spectral windows where the geometric and material properties of the human form couple maximally with the incident energy. Understanding these windows is not merely an academic exercise in biophysics; it is the foundation of occupational safety standards, aerospace design, and medical diagnostics. The historical trajectory of this research, driven by the exigencies of the Cold War—high-speed ejection seats, low-altitude turbulence, and high-power radar installations—provides the empirical backbone for our modern understanding of how the human body withstands the invisible forces of its environment.¹

2. Mechanical Biodynamics: The Low-Frequency Spectrum

The interaction of the human body with mechanical vibration is characterized by non-linear transmissibility. Unlike a rigid solid, the body is a damped, multi-degree-of-freedom system. When subjected to sinusoidal vibration, specific anatomical segments resonate, amplifying the input acceleration by factors that can exceed 200% of the driving amplitude. The research conducted by Magid, Coermann, and Ziegenruecker between 1959 and 1962 remains the definitive characterization of these low-frequency responses.⁴

2.1 Whole-Body Impedance and the Thoraco-Abdominal System

The fundamental mechanical resonance of the seated human body occurs in the frequency band of **4 to 8 Hz**. This resonance is primarily driven by the thoraco-abdominal complex—a mass-spring system comprising the visceral mass (liver, stomach, intestines), the diaphragm, and the anterior abdominal wall.

2.1.1 The Visceral Mass Oscillator

Under vertical (+Gz) vibration, the visceral mass oscillates within the abdominal cavity. At frequencies below 2 Hz, the body behaves as a relatively rigid mass; the internal organs move in phase with the skeletal frame. However, as the frequency approaches the 4–8 Hz window, the visceral mass begins to resonate. The diaphragm, acting as the upper boundary of this system, is forced upward and downward by the momentum of the abdominal contents.

This oscillation has profound physiological consequences. The movement of the diaphragm creates a "pumping" action on the lungs, forcing air in and out of the respiratory tract independent of voluntary control. This phenomenon, known as "vibration-induced hyperventilation," was documented in Magid's tolerance studies, where subjects reported dyspnea (shortness of breath) and thoracic pain.⁵ The mechanical impedance of the body—defined as the ratio of driving force to velocity—peaks sharply in this band, indicating that the body is absorbing maximum mechanical energy from the source.⁷

2.1.2 Spinal Transmissibility and Injury Mechanisms

The spinal column serves as the primary strut transmitting seat vibration to the head. In the 4–8 Hz range, the spine undergoes compression and buckling modes. The intervertebral discs and the paravertebral musculature act as dampers, but at resonance, their capacity is exceeded. The transmissibility ratio (head acceleration / seat acceleration) can reach values of 1.5 to 2.0, meaning the head is accelerating twice as violently as the vehicle floor.⁷

Secondary resonances appear in the **20 to 30 Hz** range, associated with the head-neck system. Here, the head acts as an inverted pendulum stabilized by the neck musculature. While the displacement amplitudes are smaller than in the 4–8 Hz range, the higher frequency

results in significant acceleration loads ($a = \omega^2 x$), potentially causing flexion-extension injuries to the cervical spine if the head is unsupported.

2.2 Short-Time Human Tolerance Curves

The definition of "tolerance" in biodynamics is complex, encompassing subjective discomfort, performance degradation, and tissue injury. The tolerance curves developed by Magid and Coermann (1960) for short-duration exposures (1 minute and 3 minutes) delineate the safe operational envelope for human vibration.⁴

2.2.1 The 1-Minute and 3-Minute Boundaries

The tolerance limit is lowest in the resonant band, reflecting the body's mechanical vulnerability.

- **4–8 Hz Critical Zone:** For a one-minute exposure, the tolerance limit is approximately **1.0 g** (peak). For a three-minute exposure, this drops to **0.5 g**. Subjects exposed to vibrations exceeding these limits report alarming symptoms, including substernal pain (mimicking angina), testicular pain (due to spermatic cord resonance), and profound general malaise.⁵
- **High-Frequency Robustness:** As the frequency increases beyond 10 Hz, tolerance improves significantly. At 20 Hz, the human frame can withstand nearly **3.0 g** for one minute. This increase in tolerance is due to the damping of the soft tissues, which attenuate higher-frequency inputs before they can excite large-mass resonances.⁷

The following table summarizes the subjective responses and physiological limits identified in these foundational studies:

Frequency Band	Primary Resonance Site	Subjective Symptoms	1-Min Tolerance Limit (g)
1 – 3 Hz	Whole Body (Rigid)	Motion sickness, swaying	> 2.0 g
4 – 8 Hz	Thoraco-Abdominal System	Chest pain, dyspnea, abdominal stress	0.5 – 1.0 g
10 – 12 Hz	Bladder/Pelvic Floor	Urge to defecate/urinate, back pain	1.0 – 1.5 g

15 – 20 Hz	Head-Neck System	Head vibration, speech modulation	2.0 – 3.0 g
> 20 Hz	Eyeballs (Intra-ocular)	Visual blurring, anxiety	> 3.0 g

2.3 Visual Acuity Under Sinusoidal Vibration

The degradation of visual performance during vibration is a critical factor in aerospace operations. Research by Lange and Coermann (1962) and subsequent analysis by Dennis (1965) revealed that this degradation is not linear; it is a function of the complex interplay between head transmissibility and ocular resonance.⁶

At low frequencies (< 4 Hz), the Vestibulo-Ocular Reflex (VOR) is highly effective. The semicircular canals detect head rotation and drive the extraocular muscles to counter-rotate the eyes, stabilizing the image on the retina. However, the VOR has a limited frequency response. As vibration frequency enters the 4–12 Hz resonant band, the phase lag of the VOR increases, and the gain decreases. The eyes can no longer compensate for head movement, resulting in "retinal slip" or image smear.³

At higher frequencies (> 12 Hz), the amplitude of head oscillation decreases due to body damping, but any vibration that *does* reach the head is problematic because the VOR is effectively useless at these rates. This sets the stage for the specific phenomenon investigated by Morton Ohlbaum: the resonance of the eye itself.

3. Ocular Biodynamics: The Ohlbaum Report (1976)

In 1976, Morton K. Ohlbaum of the Aerospace Medical Research Laboratory published the landmark doctoral thesis *Mechanical Resonant Frequency of the Human Eye In Vivo* (DTIC ADA030476). This document addressed a critical gap in biodynamic theory: the inadequacy of head-movement models to explain visual performance decrements at specific frequencies and viewing distances.³

3.1 The Theoretical Conflict: Tracking vs. Structure

Prior to Ohlbaum's work, the prevailing hypothesis was that visual loss was solely a failure of oculomotor tracking. If the head moved, the eye tried to compensate; if it failed, vision blurred. Ohlbaum argued this was insufficient. He hypothesized that the eye, being a mass suspended in a viscoelastic socket (the orbit), must have its own resonant frequency. If the eye were to resonate, it would oscillate within the skull, adding a secondary layer of image motion independent of the skull's motion.¹²

3.2 Methodology: Mechanical Induction and Photogrammetry

Ohlbaum's experimental design was rigorous, designed to decouple head motion from eye motion.

- **Apparatus:** Subjects were coupled to a vibration generator via a rigid bite bar to induce precise sinusoidal inputs to the skull.
- **Measurement:** High-speed cinematography and corneal reflection techniques were used to track the movement of the globe relative to the canthus (corner of the eye).
- **Stimulus:** Frequencies ranging from sub-resonant to 50 Hz were swept to identify the point of maximum amplitude magnification.³

3.3 The 18 Hz Resonance Discovery

Ohlbaum successfully identified the resonant frequency of the human eye *in vivo* to be approximately **18 Hz**.³

This finding clarified the biodynamic map. The eye acts as a secondary oscillator mounted on the primary oscillator of the head.

- **Separation of Modes:** The 18 Hz ocular resonance is distinct from the 4–8 Hz torso resonance and the 20–30 Hz head resonance. This separation implies that vibration energy must traverse a complex "mechanical filter" chain—Seat $\rightarrow$ Torso $\rightarrow$ Neck $\rightarrow$ Skull $\rightarrow$ Orbit $\rightarrow$ Eye—before exciting this mode.
- **Damping Characteristics:** Crucially, Ohlbaum found that while resonance occurred, the amplification factor (Q) was relatively low. The orbital fat and extraocular muscles provide significant damping. He concluded that the amplification factor at 18 Hz was "insufficient to account for the observed performance" decrements on its own, suggesting that the breakdown of visual acuity is a summation of VOR failure, head resonance, and ocular resonance.³

3.4 Rotational vs. Translational Dynamics

One of the most significant insights from the Ohlbaum report was the characterization of the eye's movement mode. The data demonstrated that under vertical vibration, the major component of eye movement is **rotational** rather than translational.³

This is physically intuitive but optically critical. The eye is tethered by the optic nerve and pivoted by muscles. A translational shock to the skull results in a rotation of the globe around its center of rotation. Optically, a 1-degree rotation of the eye shifts the retinal image by approximately 300 microns (assuming a 17mm nodal distance), whereas a 1mm translation shifts it far less effectively. Thus, the *rotational* resonance of the eye at 18 Hz is the primary driver of visual blurring in that frequency band.

3.5 Modern Corroboration: Corneal Natural Frequency vs. Global

Resonance

It is imperative to distinguish Ohlbaum's "global resonance" from recent findings in ophthalmology regarding "corneal resonance."

- **Global Resonance (18–30 Hz):** This is the oscillation of the *entire eyeball* within the orbital soft tissue. Recent studies using high-speed cameras confirm peaks in the 25–31 Hz range, aligning with Ohlbaum's 18 Hz finding when accounting for variations in muscle tone and measurement technique.¹¹
- **Corneal Natural Frequency (100–300 Hz):** Modern Optical Coherence Elastography (OCE) has identified the resonant frequency of the *cornea itself* (the tissue membrane) to be much higher, typically centered around **250 Hz**.¹⁴

This distinction is vital for safety analysis. Environmental vibration (vehicles, machinery) rarely contains significant energy at 250 Hz, so the cornea is unlikely to resonate structurally in a damaging way during transport. However, the 18 Hz global resonance sits squarely in the harmonic range of helicopter rotors and rough-terrain vehicles, making it a persistent operational hazard.

4. The Radiofrequency Spectrum: The Shift to Dosimetry

Leaving the domain of mechanical mass-displacement, we ascend the frequency spectrum into the electromagnetic domain. Here, the interaction is not governed by Newton's laws of motion but by Maxwell's equations. The history of this field is dominated by the transition from measuring "what is in the air" (exposure) to "what is in the body" (dose).

4.1 The USAF SAM Radiofrequency Radiation Dosimetry Handbook

In the 1970s, parallel to Ohlbaum's work, the USAF School of Aerospace Medicine recognized that existing safety standards based on incident power density (mW/cm^2) were inadequate. They failed to account for the body's variable absorption efficiency. This led to the publication of the *Radiofrequency Radiation Dosimetry Handbook*, a "living document" that evolved through four major editions between 1976 and 1986.²

This handbook introduced and standardized the metric of Specific Absorption Rate (SAR).

$$SAR = \frac{\sigma |E|^2}{\rho}$$

Where:

- σ = Tissue conductivity (S/m)
- E = Internal electric field strength (V/m)

- ρ = Tissue density (kg/m³)

This shift was revolutionary. It acknowledged that a field of 10 mW/cm² might be harmless at one frequency but lethal at another, depending entirely on how well the body coupled with that frequency. The Handbook compiled vast tables of dielectric properties and SAR calculations for various body shapes (spheres, cylinders, block models), serving as the "Bible" for RF researchers.¹⁷

4.2 The "Standard Man" Resonance: 70 MHz

The most critical finding of RF dosimetry is the existence of a **Whole-Body RF Resonance**. Just as the torso resonates mechanically at 4–8 Hz, the human body resonates electromagnetically between **30 and 100 MHz**.¹⁹

4.2.1 The Dipole Antenna Model

The physics of this resonance are best described by antenna theory. An isolated human body in free space acts as a **half-wave dipole**. Resonance occurs when the body length (L) is approximately 0.4 times the wavelength (λ) of the incident field ($L \approx 0.4\lambda$).²¹

- **Standard Adult (1.75 m):** The resonant frequency is approximately **70–80 MHz**.
- **Grounded Condition:** If the person stands on a conductive ground plane (e.g., a metal deck), the body behaves as a **quarter-wave monopole**. The resonant frequency drops by half, to approximately **35–40 MHz**.²⁰
- **Scaling Factors:** Children or shorter adults resonate at higher frequencies, approaching **100 MHz**.

At this resonant frequency, the body's RF absorption cross-section can exceed its physical cross-section. The SAR increases dramatically—by a factor of 10 or more compared to non-resonant frequencies.¹⁹ This is why RF safety standards (like FCC or ICNIRP guidelines) impose the strictest exposure limits in the 30–300 MHz band; the body is essentially a tuned antenna for these frequencies.

4.2.2 SAR Distribution and "Hot Spots"

Unlike mechanical vibration, which stresses joints, RF resonance creates thermal "hot spots" driven by current density.

- **The Ankles:** In a standing, grounded person exposed to vertical polarization, the induced current must flow to the ground. The ankles, having a small cross-sectional area, experience extremely high current densities, leading to potentially severe local heating (high local SAR) even if the whole-body average SAR is low.²¹
- **The Neck:** Similarly, the narrowing of the conductive path at the neck can create a localized SAR peak.
- **The Brain:** At higher frequencies (Microwave bands > 300 MHz), the head itself can

resonate (acting as a sphere). This can lead to "hot spots" deep within the brain due to the focusing effect of the skull's curvature and dielectric properties.²²

5. Dielectric Dispersion: The Physics of Tissue Interaction

To understand *why* the body absorbs RF energy, one must look at the microscopic level. The *Radiofrequency Radiation Dosimetry Handbook* provides extensive data on the **dielectric properties** of tissues—permittivity (ϵ) and conductivity (σ). These properties are not constant; they vary with frequency according to dispersion mechanisms characterized by Herman P. Schwan.²³

5.1 Schwan's Dispersion Regions

Schwan identified three primary dispersion regions, labeled Alpha, Beta, and Gamma. Each corresponds to a specific biophysical relaxation mechanism.

5.1.1 Alpha Dispersion (α): 10 Hz – 10 kHz

Occurring at low frequencies (audio range), α -dispersion is dominated by **counter-ion polarization** along cell membranes. Ions in the extracellular fluid are attracted to the charged surface of the cell membrane, forming an ionic cloud. Under an alternating field, this cloud distorts, creating a massive induced dipole. This results in incredibly high permittivity values (often $> 10^5$) in tissues like muscle at low frequencies.²⁴

5.1.2 Beta Dispersion (β): 10 kHz – 100 MHz

This is the "Radiofrequency" dispersion. It is caused by the **Maxwell-Wagner interfacial polarization** of the cell membrane.

- **Mechanism:** The cell membrane acts as a thin capacitive barrier separating the conductive cytoplasm from the conductive extracellular fluid. At low frequencies, the membrane impedance is high ($X_c = 1/2\pi fC$), blocking current entry into the cell. As frequency increases into the MHz range, the capacitive reactance drops. The current begins to short-circuit the membrane capacitance.
- **Transition:** By 100 MHz, the membrane is effectively transparent to the current. The tissue transitions from a heterogeneous suspension of insulating shells to a more homogeneous lossy dielectric. This is why muscle conductivity rises significantly across this band.²⁶

5.1.3 Gamma Dispersion (γ): > 1 GHz

This is the "Microwave" dispersion. At these frequencies, the large biological structures (cells, membranes) are electrically invisible. The interaction is dominated by the **dipolar relaxation of water molecules**.

- **Mechanism:** Water molecules are permanent dipoles. As the field oscillates at GHz frequencies, the molecules try to rotate to align with the field. Viscous drag prevents them from tracking the field perfectly, leading to energy loss (heat).
- **Peak:** For pure water at 37°C, this relaxation peaks around 25 GHz. In tissues, interactions with proteins broaden this peak, but the fundamental mechanism remains water rotation.²⁶

5.2 Specific Tissue Data at 100 MHz

Based on the extensive databases compiled by the IT'IS Foundation and the USAF Handbooks, we can quantify the properties of key tissues at the upper edge of the β -dispersion (100 MHz), a critical frequency for FM radio and VHF TV exposure.²⁹

Tissue Type	Relative Permittivity (ϵ_r)	Conductivity (σ)	Biophysical Characteristic
Cortical Bone	~12 - 15	~0.02 - 0.04	Low water content; acts as insulator.
Fat (Infiltrated)	~6 - 12	~0.04 - 0.08	Hydrophobic; transparent to RF relative to muscle.
Muscle (Skeletal)	~60 - 70	~0.70 - 0.90	High water/ion content; primary absorber.
Brain (Grey Matter)	~65 - 75	~0.60 - 0.75	Highly conductive; susceptible to hot spots.
Blood	~60 - 65	~1.20 - 1.40	High ionic mobility; circulates heat.

This contrast between fat ($\sigma \approx 0.05$) and muscle ($\sigma \approx 0.8$) creates a "layering effect." RF waves incident on the body undergo reflection and refraction at these interfaces. Under certain conditions, a fat layer can act as an impedance matching transformer, actually *increasing* the energy deposited into the underlying muscle, contrary to the intuition that the fat would shield it.²

6. The Acoustic Bridge: The Frey Effect

Perhaps the most fascinating interaction in the entire frequency domain is the **Microwave Auditory Effect**, or "Frey Effect." First described by Allan Frey in 1961, this phenomenon allows humans to "hear" RF energy as clicks, buzzes, or chirps, occurring without the use of ears in the traditional sense.³¹

6.1 The Thermoelastic Mechanism

For decades, the mechanism was debated (direct neural stimulation vs. radiation pressure). It is now definitively established as **thermoelastic expansion**.³²

1. **Pulse Absorption:** A short, high-peak-power RF pulse (e.g., from a radar) is absorbed by the brain tissue.
2. **Thermal Spike:** The energy absorption causes a miniscule, rapid temperature rise—on the order of 10^{-6} °C. This is too small to cause cumulative heating damage but occurs so quickly that the tissue does not have time to expand gradually.
3. **Pressure Wave Generation:** The rapid thermal expansion generates a stress wave (acoustic phonon) within the cranial volume.
4. **Bone Conduction:** This acoustic wave propagates through the skull to the cochlea.
5. **Perception:** The hair cells of the cochlea detect this mechanical wave exactly as they would a sound wave conducted through bone.

6.2 Critical Parameters: Pulse Width and Peak Power

The Frey Effect is not dependent on *average* power (which determines thermal safety) but on *peak* power and *pulse width*.

- **Pulse Width Window:** Research indicates that pulse widths between **10 and 70 microseconds** are most effective at generating the acoustic frequencies audible to humans.³¹ Pulses that are too long heat the tissue too slowly to generate a sharp pressure wave; pulses that are too short may not deliver enough total energy per pulse to cross the hearing threshold.
- **Peak Power Density:** The threshold for perception can be as low as **80 mW/cm²** (peak), provided the pulse characteristics are correct, even if the average power is negligible.³¹

This phenomenon serves as a critical bridge between the electromagnetic and mechanical domains. It demonstrates that the human body can act as a transducer, converting 1 GHz electromagnetic oscillations into 5 kHz acoustic mechanical vibrations.

7. Non-Thermal Effects and Membrane Rectification

Beyond thermal heating and thermoelasticity, there exists the contentious domain of "non-thermal" effects—biological changes induced by the electric field itself.

7.1 The Rectification Hypothesis

The primary theoretical mechanism for non-thermal interaction is **membrane rectification**. Because the cell membrane is a non-linear dielectric (its resistance changes with voltage), it could theoretically demodulate an amplitude-modulated RF signal, extracting the low-frequency envelope (e.g., extracting a 16 Hz modulation from a 900 MHz carrier).³⁵

If this demodulation occurred, the resulting low-frequency electric field across the membrane could potentially gate **Voltage-Gated Calcium Channels (VGCCs)**, leading to an influx of calcium ions (Ca^{2+}). Calcium is a ubiquitous second messenger; its dysregulation could trigger oxidative stress, DNA damage, or altered neurotransmitter release.³⁶

7.2 Biophysical Constraints

However, rigorous biophysical modeling imposes strict limits on this theory. As noted in the discussion of β -dispersion, the cell membrane capacitance "shorts out" frequencies above ~10 MHz.

- **The 10 MHz Cutoff:** Above 10 MHz, the voltage drop across the membrane becomes vanishingly small because the current passes through the capacitance rather than the resistance.
- **Field Strength Requirement:** Calculations suggest that to produce a rectified voltage sufficient to open a VGCC (millivolts), the external RF field would need to be immensely strong—likely high enough to cause thermal destruction before rectification effects became relevant.³⁵

Therefore, while the mechanism is physically plausible at lower frequencies (< 10 MHz), it is unlikely to be a primary driver of biological effects at cellular communication frequencies (GHz range), where thermal noise dominates.

8. Integrated Conclusions: The Tuned System

The synthesis of these diverse datasets—from Magid's shake tables to Ohlbaum's optics and Schwan's dispersions—reveals the human body not as a passive victim of energy, but as a highly tuned spectral system.

1. **Mechanical Vulnerability (4–30 Hz):** The body is mechanically "softest" in the infrasonic range. The 4–8 Hz torso resonance and the 18 Hz ocular resonance represent the fundamental structural weaknesses of the human frame. Safety in this domain requires stiffening the coupling (tight restraints) or decoupling the source (suspension seats).
2. **Electromagnetic Vulnerability (30–100 MHz):** The body is electrically "loudest" in the VHF band. The 70 MHz whole-body resonance represents the frequency of maximal energy capture. Safety in this domain requires strict adherence to SAR limits, as the body acts as a high-gain antenna.

3. **The Acoustic Bridge:** The Frey effect demonstrates that these domains are not isolated. High-energy RF can become mechanical sound, bypassing the ear to speak directly to the cochlea.

For the modern engineer or physiologist, the "Standard Man" is a stack of coupled oscillators overlaying a lossy dielectric map. Whether designing the suspension of a lunar rover to avoid the 4–8 Hz sickness window, or tuning the antenna of a 5G handset to avoid the 70 MHz whole-body resonance, the design constraint is the same: The human body resonates. Ignoring these resonances leads to injury, fatigue, and system failure; respecting them is the prerequisite for human-machine integration.

Works cited

1. Technical Reports – Defense Technical Information Center - dtic.mil, accessed December 23, 2025, <https://discover.dtic.mil/technical-reports/>
2. Radio Frequency Radiation Dosimetry Handbook (Fifth Edition). - Google Books, accessed December 23, 2025, https://books.google.com/books/about/Radio_Frequency_Radiation_Dosimetry_Hand.html?id=cew3nQAACAAJ
3. Mechanical Resonant Frequency of the Human Eye 'In Vivo' - DTIC, accessed December 23, 2025, <https://apps.dtic.mil/sti/tr/pdf/ADA030476.pdf>
4. Human tolerance to whole body sinusoidal vibration. Short-time, one-minute and three-minute studies - PubMed, accessed December 23, 2025, <https://pubmed.ncbi.nlm.nih.gov/13765250/>
5. Short Time Human Tolerance to Sinusoidal Vibrations - Gerd H. Ziegenruecker, Edward B. Magid - Google Books, accessed December 23, 2025, https://books.google.com/books/about/Short_Time_Human_Tolerance_to_Sinusoidal.html?id=2CumE5P5FzUC
6. Visual acuity under vibration - PubMed, accessed December 23, 2025, <https://pubmed.ncbi.nlm.nih.gov/13928576/>
7. Human Issues related to Spacecraft Vibration during Ascent, accessed December 23, 2025, https://listserv-public.bcm.tmc.edu/scripts/wa.exe?A3=ind1507&L=CSMJOURNAL_CLUB&E=base64&P=8380423&B=--_006_SN1PR0601MB16294E34A04120AC33F2AAD1D8850SN1PR0601MB1629_&T=application%2Fpdf;%20name=%22Human%20Issues%20related%20to%20Spacecraft%20Vibration%20%20during%20Ascent.pdf%22&N=Human%20Issues%20related%20to%20Spacecraft%20Vibration%20%20during%20Ascent.pdf&attachment=q&XSS=3
8. Human tolerance to whole body sinusoidal vibration. Short-time, one-minute and three-minute studies. | Semantic Scholar, accessed December 23, 2025, <https://www.semanticscholar.org/paper/Human-tolerance-to-whole-body-sinusoidal-vibration.-Magid-Coermann/e8050fc178089413c9e4d45f4271e19400b16f5e>
9. SOME EFFECTS OF Y=AXIS VIBRATION ON VISUAL ACUITY - NASA Technical Reports Server, accessed December 23, 2025, <https://ntrs.nasa.gov/api/citations/19680023621/downloads/19680023621.pdf>

10. VISUAL PERFORMANCE DURING WHOLE-BODY VIBRATION - DTIC, accessed December 23, 2025, <https://apps.dtic.mil/sti/tr/pdf/AD0427254.pdf>
11. Mechanical Resonant Frequency of the Human Eye 'In Vivo' | Semantic Scholar, accessed December 23, 2025, <https://www.semanticscholar.org/paper/Mechanical-Resonant-Frequency-of-the-Human-Eye-%27In-Ohlbaum/3ef73b1375628181ac624d5199171ccaa850e800>
12. DTIC ADA030476: Mechanical Resonant Frequency of the Human Eye 'In Vivo' : Defense Technical Information Center - Internet Archive, accessed December 23, 2025, https://archive.org/details/DTIC_ADA030476
13. Retinal tear formation after whole-body vibration training exercise - PMC - NIH, accessed December 23, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC6990570/>
14. Chirp excitation for natural frequency optical coherence elastography - Optica Publishing Group, accessed December 23, 2025, <https://opg.optica.org/boe/abstract.cfm?uri=boe-15-10-5856>
15. In vivo human corneal natural frequency quantification using dynamic optical coherence elastography: repeatability and reproducibility - PubMed Central - NIH, accessed December 23, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8127410/>
16. Observation of sound-induced corneal vibrational modes by optical coherence tomography, accessed December 23, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4574659/>
17. The Radiofrequency Radiation Dosimetry Handbook: Reminiscences - PubMed, accessed December 23, 2025, <https://pubmed.ncbi.nlm.nih.gov/10334710/>
18. Radiofrequency Radiation Dosimetry Handbook. 4th Edition - DTIC, accessed December 23, 2025, <https://apps.dtic.mil/sti/tr/pdf/ADA180678.pdf>
19. Biological effects of radiofrequency electromagnetic fields due to energy absorption and heating - BfS, accessed December 23, 2025, <https://www.bfs.de/EN/topics/emf/hff/effect/hff-established/hff-established.html>
20. RF Safety FAQ | Federal Communications Commission, accessed December 23, 2025, <https://www.fcc.gov/engineering-technology/electromagnetic-compatibility-division/radio-frequency-safety/faq/rf-safety>
21. The Human Body Antenna: Characteristics and its Application - VU Research Repository, accessed December 23, 2025, https://vuir.vu.edu.au/31012/3/KIBRET%20Behailu-thesis_nosignature.pdf
22. Chapter 7 - Radiofrequency Radiation Dosimetry Handbook, accessed December 23, 2025, <https://niremf.ifac.cnr.it/docs/HANDBOOK/chp7-1.htm>
23. accessed December 23, 2025, https://www.researchgate.net/figure/Representation-of-a-b-and-g-dispersions-in-biological-tissues_fig2_329187565#:~:text=The%20dielectric%20spectrum%20of%20a.%CE%B2%20and%20%CE%B3%20dispersions%2071%20
24. Mechanisms responsible for electrical properties of tissues and cell suspensions - PubMed, accessed December 23, 2025, <https://pubmed.ncbi.nlm.nih.gov/8052170/>
25. INTERFACE PHENOMENA AND DIELECTRIC PROPERTIES OF BIOLOGICAL TISSUE,

accessed December 23, 2025,

<https://www.mn.uio.no/fysikk/english/research/projects/bioimpedance/publications/papers/encyclop.pdf>

26. Electrical Impedance Spectroscopy Study of Biological Tissues - PMC - NIH, accessed December 23, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC2597841/>
27. Dielectric Behavior of Biological Cells and Membranes (Commemoration Issue Dedicated to Professor Tetsuya HANAI On the Occasion, accessed December 23, 2025, https://repository.kulib.kyoto-u.ac.jp/bitstream/2433/77396/1/chd069_4_459.pdf
28. Radiofrequency Radiation Dosimetry Handbook - Chapter 4.1, accessed December 23, 2025, <http://niremf.ifac.cnr.it/docs/HANDBOOK/chp4-1-1.htm>
29. Estimating Human Fat and Muscle Conductivity From 100 Hz to 1 MHz Using Measurements and Modelling - PubMed, accessed December 23, 2025, <https://pubmed.ncbi.nlm.nih.gov/39825696/>
30. and frequency-dependent dielectric properties of biological tissues within the temperature and frequency ranges typ, accessed December 23, 2025, <https://www.tandfonline.com/doi/pdf/10.3109/02656736.2013.868534>
31. Microwave auditory effect - Wikipedia, accessed December 23, 2025, https://en.wikipedia.org/wiki/Microwave_auditory_effect
32. Chapter: Appendix C: Additional Comments on Directed Radio Frequency Energy, accessed December 23, 2025, <https://www.nationalacademies.org/read/25889/chapter/11>
33. Pulsed High-Power Radio Frequency Energy Can Cause Non-Thermal Harmful Effects on the BRAIN - NIH, accessed December 23, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC10914144/>
34. The Microwave Auditory Effect - IEEE Xplore, accessed December 23, 2025, <https://ieeexplore.ieee.org/iel7/7397573/9716150/09366412.pdf>
35. Rectification of RF Fields in Load Dependent Coupled Systems: Application to Non-Invasive Electroceuticals - Scirp.org., accessed December 23, 2025, https://www.scirp.org/pdf/jbise_2016022913444940.pdf
36. Electromagnetic fields act via activation of voltage-gated calcium channels to produce beneficial or adverse effects - PubMed Central, accessed December 23, 2025, <https://pmc.ncbi.nlm.nih.gov/articles/PMC3780531/>
37. Radiofrequency Fields and Calcium Movements Into and Out of Cells | Radiation Research - Allen Press, accessed December 23, 2025, <https://meridian.allenpress.com/radiation-research/article/195/1/101/448146/Radio-frequency-Fields-and-Calcium-Movements-Into>