

The Hydrodynamics and Active Mechanics of Collective Motion: From Passive Particle Settling to Biological Swarming

The mechanics of collective motion represent a profound intersection of fluid dynamics, non-equilibrium statistical mechanics, and ethology. Whether analyzing the passive descent of frozen hydrometeors through atmospheric turbulence or the highly coordinated, self-propelled maneuvers of avian murmurations and microbial suspensions, the fundamental physical constraints remain identical. The medium through which these entities move—whether air or water—acts simultaneously as a viscous resistor and a dynamic communicative substrate. Hydrodynamic forces dictate not only the trajectory of a single body but also the emergent, macroscopic architecture of massive biological aggregations. When self-propelled entities share identical morphological and kinematic properties, their mutual interactions and fluidic responses coalesce. A massive swarm of independent agents ultimately behaves, dynamically and structurally, as a single, continuous meta-organism. This report provides an exhaustive, mathematically rigorous analysis of the mechanics governing both passive particle settling and the active, equidistant swarming of biological organisms, functioning as a comprehensive foundation for further research into fluid-mediated collective behavior.

The Passive Paradigm: Kinematics and Fluid Mechanics of Settling Particles

To comprehend the self-propelled mechanics of biological swarms, one must first establish the foundational fluid dynamics governing passive particles. The precipitation of frozen hydrometeors, such as snowflakes and plate crystals, serves as a canonical model for understanding how morphology, inertia, and ambient turbulence dictate collective spatial distribution.

The Maxey-Riley Equation and Basset History Force

The trajectory of a small, rigid spherical or near-spherical particle moving through a non-uniform fluid field is classically described by the Maxey-Riley equation. Derived from the Navier-Stokes equations, this integro-differential framework accounts for a complex balance of hydrodynamic forces, including the undisturbed fluid force, buoyancy, Stokes drag, added mass, and the critically important Basset history force¹. For a particle of mass and specific radius, moving with velocity through a fluid with kinematic viscosity, the dimensionless acceleration relies heavily on the Stokes number, which represents the ratio of the particle's momentum response time to the characteristic timescale of the fluid flow³. Historically, the Basset history force has been omitted from many computational models due to

its exceptional mathematical complexity and immense memory cost during numerical simulation¹. This force models the temporal delay in the development of the viscous boundary layer around an accelerating particle, acting as a physical memory of the particle's past kinematic states³. The history force is expressed mathematically as a convolution integral incorporating a singular kernel that decays over time. The inclusion of this history term fundamentally alters the asymptotic settling behavior of particles in a fluid. Rather than an exponential relaxation to a terminal velocity, the Basset history force slows the decay to a power-law relaxation, scaling inversely with the square root of time, proving that the fluid medium retains a persistent memory of the particle's passage and thereby alters the trajectory of any subsequent particles following in its wake¹.

Morphological Porosity, Turbulence, and Tumbling Dynamics

For highly anisotropic, non-spherical particles such as snowflakes, which are mathematically modeled as thin disks with specific diameter-to-thickness aspect ratios, morphological porosity strictly controls fluid stability⁶. Advanced fluid analysis utilizing Proper Orthogonal Decomposition and Dynamic Mode Decomposition reveals that high-porosity dendritic snowflakes permit fluid to pass through their intricate lattices, which stabilizes small-scale wake structures and promotes steady, horizontal descent⁷. In sharp contrast, low-porosity planar crystals generate massive, unstable hairpin vortices that induce highly chaotic, three-dimensional trajectories⁸.

As these anisotropic particles descend through the fluid, they transition through four primary kinematic regimes: stable, zigzag, transitional, and tumbling or spiraling⁸. Ambient air turbulence fundamentally alters these regimes. Experimental data indicates that air turbulence reduces the terminal settling velocity of snowflakes by up to thirty-five percent compared to their velocity in quiescent air⁶. This reduction is driven by cross-flow-induced nonlinear drag. As turbulence intensity increases within the medium, the flat-falling behavior is progressively eliminated, and continuous tumbling becomes the prevalent kinematic state¹⁰. Within turbulent boundary layers, the behavior of the swarm of falling particles is governed by a critical dimensionless parameter that characterizes the ratio of gravitational settling to turbulent diffusion. When this parameter drops below a specific threshold, horizontal relative motion completely dominates vertical descent¹¹. The turbulence induces phenomena known as preferential sweeping, where particles cluster rapidly in the downward-moving regions of turbulent eddies, or loitering, where particles become trapped in upward-moving regions¹⁰. These fluidic interactions fundamentally organize the passive swarm, clustering particles in a manner analogous to biological aggregation.

Microscopic Active Matter: Microbial Swarming and Hydrodynamic Coupling

Transitioning from passive settling to active matter, microscopic biological entities such as bacteria and microalgae operate entirely within the Stokes flow regime, characterized by exceedingly low Reynolds numbers¹². In this microscopic domain, viscous forces completely

overpower inertial forces. Therefore, sustained motion requires the continuous expenditure of internal chemical energy to break time-reversal symmetry, creating localized active stresses within the fluid medium¹⁴.

Pushers, Pullers, and the Stresslet Tensor

A single microbe swimming through a viscous fluid is fundamentally force-free and torque-free; the propulsive thrust it generates is instantaneously and perfectly balanced by the viscous drag exerted across its cell body¹⁶. Consequently, the leading-order disturbance flow generated by a swimming microbe is not a monopole, but a force dipole. This dipolar flow field is mathematically described by a symmetric, traceless second-rank tensor known as the stresslet¹².

The orientation of the biological propulsion mechanism dictates the sign of the stresslet strength. Bacteria such as *Escherichia coli* propel themselves by rotating helical flagella located at the rear of their bodies. Because the thrust is generated posterior to the center of drag, they push fluid outward along their primary longitudinal axis and draw fluid inward laterally. These organisms are classified hydrodynamically as pushers, possessing a negative stresslet strength¹⁶. Conversely, microalgae like *Chlamydomonas reinhardtii* utilize a coordinated breaststroke motion using anterior flagella. They draw fluid inward along their longitudinal axis and expel it laterally, operating as pullers with a positive stresslet strength¹⁶.

When massive suspensions of pushers and pullers coexist, their respective fluidic velocity fields couple mechanically¹⁹. The interaction of opposite-type active microswimmers generates spontaneous, non-equilibrium fluctuations in the fluid. The dynamic viscosity of the medium is actively altered by the stresslets. Pusher suspensions tend to decrease the effective viscosity of the fluid, a phenomenon termed active superfluidity, which leads to large-scale chaotic fluid flows known as active turbulence²⁰. In these dense states, the conservation of particle number and orientational probability is strictly governed by the Smoluchowski equation, which models the spatial and angular fluxes of the swarm as the active fluid attempts to minimize orientational entropy¹⁷.

Chemotaxis, Temporal Gradients, and the Keller-Segel Equations

The self-propelled motion of microbes is rarely a purely isotropic random walk; it is heavily biased by chemical gradients in the environment, a biological process known as chemotaxis²³. When modeling the macroscopic swarming behavior of a bacterial population, the discrete run-and-tumble kinematics of individual cells are coarse-grained into continuum partial differential equations, most notably the Patlak-Keller-Segel model²⁵.

In this mathematical framework, the bacterial population density and the chemoattractant concentration are deeply coupled²³. The population undergoes both unbiased Fickian diffusion and highly directed advection up the chemical gradient. Modern variations of the Keller-Segel model incorporate logistic growth terms to prevent non-physical infinite density blow-ups, and they directly couple the chemotactic system to the Navier-Stokes equations to account for the macroscopic fluid flow carrying the cells²⁶. Furthermore, the bacteria exert a reciprocal body force back onto the fluid. For example, if the cells are denser than the surrounding medium,

gravitational forcing is applied to the fluid momentum equations²⁷. This reciprocal mechanical interaction causes dense bacterial swarms to trigger localized fluid convection, accelerating the mixing of nutrients and structuring the swarm into stable, collective bioconvection plumes that behave as a single organism²³.

Recent advancements in continuum modeling also account for temporal gradients²⁴. Bacteria do not merely sense spatial gradients instantaneously; they possess a chemotactic memory. By evaluating the derivative of bacterial position along a gradient over a delay time, researchers use path integrals to map how temporal changes in chemical concentration modulate the base tumble rate of the bacteria, optimizing the swarm's navigation through dynamically shifting chemical landscapes²⁴.

Phase Synchronization of Flagella: The Kuramoto Model

At an even finer biological scale, the swarming behavior of a single microbe requires the absolute coordination of its own appendages. The synchronization of multiple flagella beating within a viscous fluid is an emergent mechanical phenomenon driven entirely by hydrodynamic coupling¹³. Because the flagella move through a fluid with negligible inertia, the velocity flow field generated by one flagellum exerts an immediate drag force on its neighbor, shifting its orbital phase¹³.

This phase-locking behavior is mathematically encapsulated by the Kuramoto model of coupled phase oscillators³⁰. For a specified number of interacting flagella, each possessing an intrinsic natural frequency and a dynamic phase, the rate of phase change is governed by the addition of the natural frequency and a sinusoidal coupling term that scales with the hydrodynamic coupling strength³⁰. To measure the degree to which the flagella beat in unison as a single mechanical entity, physicists utilize the Kuramoto order parameter, defined as the magnitude of the mean phase vector of the system³⁰.

When the fluidic coupling strength is below a critical threshold, the flagella beat incoherently, and the order parameter rests near zero³⁰. As the hydrodynamic interactions overcome the natural frequency dispersion, the system undergoes a continuous phase transition. The flagella lock into a synchronized rhythm, pushing the order parameter toward unity, enabling the microbe to achieve rapid directional swimming rather than spinning erratically³⁰.

Time-frequency computations based on continuous wavelet transforms demonstrate that this phase-locking mechanism is remarkably robust to external perturbations, allowing organisms to quickly resynchronize their beating patterns after experiencing severe environmental stressors, such as intense light stimuli during photoshock¹³.

Fluid-Mediated Cohesion at the Macroscale: Aquatic Schooling

At larger biological scales, where the Reynolds number ranges from ten thousand to one million, inertial forces strictly dominate over viscous forces³². Fish schools provide an exemplary model of macroscopic organisms exploiting fluid mechanics to maintain a cohesive, equidistant swarm structure while drastically minimizing the energetic cost of collective locomotion³⁴.

Kármán and Reverse Kármán Vortex Streets

When a fluid flows past a stationary bluff body, it generates a regular, alternating pattern of swirling vortices known as a Kármán vortex street³². The arrangement and shedding frequency of this wake are parameterized by the non-dimensional Strouhal number, which relates the shedding frequency and characteristic length to the overall flow velocity³².

Conversely, when an aquatic organism undulates its flexible body and tail to generate forward thrust, it sheds a highly organized wake of alternating vortices with a rotational direction perfectly opposite to that of a classical drag wake³². This propulsive flow is classified as a reverse Kármán vortex street³⁷. Fish swimming continuously adjust their tail-beat amplitude and undulation frequency to maintain a Strouhal number tightly bounded between 0.2 and 0.4, a specific kinematic window that ensures maximum propulsive efficiency and optimal thrust generation³².

Rheotaxis and the Energetics of the Kármán Gait

Fishes utilize a highly specialized sensory organ known as the lateral line system to detect minute pressure gradients and velocity changes in the fluid surrounding their bodies³⁹. This acute hydrodynamic perception allows them to engage in rheotaxis, which is the behavioral tendency to align their bodies directly into oncoming currents⁴¹.

When fish assemble into a school, trailing individuals strategically position themselves within the reverse Kármán vortex streets generated by the leading individuals³⁵. By adopting a staggered or diamond formation, a trailing fish encounters an oscillating flow field. Instead of expending energy to fight the surrounding fluid, the trailing fish actively synchronizes its own tail beats to match the phase of the oncoming vortices³⁵. This phenomenon, termed the Kármán gait, allows the follower to effectively slalom through the alternating vortices, utilizing the low-pressure vortex cores to be pulled forward hydrodynamically³⁹.

Mathematical modeling and computational fluid dynamics employing the immersed boundary method demonstrate that a flexible body positioned appropriately in a reverse Kármán street can achieve forward propulsion entirely passively, extracting kinetic energy directly from the leader's wake³⁶. The integration of advanced neural network models, such as Kolmogorov-Arnold Networks, into autonomous simulated swimmers has proven that perception and decision-making in complex fluid environments heavily prioritize finding these optimal phase-matching positions⁴¹. Through this fluidic coupling, the entire school achieves sustained high swimming speeds with significantly reduced average metabolic and locomotor costs. The trailing fish experience massive drag reduction and enhanced thrust, allowing the swarm to act synergistically as a unified hydrodynamic entity moving equidistantly through the water column³⁴.

Aerial Swarming Mechanics: Avian Formations and Murmurations

While aquatic schooling relies heavily on discrete vortex harvesting, the aerial swarming of

birds encompasses both rigid aerodynamic drafting, seen in V-formations, and highly fluid, isotropic collective maneuvers, observed in murmurations. Despite the differing functional geometries of these flight patterns, both fundamentally rely on profound spatial synchronization to unify the flock into a single mechanical structure.

Aerodynamic Drafting in V-Formations

Migratory birds, such as geese, pelicans, and cranes, arrange themselves in precise V-shaped formations to optimize group energy expenditure during grueling long-distance flights⁴³. The mechanics of this formation are deeply rooted in fixed-wing aerodynamics and lifting-line theory⁴⁵.

As the leading bird flaps its wings, the intense pressure differential between the lower and upper surfaces of the wing generates a powerful pair of trailing wingtip vortices⁴⁵. Inside the wingspan, these vortices induce a downward flow of air, known as downwash, which increases the induced drag on the bird. However, immediately outside the wingspan, the vortices induce a strong upward flow of air, known as upwash⁴³.

Trailing birds position their own wingtips precisely within the upwash region of the bird immediately ahead of them⁴³. The upwash tilts the local aerodynamic force vector forward, effectively providing free lift and substantially reducing the induced drag experienced by the follower. High-fidelity computational fluid dynamics simulations reveal that optimal positioning within this upwash zone can increase a trailing bird's aerodynamic efficiency, measured as the lift-to-drag ratio, by up to thirty-two percent⁴³.

Because the wingtip vortices undulate vertically with the flapping motion of the leader, the trailing birds must enact precise spatial and temporal wingbeat synchronization to remain within the narrow band of supportive air⁴³. The formation is dictated by a specific longitudinal wavelength separation, perfectly matching the distance the leader travels in one full wingbeat cycle⁴³. By maintaining an optimal separation distance of exactly one wavelength, the entire flock distributes the aerodynamic load, rotating leadership dynamically to prevent fatigue, and seamlessly migrating as a macroscopic, interconnected wing⁴⁴.

Murmurations, Topological Distance, and Critical Systems

In extreme contrast to the rigid, metric structures of V-formations, the murmurations of starlings exhibit incredibly dense, fluid, and rapidly shifting geometries⁴⁸. These massive swarms can comprise hundreds of thousands of individuals executing simultaneous, high-speed maneuvers to evade predators⁴⁸. The mechanics governing these evasive maneuvers represent one of the most sophisticated examples of collective animal behavior, operating on mathematical principles akin to statistical mechanics and thermodynamic phase transitions⁴⁸.

Traditional metric models of animal swarming assume that individuals interact with all neighbors located within a specific radial metric distance⁵⁰. However, high-speed stereographic three-dimensional tracking of starling flocks and jackdaw murmurations has proven the metric hypothesis incorrect⁵⁰. Instead, these avian swarms operate on topological distance. Regardless of whether the flock is tightly compressed during a predatory attack or

widely dispersed in calm flight, each bird consistently monitors and coordinates its movements with exactly its six or seven nearest neighbors⁴⁸.

Mapping the spatial position of these nearest neighbors reveals a striking anisotropy; starlings maintain closer topological distances to birds beside them than to those directly in front or behind, establishing a pancake-like geometric structure that prevents collisions during intricate maneuvers⁴⁸. Mathematical models evaluating the cost-benefit ratio of biological computation demonstrate that tracking exactly seven neighbors represents the optimal balance between maintaining absolute flock cohesion and minimizing cognitive load and energetic expenditure⁵³. By interacting strictly based on topology, the flock ensures that interaction strength remains perfectly constant even during extreme density fluctuations, preventing the swarm from splitting or individuals from becoming dangerously isolated⁵¹.

Furthermore, murmurations exhibit a remarkable physical phenomenon known as scale-free behavioral correlation⁴⁹. In standard physical systems, correlation length determines how far a local perturbation propagates through a medium before dissipating into noise⁴⁹. In most biological systems, such as bacterial swarms, this correlation length is finite, meaning distant parts of the swarm operate independently. However, in starling flocks, researchers discovered that the correlation length is not constant; rather, it scales dynamically with the linear size of the entire flock⁴⁹.

Because the behavioral correlation is scale-free, the flock behaves mathematically as a critical system—a system poised exactly on the brink of a phase transition⁴⁸. In this highly balanced critical state, a sudden change in the directional velocity of a single bird on the edge of the flock instantly affects, and is simultaneously affected by, every other bird in the network, regardless of the flock's total size⁴⁹. The information regarding directional shifts propagates through the entire flock with virtually no signal degradation, yielding a nearly noiseless macroscopic response⁴⁸. The murmuration cannot be mathematically or physically divided into independent sub-parts; it is intrinsically bound by its scale-free correlations, rendering the flock a single, highly sensitive organism reacting instantaneously as a unified whole to environmental stimuli⁴⁹.

The Mathematics of Collective Motion and Equidistant Spacing

To understand how individual agents—whether bacteria, fish, or birds—coalesce into a meta-organism that moves with uniform density and coherent motion, scientists rely on both discrete agent-based models and continuum hydrodynamic equations.

The Vicsek Model and the Toner-Tu Hydrodynamic Equations

The mathematical foundation of modern swarming mechanics is the Vicsek model, an elegant, minimal agent-based model designed to describe the onset of collective motion⁵⁸. The model assumes that point-like active particles travel at a constant magnitude of speed and, at each discrete time step, align their orientation with the average direction of all neighbors within a specified radius, subject to a random scalar noise variable⁵⁸. By tuning the noise parameter or

the particle density, the Vicsek model demonstrates a sudden, first-order kinetic phase transition. The system shifts from a disordered, gaseous state of random motion into a highly ordered, liquid-like polar phase where all particles migrate cohesively in the exact same direction⁵⁸.

While the Vicsek model flawlessly handles discrete entities, the sheer number of interacting bodies in massive biological swarms necessitates a continuum approach. By coarse-graining the discrete Vicsek rules utilizing the Chapman-Enskog method from kinetic theory, researchers derive the Toner-Tu equations. These equations form a set of stochastic partial differential equations that describe the swarm as an active, compressible fluid¹⁴. For an incompressible active fluid, the velocity field and pressure gradient are governed by a modified Navier-Stokes framework⁶¹.

Unlike passive Newtonian fluids, active swarming fluids lack Galilean invariance due to continuous momentum exchange with the substrate or the ambient fluid medium⁶¹. The defining feature of the Toner-Tu continuum equation is the Rayleigh-Helmholtz friction term, derived directly from a quartic Landau velocity potential⁶⁰. When the damping coefficients cross a critical threshold, the fluid is driven actively away from a resting state and settles into a non-vanishing, uniform collective velocity. This spontaneous symmetry breaking mathematically proves the emergence of long-range polar order in the swarm⁶¹. Extensive dynamic renormalization group analyses of the Toner-Tu equations reveal unique scaling exponents that perfectly characterize the temporal and spatial correlations of fluctuations within the flock, proving that the flocking phase avoids the restrictions of the Mermin-Wagner-Hohenberg theorem typically applied to thermal systems⁶³.

Interaction Potentials and Equidistant Spacing

The defining structural feature of any mature swarm is its ability to maintain a relatively uniform internal density—where individuals remain roughly equidistant from one another—without dispersing into the environment or collapsing catastrophically into a single, infinitesimally small point⁶⁵. This highly regulated equilibrium spacing is governed by pairwise interaction potentials defined by three fundamental biological rules: short-range repulsion to ensure collision avoidance, intermediate-range alignment for velocity matching, and long-range attraction to ensure flock cohesion⁵⁴.

Mathematically, the interactions between any two agents within the swarm are modeled using continuous, radially symmetric potential functions, most notably the Morse potential and the Quasi-Morse potential⁶⁸. The Morse potential, a function of the metric distance between individuals, is expressed analytically as the sum of exponentially decaying attractive and repulsive forces. The equation relies on specific amplitude coefficients for attraction and repulsion, alongside parameters that dictate their respective length scales⁷⁰.

For a biological swarm to achieve a stable, equidistant distribution of agents, the repulsive forces must operate over a significantly shorter spatial range than the attractive forces to maintain overall group cohesion, yet the magnitude of the repulsion must overwhelmingly dominate at close ranges to prevent physical collision⁶⁶. To rigorously guarantee that the swarm does not collapse as the total number of individuals approaches infinity, the interaction

potential must strictly satisfy the thermodynamic condition of H-stability⁶⁷. A potential is deemed H-stable if there exists a non-negative constant such that the total potential energy of the swarm remains bounded from below for any possible spatial configuration of particles⁷¹. The absolute mathematical constraint required to avoid catastrophic collapse dictates that the product of the repulsive amplitude and its characteristic spatial range raised to the power of the space dimension plus one must strictly exceed the equivalent product for the attractive forces⁶⁶. When this specific inequality is met, the active particles settle effortlessly into a stable energy minimum. At the macroscopic scale, this microscopic H-stability manifests as a perfectly constant individual distance, or a comfortable distance, between all members of the swarm⁶⁶. Consequently, the density within the swarm becomes strikingly uniform, and the physical physical boundaries of the swarm scale linearly with the number of individuals, matching empirical observations of massive biological aggregations⁶⁶. This theoretical framework seamlessly bridges the vast gap between individual, agent-based kinematics and the emergence of macroscopic continuum behavior. The precise evolutionary tuning of Morse potentials, combined with the scale-free correlations observed in topological networks, explains precisely how an aggregate of entirely independent biological agents forms a robust, highly structured lattice. Driven continuously by non-equilibrium active forces perfectly described by the Toner-Tu equations, the swarm navigates its fluid environment not as a loose, chaotic collection of colliding bodies, but as a seamlessly unified, equidistant, and highly responsive singular entity.

Works cited

1. Accurate solution method for the Maxey–Riley equation, and the effects of Basset history,
https://www.researchgate.net/publication/332364075_Accurate_solution_method_for_the_Maxey-Riley_equation_and_the_effects_of_Basset_history
2. Maxey-Riley-global-existence.pdf - George Haller,
<http://www.georgehaller.com/reprints/Maxey-Riley-global-existence.pdf>
3. On the role of the history force for inertial particles in turbulence - arXiv,
<https://arxiv.org/pdf/1501.04770>
4. The Basset–Boussinesq history force: its neglect, validity, and recent numerical developments - Frontiers,
<https://www.frontiersin.org/journals/physics/articles/10.3389/fphy.2023.1167338/full>
5. An efficient, second order method for the approximation of the Basset history force, https://ris.utwente.nl/ws/files/6725447/2011_JCompP_vHtTC.pdf
6. (PDF) Thin disks falling in air - ResearchGate,
https://www.researchgate.net/publication/370318503_Thin_disks_falling_in_air
7. Modal analysis reveals imprint of snowflake shape on wake flow structures - DORA 4RI,
[https://www.dora.lib4ri.ch/wsl/dload/wsl:41116/PDF/Tagliavini-2025-Modal_analysis_reveals_imprint_of-\(published_version\).pdf](https://www.dora.lib4ri.ch/wsl/dload/wsl:41116/PDF/Tagliavini-2025-Modal_analysis_reveals_imprint_of-(published_version).pdf)
8. (PDF) Three dimensional wakes of freely falling planar polygons - ResearchGate,

- https://www.researchgate.net/publication/333857209_Three_dimensional_wakes_of_freely_falling_planar_polygons
9. Visualisation of Euler angle convention used; an example of the... - ResearchGate, https://www.researchgate.net/figure/Visualisation-of-Euler-angle-convention-used-an-example-of-the-orientation-of-an_fig2_346247219
 10. (PDF) Turbulence effect on disk settling dynamics - ResearchGate, https://www.researchgate.net/publication/383472722_Turbulence_effect_on_disk_settling_dynamics
 11. The p.d.f.s of the snow particle vertical velocity (w_s) as measured... - ResearchGate, https://www.researchgate.net/figure/The-pdfs-of-the-snow-particle-vertical-velocity-w-s-as-measured-by-PIV-for-January_fig3_349404139
 12. Rheology of Active Fluids - David Saintillan, http://stokeslet.ucsd.edu/publications_files/S2018.pdf
 13. Tracking phase synchronization between flagella in the time-frequency domain resolves photophobic response - arXiv, <https://arxiv.org/html/2510.21393v2>
 14. Active Matter: from motility to self-organization Lectures presented at the Boulder School on Self-Organizing Matter July 2024, https://boulderschool.yale.edu/sites/default/files/files/BSS2024/Boulder_2024_Lecture_Notes_Marchetti1.pdf
 15. Energetic Variational Modeling of Active Nematics: Coupling the Toner–Tu Model with ATP Hydrolysis - arXiv, <https://arxiv.org/html/2506.23043v1>
 16. KINETIC MODELS FOR BIOLOGICALLY ACTIVE SUSPENSIONS 1. Introduction. Microorganisms are present in every part of the biosphere, r - David Saintillan, http://stokeslet.ucsd.edu/publications_files/S2012.pdf
 17. Theory of active suspensions - math.nyu.edu., <https://math.nyu.edu/~shelley/papers/SS2015.pdf>
 18. Swimmer types of optimum surface-driven active particles | Journal of Fluid Mechanics, <https://www.cambridge.org/core/journals/journal-of-fluid-mechanics/article/swimmer-types-of-optimum-surfacedriven-active-particles/AB1307D566135517F624C61B8622CA41>
 19. Mechanical Coupling of Puller and Pusher Active Microswimmers Influences Motility - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7304893/>
 20. Mechanical Coupling of Puller and Pusher Active Microswimmers Influences Motility | Langmuir - ACS Publications, <https://pubs.acs.org/doi/10.1021/acs.langmuir.9b03665>
 21. An effective and efficient model of the near-field hydrodynamic interactions for active suspensions of bacteria | PNAS, <https://www.pnas.org/doi/10.1073/pnas.2100145118>
 22. Active Stokesian dynamics - Soft Matter Group, <https://mech-aml.sites.olt.ubc.ca/files/2022/11/elfring22.pdf>
 23. On the simultaneous recovery of environmental factors in the 3D chemotaxis-Navier-Stokes models - American Institute of Mathematical Sciences, <https://www.aims sciences.org/article/doi/10.3934/cac.2024003>

24. Drift velocity of bacterial chemotaxis in dynamic chemical environments - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12423644/>
25. The Keller--Segel Model with Logistic Sensitivity Function and Small Diffusivity | SIAM Journal on Applied Mathematics, <https://epubs.siam.org/doi/10.1137/040612841>
26. Multi-Species Keller–Segel Systems: Analysis, Pattern Formation, and Emerging Mathematical Structures - arXiv, <https://arxiv.org/html/2603.04931v1>
27. Global Solutions to the Coupled Chemotaxis-Fluid Equations - ResearchGate, https://www.researchgate.net/publication/243046997_Global_Solutions_to_the_Coupled_Chemotaxis-Fluid_Equations
28. A diffuse-domain-based numerical method for a chemotaxis-fluid model - Alina Chertock, <https://chertock.wordpress.ncsu.edu/files/2023/03/WCKZ22.pdf>
29. Tracking phase synchronization between flagella in the time-frequency domain resolves photophobic response - arXiv, <https://arxiv.org/html/2510.21393v1>
30. Kuramoto model - Wikipedia, https://en.wikipedia.org/wiki/Kuramoto_model
31. Synchronization and Critical Coupling in the Classical Kuramoto Model, <https://www.journalofyoungphysicists.org/post/synchronization-and-critical-coupling-in-the-classical-kuramoto-model>
32. The role of hydrodynamics in collective motions of fish schools and bioinspired underwater robots | Journal of The Royal Society Interface, <https://royalsocietypublishing.org/rsif/article/20/207/20230357/90505/The-role-of-hydrodynamics-in-collective-motions-of>
33. Numerical Study on the Hydrodynamics of Fish Swimming with Different Morphologies in Oblique Flow - MDPI, <https://www.mdpi.com/2077-1312/12/12/2302>
34. Biomechanics and energetics of fish schooling behavior - ScienceDirect - DOI, <https://doi.org/10.1016/bs.fp.2025.06.001>
35. Hydrodynamic Efficiency and Wake Interactions in Fish School Swimming - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC13113052/>
36. In-line swimming dynamics revealed by fish interacting with a robotic mechanism - eLife, <https://elifesciences.org/articles/81392>
37. Kinematic Characteristics and Drag Reduction of a Flexible Filament behind a Flapping Foil, <https://www.scirp.org/journal/paperinformation?paperid=150809>
38. Full article: Evolvement rule and hydrodynamic effect of fluid field around fish-like model from starting to cruising - Taylor & Francis, <https://www.tandfonline.com/doi/full/10.1080/19942060.2020.1734095>
39. The role of the lateral line and vision on body kinematics and hydrodynamic preference of rainbow trout in turbulent flow - Company of Biologists journals, <https://journals.biologists.com/jeb/article/209/20/4077/16399/The-role-of-the-lateral-line-and-vision-on-body>
40. A model of a lateral line of fish for vortex sensing - Kamran Mohseni - University of Florida, <https://enstrophy.mae.ufl.edu/publications/MyPapers/BioinspirBiomim2012LateralLineModel.pdf>
41. The Karman gait: novel body kinematics of rainbow trout swimming in a vortex

street,

https://www.researchgate.net/publication/277436689_The_Karman_gait_novel_body_kinematics_of_rainbow_trout_swimming_in_a_vortex_street

42. A numerical study of fish adaption behaviors in complex environments with a deep reinforcement learning and immersed boundary–lattice Boltzmann method - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7814145/>
43. Aerodynamic mechanisms behind energy efficiency in migratory bird formations - AIP Publishing, https://pubs.aip.org/aip/pof/article-pdf/doi/10.1063/5.0252553/20404860/0252021_5.0252553.pdf
44. Aerodynamics of Flight Formations in Birds: A Quest for Energy Efficiency - ResearchGate, https://www.researchgate.net/publication/390135521_Aerodynamics_of_Flight_Formations_in_Birds_A_Quest_for_Energy_Efficiency
45. Effects of Leader's Position and Shape on Aerodynamic Performances of V Flight Formation - arXiv, <https://arxiv.org/pdf/0804.3879>
46. V-formation flight of birds - Robots in the sun -, <http://robotsinthesun.org/v-formation-flight-of-birds/>
47. Aerodynamics of Flight Formations in Birds: A Quest for Energy Efficiency - MDPI, <https://www.mdpi.com/2673-6004/6/2/15>
48. The Science Of Starling Murmurations - Bird Spot, <https://www.birdspot.co.uk/bird-behaviour/the-science-of-starling-murmurations>
49. Starling Murmuration - One Shared Myth, <https://www.onesharedmyth.com/murmuration.html>
50. Topological vs. Metric distance - CNR-ISC, <https://www.isc.cnr.it/research/topics/physical-biology/biological-systems/topological-vs-metric-distance/>
51. Interaction ruling animal collective behavior depends on topological rather than metric distance: Evidence from a field study - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC2234121/>
52. Why Birds on the Edge Stay There: Study Sheds Light on Murmuration Mysteries, <https://www.rothamsted.ac.uk/news/why-birds-edge-stay-there-study-sheds-light-murmuration-mysteries>
53. Birds of a feather ... track seven neighbors to flock together - Princeton University, <https://www.princeton.edu/news/2013/02/07/birds-feather-track-seven-neighbors-flock-together>
54. Diffusion and Topological Neighbours in Flocks of Starlings: Relating a Model to Empirical Data - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4436282/>
55. How Do Starling Flocks Create Those Mesmerizing Murmurations? - All About Birds, <https://www.allaboutbirds.org/news/how-do-starling-flocks-create-those-mesmerizing-murmurations/>
56. Scale-free correlations in starling flocks - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC2900681/>
57. (PDF) Scale-free correlations in starling flocks - ResearchGate,

- https://www.researchgate.net/publication/44673012_Scale-free_correlations_in_starling_flocks
58. Vicsek model - Wikipedia, https://en.wikipedia.org/wiki/Vicsek_model
 59. PHYS 563 term Paper The Flocking Transition : A Review of The Vicsek Model, https://guava.physics.ucsd.edu/~nigel/Courses/Web%20page%20563/Essays_2017/PDF/Chatterjee.pdf
 60. A microscopically reversible kinetic theory of flocking - arXiv, <https://arxiv.org/pdf/2508.09274>
 61. Well-Posedness and Asymptotic Stability of Solutions for the Incompressible Toner-Tu Model | SIAM Journal on Mathematical Analysis, <https://epubs.siam.org/doi/10.1137/23M1599756>
 62. An analytical and computational study of the incompressible Toner-Tu Equations - Department of Mathematics - Imperial College London, <https://www.ma.ic.ac.uk/~jdq/ITT-arx1.pdf>
 63. A new universality class describes Vicsek's flocking phase in physical dimensions - arXiv, <https://arxiv.org/html/2402.01316v1>
 64. Quantum Vicsek Model for Active Matter - arXiv, <https://arxiv.org/html/2407.09860v1>
 65. Swarm behaviour - Wikipedia, https://en.wikipedia.org/wiki/Swarm_behaviour
 66. Mutual interactions, potentials, and individual distance in a social aggregation, https://www.researchgate.net/publication/9067605_Mutual_interactions_potentials_and_individual_distance_in_a_social_aggregation
 67. An agent-based approach for modelling collective dynamics in animal groups distinguishing individual speed and orientation - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7423381/>
 68. A new interaction potential for swarming models - UC Davis Mathematics, https://www.math.ucdavis.edu/~saito/data/PSO-ACO/carrillo-et-al_interaction-potential-swarming.pdf
 69. Swarm dynamics and equilibria for a nonlocal aggregation model - Simon Fraser University, https://www.sfu.ca/~van/papers/FeHuKo_2011.pdf
 70. Numerical simulation of nonlinear continuity equations by evolving diffeomorphisms - RICAM, <https://www.ricam.oeaw.ac.at/files/reports/16/rep16-19.pdf>
 71. Self-propelled particles with soft-core interactions: patterns, stability and collapse - UCLA Mathematics, <https://www3.math.ucla.edu/camreport/cam05-53.pdf>