

Sequential Regressions for Efficient Continuous-Time Causal Inference

Johan Sebastian Ohlendorff

University of Copenhagen

- PhD student in Biostatistics at the Section of Biostatistics, University of Copenhagen.

- PhD student in Biostatistics at the Section of Biostatistics, University of Copenhagen.
- Supervisors: Thomas Alexander Gerds and Anders Munch.

- PhD student in Biostatistics at the Section of Biostatistics, University of Copenhagen.
- Supervisors: Thomas Alexander Gerds and Anders Munch.
- Work on continuous-time longitudinal causal inference using
 - Targeted learning (e.g., TMLE (van der Laan & Rubin (2006)) or double/debiased machine learning (Chernozhukov et al. (2018))).

- PhD student in Biostatistics at the Section of Biostatistics, University of Copenhagen.
- Supervisors: Thomas Alexander Gerds and Anders Munch.
- Work on continuous-time longitudinal causal inference using
 - Targeted learning (e.g., TMLE (van der Laan & Rubin (2006)) or double/debiased machine learning (Chernozhukov et al. (2018))).
- Working on a continuous-time scale is motivated by:
 - Real-world data often recorded in continuous time (e.g., electronic health records).
 - Avoiding bias due to discretization (Adams et al. (2020); Ferreira Guerra et al. (2020); Kant & Krijthe (2025); Ryalen et al. (2019); Sofrygin et al. (2019); Sun & Crawford (2023)).

- PhD student in Biostatistics at the Section of Biostatistics, University of Copenhagen.
- Supervisors: Thomas Alexander Gerds and Anders Munch.
- Work on continuous-time longitudinal causal inference using
 - Targeted learning (e.g., TMLE (van der Laan & Rubin (2006)) or double/debiased machine learning (Chernozhukov et al. (2018))).
- Working on a continuous-time scale is motivated by:
 - Real-world data often recorded in continuous time (e.g., electronic health records).
 - Avoiding bias due to discretization (Adams et al. (2020); Ferreira Guerra et al. (2020); Kant & Krijthe (2025); Ryalen et al. (2019); Sofrygin et al. (2019); Sun & Crawford (2023)).
- We follow the setting in Rytgaard et al. (2022) and are interested in the mean interventional absolute risk under a specified treatment regime in continuous time.

- PhD student in Biostatistics at the Section of Biostatistics, University of Copenhagen.
- Supervisors: Thomas Alexander Gerds and Anders Munch.
- Work on continuous-time longitudinal causal inference using
 - Targeted learning (e.g., TMLE (van der Laan & Rubin (2006)) or double/debiased machine learning (Chernozhukov et al. (2018))).
- Working on a continuous-time scale is motivated by:
 - Real-world data often recorded in continuous time (e.g., electronic health records).
 - Avoiding bias due to discretization (Adams et al. (2020); Ferreira Guerra et al. (2020); Kant & Krijthe (2025); Ryalen et al. (2019); Sofrygin et al. (2019); Sun & Crawford (2023)).
- We follow the setting in Rytgaard et al. (2022) and are interested in the mean interventional absolute risk under a specified treatment regime in continuous time.
- **Problem:** Rytgaard et al. (2022) do not provide a feasibly implementable procedure for estimation.

- **Study period:** $[0, \tau_{\text{end}}]$.

- **Study period:** $[0, \tau_{\text{end}}]$.
- **Càdlàg, jump processes**
 - Treatment: $A(t) \in \{0, 1\}$
 - Covariates: $L(t) \in \mathbb{R}^d$

- **Study period:** $[0, \tau_{\text{end}}]$.
- **Càdlàg, jump processes**
 - Treatment: $A(t) \in \{0, 1\}$
 - Covariates: $L(t) \in \mathbb{R}^d$
- **Counting processes**
 - $N^x; x \in \{a, \ell, y\}$ (treatment, covariate, death)
 - Observed until censoring time C
 - Terminal event time: T_e

- **Study period:** $[0, \tau_{\text{end}}]$.
- **Càdlàg, jump processes**
 - Treatment: $A(t) \in \{0, 1\}$
 - Covariates: $L(t) \in \mathbb{R}^d$
- **Counting processes**
 - $N^x; x \in \{a, \ell, y\}$ (treatment, covariate, death)
 - Observed until censoring time C
 - Terminal event time: T_e
- **Assumptions**
 - Jumps in $A(t)$ or $L(t)$ only when corresponding N^a or (N^ℓ or N^a) jump.
 - No simultaneous jumps + orthogonal martingales

- **Study period:** $[0, \tau_{\text{end}}]$.
- **Càdlàg, jump processes**
 - Treatment: $A(t) \in \{0, 1\}$
 - Covariates: $L(t) \in \mathbb{R}^d$
- **Counting processes**
 - $N^x; x \in \{a, \ell, y\}$ (treatment, covariate, death)
 - Observed until censoring time C
 - Terminal event time: T_e
- **Assumptions**
 - Jumps in $A(t)$ or $L(t)$ only when corresponding N^a or (N^ℓ or N^a) jump.
 - No simultaneous jumps + orthogonal martingales
- **Treatment decisions**
 - Doctor chooses treatment at jump times of N^a
 - Intervention specifies decision, not timing

- **Study period:** $[0, \tau_{\text{end}}]$.
- **Càdlàg, jump processes**
 - Treatment: $A(t) \in \{0, 1\}$
 - Covariates: $L(t) \in \mathbb{R}^d$
- **Counting processes**
 - $N^x; x \in \{a, \ell, y\}$ (treatment, covariate, death)
 - Observed until censoring time C
 - Terminal event time: T_e
- **Assumptions**
 - Jumps in $A(t)$ or $L(t)$ only when corresponding N^a or (N^ℓ or N^a) jump.
 - No simultaneous jumps + orthogonal martingales
- **Treatment decisions**
 - Doctor chooses treatment at jump times of N^a
 - Intervention specifies decision, not timing
- **Bounded events**
 - Each individual has at most K events in $[0, \tau_{\text{end}}]$

$$\mathcal{F}_t = \sigma\left(\left(A(s), L(s), N^a(s), N^\ell(s), N^y(s)\right) : s \leq t\right) \vee \sigma\left(\left(A(0), L(0)\right)\right)$$

- \mathcal{F}_t : natural filtration for the processes without censoring

$$\mathcal{F}_t = \sigma\left(\left(A(s), L(s), N^a(s), N^\ell(s), N^y(s)\right) : s \leq t\right) \vee \sigma\left(\left(A(0), L(0)\right)\right)$$

$$\mathcal{F}_t^{\text{full}} = \sigma\left(\left(A(s), L(s), N^a(s), N^\ell(s), N^y(s), N^c(s)\right) : s \leq t\right) \vee \sigma\left(\left(A(0), L(0)\right)\right)$$

- \mathcal{F}_t : natural filtration for the processes without censoring
- $\mathcal{F}_t^{\text{full}}$: natural filtration for the processes including censoring

$$\mathcal{F}_t = \sigma\left(\left(A(s), L(s), N^a(s), N^\ell(s), N^y(s)\right) : s \leq t\right) \vee \sigma((A(0), L(0)))$$

$$\mathcal{F}_t^{\text{full}} = \sigma\left(\left(A(s), L(s), N^a(s), N^\ell(s), N^y(s), N^c(s)\right) : s \leq t\right) \vee \sigma((A(0), L(0)))$$

$$\bar{\mathcal{F}}_t = \sigma\left(\left(A(s \wedge C), L(s \wedge C), N^a(s \wedge C), N^\ell(s \wedge C), N^y(s \wedge C), N^c(s \wedge T^e)\right) : s \leq t\right) \vee \sigma((A(0), L(0)))$$

- \mathcal{F}_t : natural filtration for the processes without censoring
- $\mathcal{F}_t^{\text{full}}$: natural filtration for the processes including censoring
- $\bar{\mathcal{F}}_t$: observed (natural) filtration

$$\mathcal{F}_t = \sigma\left(\left(A(s), L(s), N^a(s), N^\ell(s), N^y(s)\right) : s \leq t\right) \vee \sigma\left(\left(A(0), L(0)\right)\right)$$

$$\mathcal{F}_t^{\text{full}} = \sigma\left(\left(A(s), L(s), N^a(s), N^\ell(s), N^y(s), N^c(s)\right) : s \leq t\right) \vee \sigma\left(\left(A(0), L(0)\right)\right)$$

$$\bar{\mathcal{F}}_t = \sigma\left(\left(A(s \wedge C), L(s \wedge C), N^a(s \wedge C), N^\ell(s \wedge C), N^y(s \wedge C), N^c(s \wedge T^e)\right) : s \leq t\right) \vee \sigma\left(\left(A(0), L(0)\right)\right)$$

- \mathcal{F}_t : natural filtration for the processes without censoring
- $\mathcal{F}_t^{\text{full}}$: natural filtration for the processes including censoring
- $\bar{\mathcal{F}}_t$: observed (natural) filtration
- Data format (uncensored)

$$\left(T_{(K)}, \Delta_{(K)}, A\left(T_{(K-1)}\right), L\left(T_{(K-1)}\right), \underbrace{T_{(K-1)}}_{\text{ordered event time}}, \underbrace{\Delta_{(K-1)}}_{\substack{\in \{a, y, \ell\} \\ \text{status indicator}}}, \dots, A(0), L(0) \right)$$

- Data format (censored)

$$\left(\bar{T}_{(K)}, \bar{\Delta}_{(K)}, A\left(\bar{T}_{K-1}\right), L\left(\bar{T}_{K-1}\right), \bar{T}_{(K-1)}, \bar{\Delta}_{(K-1)}, \dots, A(0), L(0) \right)$$

- **Random measure** N_t^{a*} : random measure associated to N_a and A given by

$$N_t^{a*} = \sum_{k: \Delta_{(k)} = a} \delta_{(T_{(k)}, A(T_{(k)}))}$$

- **Random measure** N_t^{a*} : random measure associated to N_a and A given by

$$N_t^{a*} = \sum_{k: \Delta_{(k)} = a} \delta_{(T_{(k)}, A(T_{(k)}))}$$

- **Intervention**

- ▶ Modify compensator: $\Lambda_t^{a*}(\cdot) = \pi_t(\cdot) \Lambda^a(t)$
- ▶ Replace treatment mechanism
 - $\pi_t(\{x\}) = P(A(t) = x \mid \mathcal{F}_{t-})$
 - Under new law P^{G^*} , compensator of N^a is $\pi_t^*(\cdot) \Lambda^a(t)$ with respect to \mathcal{F}_t

- **Random measure** N_t^{a*} : random measure associated to N_a and A given by

$$N_t^{a*} = \sum_{k: \Delta_{(k)} = a} \delta_{(T_{(k)}, A(T_{(k)}))}$$

- **Intervention**

- ▶ Modify compensator: $\Lambda_t^{a*}(\cdot) = \pi_t(\cdot) \Lambda^a(t)$
- ▶ Replace treatment mechanism
 - $\pi_t(\{x\}) = P(A(t) = x \mid \mathcal{F}_{t-})$
 - Under new law P^{G^*} , compensator of N^a is $\pi_t^*(\cdot) \Lambda^a(t)$ with respect to \mathcal{F}_t

- **Special case**

- ▶ $\pi_t^*(\{x\}) \equiv \mathbb{1}\{x = 1\}$ (stay on treatment)

- **Random measure** N_t^{a*} : random measure associated to N_a and A given by

$$N_t^{a*} = \sum_{k: \Delta_{(k)} = a} \delta_{(T_{(k)}, A(T_{(k)}))}$$

- **Intervention**

- ▶ Modify compensator: $\Lambda_t^{a*}(\cdot) = \pi_t(\cdot) \Lambda^a(t)$
- ▶ Replace treatment mechanism
 - $\pi_t(\{x\}) = P(A(t) = x \mid \mathcal{F}_{t-})$
 - Under new law P^{G^*} , compensator of N^a is $\pi_t^*(\cdot) \Lambda^a(t)$ with respect to \mathcal{F}_t

- **Special case**

- ▶ $\pi_t^*(\{x\}) \equiv \mathbb{1}\{x = 1\}$ (stay on treatment)

- **Target parameter**

- ▶ $\Psi_\tau(P) = \mathbb{E}_P \left[\frac{dP^{G^*}}{dP}(\tau) N^y(\tau) \right] = \mathbb{E}_{P^{G^*}} [N^y(\tau)], \tau < \tau_{\text{end}}$

- **Efficient influence function (EIF)** for $\Psi_\tau(P)$ in the nonparametric model is given by (Rytgaard et al. (2022))

$$\begin{aligned}
\varphi_\tau^*(P) = & \mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \mathcal{F}_0] - \Psi_\tau(P) \\
& + \int_0^\tau \frac{dP^{G^*}}{dP}(t-) (\mathbb{E}_{P^{G^*}} [N_y(\tau) \mid L(t), N^\ell(t), \mathcal{F}_{t-}] - \mathbb{E}_{P^{G^*}} [N_y(\tau) \mid N^\ell(t), \mathcal{F}_{t-}]) \tilde{N}^\ell(dt) \\
& + \int_0^\tau \frac{dP^{G^*}}{dP}(t-) (\mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \Delta N^\ell(t) = 1, \mathcal{F}_{t-}] - \mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \Delta N^\ell(t) = 0, \mathcal{F}_{t-}]) \tilde{M}^\ell(dt) \\
& + \int_0^\tau \frac{dP^{G^*}}{dP}(t-) (\mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \Delta N^a(t) = 1, \mathcal{F}_{t-}] - \mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \Delta N^a(t) = 0, \mathcal{F}_{t-}]) \tilde{M}^a(dt) \\
& + \int_0^\tau \frac{dP^{G^*}}{dP}(t-) (1 - \mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \Delta N^y(t) = 0, \mathcal{F}_{t-}]) \tilde{M}^y(dt).
\end{aligned}$$

- $\tilde{M}^x(t) = \tilde{N}^x(t) - \Lambda^x(t)$ is the P - $\bar{\mathcal{F}}_t$ martingale for $\tilde{N}^x(t) = N^x(t \wedge C)$.

5.1. Efficient influence function (continued) 5. Efficient influence function (Rytgaard et al. (2022))

- To work within the targeted learning framework, we need the efficient influence function.

5.1. Efficient influence function (continued) 5. Efficient influence function (Rytgaard et al. (2022))


- To work within the targeted learning framework, we need the efficient influence function.
- Rytgaard et al. (2022) propose sequential regressions for estimating terms in $\varphi_{\tau}^*(P)$.

5.1. Efficient influence function (continued) 5. Efficient influence function (Rytgaard et al. (2022))


- To work within the targeted learning framework, we need the efficient influence function.
- Rytgaard et al. (2022) propose sequential regressions for estimating terms in $\varphi_{\tau}^*(P)$.
- Implementation is unclear and may require thousands of iterations (iterate through all unique event times in the sample).

- To work within the targeted learning framework, we need the efficient influence function.
- Rytgaard et al. (2022) propose sequential regressions for estimating terms in $\varphi_{\tau}^*(P)$.
- Implementation is unclear and may require thousands of iterations (iterate through all unique event times in the sample).
- **Idea:** Replace \mathcal{F}_{t-} with simpler histories:
 - $\mathcal{F}_{T_{(k)}} = \sigma\left(A\left(T_{(j)}\right), L\left(T_{(j)}\right), T_{(j)}, \Delta_{(j)} : j \leq k\right) \vee \sigma((A(0), L(0)))$
 - Censored versions: $\mathcal{F}_{\bar{T}_{(k)}} = \sigma\left(A\left(\bar{T}_j\right), L\left(\bar{T}_j\right), \bar{T}_{(j)}, \bar{\Delta}_{(j)} : j \leq k\right) \vee \sigma((A(0), L(0)))$


Rytgaard et al. (2022):



Rytgaard et al. (2022):



ICE-IPCW (Ohlendorff et al. (2025)):



- **Propensity score:**

- $\pi_k(t, L(T_{(k)}), \mathcal{F}_{T_{(k-1)}}): P(A(T_{(k)}) = 1 \mid L(T_{(k)}), T_{(k)} = t, \Delta_{(k)} = a, \mathcal{F}_{T_{(k-1)}})$

- **Hazard measures:**

- $\tilde{\Lambda}_k^c(t \mid \mathcal{F}_{\bar{T}_{(k-1)}}):$ hazard measure for $(\bar{T}_{(k)}, \mathbb{1}\{\bar{\Delta}_{(k)} = c\})$ given $\mathcal{F}_{\bar{T}_{(k-1)}}$
 - $\Lambda_k^x(t, \mathcal{F}_{T_{(k-1)}}):$ hazard measure of $(T_{(k)}, \mathbb{1}\{\Delta_{(k)} = x\})$ given $\mathcal{F}_{T_{(k-1)}}$ for $x \in \{a, \ell, y\}$

- **Propensity score:**

- $\pi_k(t, L(T_{(k)}), \mathcal{F}_{T_{(k-1)}}): P(A(T_{(k)}) = 1 \mid L(T_{(k)}), T_{(k)} = t, \Delta_{(k)} = a, \mathcal{F}_{T_{(k-1)}})$

- **Hazard measures:**

- $\tilde{\Lambda}_k^c(t \mid \mathcal{F}_{\bar{T}_{(k-1)}}):$ hazard measure for $(\bar{T}_{(k)}, \mathbb{1}\{\bar{\Delta}_{(k)} = c\})$ given $\mathcal{F}_{\bar{T}_{(k-1)}}$
 - $\Lambda_k^x(t, \mathcal{F}_{T_{(k-1)}}):$ hazard measure of $(T_{(k)}, \mathbb{1}\{\Delta_{(k)} = x\})$ given $\mathcal{F}_{T_{(k-1)}}$ for $x \in \{a, \ell, y\}$

- \prod : product integral (Gill & Johansen (1990))

- **Propensity score:**

- $\pi_k(t, L(T_{(k)}), \mathcal{F}_{T_{(k-1)}}): P(A(T_{(k)}) = 1 \mid L(T_{(k)}), T_{(k)} = t, \Delta_{(k)} = a, \mathcal{F}_{T_{(k-1)}})$

- **Hazard measures:**

- $\tilde{\Lambda}_k^c(t \mid \mathcal{F}_{\bar{T}_{(k-1)}}):$ hazard measure for $(\bar{T}_{(k)}, \mathbb{1}\{\bar{\Delta}_{(k)} = c\})$ given $\mathcal{F}_{\bar{T}_{(k-1)}}$
 - $\Lambda_k^x(t, \mathcal{F}_{T_{(k-1)}}):$ hazard measure of $(T_{(k)}, \mathbb{1}\{\Delta_{(k)} = x\})$ given $\mathcal{F}_{T_{(k-1)}}$ for $x \in \{a, \ell, y\}$

- \prod : product integral (Gill & Johansen (1990))

- **Survival functions:**

- $\tilde{S}^c(t \mid \mathcal{F}_{\bar{T}_{(k-1)}}) = \prod_{s \in (\bar{T}_{(k-1)}, t]} \left(1 - d\tilde{\Lambda}_k^c(s \mid \mathcal{F}_{\bar{T}_{(k-1)}})\right)$
 - $S(t \mid \mathcal{F}_{T_{(k-1)}}) = \prod_{s \in (T_{(k-1)}, t]} \left(1 - \sum_{x=a, \ell, y, d} d\Lambda_k^x(s \mid \mathcal{F}_{T_{(k-1)}})\right)$

- **Propensity score:**

- $\pi_k(t, L(T_{(k)}), \mathcal{F}_{T_{(k-1)}}): P(A(T_{(k)}) = 1 \mid L(T_{(k)}), T_{(k)} = t, \Delta_{(k)} = a, \mathcal{F}_{T_{(k-1)}})$

- **Hazard measures:**

- $\tilde{\Lambda}_k^c(t \mid \mathcal{F}_{\bar{T}_{(k-1)}}):$ hazard measure for $(\bar{T}_{(k)}, \mathbb{1}\{\bar{\Delta}_{(k)} = c\})$ given $\mathcal{F}_{\bar{T}_{(k-1)}}$
 - $\Lambda_k^x(t, \mathcal{F}_{T_{(k-1)}}):$ hazard measure of $(T_{(k)}, \mathbb{1}\{\Delta_{(k)} = x\})$ given $\mathcal{F}_{T_{(k-1)}}$ for $x \in \{a, \ell, y\}$

- \prod : product integral (Gill & Johansen (1990))

- **Survival functions:**

- $\tilde{S}^c(t \mid \mathcal{F}_{\bar{T}_{(k-1)}}) = \prod_{s \in (\bar{T}_{(k-1)}, t]} \left(1 - d\tilde{\Lambda}_k^c(s \mid \mathcal{F}_{\bar{T}_{(k-1)}})\right)$
 - $S(t \mid \mathcal{F}_{T_{(k-1)}}) = \prod_{s \in (T_{(k-1)}, t]} \left(1 - \sum_{x=a, \ell, y, d} d\Lambda_k^x(s \mid \mathcal{F}_{T_{(k-1)}})\right)$

- **Random measure:**

- $N = \sum_k \delta_{(T_{(k)}, \Delta_{(k)}, A(T_{(k)}), L(T_{(k)}))}$
 - Turns out that natural filtration of N is $\mathcal{F}_t^{\text{full}}$ (Chapter 2.5 of Last & Brandt (1995))

7.1. Independent censoring conditions

- Let $\bar{Q}_{K,\tau}^g : (a_k, h_k) \mapsto 0$

7. Consistency of ICE-IPCW (right-censoring)

7.1. Independent censoring conditions

- Let $\bar{Q}_{K,\tau}^g : (a_k, h_k) \mapsto 0$
- Define recursively, for $k = K, \dots, 0$,

$$\begin{aligned} \bar{Z}_{k,\tau}^a(u) = & \frac{1}{\tilde{S}^c(\bar{T}_{(k)} - | A(\bar{T}_{k-1}), \bar{H}_{k-1})} \left(\mathbb{1}\{\bar{T}_{(k)} \leq u, \bar{T}_{(k)} < \tau, \bar{\Delta}_{(k)} = a\} \bar{Q}_{k,\tau}^g(1, \bar{H}_k) \right. \\ & + \mathbb{1}\{\bar{T}_{(k)} \leq u, \bar{T}_{(k)} < \tau, \bar{\Delta}_{(k)} = \ell\} \bar{Q}_{k,\tau}^g(A(\bar{T}_k), \bar{H}_k) \\ & \left. + \mathbb{1}\{\bar{T}_{(k)} \leq u, \bar{\Delta}_{(k)} = y\} \right), \end{aligned}$$

and

$$\bar{Q}_{k,\tau}^g : (u, a_k, h_k) \mapsto \mathbb{E}_P \left[\bar{Z}_{k+1,\tau}^a(u) \mid A(\bar{T}_k) = a_k, \bar{H}_k = h_k \right], \quad u \leq \tau$$

where $h_k = (a_k, l_k, t_k, d_k, \dots, a_0, l_0)$.

Theorem 7.2.1 Assume that the $P\text{-}\mathcal{F}_t^{\text{full}}$ compensator Λ of N is also the $P\text{-}\mathcal{F}_t$ compensator of N .

Theorem 7.2.1 Assume that the P - $\mathcal{F}_t^{\text{full}}$ compensator Λ of N is also the P - \mathcal{F}_t compensator of N .

If

1. $\Delta \tilde{\Lambda}_k^c(\cdot, \bar{\mathcal{F}}_{\bar{T}_{(k-1)}}) \Delta \Lambda_k^x(\cdot, \mathcal{F}_{T_{(k-1)}}) \equiv 0$ for $x \in \{a, \ell, y\}$ and $k \in \{1, \dots, K\}$.
2. $\tilde{S}^c(t \mid \bar{\mathcal{F}}_{\bar{T}_{(k-1)}}) > \eta$ for all $t \in (0, \tau]$ and $k \in \{1, \dots, K\}$ P -a.s. for some $\eta > 0$.

Theorem 7.2.1 Assume that the P - $\mathcal{F}_t^{\text{full}}$ compensator Λ of N is also the P - \mathcal{F}_t compensator of N .

If

1. $\Delta \tilde{\Lambda}_k^c(\cdot, \bar{\mathcal{F}}_{\bar{T}_{(k-1)}}) \Delta \Lambda_k^x(\cdot, \mathcal{F}_{T_{(k-1)}}) \equiv 0$ for $x \in \{a, \ell, y\}$ and $k \in \{1, \dots, K\}$.
2. $\tilde{S}^c(t \mid \bar{\mathcal{F}}_{\bar{T}_{(k-1)}}) > \eta$ for all $t \in (0, \tau]$ and $k \in \{1, \dots, K\}$ P -a.s. for some $\eta > 0$.

It holds that

$$\Psi_\tau^g(P) = \mathbb{E}_P[\bar{Q}_{0,\tau}^g(\tau, 1, L(0))].$$

Theorem 7.2.1 Assume that the P - $\mathcal{F}_t^{\text{full}}$ compensator Λ of N is also the P - \mathcal{F}_t compensator of N .

If

1. $\Delta \tilde{\Lambda}_k^c(\cdot, \bar{\mathcal{F}}_{\bar{T}_{(k-1)}}) \Delta \Lambda_k^x(\cdot, \mathcal{F}_{T_{(k-1)}}) \equiv 0$ for $x \in \{a, \ell, y\}$ and $k \in \{1, \dots, K\}$.
2. $\tilde{S}^c(t \mid \bar{\mathcal{F}}_{\bar{T}_{(k-1)}}) > \eta$ for all $t \in (0, \tau]$ and $k \in \{1, \dots, K\}$ P -a.s. for some $\eta > 0$.

It holds that

$$\Psi_{\tau}^g(P) = \mathbb{E}_P [\bar{Q}_{0,\tau}^g(\tau, 1, L(0))].$$

- We make *explicit* use of the fact that the compensator can be explicitly written in terms of the regular conditional distributions of the variables $(\bar{T}_{(k)}, \bar{\Delta}_{(k)}, A(\bar{T}_k), L(\bar{T}_k))$, $k = 1, \dots, K$ and $(A(0), L(0))$.

Theorem 8.1 Under suitable regularity conditions, $\varphi_\tau^*(P)$ can be rewritten as

$$\begin{aligned}
\varphi_\tau^*(P) = & \frac{\mathbb{1}\{A(0) = 1\}}{\pi_0(L(0))} \sum_{k=1}^K \prod_{j=1}^{k-1} \left(\frac{\mathbb{1}\{A(\bar{T}_j) = 1\}}{\pi_j(\bar{T}_{(j)}, L(\bar{T}_j), \bar{\mathcal{F}}_{\bar{T}_{(j-1)}})} \right)^{\mathbb{1}\{\bar{\Delta}_{(j)}=a\}} \frac{1}{\prod_{j=1}^{k-1} \tilde{S}^c(\bar{T}_{(j)} - | \bar{\mathcal{F}}_{\bar{T}_{(j-1)}})} \\
& \times \mathbb{1}\{\bar{\Delta}_{(k-1)} \in \{\ell, a\}, \bar{T}_{(k-1)} < \tau\} \left((\bar{Z}_{k,\tau}^a(\tau) - \bar{Q}_{k-1,\tau}^g(\tau)) \right. \\
& \quad \left. + \int_{\bar{T}_{(k-1)}}^{\tau \wedge \bar{T}_{(k)}} (\bar{Q}_{k-1,\tau}^g(\tau) - \bar{Q}_{k-1,\tau}^g(u)) \frac{1}{\tilde{S}^c(u | \bar{\mathcal{F}}_{\bar{T}_{(k-1)}}) S(u - | \bar{\mathcal{F}}_{\bar{T}_{(k-1)}})} \tilde{M}^c(du) \right) \\
& + \bar{Q}_{0,\tau}^g(\tau) - \Psi_\tau^g(P).
\end{aligned}$$

- **Estimator**
 - One-step estimator based on the efficient influence function

- **Estimator**
 - One-step estimator based on the efficient influence function
- **Simulations**
 - Lower bias than LTMLE (van der Laan & Gruber (2012)) and good CI coverage.
 - Mean squared errors are however close to being the same.
- **Application**: Apply to real-world data with Danish registry data:
 - EIF provides confidence intervals comparable to bootstrap CIs.

- **Estimator**
 - One-step estimator based on the efficient influence function
- **Simulations**
 - Lower bias than LTMLE (van der Laan & Gruber (2012)) and good CI coverage.
 - Mean squared errors are however close to being the same.
- **Application**: Apply to real-world data with Danish registry data:
 - EIF provides confidence intervals comparable to bootstrap CIs.
- **Challenges**
 - Variance estimation is difficult (censoring martingale term)
 - Possible approaches:
 - Undersmooth estimation of censoring compensator
 - Machine learning (e.g., neural networks) methods for multivariate outcomes (to $\bar{Q}_{k,\tau}^g(u)$)

- **Estimator**
 - One-step estimator based on the efficient influence function
- **Simulations**
 - Lower bias than LTMLE (van der Laan & Gruber (2012)) and good CI coverage.
 - Mean squared errors are however close to being the same.
- **Application:** Apply to real-world data with Danish registry data:
 - EIF provides confidence intervals comparable to bootstrap CIs.
- **Challenges**
 - Variance estimation is difficult (censoring martingale term)
 - Possible approaches:
 - Undersmooth estimation of censoring compensator
 - Machine learning (e.g., neural networks) methods for multivariate outcomes (to $\bar{Q}_{k,\tau}^g(u)$)
- **Next steps**
 - Empirical process & remainder term conditions not yet addressed (ongoing work)
 - Consider TMLE instead of one-step \Rightarrow ensures estimates in $[0, 1]$
 - Apply flexible, data-adaptive estimators for nuisance parameters
 - Clarify causal interpretation of target parameter (identifiability)

Bibliography

- Adams, R., Saria, S., & Rosenblum, M. (2020). The impact of time series length and discretization on longitudinal causal estimation methods. *Arxiv Preprint Arxiv:2011.15099*.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., Newey, W., & Robins, J. (2018). Double/debiased machine learning for treatment and structural parameters. *The Econometrics Journal*, 21(1), C1–C68. <https://doi.org/10.1111/ectj.12097>
- Ferreira Guerra, S., Schnitzer, M. E., Forget, A., & Blais, L. (2020). Impact of discretization of the timeline for longitudinal causal inference methods. *Statistics in Medicine*, 39(27), 4069–4085. <https://doi.org/https://doi.org/10.1002/sim.8710>
- Gill, R. D., & Johansen, S. (1990). A survey of product-integration with a view toward application in survival analysis. *The Annals of Statistics*, 1501–1555.
- Kant, W. M., & Krijthe, J. H. (2025). Irregular measurement times in estimating time-varying treatment effects: Categorizing biases and comparing adjustment methods. *Arxiv Preprint Arxiv:2501.11449*.
- Last, G., & Brandt, A. (1995). *Marked Point Processes on the Real Line: The Dynamical Approach*. Springer. <https://link.springer.com/book/9780387945477>

- Ohlendorff, J. S., Munch, A., & Gerds, T. A. (2025). *A Novel Sequential Regression Approach for Efficient Continuous-Time Causal Inference*.
- Ryalen, P. C., Stensrud, M. J., & Røysland, K. (2019). The additive hazard estimator is consistent for continuous-time marginal structural models. *Lifetime Data Analysis*, 25, 611–638.
- Rytgaard, H. C., Gerds, T. A., & van der Laan, M. J. (2022). Continuous-Time Targeted Minimum Loss-Based Estimation of Intervention-Specific Mean Outcomes. *The Annals of Statistics*, 50(5), 2469–2491. <https://doi.org/10.1214/21-AOS2114>
- Sofrygin, O., Zhu, Z., Schmittdiel, J. A., Adams, A. S., Grant, R. W., van der Laan, M. J., & Neugebauer, R. (2019). Targeted learning with daily EHR data. *Statistics in Medicine*, 38(16), 3073–3090.
- Sun, J., & Crawford, F. W. (2023). The role of discretization scales in causal inference with continuous-time treatment. *Arxiv Preprint Arxiv:2306.08840*.
- van der Laan, M. J., & Gruber, S. (2012). *The International Journal of Biostatistics*, 8(1). <https://doi.org/doi:10.1515/1557-4679.1370>
- van der Laan, M. J., & Rubin, D. (2006). Targeted Maximum Likelihood Learning. *The International Journal of Biostatistics*, 2(1). <https://doi.org/10.2202/1557-4679.1043>