

Sequential Regressions for Efficient Continuous-Time Causal Inference

Johan Sebastian Ohlendorff

2025-09-02

- PhD student in Biostatistics at the Section of Biostatistics, University of Copenhagen.

- PhD student in Biostatistics at the Section of Biostatistics, University of Copenhagen.
- Supervisors: Thomas Alexander Gerds and Anders Munch.

- PhD student in Biostatistics at the Section of Biostatistics, University of Copenhagen.
- Supervisors: Thomas Alexander Gerds and Anders Munch.
- Work on continuous-time longitudinal causal inference using
 - Targeted learning (e.g., TMLE Rytgaard et al. (2022) or one-step estimation).

- PhD student in Biostatistics at the Section of Biostatistics, University of Copenhagen.
- Supervisors: Thomas Alexander Gerds and Anders Munch.
- Work on continuous-time longitudinal causal inference using
 - Targeted learning (e.g., TMLE Rytgaard et al. (2022) or one-step estimation).
- Working on a continuous-time scale is motivated by:
 - Real-world data often recorded in continuous time (e.g., electronic health records).
 - Avoiding bias due to discretization (Adams et al. (2020); Ferreira Guerra et al. (2020); Kant & Krijthe (2025); Ryalen et al. (2019); Sofrygin et al. (2019); Sun & Crawford (2023))

- We observe a càdlàg, jump process for the treatment $(A(t))_{t \in [0, \tau_{\text{end}}]} \in \{0, 1\}$ and a covariate process $(L(t))_{t \in [0, \tau_{\text{end}}]}$, such that $L(t)$ almost surely takes values some finite subset of \mathbb{R}^d .

- We observe a càdlàg, jump process for the treatment $(A(t))_{t \in [0, \tau_{\text{end}}]} \in \{0, 1\}$ and a covariate process $(L(t))_{t \in [0, \tau_{\text{end}}]}$, such that $L(t)$ almost surely takes values some finite subset of \mathbb{R}^d .
- Assume that we observe the counting processes N^x , $x = a, \ell, y$ (treatment, covariate, death, censoring) up to a right-censoring time C which is distinct from all event times with probability 1. Terminal event time is denoted by T^e .

- We observe a càdlàg, jump process for the treatment $(A(t))_{t \in [0, \tau_{\text{end}}]} \in \{0, 1\}$ and a covariate process $(L(t))_{t \in [0, \tau_{\text{end}}]}$, such that $L(t)$ almost surely takes values some finite subset of \mathbb{R}^d .
- Assume that we observe the counting processes N^x , $x = a, \ell, y$ (treatment, covariate, death, censoring) up to a right-censoring time C which is distinct from all event times with probability 1. Terminal event time is denoted by T^e .
- Assume that $\Delta A(t) \neq 0$ only if $\Delta N^a(t) \neq 0$ and $\Delta L(t) \neq 0$ only if $\Delta N^\ell(t) \neq 0$ or $\Delta N^a(t) \neq 0$.

- We observe a càdlàg, jump process for the treatment $(A(t))_{t \in [0, \tau_{\text{end}}]} \in \{0, 1\}$ and a covariate process $(L(t))_{t \in [0, \tau_{\text{end}}]}$, such that $L(t)$ almost surely takes values some finite subset of \mathbb{R}^d .
- Assume that we observe the counting processes N^x , $x = a, \ell, y$ (treatment, covariate, death, censoring) up to a right-censoring time C which is distinct from all event times with probability 1. Terminal event time is denoted by T^e .
- Assume that $\Delta A(t) \neq 0$ only if $\Delta N^a(t) \neq 0$ and $\Delta L(t) \neq 0$ only if $\Delta N^\ell(t) \neq 0$ or $\Delta N^a(t) \neq 0$. It then seems reasonable to assume that $\Delta N^a \Delta N^\ell \equiv 0$ and that, in fact, every counting process does not jump at the same time as any other counting process. In addition, all considered counting processes have orthogonal martingales.

- We observe a càdlàg, jump process for the treatment $(A(t))_{t \in [0, \tau_{\text{end}}]} \in \{0, 1\}$ and a covariate process $(L(t))_{t \in [0, \tau_{\text{end}}]}$, such that $L(t)$ almost surely takes values some finite subset of \mathbb{R}^d .
- Assume that we observe the counting processes N^x , $x = a, \ell, y$ (treatment, covariate, death, censoring) up to a right-censoring time C which is distinct from all event times with probability 1. Terminal event time is denoted by T^e .
- Assume that $\Delta A(t) \neq 0$ only if $\Delta N^a(t) \neq 0$ and $\Delta L(t) \neq 0$ only if $\Delta N^\ell(t) \neq 0$ or $\Delta N^a(t) \neq 0$. It then seems reasonable to assume that $\Delta N^a \Delta N^\ell \equiv 0$ and that, in fact, every counting process does not jump at the same time as any other counting process. In addition, all considered counting processes have orthogonal martingales.
- The doctor may decide treatment based at times at which $\Delta N^a(t) \neq 0$. The intervention in which we are interested specifies what this decision should be (or the probability of being treated), but does not naturally intervene on when the doctor decides to do so.

- We observe a càdlàg, jump process for the treatment $(A(t))_{t \in [0, \tau_{\text{end}}]} \in \{0, 1\}$ and a covariate process $(L(t))_{t \in [0, \tau_{\text{end}}]}$, such that $L(t)$ almost surely takes values some finite subset of \mathbb{R}^d .
- Assume that we observe the counting processes N^x , $x = a, \ell, y$ (treatment, covariate, death, censoring) up to a right-censoring time C which is distinct from all event times with probability 1. Terminal event time is denoted by T^e .
- Assume that $\Delta A(t) \neq 0$ only if $\Delta N^a(t) \neq 0$ and $\Delta L(t) \neq 0$ only if $\Delta N^\ell(t) \neq 0$ or $\Delta N^a(t) \neq 0$. It then seems reasonable to assume that $\Delta N^a \Delta N^\ell \equiv 0$ and that, in fact, every counting process does not jump at the same time as any other counting process. In addition, all considered counting processes have orthogonal martingales.
- The doctor may decide treatment based at times at which $\Delta N^a(t) \neq 0$. The intervention in which we are interested specifies what this decision should be (or the probability of being treated), but does not naturally intervene on when the doctor decides to do so.
- Each individual has at most K events in $[0, \tau_{\text{end}}]$, i.e., $\sum_{x=a,y,c,\ell} N^x(\tau_{\text{end}}) \leq K$ almost surely.

$$\mathcal{F}_t = \sigma\big((A(s), L(s), N^a(s), N^\ell(s), N^y(s)) : s \leq t\big)$$

- \mathcal{F}_t is the natural filtration for the processes without censoring.

$$\mathcal{F}_t = \sigma\big((A(s), L(s), N^a(s), N^\ell(s), N^y(s)) : s \leq t\big)$$

$$\mathcal{F}_t^\beta = \sigma\big((A(s), L(s), N^a(s), N^\ell(s), N^y(s), N^c(s)) : s \leq t\big)$$

- \mathcal{F}_t is the natural filtration for the processes without censoring.
- \mathcal{F}_t^β is the natural filtration for the processes including censoring.

$$\begin{aligned}\mathcal{F}_t &= \sigma\big((A(s), L(s), N^a(s), N^\ell(s), N^y(s)) : s \leq t\big) \\ \mathcal{F}_t^\beta &= \sigma\big((A(s), L(s), N^a(s), N^\ell(s), N^y(s), N^c(s)) : s \leq t\big) \\ \mathcal{F}_t^{\tilde{\beta}} &= \sigma\big((A(s \wedge C), L(s \wedge C), N^a(s \wedge C), N^\ell(s \wedge C), \\ &\quad N^y(s \wedge C), N^c(s \wedge T^e)) : s \leq t\big)\end{aligned}$$

- \mathcal{F}_t is the natural filtration for the processes without censoring.
- \mathcal{F}_t^β is the natural filtration for the processes including censoring.
- $\mathcal{F}_t^{\tilde{\beta}}$ is the observed filtration, i.e., the natural filtration stopped by death and censoring.

$$\begin{aligned}\mathcal{F}_t &= \sigma\big((A(s), L(s), N^a(s), N^\ell(s), N^y(s)) : s \leq t\big) \\ \mathcal{F}_t^\beta &= \sigma\big((A(s), L(s), N^a(s), N^\ell(s), N^y(s), N^c(s)) : s \leq t\big) \\ \mathcal{F}_t^{\tilde{\beta}} &= \sigma\big((A(s \wedge C), L(s \wedge C), N^a(s \wedge C), N^\ell(s \wedge C), \\ &\quad N^y(s \wedge C), N^c(s \wedge T^e)) : s \leq t\big)\end{aligned}$$

- \mathcal{F}_t is the natural filtration for the processes without censoring.
- $\mathcal{F}_{t_{\sim}}^\beta$ is the natural filtration for the processes including censoring.
- $\mathcal{F}_t^{\tilde{\beta}}$ is the observed filtration, i.e., the natural filtration stopped by death and censoring.
- Data format (uncensored)

$$(T_{(K)}, \Delta_{(K)}, A(T_{(K-1)}), L(T_{(K-1)}), T_{(K-1)}, \Delta_{(K-1)}, \dots, A(0), L(0))$$

- Data format (censored)

$$(\bar{T}_{(K)}, \bar{\Delta}_{(K)}, A(\bar{T}_{K-1}), L(\bar{T}_{K-1}), \bar{T}_{(K-1)}, \bar{\Delta}_{(K-1)}, \dots, A(0), L(0))$$

- Let $N_t^a(\cdot)$ denote the random measure associated with N^a and $A(\cdot)$,

$$N_t^a(A) = \sum_{k: \Delta_{(k)}=a} \delta_{(T_{(k)}, A(T_{(k)}))}(A).$$

- Let $N_t^a(\cdot)$ denote the random measure associated with N^a and $A(\cdot)$,

$$N_t^a(A) = \sum_{k: \Delta_{(k)}=a} \delta_{(T_{(k)}, A(T_{(k)}))}(A).$$

- Interested on “intervening” on the compensator of $N^a(\cdot)$, $\Lambda_t^a(A) = \pi_t(A)\Lambda^a(t)$, replacing the treatment mechanism $\pi_t(\{x\}) = P(A(t) = x \mid \mathcal{F}_{t-})$ with a specified treatment mechanism $\pi_t^*(A)$. We denote by P^{G^*} the probability measure in which the P^{G^*} - \mathcal{F}_t compensator of $N^a(\cdot)$ is $\pi_t^*(A)\Lambda^a(t)$.

- Let $N_t^a(\cdot)$ denote the random measure associated with N^a and $A(\cdot)$,

$$N_t^a(A) = \sum_{k: \Delta_{(k)}=a} \delta_{(T_{(k)}, A(T_{(k)}))}(A).$$

- Interested on “intervening” on the compensator of $N^a(\cdot)$, $\Lambda_t^a(A) = \pi_t(A)\Lambda^a(t)$, replacing the treatment mechanism $\pi_t(\{x\}) = P(A(t) = x \mid \mathcal{F}_{t-})$ with a specified treatment mechanism $\pi_t^*(A)$. We denote by P^{G^*} the probability measure in which the P^{G^*} - \mathcal{F}_t compensator of $N^a(\cdot)$ is $\pi_t^*(A)\Lambda^a(t)$.
- Focus on the case $\pi_t^*(\{x\}) \equiv \mathbb{1}\{x = 1\}$.

- Let $N_t^a(\cdot)$ denote the random measure associated with N^a and $A(\cdot)$,

$$N_t^a(A) = \sum_{k: \Delta_{(k)}=a} \delta_{(T_{(k)}, A(T_{(k)}))}(A).$$

- Interested on “intervening” on the compensator of $N^a(\cdot)$, $\Lambda_t^a(A) = \pi_t(A)\Lambda^a(t)$, replacing the treatment mechanism $\pi_t(\{x\}) = P(A(t) = x \mid \mathcal{F}_{t-})$ with a specified treatment mechanism $\pi_t^*(A)$. We denote by P^{G^*} the probability measure in which the P^{G^*} - \mathcal{F}_t compensator of $N^a(\cdot)$ is $\pi_t^*(A)\Lambda^a(t)$.
- Focus on the case $\pi_t^*(\{x\}) \equiv \mathbb{1}\{x = 1\}$.
- We are then interested (are we?) in

$$\Psi_\tau(P) = \mathbb{E}_P \left[\frac{dP^{G^*}}{dP}(\tau) N^y(\tau) \right] = \mathbb{E}_{P^{G^*}}[N^y(\tau)]$$

- With $W^g(t) = \frac{dP^{G^*}}{dP}(t)$, Rytgaard et al. (2022) claims that the following is the EIF:

$$\begin{aligned}
\varphi_{\tau}^*(P) &= \mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \mathcal{F}_0] - \Psi_{\tau}(P) \\
&+ \int_0^{\tau} W^g(t-) (\mathbb{E}_{P^{G^*}} [N_y(\tau) \mid L(t), N^{\ell}(t), \mathcal{F}_{t-}] - \mathbb{E}_{P^{G^*}} [N_y(\tau) \mid N^{\ell}(t), \mathcal{F}_{t-}]) \tilde{N}^{\ell}(dt) \\
&+ \int_0^{\tau} W^g(t-) (\mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \Delta N^{\ell}(t) = 1, \mathcal{F}_{t-}] - \mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \Delta N^{\ell}(t) = 0, \mathcal{F}_{t-}]) \tilde{M}^{\ell}(dt) \\
&+ \int_0^{\tau} W^g(t-) (\mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \Delta N^a(t) = 1, \mathcal{F}_{t-}] - \mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \Delta N^a(t) = 0, \mathcal{F}_{t-}]) \tilde{M}^a(dt) \\
&+ \int_0^{\tau} W^g(t-) (1 - \mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \Delta N^y(t) = 0, \mathcal{F}_{t-}]) \tilde{M}^y(dt).
\end{aligned}$$

- With $W^g(t) = \frac{dP^{G^*}}{dP}(t)$, Rytgaard et al. (2022) claims that the following is the EIF:


$$\begin{aligned}
\varphi_\tau^*(P) &= \mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \mathcal{F}_0] - \Psi_\tau(P) \\
&+ \int_0^\tau W^g(t-) (\mathbb{E}_{P^{G^*}} [N_y(\tau) \mid L(t), N^\ell(t), \mathcal{F}_{t-}] - \mathbb{E}_{P^{G^*}} [N_y(\tau) \mid N^\ell(t), \mathcal{F}_{t-}]) \tilde{N}^\ell(dt) \\
&+ \int_0^\tau W^g(t-) (\mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \Delta N^\ell(t) = 1, \mathcal{F}_{t-}] - \mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \Delta N^\ell(t) = 0, \mathcal{F}_{t-}]) \tilde{M}^\ell(dt) \\
&+ \int_0^\tau W^g(t-) (\mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \Delta N^a(t) = 1, \mathcal{F}_{t-}] - \mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \Delta N^a(t) = 0, \mathcal{F}_{t-}]) \tilde{M}^a(dt) \\
&+ \int_0^\tau W^g(t-) (1 - \mathbb{E}_{P^{G^*}} [N_y(\tau) \mid \Delta N^y(t) = 0, \mathcal{F}_{t-}]) \tilde{M}^y(dt).
\end{aligned}$$



- Here $\tilde{M}^x(t) = \tilde{N}^x(t) - \Lambda^x(t)$ is the martingale for $\tilde{N}^x(t) = N^x(t \wedge C)$.

- It is unclear how to estimate $\mathbb{E}_{PG^*} [N_y(\tau) \mid \Delta N^x(t), \mathcal{F}_{t-}]$.

- It is unclear how to estimate $\mathbb{E}_{PG^*} [N_y(\tau) \mid \Delta N^x(t), \mathcal{F}_{t-}]$.
- Sequential regression not clear how to implement.
- Rytgaard et al. (2022) iterative procedure requires 1000s of iterative steps.
 - Assume that $n = 1000$; if all registrations in the sample are unique and each person has 10 events on average, then we need to fit 10,000 regressions.
- Hard to work with \mathcal{F}_{t-} .

- It is unclear how to estimate $\mathbb{E}_{PG^*} [N_y(\tau) \mid \Delta N^x(t), \mathcal{F}_{t-}]$.
- Sequential regression not clear how to implement.
- Rytgaard et al. (2022) iterative procedure requires 1000s of iterative steps.
 - Assume that $n = 1000$; if all registrations in the sample are unique and each person has 10 events on average, then we need to fit 10,000 regressions.
- Hard to work with \mathcal{F}_{t-} .
- My idea: Can we work with $\mathcal{F}_{T_{(k)}} = \sigma(A(T_{(j)}), L(T_{(j)}), T_{(j)}, \Delta_{(j)} : j \leq k) \vee \sigma((A(0), L(0)))$ instead and more generally $\mathcal{F}_{\bar{T}_{(k)}}^{\bar{\beta}} = \sigma(A(\bar{T}_j), L(\bar{T}_j), \bar{T}_{(j)}, \bar{\Delta}_{(j)} : j \leq k) \vee \sigma((A(0), L(0)))$ and regress back on that information instead of \mathcal{F}_{t-} ?





Let

$$\begin{aligned} \bar{Z}_{k,\tau}^a(u) = & \frac{1}{\tilde{S}^c(\bar{T}_{(k)} - | A(\bar{T}_{k-1}), \bar{H}_{k-1})} \left(\mathbb{1}\{\bar{T}_{(k)} \leq u, \bar{T}_{(k)} < \tau, \bar{\Delta}_{(k)} = a\} \bar{Q}_{k,\tau}^g(1, \bar{H}_k) \right. \\ & + \mathbb{1}\{\bar{T}_{(k)} \leq u, \bar{T}_{(k)} < \tau, \bar{\Delta}_{(k)} = \ell\} \bar{Q}_{k,\tau}^g(A(\bar{T}_k), \bar{H}_k) \\ & \left. + \mathbb{1}\{\bar{T}_{(k)} \leq u, \bar{\Delta}_{(k)} = y\} \right). \end{aligned}$$

Let

$$\begin{aligned}
 \bar{Z}_{k,\tau}^a(u) &= \frac{1}{\tilde{S}^c(\bar{T}_{(k)} - | A(\bar{T}_{k-1}), \bar{H}_{k-1})} \left(\mathbb{1}\{\bar{T}_{(k)} \leq u, \bar{T}_{(k)} < \tau, \bar{\Delta}_{(k)} = a\} \bar{Q}_{k,\tau}^g(1, \bar{H}_k) \right. \\
 &\quad \left. + \mathbb{1}\{\bar{T}_{(k)} \leq u, \bar{T}_{(k)} < \tau, \bar{\Delta}_{(k)} = \ell\} \bar{Q}_{k,\tau}^g(A(\bar{T}_k), \bar{H}_k) \right. \\
 &\quad \left. + \mathbb{1}\{\bar{T}_{(k)} \leq u, \bar{\Delta}_{(k)} = y\} \right). \\
 \bullet \quad \tilde{S}^c\left(t \mid \mathcal{F}_{\bar{T}_{(k-1)}}^{\tilde{\beta}}\right) &= \prod_{s \in (\bar{T}_{(k-1)}, t]} \left(1 - d\tilde{\Lambda}_k^c\left(s \mid \mathcal{F}_{\bar{T}_{(k-1)}}^{\tilde{\beta}}\right) \right).
 \end{aligned}$$

Let

$$\begin{aligned} \bar{Z}_{k,\tau}^a(u) = & \frac{1}{\tilde{S}^c(\bar{T}_{(k)} - | A(\bar{T}_{k-1}), \bar{H}_{k-1})} \left(\mathbb{1}\{\bar{T}_{(k)} \leq u, \bar{T}_{(k)} < \tau, \bar{\Delta}_{(k)} = a\} \bar{Q}_{k,\tau}^g(1, \bar{H}_k) \right. \\ & + \mathbb{1}\{\bar{T}_{(k)} \leq u, \bar{T}_{(k)} < \tau, \bar{\Delta}_{(k)} = \ell\} \bar{Q}_{k,\tau}^g(A(\bar{T}_k), \bar{H}_k) \\ & \left. + \mathbb{1}\{\bar{T}_{(k)} \leq u, \bar{\Delta}_{(k)} = y\} \right). \end{aligned}$$

- $\tilde{S}^c\left(t \mid \mathcal{F}_{\bar{T}_{(k-1)}}^{\tilde{\beta}}\right) = \prod_{s \in (\bar{T}_{(k-1)}, t]} \left(1 - d\tilde{\Lambda}_k^c\left(s \mid \mathcal{F}_{\bar{T}_{(k-1)}}^{\tilde{\beta}}\right)\right).$
- $\tilde{\Lambda}_k^c\left(t \mid \mathcal{F}_{\bar{T}_{(k-1)}}^{\tilde{\beta}}\right)$ denotes the hazard measure of $(\bar{T}_{(k)}, \mathbb{1}\{\bar{\Delta}_{(k)} = c\})$ given $\mathcal{F}_{\bar{T}_{(k-1)}}^{\tilde{\beta}}$ and
- $\Lambda_k^x\left(t, \mathcal{F}_{T_{(k-1)}}\right)$ denotes the hazard measure of $(T_{(k)}, \mathbb{1}\{\Delta_{(k)} = x\})$ given $\mathcal{F}_{T_{(k-1)}}$ for $x \in \{a, \ell, y, d\}.$

Theorem 1 Assume that the compensator Λ^α of N^α with respect to the filtration \mathcal{F}_t^β is also the compensator with respect to the filtration \mathcal{F}_t . Let $\bar{Q}_{K,\tau}^g : (a_k, h_k) \mapsto 0$.

Theorem 1 Assume that the compensator Λ^α of N^α with respect to the filtration \mathcal{F}_t^β is also the compensator with respect to the filtration \mathcal{F}_t . Let $\bar{Q}_{K,\tau}^g : (a_k, h_k) \mapsto 0$. If

1. $\Delta \tilde{\Lambda}_k^c \left(\cdot, \mathcal{F}_{\bar{T}_{(k-1)}}^{\tilde{\beta}} \right) \Delta \Lambda_k^x \left(\cdot, \mathcal{F}_{T_{(k-1)}} \right) \equiv 0$ for $x \in \{a, \ell, y, d\}$ and $k \in \{1, \dots, K\}$.
2. $\tilde{S}^c \left(t \mid \mathcal{F}_{\bar{T}_{(k-1)}}^{\tilde{\beta}} \right) > \eta$ for all $t \in (0, \tau]$ and $k \in \{1, \dots, K\}$ P -a.s. for some $\eta > 0$.

Theorem 1 Assume that the compensator Λ^α of N^α with respect to the filtration \mathcal{F}_t^β is also the compensator with respect to the filtration \mathcal{F}_t . Let $\bar{Q}_{K,\tau}^g : (a_k, h_k) \mapsto 0$. If

1. $\Delta \tilde{\Lambda}_k^c \left(\cdot, \mathcal{F}_{\bar{T}_{(k-1)}}^{\tilde{\beta}} \right) \Delta \Lambda_k^x \left(\cdot, \mathcal{F}_{T_{(k-1)}} \right) \equiv 0$ for $x \in \{a, \ell, y, d\}$ and $k \in \{1, \dots, K\}$.
2. $\tilde{S}^c \left(t \mid \mathcal{F}_{\bar{T}_{(k-1)}}^{\tilde{\beta}} \right) > \eta$ for all $t \in (0, \tau]$ and $k \in \{1, \dots, K\}$ P -a.s. for some $\eta > 0$.

With $h_k = (a_k, l_k, t_k, d_k, \dots, a_0, l_0)$,

$$\bar{Q}_{k,\tau}^g : (u, a_k, h_k) \mapsto \mathbb{E}_P \left[\bar{Z}_{k+1,\tau}^a(u) \mid A(\bar{T}_k) = a_k, \bar{H}_k = h_k \right],$$

for $u \leq \tau$, it holds that

$$\Psi_\tau^g(P) = \mathbb{E}_P \left[\bar{Q}_{0,\tau}^g(\tau, 1, L(0)) \right].$$

Rewriting the efficient influence function

Independent censoring conditions

Let $\tilde{M}^c(t) = \tilde{N}^c(t) - \tilde{\Lambda}^c(t)$ and $S(t \mid \mathcal{F}_{T_{(k-1)}}) = \prod_{s \in (0, t]} \left(1 - \sum_{x=a, \ell, y, d} \Lambda_k^x(ds \mid \mathcal{F}_{T_{(k-1)}})\right)$. Under suitable regularity conditions, the efficient influence function can be written as

$$\begin{aligned} \varphi_\tau^*(P) = & \frac{\mathbb{1}\{A(0) = 1\}}{\pi_0(L(0))} \sum_{k=1}^K \prod_{j=1}^{k-1} \left(\frac{\mathbb{1}\{A(\bar{T}_j) = 1\}}{\pi_j(\bar{T}_{(j)}, L(\bar{T}_j), \mathcal{F}_{\bar{T}_{(j-1)}}^{\tilde{\beta}})} \right)^{\mathbb{1}\{\bar{\Delta}_{(j)}=a\}} \frac{1}{\prod_{j=1}^{k-1} \tilde{S}^c(\bar{T}_{(j)} - \mid \mathcal{F}_{\bar{T}_{(j-1)}}^{\tilde{\beta}})} \\ & \times \mathbb{1}\{\bar{\Delta}_{(k-1)} \in \{\ell, a\}, \bar{T}_{(k-1)} < \tau\} \left(\left(\bar{Z}_{k, \tau}^a(\tau) - \bar{Q}_{k-1, \tau}^g(\tau) \right) \right. \\ & \left. + \int_{\bar{T}_{(k-1)}}^{\tau \wedge \bar{T}_{(k)}} \left(\bar{Q}_{k-1, \tau}^g(\tau) - \bar{Q}_{k-1, \tau}^g(u) \right) \frac{1}{\tilde{S}^c(u \mid \mathcal{F}_{\bar{T}_{(k-1)}}^{\tilde{\beta}}) S(u - \mid \mathcal{F}_{\bar{T}_{(k-1)}}^{\tilde{\beta}})} \tilde{M}^c(du) \right) \\ & + \bar{Q}_{0, \tau}^g(\tau) - \Psi_\tau^g(P), \end{aligned}$$

- We consider a one-step estimator based on the EIF.

- We consider a one-step estimator based on the EIF.
- Simulation studies demonstrate favorable performance of the proposed procedure – lower bias than discrete-time procedures and good coverage of confidence intervals.
- However, variance estimation is challenging due to the censoring martingale term.

- We consider a one-step estimator based on the EIF.
- Simulation studies demonstrate favorable performance of the proposed procedure – lower bias than discrete-time procedures and good coverage of confidence intervals.
- However, variance estimation is challenging due to the censoring martingale term.
- Estimating the censoring martingale term
 - Undersmoothing of the estimation of the censoring compensator to avoid estimation altogether.
 - Using a machine learning methods that can handle multivariate outcomes.

- We consider a one-step estimator based on the EIF.
- Simulation studies demonstrate favorable performance of the proposed procedure – lower bias than discrete-time procedures and good coverage of confidence intervals.
- However, variance estimation is challenging due to the censoring martingale term.
- Estimating the censoring martingale term
 - Undersmoothing of the estimation of the censoring compensator to avoid estimation altogether.
 - Using a machine learning methods that can handle multivariate outcomes.
- For simplicity, empirical process conditions and remainder term conditions are not considered in this work (work in progress).

- We consider a one-step estimator based on the EIF.
- Simulation studies demonstrate favorable performance of the proposed procedure – lower bias than discrete-time procedures and good coverage of confidence intervals.
- However, variance estimation is challenging due to the censoring martingale term.
- Estimating the censoring martingale term
 - Undersmoothing of the estimation of the censoring compensator to avoid estimation altogether.
 - Using a machine learning methods that can handle multivariate outcomes.
- For simplicity, empirical process conditions and remainder term conditions are not considered in this work (work in progress).
- Using a TMLE approach instead of one-step approach \Rightarrow better because we want estimates in $[0, 1]$.

- We consider a one-step estimator based on the EIF.
- Simulation studies demonstrate favorable performance of the proposed procedure – lower bias than discrete-time procedures and good coverage of confidence intervals.
- However, variance estimation is challenging due to the censoring martingale term.
- Estimating the censoring martingale term
 - Undersmoothing of the estimation of the censoring compensator to avoid estimation altogether.
 - Using a machine learning methods that can handle multivariate outcomes.
- For simplicity, empirical process conditions and remainder term conditions are not considered in this work (work in progress).
- Using a TMLE approach instead of one-step approach \Rightarrow better because we want estimates in $[0, 1]$.
- Does the target parameter have a causal interpretation? (Identifiability)

- We consider a one-step estimator based on the EIF.
- Simulation studies demonstrate favorable performance of the proposed procedure – lower bias than discrete-time procedures and good coverage of confidence intervals.
- However, variance estimation is challenging due to the censoring martingale term.
- Estimating the censoring martingale term
 - Undersmoothing of the estimation of the censoring compensator to avoid estimation altogether.
 - Using a machine learning methods that can handle multivariate outcomes.
- For simplicity, empirical process conditions and remainder term conditions are not considered in this work (work in progress).
- Using a TMLE approach instead of one-step approach \Rightarrow better because we want estimates in $[0, 1]$.
- Does the target parameter have a causal interpretation? (Identifiability)
- Other target parameters (e.g., recurrent events, restricted mean survival time, etc.).

Bibliography

- Adams, R., Saria, S., & Rosenblum, M. (2020). The impact of time series length and discretization on longitudinal causal estimation methods. *Arxiv Preprint Arxiv:2011.15099*.
- Ferreira Guerra, S., Schnitzer, M. E., Forget, A., & Blais, L. (2020). Impact of discretization of the timeline for longitudinal causal inference methods. *Statistics in Medicine*, 39(27), 4069–4085. <https://doi.org/https://doi.org/10.1002/sim.8710>
- Kant, W. M., & Krijthe, J. H. (2025). Irregular measurement times in estimating time-varying treatment effects: Categorizing biases and comparing adjustment methods. *Arxiv Preprint Arxiv:2501.11449*.
- Ryalen, P. C., Stensrud, M. J., & Røysland, K. (2019). The additive hazard estimator is consistent for continuous-time marginal structural models. *Lifetime Data Analysis*, 25, 611–638.
- Rytgaard, H. C., Gerds, T. A., & van der Laan, M. J. (2022). Continuous-Time Targeted Minimum Loss-Based Estimation of Intervention-Specific Mean Outcomes. *The Annals of Statistics*, 50(5), 2469–2491. <https://doi.org/10.1214/21-AOS2114>
- Sofrygin, O., Zhu, Z., Schmittdiel, J. A., Adams, A. S., Grant, R. W., van der Laan, M. J., & Neugebauer, R. (2019). Targeted learning with daily EHR data. *Statistics in Medicine*, 38(16), 3073–3090.

Sun, J., & Crawford, F. W. (2023). The role of discretization scales in causal inference with continuous-time treatment. *Arxiv Preprint Arxiv:2306.08840*.