

2.exploration

Joao Lopes

2024-04-18

Contents

1. VARIATION	3
1.1. TYPICAL VALUES	3
1.2. UNUSUAL VALUES	3
2. MISSING VALUES	5
2.1. EXPLICIT MISSING VALUES	5
Last observation carried forward	5
Fixed values	5
Numeric value represents NA	5
NaN	5
2.2. IMPLICIT MISSING VALUES	6
Pivoting	6
Complete	6
Joins	6
2.3. FACTORS AND EMPTY GROUPS	6
3. COVARIATION	8
3.1. CATEGORICAL AND NUMERICAL VARIABLES	8
3.2. TWO CATEGORICAL VARIABLES	8
3.3. TWO NUMERICAL VARIABLES	8
4. PATTERNS AND MODELS	10

1. VARIATION

[from <https://r4ds.hadley.nz/eda#variation>]

```
library("tidyverse")
library("nycflights13")
```

1.1. TYPICAL VALUES

```
#visualize distribution
ggplot(diamonds, aes(x = carat)) +
  geom_histogram(binwidth = 0.5)

#visualize distribution in detail
diamonds |>
  filter(carat < 3) |> #filter in smaller diamonds
  ggplot(aes(x = carat)) +
  geom_histogram(binwidth = 0.01) +
  geom_vline(xintercept=c(seq(0,3,0.5)), linetype = "dashed", color="red")

#visualize distribution
ggplot(diamonds, aes(x = y)) +
  geom_histogram(binwidth = 0.5)

#visualize distribution in detail
ggplot(diamonds, aes(x = y)) +
  geom_histogram(binwidth = 0.5) +
  coord_cartesian(ylim = c(0, 50))
```

1.2. UNUSUAL VALUES

```
#look at the data
diamonds |>
  filter(between(y, 3, 20)) |> #filter out unusual values
  select(price, x, y, z) |>
  arrange(y)

#drop entire row of data
diamonds2 <- diamonds |>
  filter(between(y, 3, 20))

#replacing with missing values
diamonds2 <- diamonds |>
  mutate(y = if_else(y < 3 | y > 20, NA, y))

ggplot(diamonds2, aes(x = x, y = y)) +
  geom_point()

ggplot(diamonds2, aes(x = x, y = y)) +
  geom_point(na.rm = TRUE)

#meaningful missing values
flights |>
```

```
mutate(  
  cancelled = is.na(dep_time),  
  sched_hour = sched_dep_time %/% 100,  
  sched_min = sched_dep_time %% 100,  
  sched_dep_time = sched_hour + (sched_min / 60)  
) |>  
ggplot(aes(x = sched_dep_time)) +  
geom_density(aes(color = cancelled), bw = 1/4)
```

2. MISSING VALUES

[from <https://r4ds.hadley.nz/missing-values>]

```
library("tidyverse")
library("nycflights13")
```

2.1. EXPLICIT MISSING VALUES

```
treatment <- tribble(
  ~person,      ~treatment, ~response,
  "Derrick Whitmore", 1,      7,
  NA,              2,      10,
  NA,              3,      NA,
  "Katherine Burke", 1,      4
)
```

Last observation carried forward

```
treatment |>
  fill(everything())
```

Fixed values

```
treatment |>
  mutate(response = coalesce(response, 0)) |>
  fill(everything())
```

Numeric value represents NA

```
csv <- "
person,      treatment, response
Derrick Whitmore, 1,      7
Derrick Whitmore, 2,      10
Derrick Whitmore, 3,      99
Katherine Burke, 1,      4"
read_csv(csv, na = "99")

read_csv(csv) |>
  mutate(response = na_if(response, 99))
```

NaN

```
v <- c(NA, NaN)
v * 10
v == 1
is.na(v)
is.nan(v)

0 / 0
0 * Inf
```

```
Inf - Inf
sqrt(-1)
```

2.2. IMPLICIT MISSING VALUES

```
stocks <- tibble(
  year = c(2020, 2020, 2020, 2020, 2021, 2021, 2021),
  qtr  = c( 1,   2,   3,   4,   2,   3,   4),
  price = c(1.88, 0.59, 0.35, NA, 0.92, 0.17, 2.66)
)
#1. Price in 2020 Q4 is explicitly missing
#2. Price in 2021 Q1 is implicitly missing
```

Pivoting

```
stocks |>
  pivot_wider(
    names_from = qtr,
    values_from = price
  ) |>
  pivot_longer(
    cols = -year,
    names_to = "qtr",
    values_to = "price"
  )
```

Complete

```
stocks |>
  complete(year, qtr)

stocks |>
  complete(year = 2019:2021, qtr)
```

Joins

```
flights |>
  distinct(faa = dest) |>           #calculate the distinct "dest"
  anti_join(airports)              #obtain absent "dest"

flights |>
  distinct(tailnum) |>             #calculate the distinct "dest"
  anti_join(planes)                #obtain absent "dest"
```

2.3. FACTORS AND EMPTY GROUPS

```
health <- tibble(
  name = c("Ikaia", "Oletta", "Leriah", "Dashay", "Tresaun"),
  smoker = factor(c("no", "no", "no", "no", "no"), levels = c("yes", "no")),
  age = c(34, 88, 75, 47, 56),
)
```

```

#empty groups with count
health |> count(smoker)
health |> count(smoker, .drop = FALSE)

#empty groups with ggplot
ggplot(health, aes(x = smoker)) +
  geom_bar() +
  scale_x_discrete()
ggplot(health, aes(x = smoker)) +
  geom_bar() +
  scale_x_discrete(drop = FALSE)

#empty groups with group_by()
health |>
  group_by(smoker, .drop = FALSE) |>
  summarize(
    n = n(),
    mean_age = mean(age),
    min_age = min(age),
    max_age = max(age),
    sd_age = sd(age)
  )
health |>
  group_by(smoker) |>
  summarize(
    n = n(),
    mean_age = mean(age),
    min_age = min(age),
    max_age = max(age),
    sd_age = sd(age)
  ) |>
  complete(smoker)

```

3. COVARIATION

[from <https://r4ds.hadley.nz/eda#covariation>]

```
library("tidyverse")
library("hexbin")
```

3.1. CATEGORICAL AND NUMERICAL VARIABLES

```
#use geom_freqpoly()
ggplot(diamonds, aes(x = price, color = cut)) +
  geom_freqpoly(binwidth = 500, linewidth = 0.75)

#use geom_freqpoly() with densities
ggplot(diamonds, aes(x = price, y = after_stat(density), color = cut)) +
  geom_freqpoly(binwidth = 500, linewidth = 0.75)

#use geom_density()
ggplot(diamonds, aes(x = price, color = cut)) +
  geom_density(bw = 500, linewidth = 0.75)

#use geom_boxplot()
ggplot(diamonds, aes(x = cut, y = price)) +
  geom_boxplot()
```

3.2. TWO CATEGORICAL VARIABLES

```
#geom_count()
ggplot(diamonds, aes(x = cut, y = color)) +
  geom_count()

#geom_tile()
diamonds |>
  count(color, cut) |>
  ggplot(aes(x = color, y = cut)) +
  geom_tile(aes(fill = n))
```

3.3. TWO NUMERICAL VARIABLES

```
smaller <- diamonds |>
  filter(carat < 3)                                #filter in smaller diamonds

#geom_point()
ggplot(smaller, aes(x = carat, y = price)) +
  geom_point()

#geom_point() with alpha
ggplot(smaller, aes(x = carat, y = price)) +
  geom_point(alpha = 0.01)

#geom_bin2d()
ggplot(smaller, aes(x = carat, y = price)) +
  geom_bin2d()
```



```
#geom_hex()  
ggplot(smaller, aes(x = carat, y = price)) +  
  geom_hex()  
  
#geom_boxplot()  
ggplot(smaller, aes(x = carat, y = price)) +  
  geom_boxplot(aes(group = cut_width(carat, 0.1)))
```

4. PATTERNS AND MODELS

[from <https://r4ds.hadley.nz/eda#patterns-and-models>]

```
library("tidyverse")
library("tidymodels")

diamonds <- diamonds |>
  mutate(
    log_price = log(price),
    log_carat = log(carat)
  )

#functions linear_reg() and fit()
diamonds_fit <- linear_reg() |>
  fit(log_price ~ log_carat, data = diamonds)

#function augment()
diamonds_aug <- augment(diamonds_fit, new_data = diamonds) |>
  mutate(.resid = exp(.resid))

#unexplained variation in price using carat
ggplot(diamonds_aug, aes(x = carat, y = .resid)) +
  geom_point()

#modelling unexplained variation in price using cut
ggplot(diamonds_aug, aes(x = cut, y = .resid)) +
  geom_boxplot()
```