

Joseph Specht  
jspecht3

## HW #1

- 1) I understand plasmas to be ionized states of matter, whether that be solid, liquid, or gas. However, I have some issue with describing plasmas as another state of matter. Personally, I believe ionization percentage to be a property of the material. Even if it is another state of matter, I do not believe it should be given the same classification level as solid, liquid, & gas.

Now actually about plasmas; the most universally abundant matter we understand is plasma. Plasma is a cluster of particles where the atoms have been ionized & the electrons separate. This gives rise to a locally charged, but globally quasineutral state. This bulk matter is now subject to electromagnetic forces as it can be approximated as two fluids occupying the same space w/ equal & opposite charges.


There are abundant usages of plasma on earth, which include, but are not limited to; keeping the earth's magnetosphere w/ a plasma core, exotic/high energy chemistry, glue that can bond metals together, transistor fabrication, lightning, aurora borealis/australis, welding, neon signs, certain lights, and the list goes on. Things are different for the universe as plasmas are found as stars, nebulae, supernovae, and quasars are what come to mind.



Joseph specht  
jspecht3

HW#1 - cont

2)



$$\vec{E} = \left( \frac{kq}{l^2} \left[1 - \frac{\sqrt{2}}{2}\right], \frac{kq}{l^2} \left[3 - \frac{\sqrt{2}}{2}\right] \right) = \frac{kq}{l^2} \left( \frac{1 - \sqrt{2}}{2}, \frac{3 - \sqrt{2}}{2} \right)$$

$$|\vec{E}| = \frac{kq}{l^2} \sqrt{\left(1 - \frac{\sqrt{2}}{2}\right)^2 + \left(3 - \frac{\sqrt{2}}{2}\right)^2}$$

$$\left(1 - \frac{\sqrt{2}}{2}\right) \left(1 - \frac{\sqrt{2}}{2}\right) = 1 - \sqrt{2} + \frac{2}{4} = \frac{3 - \sqrt{2}}{2}$$

$$\left(3 - \frac{\sqrt{2}}{2}\right) \left(3 - \frac{\sqrt{2}}{2}\right) = 9 - 3\sqrt{2} + \frac{2}{4} = \frac{19 - 3\sqrt{2}}{2}$$

$$\left. \begin{array}{l} \frac{3 - \sqrt{2}}{2} \\ \frac{19 - 3\sqrt{2}}{2} \end{array} \right\} + = \frac{11 - 4\sqrt{2}}{2}$$


$$\therefore |\vec{E}| = \frac{kq}{l^2} \sqrt{11 - 4\sqrt{2}} \quad \& \quad F = q_i E_i, \text{ so}$$

$$|\vec{F}| = \frac{kq^2}{l^2} \sqrt{11 - 4\sqrt{2}} = \frac{[Nm^3/C^2] [C]^2}{[m]^2} = \frac{-9e^9 \times (1.0e^{-6})^2 \sqrt{11 - 4\sqrt{2}}}{[0.05]^2}$$

$$|\vec{F}| = 8.321 \text{ N}$$

Joseph Specht  
jspecht3

HW#1 - cont



$$a) \vec{E} = \frac{\rho r \hat{r}}{3\epsilon_0} = \frac{2.4 \times 0.005 \hat{r}}{3\epsilon_0} = 4.5198 \times 10^8 \hat{r} \text{ N/C @ } r = 0.005 \text{ m}$$

$$b) \vec{E} = \frac{\rho r \hat{r}}{3\epsilon_0} = \frac{2.4 \times 0.01 \hat{r}}{3\epsilon_0} = 9.0395 \times 10^8 \hat{r} \text{ N/C @ } r = 0.01 \text{ m}$$

$$c) \vec{E} = \frac{1}{4\pi\epsilon_0} \cdot \frac{Q}{r^2} = \frac{1}{4\pi\epsilon_0} \cdot \frac{\frac{4}{3}\pi r_0^3 \rho}{r^2} = \frac{\rho r_0^3 \hat{r}}{3\epsilon_0 r^2} = 3.6158 \times 10^7 \hat{r} \text{ N/C @ } r = 0.05 \text{ m}$$



Joseph Specht  
jspecht3

HW#1 - cont

4a) ctt will move in a circular path until the  $B$  field is gone.

b) gyro-frequency  $\omega = \frac{qB}{m}$

$$\text{where } q = 1.602 \times 10^{-19} \text{ C}$$

$$B = 1 \text{ T}$$

$$m = 9.11 \times 10^{-31} \text{ kg}$$

$$\therefore \omega = 1.7585 \times 10^{11} \text{ rad/s}$$

c) know  $F = qvB$ , but also  $F = ma$ . However,  $a$  is circular, so  $F = \frac{mv^2}{R}$ , so...

$$qvB = \frac{mv^2}{R} \Rightarrow R = \frac{mv}{qB}, \text{ but need to find } v, \text{ so use}$$

$$E = \frac{1}{2}mv^2 = k_B T \Rightarrow v^2 = \frac{2k_B T}{m} \Rightarrow v = \sqrt{\frac{2k_B T}{m}}$$

$$\therefore R = \frac{m}{qB} \sqrt{\frac{2k_B T}{m}} = \frac{1}{qB} \sqrt{2mk_B T}$$

$$\text{where } k_B = 1.38 \times 10^{-23} \frac{\text{J}}{\text{K}}$$

$$T = 100,000 \text{ K}$$

$$m = 9.109 \times 10^{-31} \text{ kg}$$

$$B = 1 \text{ T}$$

$$q = 1.602 \times 10^{-19} \text{ C}$$

$$\Rightarrow R = 9.29775 \times 10^{-6} \text{ m}$$

Joseph Specht  
jspecht3

HW#1 - cont

$$5) F = q(E + v \times B) \quad \& \quad \nabla \times B = \mu_0 J + \mu_0 \epsilon_0 \frac{\partial E}{\partial t}$$

However,  $E = 0$ , so  $F = qv \times B$  &  $\frac{\partial E}{\partial t} = 0$

$$\Rightarrow F = qv \times B \quad \& \quad \nabla \times B = \mu_0 J$$

using the relation

$$\Rightarrow \underbrace{(B \cdot \nabla) B}_2 = \nabla \left( \frac{1}{2} B^2 \right) - B \times (\nabla \times B) \quad , \text{ but } \nabla \times B = \mu_0 J, \text{ so}$$

$$\Rightarrow \frac{1}{2} \nabla B^2 = B \overset{\text{a, maxwell's law}}{\times} \nabla B + B \times (\nabla \times B) \quad ; \quad \frac{1}{2} \nabla B^2 = B \times \mu_0 J$$

$$\Rightarrow J \times B = \frac{1}{2\mu_0} \nabla B^2 = \nabla \left( \frac{B^2}{2\mu_0} \right) \quad \text{where } J \times B \text{ is Lorentz Force/unit \& is}$$

$$\Rightarrow F/A = \nabla \left( \frac{B^2}{2\mu_0} \right) \quad \Rightarrow \int F/A \, d\vec{x} = \int \nabla \frac{B^2}{2\mu_0} \, d\vec{x}$$

↑  
take a spatial integral & assume  $B$  is only function of  $x$ , integral of  $A$  is area, so...

$$\Rightarrow F/A = \frac{B^2}{2\mu_0} \rightarrow W/ \overset{P}{\parallel} \quad F/A \text{ being pressure, we get.}$$

$$p = \frac{B^2}{2\mu_0} \quad \text{no need for } \parallel \text{ as } B \cdot B \text{ is always positive}$$