

Figure 1: Major lake and watershed factors affecting lake P retention. Shaded symbols indicate factors typically considered in P retention models whereas open symbols indicate additional factors specific to the present study. Dashed lines indicate inferred relationships, which cannot be tested with available data, but are otherwise discussed herein.




Figure 2: Connectivity metric definitions along with simplified examples of high and low connectivity lakes that might arise from a binary classification. Here, high connectivity lakes with respect to both lake-stream-based and stream-based metrics are associated with lower barriers to in-stream transport. For stream-based metrics, high connectivity lakes are associated with more direct transport of P from terrestrial runoff to streams. We use the term “first order stream” to describe a headwater stream.




### A. LWS

Average  
link length

Closest  
lake distance

Stream  
density



Baseflow

Stream  
order ratio



### B. NWS

Average  
link length

Closest  
lake distance

Stream  
density



Upstream  
lake area

Baseflow

Stream  
order ratio




Figure 5: Locations of lakes of different connectivity classes at the (A) lake watershed and (B) lake network watershed extent. Low connectivity lakes are represented by green symbols, and high connectivity lakes are represented by purple symbols. NWS is the lake network watershed extent. LWS is the lake watershed extent. Connectivity metrics are defined in Figure 2.




Figure 6: Residence time (yr) versus P retention for the NES dataset and the global model fit to the data ( $R^2 = 0.34$ ,  $n = 129$ ) where the solid line and shaded interval represents the median and central 95% interval estimates respectively (A). As above except that solid lines and shaded interval estimates represent hierarchical model fits to the data ( $R^2 = 0.41$ ,  $n = 129$ ) based on network watershed average link length, which had the strongest association of any connectivity metric relative to P retention (B). Equations for these lines at median water residence time for low and high connectivity lakes are:  $R_p = 1 - (1 / (1 + 1.32 \tau^{0.4}))$  and  $R_p = 1 - (1 / (1 + 1.08 \tau^{0.4}))$  respectively.




Figure 7: Distribution of the  $k$  parameter from Eq. 1 in lakes of differing connectivity, depth, or baseflow at the (A) lake watershed and (B) lake network watershed extents. Green lines indicate lower connectivity lakes while purple lines indicate higher connectivity lakes. For maximum depth, green lines indicate deeper lakes. For baseflow, green lines indicate higher baseflow. Connectivity metrics are defined in Figure 2. Labels associated with models where differences in  $k$  translated to significant differences in P retention are bolded and starred.