




Figure 8: Sensitivity analysis for the Bayesian model-averaged meta-analysis applied to the Bem (2011) data: reduced appendix display of the supplementary density estimators. Rows correspond to the two encompassing strategies (top: bridge sampling; bottom: product-space). Left column: inclusion Bayes factor BF_{incl} as a function of the prior standard deviation for the overall effect, σ_μ ; colored lines show logspline density estimation, the log-concave maximum likelihood estimator, the truncated normal approximation, the truncated asymmetric power distribution (APD), and the Bayesian nonparametric mixture estimator from BNPmix. Right column: approximation ratio $BF_{\text{approx}}/BF_{\text{exact}}$ at the validation points. Filled circles mark the anchor at $\sigma_\mu = 1$; crosses show validation refits of the full Bayesian model-averaged meta-analysis obtained by bridge sampling.




Figure 9: Sensitivity analysis for the robust Bayesian meta-analysis applied to the Bem (2011) data: reduced appendix display of the supplementary density estimators. Left panel: inclusion Bayes factor BF_{incl} as a function of the prior standard deviation for the overall effect, σ_μ ; colored lines show logspline density estimation, the log-concave maximum likelihood estimator, the truncated normal approximation, the truncated asymmetric power distribution (APD), and the Bayesian nonparametric mixture estimator from BNPmix. Right panel: approximation ratio $BF_{\text{approx}}/BF_{\text{exact}}$ at the validation points. The product-space approach is used throughout; the filled circle marks the anchor at $\sigma_\mu = 1$, and crosses show validation refits of the full 36-model robust Bayesian meta-analysis obtained by bridge sampling.