Conjectures that should be true*

May 3, 2016

1 Edges in list-critical graphs

A graph G is k-list-critical if G is not (k-1)-choosable, but every proper subgraph of G is (k-1)-choosable. Replace '(k-1)-' with 'online (k-1)-' and 'k-' with 'online k-' in the previous sentence and read it.

Conjecture 1. Every incomplete k-list-critical graph has average degree at least

$$k - 1 + \frac{k - 3}{(k - 1)^2}.$$

Background. The connected graphs in which each block is a complete graph or an odd cycle are called *Gallai trees*. Gallai [7] proved that in a k-critical graph, the vertices of degree k-1 induce a disjoint union of Gallai trees. The same is true for k-list-critical graphs [1, 6]. This quickly implies a lower bound on the average degree of k-list-critical graphs of

$$k - 1 + \frac{k - 3}{k^2 - 3}.$$

In [14], R. improved this to

$$k - 1 + \frac{k - 3}{k^2 - 2k + 2}$$

using a lemma from Kierstead and R. [9] that generalizes a kernel technique of Kostochka and Yancey [10]. As noted at the end of [14], a small improvement to the argument would yield Conjecture 1. \Box

Conjecture 2. Every incomplete online k-list-critical graph G has

$$2 ||G|| \ge (k-1) |G| + k - 3.$$

Background. Dirac [5] proved this for k-critical graphs. Kostochka and Stiebitz [11] proved it for k-list-critical graphs. Their proof does not seem to generalize. When |G| is large compared with k, the conjecture holds by Gallai-type bounds on the average degree of online k-list-critical graphs [8, 2].

 $^{*(}dis)proofs \Rightarrow landon.rabern@gmail.com$

2 Maximum degree, clique number and colorings

2.1 Around Borodin-Kostochka

Conjecture 3. Every graph with $\chi \geq \Delta \geq 8$ contains a $K_3 \vee H$ where H is some graph on $\Delta - 3$ vertices.

Background. By results in [4], for $\Delta \geq 9$ the existence of $K_3 \vee H$ implies the existence of K_Δ . So, this (seemingly weaker) conjecture for $\Delta \geq 9$ implies the Borodin-Kostochka conjecture. The one known connected counterexample to the Borodin-Kostochka conjecture for $\Delta = 8$ is a 5-cycle with each vertex blown up to a triangle. This graph is not a counterexample to Conjecture 3.

Conjecture 4. Every graph with $\chi \geq \Delta$ contains $K_{\Delta-3}$.

Background. Results in [3] show that this holds with $K_{\Delta-4}$ instead of $K_{\Delta-3}$. Moreover, [3] proves the conjecture for all but $\Delta \in \{6, 8, 9, 11, 12\}$.

Conjecture 5. Every graph with $\chi \geq \Delta$ either contains K_{Δ} or contains a $K_{\Delta-4}$ with all Δ -vertices.

Background. Results in [3] show that this holds with $K_{\Delta-5}$ instead of $K_{\Delta-4}$. For $\Delta \leq 7$, the conjecture holds by [13, 12]. Also by [3], it holds when $\Delta = 3r + 1$ for $r \geq 3$.

Conjecture 6. Every graph with $\Delta \geq 8$ and $\omega < \Delta$ is 2-fold $(2\Delta - 1)$ -colorable.

Background. The one known connected counterexample to the Borodin-Kostochka conjecture for $\Delta = 8$ is a 5-cycle with each vertex blown up to a triangle. This graph is not a counterexample to Conjecture 6.

Conjecture 7. Every graph with $\theta \geq 10$ and $\omega \leq \frac{\theta}{2}$ is $\lfloor \frac{\theta}{2} \rfloor$ -choosable.

Background.

Conjecture 8. Every claw-free graph with $\Delta \geq 9$ and $\omega < \Delta$ is $(\Delta - 1)$ -choosable.

Background.

Conjecture 9. There is a polynomial time graph algorithm that finds either a $(\Delta - 1)$ coloring or a $K_{\Delta-3}$.

Background.

2.2 The $\frac{5}{6}$ bound

Conjecture 10. If G is vertex-transitive, then $\chi(G) \leq \max \left\{ \omega(G), \left\lceil \frac{5\Delta(G)+3}{6} \right\rceil \right\}$.

Conjecture 11. If G is the line graph of a multigraph, then $\chi(G) \leq \max \left\{ \omega(G), \left\lceil \frac{5\Delta(G)+3}{6} \right\rceil \right\}$.

Background.

3 Planar graphs

Conjecture 12. Every planar graph with no K_5 -subdivision is 2-fold 9-colorable. Background.

References

[1] O.V. Borodin, Criterion of chromaticity of a degree prescription, Abstracts of IV All-Union Conf. on Th. Cybernetics, 1977, pp. 127–128.

- [2] D. Cranston and L. Rabern, Edge lower bounds for list critical graphs, via discharging, arXiv:1602.02589 (2016).
- [3] Daniel W. Cranston and Landon Rabern, Graphs with $\chi = \Delta$ Have Big Cliques, SIAM Journal on Discrete Mathematics 29 (2015), no. 4, 1792–1814.
- [4] D.W. Cranston and L. Rabern, Conjectures equivalent to the Borodin-Kostochka Conjecture that appear weaker, Arxiv preprint arXiv:1203.5380 (2012).
- [5] G.A. Dirac, A theorem of R.L. Brooks and a conjecture of H. Hadwiger, Proceedings of the London Mathematical Society 3 (1957), no. 1, 161–195.
- [6] P. Erdős, A.L. Rubin, and H. Taylor, *Choosability in graphs*, Proc. West Coast Conf. on Combinatorics, Graph Theory and Computing, Congressus Numerantium, vol. 26, 1979, pp. 125–157.
- [7] T. Gallai, Kritische Graphen I., Publ. Math. Inst. Hungar. Acad. Sci 8 (1963), 165–192 (in German).
- [8] H.A. Kierstead and L. Rabern, Improved lower bounds on the number of edges in list critical and online list critical graphs, arXiv preprint arXiv:1406.7355 (2014).
- [9] _____, Extracting list colorings from large independent sets, arXiv:1512.08130 (2015).
- [10] Alexandr Kostochka and Matthew Yancey, Ore's conjecture on color-critical graphs is almost true, J. Combin. Theory Ser. B 109 (2014), 73–101. MR 3269903
- [11] Alexandr V Kostochka and Michael Stiebitz, A list version of dirac's theorem on the number of edges in colour-critical graphs, Journal of Graph Theory 39 (2002), no. 3, 165-177.
- [12] A.V. Kostochka, L. Rabern, and M. Stiebitz, Graphs with chromatic number close to maximum degree, Discrete Mathematics 312 (2012), no. 6, 1273–1281.
- [13] L. Rabern, Δ -critical graphs with small high vertex cliques, Journal of Combinatorial Theory, Series B **102** (2012), no. 1, 126–130.
- [14] Landon Rabern, A better lower bound on average degree of 4-list-critical graphs, arXiv:1602.08532 (2016).