

# Hitting maximum cliques

Landon Rabern

landon.rabern@gmail.com

July 7, 2011

## ① Introduction

What's it good for?

An outline of the proof

## ② The common lemmas

Hajnal's clique collection lemma

Kostochka's lemma

## ③ Proving the main results

Independent transversals

Putting it all together

Improving the constant

Introduction

What's it good  
for?

An outline of the  
proof

The common  
lemmas

Proving the  
main results

# Introduction

Finding a stable set hitting every maximum clique in a graph can be very useful for coloring problems.

# Introduction

Finding a stable set hitting every maximum clique in a graph can be very useful for coloring problems.

## Observation

A graph is perfect iff every induced subgraph has a stable set hitting every maximum clique.

# Introduction

## Introduction

What's it good for?

An outline of the proof

The common lemmas

Proving the main results

Finding a stable set hitting every maximum clique in a graph can be very useful for coloring problems.

## Observation

A graph is perfect iff every induced subgraph has a stable set hitting every maximum clique.

Kostochka [8] gave the following sufficient condition.

## Lemma (Kostochka 1980)

*A graph satisfying  $\omega \geq \Delta + \frac{3}{2} - \sqrt{\Delta}$  has a stable set hitting every maximum clique.*

## Introduction

What's it good  
for?

An outline of the  
proof

The common  
lemmas

Proving the  
main results

In [10], we improved this as follows.

## Lemma (Rabern 2009)

*There exists a positive constant  $c < 1$  such that every graph satisfying  $\omega > c(\Delta + 1)$  has a stable set hitting every maximum clique.*

# What's it good for?

- removing a stable set which hits every maximum clique decreases  $\omega$

Introduction

What's it good for?

An outline of the proof

The common lemmas

Proving the main results

## What's it good for?

- removing a stable set which hits every maximum clique decreases  $\omega$
- say you are trying to prove Brooks' theorem and start by taking a counterexample minimizing  $\Delta$



## What's it good for?

- removing a stable set which hits every maximum clique decreases  $\omega$
- say you are trying to prove Brooks' theorem and start by taking a counterexample minimizing  $\Delta$
- suppose  $\Delta \geq 4$

## What's it good for?

- removing a stable set which hits every maximum clique decreases  $\omega$
- say you are trying to prove Brooks' theorem and start by taking a counterexample minimizing  $\Delta$
- suppose  $\Delta \geq 4$
- then  $\omega = 4$ , hence by the fact that  $c = \frac{3}{4}$  works in the above lemma, we have a stable set hitting every maximum clique

## What's it good for?

- removing a stable set which hits every maximum clique decreases  $\omega$
- say you are trying to prove Brooks' theorem and start by taking a counterexample minimizing  $\Delta$
- suppose  $\Delta \geq 4$
- then  $\omega = 4$ , hence by the fact that  $c = \frac{3}{4}$  works in the above lemma, we have a stable set hitting every maximum clique
- expanding it to a maximal stable set and removing it leaves a graph with  $\omega \leq 3$ ,  $\chi = 4$  and  $\Delta \leq 3$ , contradicting minimality of  $\Delta$

## What's it good for?

- removing a stable set which hits every maximum clique decreases  $\omega$
- say you are trying to prove Brooks' theorem and start by taking a counterexample minimizing  $\Delta$
- suppose  $\Delta \geq 4$
- then  $\omega = 4$ , hence by the fact that  $c = \frac{3}{4}$  works in the above lemma, we have a stable set hitting every maximum clique
- expanding it to a maximal stable set and removing it leaves a graph with  $\omega \leq 3$ ,  $\chi = 4$  and  $\Delta \leq 3$ , contradicting minimality of  $\Delta$
- thus a counterexample to Brooks' theorem minimizing  $\Delta$  must have  $\Delta = 3$

# What's it good for?

## Introduction

What's it good  
for?

An outline of the  
proof

The common  
lemmas

Proving the  
main results

- Borodin and Kostochka [3] conjecture that graphs with  $\Delta \geq 9$  containing no  $K_\Delta$  are  $(\Delta - 1)$ -colorable.

## What's it good for?

### Introduction

#### What's it good for?

An outline of the proof

#### The common lemmas

#### Proving the main results

- Borodin and Kostochka [3] conjecture that graphs with  $\Delta \geq 9$  containing no  $K_\Delta$  are  $(\Delta - 1)$ -colorable.
- as shown by Kostochka, almost identically to the case of Brooks' theorem, we may assume that  $\Delta = 9$

## What's it good for?

### Introduction

#### What's it good for?

An outline of the proof

#### The common lemmas

#### Proving the main results

- Borodin and Kostochka [3] conjecture that graphs with  $\Delta \geq 9$  containing no  $K_\Delta$  are  $(\Delta - 1)$ -colorable.
- as shown by Kostochka, almost identically to the case of Brooks' theorem, we may assume that  $\Delta = 9$
- Reed conjectures that every graph satisfies  $\chi \leq \left\lceil \frac{\omega + \Delta + 1}{2} \right\rceil$

# What's it good for?

## Introduction

### What's it good for?

An outline of the proof

### The common lemmas

### Proving the main results

- Borodin and Kostochka [3] conjecture that graphs with  $\Delta \geq 9$  containing no  $K_\Delta$  are  $(\Delta - 1)$ -colorable.
- as shown by Kostochka, almost identically to the case of Brooks' theorem, we may assume that  $\Delta = 9$
- Reed conjectures that every graph satisfies  $\chi \leq \left\lceil \frac{\omega + \Delta + 1}{2} \right\rceil$
- when proving this conjecture for a hereditary class of graphs, a minimum counterexample must have  $\omega \leq \frac{3}{4}(\Delta + 1)$



# What's it good for?

## Introduction

### What's it good for?

An outline of the proof

### The common lemmas

### Proving the main results

- Borodin and Kostochka [3] conjecture that graphs with  $\Delta \geq 9$  containing no  $K_\Delta$  are  $(\Delta - 1)$ -colorable.
- as shown by Kostochka, almost identically to the case of Brooks' theorem, we may assume that  $\Delta = 9$
- Reed conjectures that every graph satisfies  $\chi \leq \left\lceil \frac{\omega + \Delta + 1}{2} \right\rceil$
- when proving this conjecture for a hereditary class of graphs, a minimum counterexample must have  $\omega \leq \frac{3}{4}(\Delta + 1)$
- this was used in [10] to simplify the proof of Reed's conjecture for line graphs given in [7]

Hitting  
maximum  
cliques

Landon  
Rabern

Introduction

What's it good  
for?

An outline of the  
proof

The common  
lemmas

Proving the  
main results

# An outline of the proof

PICTURE

# An outline of the proof

## PICTURE

- use lemmas of Hajnal and Kostochka to show that each component of the maximum clique graph has many universal vertices

# An outline of the proof

## Introduction

What's it good  
for?

An outline of the  
proof

The common  
lemmas

Proving the  
main results

## PICTURE

- use lemmas of Hajnal and Kostochka to show that each component of the maximum clique graph has many universal vertices
- consider the subgraph induced on these universal vertices and partition it by component

# An outline of the proof

## Introduction

What's it good  
for?

An outline of the  
proof

The common  
lemmas

Proving the  
main results

## PICTURE

- use lemmas of Hajnal and Kostochka to show that each component of the maximum clique graph has many universal vertices
- consider the subgraph induced on these universal vertices and partition it by component
- find an independent transversal

# An outline of the proof

## Introduction

What's it good  
for?

An outline of the  
proof

The common  
lemmas

Proving the  
main results

## PICTURE

- use lemmas of Hajnal and Kostochka to show that each component of the maximum clique graph has many universal vertices
- consider the subgraph induced on these universal vertices and partition it by component
- find an independent transversal
- this is our desired stable set hitting all maximum cliques

# Hajnal's clique collection lemma

## Lemma (Hajnal 1965)

*For a collection  $\mathcal{Q}$  of maximum cliques in a graph  $G$  we have*

$$|\bigcup \mathcal{Q}| + |\bigcap \mathcal{Q}| \geq 2\omega(G).$$

# Hajnal's clique collection lemma

## Lemma (Hajnal 1965)

*For a collection  $\mathcal{Q}$  of maximum cliques in a graph  $G$  we have*

$$|\bigcup \mathcal{Q}| + |\bigcap \mathcal{Q}| \geq 2\omega(G).$$

- let  $\mathcal{Q}$  be a counterexample with  $|\mathcal{Q}|$  minimal



# Hajnal's clique collection lemma

## Lemma (Hajnal 1965)

*For a collection  $\mathcal{Q}$  of maximum cliques in a graph  $G$  we have*

$$|\bigcup \mathcal{Q}| + |\bigcap \mathcal{Q}| \geq 2\omega(G).$$

- let  $\mathcal{Q}$  be a counterexample with  $|\mathcal{Q}|$  minimal
- put  $r := |\mathcal{Q}|$ , say  $\mathcal{Q} = \{Q_1, \dots, Q_r\}$

# Hajnal's clique collection lemma

## Lemma (Hajnal 1965)

*For a collection  $\mathcal{Q}$  of maximum cliques in a graph  $G$  we have*

$$|\bigcup \mathcal{Q}| + |\bigcap \mathcal{Q}| \geq 2\omega(G).$$

- let  $\mathcal{Q}$  be a counterexample with  $|\mathcal{Q}|$  minimal
- put  $r := |\mathcal{Q}|$ , say  $\mathcal{Q} = \{Q_1, \dots, Q_r\}$
- consider  $W := (Q_1 \cap \bigcup_{i=2}^r Q_i) \cup \bigcap_{i=2}^r Q_i$

# Hajnal's clique collection lemma

## Lemma (Hajnal 1965)

*For a collection  $\mathcal{Q}$  of maximum cliques in a graph  $G$  we have*

$$|\bigcup \mathcal{Q}| + |\bigcap \mathcal{Q}| \geq 2\omega(G).$$

- let  $\mathcal{Q}$  be a counterexample with  $|\mathcal{Q}|$  minimal
- put  $r := |\mathcal{Q}|$ , say  $\mathcal{Q} = \{Q_1, \dots, Q_r\}$
- consider  $W := (Q_1 \cap \bigcup_{i=2}^r Q_i) \cup \bigcap_{i=2}^r Q_i$
- $W$  is a clique, so the following machinations give a contradiction

$$\begin{aligned}
 \omega(G) &\geq |W| \\
 &= |(Q_1 \cap \bigcup_{i=2}^r Q_i) \cup \bigcap_{i=2}^r Q_i| \\
 &= |Q_1 \cap \bigcup_{i=2}^r Q_i| + |\bigcap_{i=2}^r Q_i| - |\bigcap_{i=1}^r Q_i \cap \bigcup_{i=2}^r Q_i| \\
 &= |Q_1| + |\bigcup_{i=2}^r Q_i| - |\bigcup_{i=1}^r Q_i| + |\bigcap_{i=2}^r Q_i| - |\bigcap_{i=1}^r Q_i| \\
 &= \omega(G) + |\bigcup_{i=2}^r Q_i| + |\bigcap_{i=2}^r Q_i| - |\bigcup_{i=1}^r Q_i| - |\bigcap_{i=1}^r Q_i| \\
 &\geq \omega(G) + 2\omega(G) - (|\bigcup_{i=1}^r Q_i| + |\bigcap_{i=1}^r Q_i|) \\
 &> \omega(G).
 \end{aligned}$$

# The clique graph

## Introduction

## The common lemmas

Hajnal's clique  
collection lemma  
Kostochka's  
lemma

## Proving the main results

In [9] we gave the following simple proof of Kostochka's lemma from [8]. First a definition.

# The clique graph

Introduction

The common  
lemmas

Hajnal's clique  
collection lemma  
Kostochka's  
lemma

Proving the  
main results

In [9] we gave the following simple proof of Kostochka's lemma from [8]. First a definition.

## Clique graph

For a collection of cliques  $\mathcal{Q}$  in a graph, let  $X_{\mathcal{Q}}$  be the intersection graph of  $\mathcal{Q}$ ; that is, the vertex set of  $X_{\mathcal{Q}}$  is  $\mathcal{Q}$  and there is an edge between  $Q_1 \neq Q_2 \in \mathcal{Q}$  iff  $Q_1$  and  $Q_2$  intersect.

# Kostochka's lemma

## Lemma (Kostochka 1980)

*Let  $G$  be a graph satisfying  $\omega > \frac{2}{3}(\Delta + 1)$ . If  $\mathcal{Q}$  is a collection of maximum cliques in  $G$  such that  $X_{\mathcal{Q}}$  is connected, then  $\cap \mathcal{Q} \neq \emptyset$ .*

# Kostochka's lemma

## Lemma (Kostochka 1980)

*Let  $G$  be a graph satisfying  $\omega > \frac{2}{3}(\Delta + 1)$ . If  $\mathcal{Q}$  is a collection of maximum cliques in  $G$  such that  $X_{\mathcal{Q}}$  is connected, then  $\cap \mathcal{Q} \neq \emptyset$ .*

- let  $\mathcal{Q}$  be a counterexample with  $|\mathcal{Q}|$  minimal



# Kostochka's lemma

## Lemma (Kostochka 1980)

*Let  $G$  be a graph satisfying  $\omega > \frac{2}{3}(\Delta + 1)$ . If  $\mathcal{Q}$  is a collection of maximum cliques in  $G$  such that  $X_{\mathcal{Q}}$  is connected, then  $\cap \mathcal{Q} \neq \emptyset$ .*

- let  $\mathcal{Q}$  be a counterexample with  $|\mathcal{Q}|$  minimal
- let  $A \in \mathcal{Q}$  be a non-cutvertex in  $X_{\mathcal{Q}}$  and  $B$  a neighbor of  $A$

# Kostochka's lemma

## Lemma (Kostochka 1980)

*Let  $G$  be a graph satisfying  $\omega > \frac{2}{3}(\Delta + 1)$ . If  $\mathcal{Q}$  is a collection of maximum cliques in  $G$  such that  $X_{\mathcal{Q}}$  is connected, then  $\cap \mathcal{Q} \neq \emptyset$ .*

- let  $\mathcal{Q}$  be a counterexample with  $|\mathcal{Q}|$  minimal
- let  $A \in \mathcal{Q}$  be a non-cutvertex in  $X_{\mathcal{Q}}$  and  $B$  a neighbor of  $A$
- put  $\mathcal{Z} := \mathcal{Q} - \{A\}$

# Kostochka's lemma

## Lemma (Kostochka 1980)

*Let  $G$  be a graph satisfying  $\omega > \frac{2}{3}(\Delta + 1)$ . If  $\mathcal{Q}$  is a collection of maximum cliques in  $G$  such that  $X_{\mathcal{Q}}$  is connected, then  $\cap \mathcal{Q} \neq \emptyset$ .*

- let  $\mathcal{Q}$  be a counterexample with  $|\mathcal{Q}|$  minimal
- let  $A \in \mathcal{Q}$  be a non-cutvertex in  $X_{\mathcal{Q}}$  and  $B$  a neighbor of  $A$
- put  $\mathcal{Z} := \mathcal{Q} - \{A\}$
- then  $X_{\mathcal{Z}}$  is connected and hence by minimality of  $|\mathcal{Q}|$ ,  
 $\cap \mathcal{Z} \neq \emptyset$

# Kostochka's lemma

## Lemma (Kostochka 1980)

*Let  $G$  be a graph satisfying  $\omega > \frac{2}{3}(\Delta + 1)$ . If  $\mathcal{Q}$  is a collection of maximum cliques in  $G$  such that  $X_{\mathcal{Q}}$  is connected, then  $\cap \mathcal{Q} \neq \emptyset$ .*

- let  $\mathcal{Q}$  be a counterexample with  $|\mathcal{Q}|$  minimal
- let  $A \in \mathcal{Q}$  be a non-cutvertex in  $X_{\mathcal{Q}}$  and  $B$  a neighbor of  $A$
- put  $\mathcal{Z} := \mathcal{Q} - \{A\}$
- then  $X_{\mathcal{Z}}$  is connected and hence by minimality of  $|\mathcal{Q}|$ ,  $\cap \mathcal{Z} \neq \emptyset$
- in particular  $|\cup \mathcal{Z}| \leq \Delta(G) + 1$

# Kostochka's lemma

## Lemma (Kostochka 1980)

*Let  $G$  be a graph satisfying  $\omega > \frac{2}{3}(\Delta + 1)$ . If  $\mathcal{Q}$  is a collection of maximum cliques in  $G$  such that  $X_{\mathcal{Q}}$  is connected, then  $\cap \mathcal{Q} \neq \emptyset$ .*

- let  $\mathcal{Q}$  be a counterexample with  $|\mathcal{Q}|$  minimal
- let  $A \in \mathcal{Q}$  be a non-cutvertex in  $X_{\mathcal{Q}}$  and  $B$  a neighbor of  $A$
- put  $\mathcal{Z} := \mathcal{Q} - \{A\}$
- then  $X_{\mathcal{Z}}$  is connected and hence by minimality of  $|\mathcal{Q}|$ ,  $\cap \mathcal{Z} \neq \emptyset$
- in particular  $|\cup \mathcal{Z}| \leq \Delta(G) + 1$
- thus  $|\cup \mathcal{Q}| \leq |A - B| + |\cup \mathcal{Z}| \leq 2(\Delta(G) + 1) - \omega(G) < 2\omega(G)$   
contradicting Hajnal

# Independent transversals

## Definition

An *independent transversal* of a partition  $\{V_1, \dots, V_r\}$  of a the vertex set of a graph  $G$  is a stable set  $\{v_1, \dots, v_r\} \subseteq V(G)$  such that  $v_i \in V_i$  for each  $i$ .

# Independent transversals

Introduction

The common  
lemmas

Proving the  
main results

Independent  
transversals

Putting it all  
together

Improving the  
constant

## Definition

An *independent transversal* of a partition  $\{V_1, \dots, V_r\}$  of a the vertex set of a graph  $G$  is a stable set  $\{v_1, \dots, v_r\} \subseteq V(G)$  such that  $v_i \in V_i$  for each  $i$ .

Alon [2] proved a simple sufficient condition probabilistically.

# Independent transversals

## Definition

An *independent transversal* of a partition  $\{V_1, \dots, V_r\}$  of a the vertex set of a graph  $G$  is a stable set  $\{v_1, \dots, v_r\} \subseteq V(G)$  such that  $v_i \in V_i$  for each  $i$ .

Alon [2] proved a simple sufficient condition probabilistically.

## Lemma (Alon 1988)

A partition  $\{V_1, \dots, V_r\}$  of the vertex set of a graph  $G$  has an *independent transversal* if  $|V_i| \geq 2e\Delta(G)$  for each  $i$ .



# Alon's proof

- put  $\Delta = \Delta(G)$

Introduction

The common  
lemmas

Proving the  
main results

**Independent  
transversals**

Putting it all  
together

Improving the  
constant

## Alon's proof

Introduction

The common  
lemmas

Proving the  
main results

Independent  
transversals

Putting it all  
together

Improving the  
constant

- put  $\Delta = \Delta(G)$
- without loss of generality we may suppose that  $|V_i| = k \geq 2e\Delta$  for each  $i$

## Alon's proof

Introduction

The common  
lemmas

Proving the  
main results

Independent  
transversals

Putting it all  
together

Improving the  
constant

- put  $\Delta = \Delta(G)$
- without loss of generality we may suppose that  $|V_i| = k \geq 2e\Delta$  for each  $i$
- randomly select one vertex from each  $V_i$

## Alon's proof

Introduction

The common  
lemmas

Proving the  
main results

Independent  
transversals

Putting it all  
together

Improving the  
constant

- put  $\Delta = \Delta(G)$
- without loss of generality we may suppose that  $|V_i| = k \geq 2e\Delta$  for each  $i$
- randomly select one vertex from each  $V_i$
- for each edge  $e$  of  $G$ , let  $B_e$  be the event that both ends of  $e$  get selected

## Alon's proof

Introduction

The common  
lemmas

Proving the  
main results

Independent  
transversals

Putting it all  
together

Improving the  
constant

- put  $\Delta = \Delta(G)$
- without loss of generality we may suppose that  $|V_i| = k \geq 2e\Delta$  for each  $i$
- randomly select one vertex from each  $V_i$
- for each edge  $e$  of  $G$ , let  $B_e$  be the event that both ends of  $e$  get selected
- $\mathcal{P}(B_e) \leq k^{-2}$

## Alon's proof

Introduction

The common  
lemmas

Proving the  
main results

Independent  
transversals

Putting it all  
together

Improving the  
constant

- put  $\Delta = \Delta(G)$
- without loss of generality we may suppose that  $|V_i| = k \geq 2e\Delta$  for each  $i$
- randomly select one vertex from each  $V_i$
- for each edge  $e$  of  $G$ , let  $B_e$  be the event that both ends of  $e$  get selected
- $\mathcal{P}(B_e) \leq k^{-2}$
- each  $B_e$  is independent of all but at most  $d := 2k(\Delta - 1) < 2k\Delta - 1 \leq k^2e^{-1} - 1$  other events

# Alon's proof

## Introduction

## The common lemmas

## Proving the main results

## Independent transversals

## Putting it all together Improving the constant

- put  $\Delta = \Delta(G)$
- without loss of generality we may suppose that  $|V_i| = k \geq 2e\Delta$  for each  $i$
- randomly select one vertex from each  $V_i$
- for each edge  $e$  of  $G$ , let  $B_e$  be the event that both ends of  $e$  get selected
- $\mathcal{P}(B_e) \leq k^{-2}$
- each  $B_e$  is independent of all but at most  $d := 2k(\Delta - 1) < 2k\Delta - 1 \leq k^2e^{-1} - 1$  other events
- since  $e\mathcal{P}(B_e)(d + 1) < 1$ , the Lovász Local Lemma implies that the probability that none of the  $B_e$  occur is positive

- put  $\Delta = \Delta(G)$
- without loss of generality we may suppose that  $|V_i| = k \geq 2e\Delta$  for each  $i$
- randomly select one vertex from each  $V_i$
- for each edge  $e$  of  $G$ , let  $B_e$  be the event that both ends of  $e$  get selected
- $\mathcal{P}(B_e) \leq k^{-2}$
- each  $B_e$  is independent of all but at most  $d := 2k(\Delta - 1) < 2k\Delta - 1 \leq k^2e^{-1} - 1$  other events
- since  $e\mathcal{P}(B_e)(d + 1) < 1$ , the Lovász Local Lemma implies that the probability that none of the  $B_e$  occur is positive
- hence an independent transversal exists



# Putting it all together

## Lemma (Rabern 2009)

*A graph satisfying  $\omega \geq \frac{12}{13}(\Delta + 1)$  has a stable set hitting every maximum clique.*

# Putting it all together

## Lemma (Rabern 2009)

*A graph satisfying  $\omega \geq \frac{12}{13}(\Delta + 1)$  has a stable set hitting every maximum clique.*

- let  $G$  be a graph satisfying  $\omega \geq \frac{12}{13}(\Delta + 1)$

# Putting it all together

## Lemma (Rabern 2009)

*A graph satisfying  $\omega \geq \frac{12}{13}(\Delta + 1)$  has a stable set hitting every maximum clique.*

- let  $G$  be a graph satisfying  $\omega \geq \frac{12}{13}(\Delta + 1)$
- let  $\mathcal{Q}$  be the maximum cliques in  $G$  and  $Q_1, \dots, Q_t$  the vertex sets of the components of  $X_{\mathcal{Q}}$

# Putting it all together

## Lemma (Rabern 2009)

*A graph satisfying  $\omega \geq \frac{12}{13}(\Delta + 1)$  has a stable set hitting every maximum clique.*

- let  $G$  be a graph satisfying  $\omega \geq \frac{12}{13}(\Delta + 1)$
- let  $\mathcal{Q}$  be the maximum cliques in  $G$  and  $Q_1, \dots, Q_t$  the vertex sets of the components of  $X_{\mathcal{Q}}$
- put  $K_i = \cap Q_i$

# Putting it all together

## Lemma (Rabern 2009)

*A graph satisfying  $\omega \geq \frac{12}{13}(\Delta + 1)$  has a stable set hitting every maximum clique.*

- let  $G$  be a graph satisfying  $\omega \geq \frac{12}{13}(\Delta + 1)$
- let  $\mathcal{Q}$  be the maximum cliques in  $G$  and  $Q_1, \dots, Q_t$  the vertex sets of the components of  $X_{\mathcal{Q}}$
- put  $K_i = \cap Q_i$
- by Kostochka's lemma,  $K_i \neq \emptyset$  for each  $i$

# Putting it all together

## Lemma (Rabern 2009)

*A graph satisfying  $\omega \geq \frac{12}{13}(\Delta + 1)$  has a stable set hitting every maximum clique.*

- let  $G$  be a graph satisfying  $\omega \geq \frac{12}{13}(\Delta + 1)$
- let  $\mathcal{Q}$  be the maximum cliques in  $G$  and  $Q_1, \dots, Q_t$  the vertex sets of the components of  $X_{\mathcal{Q}}$
- put  $K_i = \cap Q_i$
- by Kostochka's lemma,  $K_i \neq \emptyset$  for each  $i$
- in particular,  $|\cup Q_i| \leq \Delta(G) + 1$

# Putting it all together

- put  $k := \min_i |K_i|$

# Putting it all together

Introduction

The common  
lemmas

Proving the  
main results

Independent  
transversals

**Putting it all  
together**

Improving the  
constant

- put  $k := \min_i |K_i|$
- by Hajnal's lemma,  $k \geq 2\omega(G) - (\Delta(G) + 1) \geq \frac{11}{13}(\Delta + 1)$



## Putting it all together

Introduction

The common  
lemmas

Proving the  
main results

Independent  
transversals

Putting it all  
together

Improving the  
constant

- put  $k := \min_i |K_i|$
- by Hajnal's lemma,  $k \geq 2\omega(G) - (\Delta(G) + 1) \geq \frac{11}{13}(\Delta + 1)$
- consider the graph  $H$  with vertex set  $\cup_i K_i$  and edge set  $\{xy \in E(G) \mid x \in K_i, y \in K_j, i \neq j\}$

## Putting it all together

- put  $k := \min_i |K_i|$
- by Hajnal's lemma,  $k \geq 2\omega(G) - (\Delta(G) + 1) \geq \frac{11}{13}(\Delta + 1)$
- consider the graph  $H$  with vertex set  $\cup_i K_i$  and edge set  $\{xy \in E(G) \mid x \in K_i, y \in K_j, i \neq j\}$
- $\Delta(H) \leq \Delta(G) + 1 - k \leq \frac{2}{13}(\Delta(G) + 1)$

## Putting it all together

- put  $k := \min_i |K_i|$
- by Hajnal's lemma,  $k \geq 2\omega(G) - (\Delta(G) + 1) \geq \frac{11}{13}(\Delta + 1)$
- consider the graph  $H$  with vertex set  $\cup_i K_i$  and edge set  $\{xy \in E(G) \mid x \in K_i, y \in K_j, i \neq j\}$
- $\Delta(H) \leq \Delta(G) + 1 - k \leq \frac{2}{13}(\Delta(G) + 1)$
- hence  $k \geq \frac{11}{13}(\Delta(G) + 1) \geq \frac{4e}{13}(\Delta(G) + 1) \geq 2e\Delta(H)$

# Putting it all together

- put  $k := \min_i |K_i|$
- by Hajnal's lemma,  $k \geq 2\omega(G) - (\Delta(G) + 1) \geq \frac{11}{13}(\Delta + 1)$
- consider the graph  $H$  with vertex set  $\cup_i K_i$  and edge set  $\{xy \in E(G) \mid x \in K_i, y \in K_j, i \neq j\}$
- $\Delta(H) \leq \Delta(G) + 1 - k \leq \frac{2}{13}(\Delta(G) + 1)$
- hence  $k \geq \frac{11}{13}(\Delta(G) + 1) \geq \frac{4e}{13}(\Delta(G) + 1) \geq 2e\Delta(H)$
- by Alon's lemma, we have an independent transversal through the  $K_i$  and this is the desired stable set hitting every maximum clique

# Improving the constant

In [10] we proved that  $c = \frac{3}{4}$  works in the same way as above using the following lemma of Haxell [5].

# Improving the constant

In [10] we proved that  $c = \frac{3}{4}$  works in the same way as above using the following lemma of Haxell [5].

## Lemma (Haxell 2001)

*A partition  $\{V_1, \dots, V_r\}$  of the vertex set of a graph  $G$  has an independent transversal if  $|V_i| \geq 2\Delta(G)$  for each  $i$ .*

## Improving the constant

In [10] we proved that  $c = \frac{3}{4}$  works in the same way as above using the following lemma of Haxell [5].

### Lemma (Haxell 2001)

*A partition  $\{V_1, \dots, V_r\}$  of the vertex set of a graph  $G$  has an independent transversal if  $|V_i| \geq 2\Delta(G)$  for each  $i$ .*

- Haxell's proof is elementary and uses some somewhat delicate induction

# Improving the constant

In [10] we proved that  $c = \frac{3}{4}$  works in the same way as above using the following lemma of Haxell [5].

## Lemma (Haxell 2001)

*A partition  $\{V_1, \dots, V_r\}$  of the vertex set of a graph  $G$  has an independent transversal if  $|V_i| \geq 2\Delta(G)$  for each  $i$ .*

- Haxell's proof is elementary and uses some somewhat delicate induction
- there are also proofs based on topological connectivity of the independent set complex



# Improving the constant

Introduction

The common  
lemmas

Proving the  
main results

Independent  
transversals

Putting it all  
together

Improving the  
constant

Building on observations by Aharoni, Berger and Ziv [1] about the proof of Haxell's lemma, King [6] proved the following lopsided version of Haxell's lemma. Using this, he proved that  $c = \frac{2}{3}$  works.

## Improving the constant

## Introduction

The common  
lemmasProving the  
main resultsIndependent  
transversalsPutting it all  
togetherImproving the  
constant

Building on observations by Aharoni, Berger and Ziv [1] about the proof of Haxell's lemma, King [6] proved the following lopsided version of Haxell's lemma. Using this, he proved that  $c = \frac{2}{3}$  works.

## Lemma (King 2009)

*A partition  $\{V_1, \dots, V_r\}$  of the vertex set of a graph  $G$  has an independent transversal if there exists a positive integer  $k$  such that for each  $i$  we have  $\min\{k, |V_i| - k\} \geq \max_{v \in V_i} d(v)$ .*

# Tightness of King's lemma

Introduction

The common  
lemmas

Proving the  
main results

Independent

transversals

Putting it all  
together

Improving the  
constant

- let  $G_r$  be line graph of a 5-cycle where each edge has multiplicity  $r$ ; that is,  $G_r := L(r \cdot C_5)$

# Tightness of King's lemma

Introduction

The common  
lemmas

Proving the  
main results

Independent  
transversals  
Putting it all  
together

Improving the  
constant

- let  $G_r$  be line graph of a 5-cycle where each edge has multiplicity  $r$ ; that is,  $G_r := L(r \cdot C_5)$
- then  $\omega(G_r) = 2r = \frac{2}{3}(\Delta(G_r) + 1)$  and no stable set hits every maximum clique

# Tightness of King's lemma

Introduction

The common  
lemmas

Proving the  
main results

Independent  
transversals  
Putting it all  
together

Improving the  
constant

- let  $G_r$  be line graph of a 5-cycle where each edge has multiplicity  $r$ ; that is,  $G_r := L(r \cdot C_5)$
- then  $\omega(G_r) = 2r = \frac{2}{3}(\Delta(G_r) + 1)$  and no stable set hits every maximum clique
- thus King's lemma is tight

# Tightness of King's lemma

## Introduction

## The common lemmas

## Proving the main results

## Independent transversals Putting it all together

## Improving the constant

- let  $G_r$  be line graph of a 5-cycle where each edge has multiplicity  $r$ ; that is,  $G_r := L(r \cdot C_5)$
- then  $\omega(G_r) = 2r = \frac{2}{3}(\Delta(G_r) + 1)$  and no stable set hits every maximum clique
- thus King's lemma is tight

PICTURE GOES HERE

## Hitting maximum cliques

Landon Rabern

### Introduction

### The common lemmas

### Proving the main results

Independent transversals  
Putting it all together  
Improving the constant



R. Aharoni, E. Berger, and R. Ziv, *Independent systems of representatives in weighted graphs*, *Combinatorica* **27** (2007), no. 3, 253–267.



N. Alon, *The linear arboricity of graphs*, *Israel Journal of Mathematics* **62** (1988), no. 3, 311–325.



O.V. Borodin and A.V. Kostochka, *On an upper bound of a graph's chromatic number, depending on the graph's degree and density*, *Journal of Combinatorial Theory, Series B* **23** (1977), no. 2-3, 247–250.



A. Hajnal, *A theorem on  $k$ -saturated graphs*, *Canadian Journal of Mathematics* (1965), 720.



P.E. Haxell, *A note on vertex list colouring*, *Combinatorics, Probability and Computing* **10** (2001), no. 4, 345–347.



A.D. King, *Hitting all maximum cliques with a stable set using lopsided independent transversals*, *CoRR* [abs/0911.1741](https://arxiv.org/abs/0911.1741) (2009).



A.D. King, B.A. Reed, and A. Vetta, *An upper bound for the chromatic number of line graphs*, *European Journal of Combinatorics* **28** (2007), no. 8, 2182–2187.



A.V. Kostochka, *Degree, density, and chromatic number*, *Metody Diskret. Anal.* **35** (1980), 45–70.



L. Rabern, *On hitting all maximum cliques with an independent set*, *ArXiv e-prints* (2009).



———, *On hitting all maximum cliques with an independent set*, *Journal of Graph Theory* **66** (2011), no. 1, 32–37.