

graph coloring tools

london rabern

Contents

Preface	v
graphs	1
coloring vertices	3
basic estimates	3
edge coloring	7
Hall's theorem	7
a Hall's theorem game	7
Vizing's theorem	7
hardness	7
vertex coloring, again	9
list coloring	9
online list coloring	9
kernel tools	9
polynomial tools	9
edge coloring, again	11
fans as a greedy strategy	11
Kierstead paths	11
Tashkinov trees	11
edge list coloring	11
Lovász-Catlin shuffle tool	13
destroying non-complete	13
Mozhan basics	13
a hint of Kostochka's technique	13
independent transversals	15
randomly	15
Haxell's tool	15
Hajnal and Kostochka maximum clique tool	15
King's lemma	15
vertex transitive graphs	17
strong coloring	17
medium clique implies big clique	17
Kostochka-Yancey potential tool	19

Preface

This is the preface.

graphs

A *graph* is a collection of dots we call *vertices* some of which are connected by curves we call *edges*. The relative location of the dots and the shape of the curves are not relevant, we are only concerned with whether or not a given pair of dots is connected by a curve. Initially, we forbid edges from a vertex to itself and multiple edges between two vertices. If G is a graph, then $V(G)$ is its set of vertices and $E(G)$ its set of edges. We write $|G|$ for the number of vertices in $V(G)$ and $\|G\|$ for the number of edges in $E(G)$. Two vertices are *adjacent* if they are connected by an edge. The set of vertices to which v is adjacent is its *neighborhood*, written $N(v)$. For the size of v 's neighborhood $|N(v)|$, we write $d(v)$ and call this the *degree* of v .

[ADD PICTURES]

vertices
edges
 $V(G)$, $E(G)$
 $|G|$, $\|G\|$
adjacent
neighborhood
 $N(v)$
 $d(v)$, degree

coloring vertices

The entire book concerns one simple task: we want to color the vertices of a given graph so that adjacent vertices receive different colors. With no preferences about what the coloring should look like, this is easy, we just give each vertex a different color. Things get interesting when we ask how few colors we can use. We are definitely going to need at least zero colors and that will only do for the graph with no vertices at all. Given one color, we can handle all graphs with no edges. With two colors, we can do any path and any cycle with an even number of vertices [PICTURE]. But, we can't handle a triangle or any other cycle with an odd number of vertices [PICTURE]. In fact, odd cycles are really the only thing that will prevent us from using two colors. A graph H is a *subgraph* of a graph G , written $H \subseteq G$ if $V(H) \subseteq V(G)$ and $E(H) \subseteq E(G)$. When $H \subseteq G$, we say that G *contains* H . If $v \in V(G)$, then $G - v$ is the graph we get by removing v and all edges incident to v from G . A graph is k -colorable if we can color its vertices with (at most) k colors such that adjacent vertices receive different colors.

subgraph, \subseteq
contains
 $G - v$
 k -colorable

THEOREM 1. *A graph is 2-colorable just in case it contains no odd cycle.*

PROOF. A graph containing an odd cycle clearly can't be 2-colored. For the other implication, suppose there is a graph that is not 2-colorable and doesn't contain an odd cycle. Then we may pick such a graph G with $|G|$ as small as possible. Surely, $|G| > 0$, so we may pick $v \in V(G)$. If $x, y \in N(v)$, then x is not adjacent to y since then xyz would be an odd cycle. So we can construct a graph H from G by removing v and identifying all of $N(v)$ to a new vertex x_v . Any odd cycle in H would contain x_v and hence give rise to an odd cycle in G . So H contains no odd cycle. Since $|H| < |G|$, applying the theorem to H gives a 2-coloring of H , say with red and blue where x_v gets colored red. But this gives a 2-coloring of G by coloring all vertices in $N(v)$ red and v blue, a contradiction. \square

Well, this is embarrassing, coloring appears to be easy. Fortunately, things get more interesting when we move up to three colors.

THEOREM 2. *3-coloring is hard supposing other things we think are hard are actually hard.*

PROOF. reduce 3-SAT to 3-coloring. \square

basic estimates

Even though finding the minimum number of colors needed to color a graph is hard in general (supposing it is), we can still look for lower and upper bounds on this value. The *chromatic number* $\chi(G)$ of a graph G is the smallest k for which G is k -colorable. The simplest thing we can do is give each vertex a different color.

chromatic number
 $\chi(G)$

THEOREM 3. *If G is a graph, then $\chi(G) \leq |G|$.*

The only graphs that attain the upper bound in Theorem 3 are the *complete* graphs; those in which any two vertices are adjacent. We can usually do much better by just arbitrarily coloring vertices, reusing colors when we can. The *maximum degree* $\Delta(G)$ of a graph G is the largest degree of any vertex in G ; that is

$$\Delta(G) := \max_{v \in V(G)} d(v).$$

THEOREM 4. *If G is a graph, then $\chi(G) \leq \Delta(G) + 1$.*

PROOF. Suppose there is a graph G that is not $(\Delta(G) + 1)$ -colorable. Then we may pick such a graph G with $|G|$ as small as possible. Surely, $|G| > 0$, so we may pick $v \in V(G)$. Then $|G - v| < |G|$ and $\Delta(G - v) \leq \Delta(G)$, so applying the theorem to $G - v$ gives a $(\Delta(G - v) + 1)$ -coloring of $G - v$. But v has at most $\Delta(G)$ neighbors, so there is some color, say red, not used on $N(v)$, coloring v red gives a $(\Delta(G) + 1)$ -coloring of G , a contradiction. \square

Both complete graphs and odd cycles attain the upper bound in Theorem 4. Theorem 1 says we can do better for graphs that don't contain odd cycles. We can also do better for graphs that don't contain large complete subgraphs. A set of vertices S in a graph G is a *clique* if the vertices in S are pairwise adjacent. The *clique number* of a graph G , written $\omega(G)$, is the number of vertices in a largest clique in G . A set of vertices S in a graph G is *independent* if the vertices in S are pairwise non-adjacent. The *independence number* of a graph G , written $\alpha(G)$, is the number of vertices in a largest independent set in G .

THEOREM 5 (Brooks' theorem). *If G is a graph with $\Delta(G) \geq 3$ and $\omega(G) \leq \Delta(G)$, then $\chi(G) \leq \Delta(G)$.*

PROOF. Suppose there is a graph G with $\Delta(G) \geq 3$ and $\omega(G) \leq \Delta(G)$ that is not $\Delta(G)$ -colorable. Then we may pick such a graph G with $|G|$ as small as possible. Let S be a maximal independent set in G . Since S is maximal, every vertex in $G - S$ has a neighbor in S , so $\Delta(G) > \Delta(G - S)$. If red is an unused color in a $\chi(G - S)$ -coloring of $G - S$, then by coloring all vertices in S red we get a $(\chi(G - S) + 1)$ -coloring of G . So, $\Delta(G) + 1 \leq \chi(G) \leq \chi(G - S) + 1$. We conclude $\chi(G - S) > \Delta(G - S)$ and thus $\Delta(G) = \chi(G - S) = \Delta(G - S) + 1$ by Theorem 4. Since $|G - S| < |G|$, applying the theorem to $G - S$ shows that $\Delta(G - S) < 3$ or $\Delta(G - S) < \omega(G - S)$. So, either $\chi(G - S) = \Delta(G) = 3$ or $\omega(G - S) \geq \Delta(G)$. In the former case, let X be the vertex set of an odd cycle in $G - S$ guaranteed by Theorem 1. In the latter case, let X be a $\Delta(G)$ -clique in $G - S$.

Since S is maximal and $\omega(G) \leq \Delta(G)$, there are $x_1, x_2 \in X$ and $y_1, y_2 \in S$ such that x_1 is adjacent to y_1 and x_2 is adjacent to y_2 . Construct a graph H from $G - X$ by adding the edge $y_1 y_2$. Since $|H| < |G|$, applying the theorem to H shows that $\omega(H) > \Delta(G)$ or $\chi(H) \leq \Delta(G)$. Suppose $\chi(H) \leq \Delta(G)$. Then there is a $\Delta(G)$ -coloring of $G - X$ where y_1 and y_2 receive different colors, say red and blue respectively. Pick the first vertex z in a shortest path P from x_1 to x_2 in X that has a blue colored neighbor in $V(H)$. Each vertex in X has $\Delta(G) - 1$ neighbors in X and hence at most one neighbor in $V(H)$. So, $z \neq x_1$ since x_1 already has a red colored neighbor in $V(H)$. Let w be the vertex preceding z on P . Then w has no blue colored neighbor. Since X is the vertex set of a cycle or a complete graph, there is a path Q from w to z passing through every vertex of X . Color w

blue and then proceed along Q , coloring one vertex at a time. Since each vertex we encounter before we get to z has at most $\Delta(G) - 1$ colored neighbors, we always have an available color to use. But, z is adjacent to both w and another blue colored vertex in $V(H)$, so there is an available color for z as well. This gives a $\Delta(G)$ -coloring of G , a contradiction.

So, $\omega(H) > \Delta(G)$. In particular, y_1 and y_2 each have exactly one neighbor in X and $\Delta(G) - 1$ neighbors in the same $\Delta(G) - 1$ clique A in $G - X$. Since S is maximal and $|X| \geq 3$, there must be adjacent $x_3 \in X \setminus \{x_1, x_2\}$ and $y_3 \in S \setminus \{y_1, y_2\}$. Applying the same argument with x_3, y_3 in place of x_2, y_2 shows that y_1 and y_3 each have exactly one neighbor in X and $\Delta(G) - 1$ neighbors in the same $\Delta(G) - 1$ clique B in $G - X$. But $|A \cap B| = |A| + |B| - |A \cup B| \geq 2(\Delta(G) - 1) - d(y_1) \geq \Delta(G) - 2 > 0$. But there can't be a vertex in $A \cap B$ since it would be adjacent to y_1, y_2, y_3 as well as $\Delta(G) - 2$ vertices in A and thus have degree greater than $\Delta(G)$, a contradiction.

edge coloring

two-coloring

Hall's theorem

a Hall's theorem game

Vizing's theorem

hardness

vertex coloring, again

list coloring

online list coloring

kernel tools

Brooks again.

maximum independent covers.

polynomial tools

combinatorial nullstellensatz.

coefficient formulae.

a combinatorial interpretation.

edge coloring, again

fans as a greedy strategy

Kierstead paths

Tashkinov trees

edge list coloring

more kernel method.

2-edge-coloring.

improved Brooks' theorem.

a hint of quasiline and claw-free graphs.

Lovász-Catlin shuffle tool

destroying non-complete

Mozhan basics

a hint of Kostochka's technique

independent transversals

randomly

Haxell's tool

a hint of Sperner's lemma.

Hajnal and Kostochka maximum clique tool

King's lemma

vertex transitive graphs

strong coloring

medium clique implies big clique

Kostochka-Yancey potential tool