

A common generalization of Hall's theorem and Vizing's edge-coloring theorem

landon rabern

LBD Data

Miami University Colloquium
November 6, 2014

Hall's theorem

- given finite sets A_1, A_2, \dots, A_n

Hall's theorem

- given finite sets A_1, A_2, \dots, A_n
- a **system of distinct representatives** (SDR) is a choice of $a_i \in A_i$ for all i where $a_i \neq a_j$ for $i \neq j$

Hall's theorem

- given finite sets A_1, A_2, \dots, A_n
- a **system of distinct representatives** (SDR) is a choice of $a_i \in A_i$ for all i where $a_i \neq a_j$ for $i \neq j$
- when can we pick an SDR?

Hall's theorem

- given finite sets A_1, A_2, \dots, A_n
- a **system of distinct representatives** (SDR) is a choice of $a_i \in A_i$ for all i where $a_i \neq a_j$ for $i \neq j$
- when can we pick an SDR?
- if k of the sets together have fewer than k elements, we can't

Hall's theorem

- given finite sets A_1, A_2, \dots, A_n
- a **system of distinct representatives** (SDR) is a choice of $a_i \in A_i$ for all i where $a_i \neq a_j$ for $i \neq j$
- when can we pick an SDR?
- if k of the sets together have fewer than k elements, we can't
 - $A_1 = \{1, 2\}, A_2 = \{1, 2\}, A_3 = \{1, 2\}$

Hall's theorem






- given finite sets A_1, A_2, \dots, A_n
- a **system of distinct representatives** (SDR) is a choice of $a_i \in A_i$ for all i where $a_i \neq a_j$ for $i \neq j$
- when can we pick an SDR?
- if k of the sets together have fewer than k elements, we can't
 - $A_1 = \{1, 2\}, A_2 = \{1, 2\}, A_3 = \{1, 2\}$
- **Hall's theorem: this is the only thing that can go wrong**

$$\text{SDR exists} \Leftrightarrow \left| \bigcup_{i \in I} A_i \right| \geq |I| \text{ for all } I \subseteq \{1, \dots, n\}$$

some card games

the simplest variation



- two players, Dealer and Player



some card games

the simplest variation



- two players, Dealer and Player
- the deck has just many copies of the high spade cards



some card games

the simplest variation



- two players, Dealer and Player
- the deck has just many copies of the high spade cards
- Dealer makes 5 stacks of cards with no duplicates, all cards face-up



some card games



the simplest variation

- two players, Dealer and Player
- the deck has just many copies of the high spade cards
- Dealer makes 5 stacks of cards with no duplicates, all cards face-up
- Player wins if he can pick a Royal Flush, one card from each stack





some card games

example, a Player win




some card games

example, a Player win



some card games

example, a Dealer win



some card games

winning condition

- Player cannot win if there is a set of k stacks that together have fewer than k different cards

some card games

winning condition

- Player cannot win if there is a set of k stacks that together have fewer than k different cards



some card games

winning condition

- Player cannot win if there is a set of k stacks that together have fewer than k different cards
- Hall's theorem says: **Player wins otherwise**



some card games

making things harder for Dealer

- this isn't a fun game, far too easy for Dealer to win

some card games

making things harder for Dealer

- this isn't a fun game, far too easy for Dealer to win
- to make a better game, we allow Player to modify some of the stacks

some card games

making things harder for Dealer

- this isn't a fun game, far too easy for Dealer to win
- to make a better game, we allow Player to modify some of the stacks

Player's Move

Player can pick any card A from the deck and swap it for another card B in one stack (not containing A).

some card games

making things harder for Dealer

- this isn't a fun game, far too easy for Dealer to win
- to make a better game, we allow Player to modify some of the stacks

Player's Move

Player can pick any card A from the deck and swap it for another card B in one stack (not containing A).

Dealer's Move

Dealer can (i) do nothing or (ii) swap A and B in at most one other stack.

some card games

making things harder for Dealer

- this isn't a fun game, far too easy for Dealer to win
- to make a better game, we allow Player to modify some of the stacks

Player's Move

Player can pick any card A from the deck and swap it for another card B in one stack (not containing A).

Dealer's Move


Dealer can (i) do nothing or (ii) swap A and B in at most one other stack.

Winning

Player wins if he can pick a Royal Flush at the start of one of his turns, otherwise Dealer wins.

some card games


example, a Player win



some card games

example, a Player win


- Player picks a King from the deck and swaps it for a Queen in the first stack



some card games

example, a Player win


- Player picks a King from the deck and swaps it for a Queen in the first stack



some card games

example, a Player win

- Player picks a King from the deck and swaps it for a Queen in the first stack
- Dealer can swap a King and Queen in one of the other stacks



some card games

example, a Player win


- Player picks a King from the deck and swaps it for a Queen in the first stack
- Dealer can swap a King and Queen in one of the other stacks



some card games


example, a Player win

- Player picks a King from the deck and swaps it for a Queen in the first stack
- Dealer can swap a King and Queen in one of the other stacks
- Player wins no matter what Dealer does





some card games

example, a Dealer win




some card games

example, a Dealer win






some card games

example, a Dealer win




some card games

example, a Dealer win



some card games


what was the difference?



some card games

what was the difference?


- in the top game, Dealer can prevent Player from increasing the number of different cards in the first two stacks



some card games

what was the difference?

- in the top game, Dealer can prevent Player from increasing the number of different cards in the first two stacks
- in the bottom game, Dealer cannot prevent Player from increasing the number of different cards in the first three stacks



some card games

necessary condition

- if the same card appears on three stacks, Player can force the addition of a new card to these stacks

some card games

necessary condition

- if the same card appears on three stacks, Player can force the addition of a new card to these stacks
- it is not hard to show that this is essentially all Player can do

some card games

necessary condition

- if the same card appears on three stacks, Player can force the addition of a new card to these stacks
- it is not hard to show that this is essentially all Player can do
- this suggests a necessary condition

some card games

necessary condition

- if the same card appears on three stacks, Player can force the addition of a new card to these stacks
- it is not hard to show that this is essentially all Player can do
- this suggests a necessary condition

Degree

The *degree* of a card C in a set of stacks S is the number of times C appears in S . We write $d_S(C)$ for this quantity.

some card games

necessary condition

- if the same card appears on three stacks, Player can force the addition of a new card to these stacks
- it is not hard to show that this is essentially all Player can do
- this suggests a necessary condition

Degree

The *degree* of a card C in a set of stacks S is the number of times C appears in S . We write $d_S(C)$ for this quantity.

Necessary Condition

If Player has a winning strategy, then for every set of stacks S we must have

$$\sum_{C \in \cup S} \left\lceil \frac{d_S(C)}{2} \right\rceil \geq |S|.$$

some card games

intuition

Degree

The *degree* of a card C in a set of stacks S is the number of times C appears in S . We write $d_S(C)$ for this quantity.

Necessary Condition

If Player has a winning strategy, then for every set of stacks S we must have

$$\sum_{C \in \bigcup S} \left\lceil \frac{d_S(C)}{2} \right\rceil \geq |S|.$$

- in Hall's theorem, each C is 'worth' 1 in $\sum_{C \in \bigcup S} 1 = \left| \bigcup S \right| \geq |S|$

some card games

intuition

Degree

The *degree* of a card C in a set of stacks S is the number of times C appears in S . We write $d_S(C)$ for this quantity.

Necessary Condition

If Player has a winning strategy, then for every set of stacks S we must have

$$\sum_{C \in \bigcup S} \left\lceil \frac{d_S(C)}{2} \right\rceil \geq |S|.$$

- in Hall's theorem, each C is 'worth' 1 in $\sum_{C \in \bigcup S} 1 = \left| \bigcup S \right| \geq |S|$
- Player can turn $2t + 1$ of the same card into $t + 1$ different cards, so C is 'worth' $\left\lceil \frac{d_S(C)}{2} \right\rceil$

some card games

Dealer's strategy

- given a set of stacks S with $\sum_{C \in \cup S} \left\lceil \frac{d_S(C)}{2} \right\rceil < |S|$

some card games

Dealer's strategy

- given a set of stacks S with $\sum_{C \in \cup S} \left\lceil \frac{d_S(C)}{2} \right\rceil < |S|$
- Dealer's strategy: **maintain this invariant**

some card games

Dealer's strategy

- given a set of stacks S with $\sum_{C \in \bigcup S} \left\lceil \frac{d_S(C)}{2} \right\rceil < |S|$
- Dealer's strategy: **maintain this invariant**
 - this is good enough since then $|\bigcup S| \leq \sum_{C \in \bigcup S} \left\lceil \frac{d_S(C)}{2} \right\rceil < |S|$ always

some card games

Dealer's strategy

- given a set of stacks S with $\sum_{C \in \bigcup S} \left\lceil \frac{d_S(C)}{2} \right\rceil < |S|$
- Dealer's strategy: **maintain this invariant**
 - this is good enough since then $|\bigcup S| \leq \sum_{C \in \bigcup S} \left\lceil \frac{d_S(C)}{2} \right\rceil < |S|$ always
 - if Player's swaps A in for B , increasing $\left\lceil \frac{d_S(A)}{2} \right\rceil$, then $d_S(A)$ and $d_S(B)$ both changed from even to odd

some card games

Dealer's strategy

- given a set of stacks S with $\sum_{C \in \cup S} \left\lceil \frac{d_S(C)}{2} \right\rceil < |S|$
- Dealer's strategy: **maintain this invariant**
 - this is good enough since then $|\cup S| \leq \sum_{C \in \cup S} \left\lceil \frac{d_S(C)}{2} \right\rceil < |S|$ always
 - if Player's swaps A in for B , increasing $\left\lceil \frac{d_S(A)}{2} \right\rceil$, then $d_S(A)$ and $d_S(B)$ both changed from even to odd
 - Dealer can swap A for B somewhere else, decreasing $\left\lceil \frac{d_S(B)}{2} \right\rceil$ and not increasing $\left\lceil \frac{d_S(A)}{2} \right\rceil$

some card games

Dealer's strategy

- given a set of stacks S with $\sum_{C \in \cup S} \left\lceil \frac{d_S(C)}{2} \right\rceil < |S|$
- Dealer's strategy: **maintain this invariant**
 - this is good enough since then $|\cup S| \leq \sum_{C \in \cup S} \left\lceil \frac{d_S(C)}{2} \right\rceil < |S|$ always
 - if Player's swaps A in for B , increasing $\left\lceil \frac{d_S(A)}{2} \right\rceil$, then $d_S(A)$ and $d_S(B)$ both changed from even to odd
 - Dealer can swap A for B somewhere else, decreasing $\left\lceil \frac{d_S(B)}{2} \right\rceil$ and not increasing $\left\lceil \frac{d_S(A)}{2} \right\rceil$
 - Dealer has maintained $\sum_{C \in \cup S} \left\lceil \frac{d_S(C)}{2} \right\rceil < |S|$

some card games

winning condition

- **this necessary condition is also sufficient**

some card games

winning condition

- **this necessary condition is also sufficient**

Winning Condition


Player has a winning strategy if and only if for every set of stacks S we have

$$\sum_{C \in \cup S} \left\lceil \frac{d_S(C)}{2} \right\rceil \geq |S|.$$

some card games

proof idea


- 1 Player looks for a set of card types that give a system of distinct representatives of all the stacks containing them



some card games

proof idea


- 1 Player looks for a set of card types that give a system of distinct representatives of all the stacks containing them



some card games

proof idea


- 1 Player looks for a set of card types that give a system of distinct representatives of all the stacks containing them



some card games

proof idea


- 1 Player looks for a set of card types that give a system of distinct representatives of all the stacks containing them



some card games

proof idea


- 1 Player looks for a set of card types that give a system of distinct representatives of all the stacks containing them
- 2 Player calls those stacks done and never plays with those card types again



some card games

proof idea


- ③ if no such set of card types exists, then Hall's theorem shows that there is at least one card appearing on none of the remaining stacks



some card games

proof idea


- 3 if no such set of card types exists, then Hall's theorem shows that there is at least one card appearing on none of the remaining stacks
- 4 but then some card appears at least thrice, so Player can increase the number of card types in the stacks



some card games

proof idea


- 3 if no such set of card types exists, then Hall's theorem shows that there is at least one card appearing on none of the remaining stacks
- 4 but then some card appears at least thrice, so Player can increase the number of card types in the stacks
- 5 goto step 1



some card games

proof idea


- ③ if no such set of card types exists, then Hall's theorem shows that there is at least one card appearing on none of the remaining stacks
- ④ but then some card appears at least thrice, so Player can increase the number of card types in the stacks
- ⑤ goto step 1



some card games

proof idea


- 3 if no such set of card types exists, then Hall's theorem shows that there is at least one card appearing on none of the remaining stacks
- 4 but then some card appears at least thrice, so Player can increase the number of card types in the stacks
- 5 goto step 1



some card games

proof idea

- 3 if no such set of card types exists, then Hall's theorem shows that there is at least one card appearing on none of the remaining stacks
- 4 but then some card appears at least thrice, so Player can increase the number of card types in the stacks
- 5 goto step 1



A generalization of Hall's theorem

making it harder for Player

- allow Dealer to make more swaps in response to Player's move

A generalization of Hall's theorem

making it harder for Player

- allow Dealer to make more swaps in response to Player's move
- for each $t \geq 1$, the t -game allows Dealer to make up to t swaps

A generalization of Hall's theorem

making it harder for Player

- allow Dealer to make more swaps in response to Player's move
- for each $t \geq 1$, the t -game allows Dealer to make up to t swaps

Winning Condition

Player has a winning strategy in the t -game if and only if for every set of stacks S we have

$$\sum_{C \in \cup S} \left\lceil \frac{d_S(C)}{t+1} \right\rceil \geq |S|.$$

A generalization of Hall's theorem

making it harder for Player

- allow Dealer to make more swaps in response to Player's move
- for each $t \geq 1$, the t -game allows Dealer to make up to t swaps

Winning Condition

Player has a winning strategy in the t -game if and only if for every set of stacks S we have

$$\sum_{C \in \cup S} \left\lceil \frac{d_S(C)}{t+1} \right\rceil \geq |S|.$$

- Hall's theorem is the winning condition in the $(k-1)$ -game when there are k total stacks:

A generalization of Hall's theorem

making it harder for Player

- allow Dealer to make more swaps in response to Player's move
- for each $t \geq 1$, the t -game allows Dealer to make up to t swaps

Winning Condition

Player has a winning strategy in the t -game if and only if for every set of stacks S we have

$$\sum_{C \in \cup S} \left\lceil \frac{d_S(C)}{t+1} \right\rceil \geq |S|.$$

- Hall's theorem is the winning condition in the $(k-1)$ -game when there are k total stacks:
 - $1 \leq d_S(C) \leq k$, so $\left\lceil \frac{d_S(C)}{t+1} \right\rceil = 1$

A generalization of Hall's theorem

making it harder for Player

- allow Dealer to make more swaps in response to Player's move
- for each $t \geq 1$, the t -game allows Dealer to make up to t swaps

Winning Condition

Player has a winning strategy in the t -game if and only if for every set of stacks S we have

$$\sum_{C \in \bigcup S} \left\lceil \frac{d_S(C)}{t+1} \right\rceil \geq |S|.$$

- Hall's theorem is the winning condition in the $(k-1)$ -game when there are k total stacks:
 - $1 \leq d_S(C) \leq k$, so $\left\lceil \frac{d_S(C)}{t+1} \right\rceil = 1$
 - so, the sum equals $|\bigcup S|$

A generalization of Hall's theorem

making it harder for Player

- allow Dealer to make more swaps in response to Player's move
- for each $t \geq 1$, the t -game allows Dealer to make up to t swaps

Winning Condition

Player has a winning strategy in the t -game if and only if for every set of stacks S we have

$$\sum_{C \in \bigcup S} \left\lceil \frac{d_S(C)}{t+1} \right\rceil \geq |S|.$$

- Hall's theorem is the winning condition in the $(k-1)$ -game when there are k total stacks:
 - $1 \leq d_S(C) \leq k$, so $\left\lceil \frac{d_S(C)}{t+1} \right\rceil = 1$
 - so, the sum equals $|\bigcup S|$
 - Player's moves are useless

edge coloring

setup

- assign colors to the edges of a graph so that incident edges get different colors

edge coloring

setup

- assign colors to the edges of a graph so that incident edges get different colors
- how few colors can we use?

edge coloring

setup

- assign colors to the edges of a graph so that incident edges get different colors
- how few colors can we use?


Vizing's theorem

Any simple graph can be edge-colored using at most one more color than its maximum degree.

edge coloring

proof of Vizing's theorem


- proceed by induction on the number of vertices



edge coloring

proof of Vizing's theorem


- proceed by induction on the number of vertices
- remove a vertex and edge-color the rest with one more color than its maximum degree



edge coloring

proof of Vizing's theorem


- proceed by induction on the number of vertices
- remove a vertex and edge-color the rest with one more color than its maximum degree



edge coloring

proof of Vizing's theorem


- proceed by induction on the number of vertices
- remove a vertex and edge-color the rest with one more color than its maximum degree



edge coloring

proof of Vizing's theorem


- proceed by induction on the number of vertices
- remove a vertex and edge-color the rest with one more color than its maximum degree



edge coloring

proof of Vizing's theorem


- proceed by induction on the number of vertices
- remove a vertex and edge-color the rest with one more color than its maximum degree



edge coloring

proof of Vizing's theorem


- proceed by induction on the number of vertices
- remove a vertex and edge-color the rest with one more color than its maximum degree



edge coloring

proof of Vizing's theorem

- proceed by induction on the number of vertices
- remove a vertex and edge-color the rest with one more color than its maximum degree



edge coloring

proof of Vizing's theorem

- **exchanging colors on a two-colored path is just a Player move followed by a Dealer move**

edge coloring

proof of Vizing's theorem

- **exchanging colors on a two-colored path is just a Player move followed by a Dealer move**
- we can make any of Player's legal moves this way, so if the winning conditions are satisfied, Vizing's theorem is true

edge coloring

proof of Vizing's theorem

- **exchanging colors on a two-colored path is just a Player move followed by a Dealer move**
- we can make any of Player's legal moves this way, so if the winning conditions are satisfied, Vizing's theorem is true
- each stack has at least two colors, so counting the 'cards' in two ways we get for each set of stacks S ,

$$\sum_{C \in \cup S} d_S(C) \geq 2|S|$$

edge coloring

proof of Vizing's theorem

- **exchanging colors on a two-colored path is just a Player move followed by a Dealer move**
- we can make any of Player's legal moves this way, so if the winning conditions are satisfied, Vizing's theorem is true
- each stack has at least two colors, so counting the 'cards' in two ways we get for each set of stacks S ,

$$\sum_{C \in \cup S} d_S(C) \geq 2|S|$$

- so, we have the desired winning condition

$$\sum_{C \in \cup S} \frac{d_S(C)}{2} \geq |S|$$

summary

- we introduced a simple card game

summary

- we introduced a simple card game
 - Player can pick any card A from the deck and swap it for another card B in one stack (not containing A)

- we introduced a simple card game
 - Player can pick any card A from the deck and swap it for another card B in one stack (not containing A)
 - Dealer can (i) do nothing or (ii) swap A and B in at most one other stack

- we introduced a simple card game
 - Player can pick any card A from the deck and swap it for another card B in one stack (not containing A)
 - Dealer can (i) do nothing or (ii) swap A and B in at most one other stack
 - Player wins if he can pick a Royal Flush at the start of one of his turns, otherwise Dealer wins

- we introduced a simple card game
 - Player can pick any card A from the deck and swap it for another card B in one stack (not containing A)
 - Dealer can (i) do nothing or (ii) swap A and B in at most one other stack
 - Player wins if he can pick a Royal Flush at the start of one of his turns, otherwise Dealer wins
- Player has a winning strategy exactly when a Hall-like condition is satisfied

- we introduced a simple card game
 - Player can pick any card A from the deck and swap it for another card B in one stack (not containing A)
 - Dealer can (i) do nothing or (ii) swap A and B in at most one other stack
 - Player wins if he can pick a Royal Flush at the start of one of his turns, otherwise Dealer wins
- Player has a winning strategy exactly when a Hall-like condition is satisfied
- Vizing's edge-coloring theorem is an easy corollary

- we introduced a simple card game
 - Player can pick any card A from the deck and swap it for another card B in one stack (not containing A)
 - Dealer can (i) do nothing or (ii) swap A and B in at most one other stack
 - Player wins if he can pick a Royal Flush at the start of one of his turns, otherwise Dealer wins
- Player has a winning strategy exactly when a Hall-like condition is satisfied
- Vizing's edge-coloring theorem is an easy corollary
- most other classical edge-coloring results follow easily as well

- we introduced a simple card game
 - Player can pick any card A from the deck and swap it for another card B in one stack (not containing A)
 - Dealer can (i) do nothing or (ii) swap A and B in at most one other stack
 - Player wins if he can pick a Royal Flush at the start of one of his turns, otherwise Dealer wins
- Player has a winning strategy exactly when a Hall-like condition is satisfied
- Vizing's edge-coloring theorem is an easy corollary
- most other classical edge-coloring results follow easily as well
- generalizes easily to multigraphs

- we introduced a simple card game
 - Player can pick any card A from the deck and swap it for another card B in one stack (not containing A)
 - Dealer can (i) do nothing or (ii) swap A and B in at most one other stack
 - Player wins if he can pick a Royal Flush at the start of one of his turns, otherwise Dealer wins
- Player has a winning strategy exactly when a Hall-like condition is satisfied
- Vizing's edge-coloring theorem is an easy corollary
- most other classical edge-coloring results follow easily as well
- generalizes easily to multigraphs
- a more general game unifies much of edge-coloring theory