

# Part II — Asymptotic Methods

Based on lectures by E. S. Titi

Notes taken by Joseph Tedds using Dexter Chua's header and Gilles Castel's snippets.

Lent 2020

These notes are not endorsed by the lecturers, and I have modified them (often significantly) after lectures. They are nowhere near accurate representations of what was actually lectured, and in particular, all errors are almost surely mine.

## **Asymptotic expansion**

Definition (Poincare) of  $\phi(z) \sim \sum a_n z^{-n}$ ; examples; elementary properties; uniqueness; Stokes' phenomenon [4]

## **Asymptotics behaviour of functions defined by integrals**

Integration by parts. Watson's lemma and Laplace's method. Riemann-Lebesgue lemma and method of stationary phase. The method of steepest descent (including derivation of higher order terms). Airy function, \*and application to wave theory of a rainbow\*. [7]

## **Asymptotic behaviour of solutions of differential equations**

Asymptotic solution of second-order linear differential equations, including Liouville-Green functions (proof that they are asymptotic not required) and WKBJ with the quantum harmonic oscillator as an example. [4]

## **Recent developments**

Further discussion of Stokes phenomenon. \*Asymptotics beyond all orders\*. [1]

## Contents

<b>0</b>	<b>Introduction</b>	<b>3</b>
<b>1</b>	<b>Asymptotic Expansions / Series</b>	<b>4</b>

## 0 Introduction

Consider  $f(x) = x + 3 + \frac{1}{x^2}$  (1), from the sketch we see that  $f(x) - (x + 3) = \frac{1}{x^2} \xrightarrow{x \rightarrow \infty} 0$ . Now, for  $g(x) = x^2 + x + 3\frac{1}{x^3}$  we obtain the asymptote  $y = x^2 + x + 3$  from  $g(x) - (x^2 + x + 3) \xrightarrow{x \rightarrow \infty} 0$ . In this course, we will study

- (i) Functions
- (ii) Solutions to differential equations
- (iii) Integrals

Important definitions:

**Definition (Big  $\mathcal{O}$ ).** –  $f : (a, \infty) \rightarrow \mathbb{C} \setminus \mathbb{R}$ ,  $g : (a, \infty) \rightarrow \mathbb{R}$ . Assume that  $g(x) > 0$  for  $x \geq A \geq a$ . We say that  $f(x) = \mathcal{O}(g(x))$  as  $x \rightarrow \infty$  if  $\exists M > 0, B > 0$  such that

$$|f(x)| \leq Mg(x) \quad x \geq B > A.$$

(2)

- We say  $f(x) = \mathcal{O}(g(x))$  as  $x \rightarrow x_0$  if  $\exists M, \delta > 0$  such that

$$\frac{|f(x)|}{g(x)} \leq M; \quad 0 < |x - x_0| < \delta.$$

(3) i.e.

$$\limsup_{x \rightarrow x_0} \frac{|f(x)|}{g(x)} < \infty.$$

Observation If  $f(x) = \mathcal{O}(g(x))$  as  $x \rightarrow x_0$ , then  $cf = \mathcal{O}(g(x))$ ,  $c \in \mathbb{R}$

**Example.** Let  $f(x) = \frac{1}{x^2} \sin \frac{1}{x}$  as  $x \rightarrow x_0$ ,  $f(x) = \mathcal{O}(\frac{1}{x^2})$

$$\limsup \frac{|f(x)|}{g(x)} = \limsup_{x \rightarrow 0} \left| \sin \frac{1}{x} \right| = 1 < \infty.$$

**Definition (Little  $o$ ).** We say  $f(x) = o(g(x))$  as  $x \rightarrow x_0$  if

$$\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = 0.$$

More rigorously,  $\forall \varepsilon > 0 \exists \delta(\varepsilon)$  such that  $0 < |x - x_0| < \delta$  then  $\left| \frac{f(x)}{g(x)} \right| < \varepsilon$

**Example.**  $f(x) = \frac{1}{x^2} \sin \frac{1}{x}$  and  $g(x) = \frac{1}{x^3}$  as  $x \rightarrow 0$ . We claim  $f(x) = o(g(x))$

$$\frac{f(x)}{g(x)} = \sin \frac{1}{x} \cdot x \xrightarrow{x \rightarrow 0} 0.$$

**Example.**  $f(x) = 4x^2$  as  $x \rightarrow \infty$

- (i)  $f(x) = \mathcal{O}(x^3), o(x^3)$  but not  $o(x^2)$

**Notation.**

$$f(x) = o(g(x)) \iff f(x) \ll g(x).$$

# 1 Asymptotic Expansions / Series

**Definition** (Asymptotic sequence).  $\phi_n : D \subset \mathbb{C} \mapsto \mathbb{C} \setminus \mathbb{R} \ n = 0, 1, 2, \dots$  is called an asymptotic sequence as  $z \rightarrow z_0 \in D$  if

$$\phi_n(z) = o(\phi_m(z)) \text{ as } z \rightarrow z_0$$

for all

$$n > m \iff \frac{\phi_n(z)}{\phi_m(z)} \xrightarrow{z \rightarrow z_0} 0 \ \forall \ n > m.$$

**Example.** (i)

$$\frac{1}{x^3}, \frac{1}{x^2}, \frac{1}{x}, 1, x, x^2, \dots, x^n, \dots$$

$$x = 0, \frac{x^n}{x^m} = x^{n-m} \xrightarrow{x \rightarrow 0} 0$$

$$(ii) \ \frac{1}{(x-5)^k}, k = 0, 1, \dots, \text{ as } x \rightarrow \infty$$

$$(iii) \ \phi_n(x) = \frac{1}{x^2} \cos nx \text{ for } n = 0, 1, 2, \dots, \text{ consider}$$

$$\frac{\phi_n(x)}{\phi_m(x)} = \frac{\cos nx}{\cos mx}.$$

But  $\limsup \frac{\cos nx}{\cos mx} = 1 \neq 0$  hence is not an asymptotic sequence.

**Definition** (Asymptotic Expansion). Let  $\phi_n : D \subset \mathbb{C} \rightarrow \mathbb{C} \setminus \mathbb{R}$  be an asymptotic sequence about  $z \rightarrow z_0$ . We say the sum  $\sum_{n=0}^{\infty} a_n \phi_n$  is an *asymptotic expansion* of  $f(z)$  as  $z \rightarrow z_0$  if

$$f(z) - \sum_{n=0}^N a_n \phi_n(z) = o(\phi_N(z)) \text{ as } z \rightarrow z_0 \ \forall \ N = 0, 1, \dots$$

Or

$$\frac{f(z) - \sum_{n=0}^N a_n \phi_n(z)}{\phi_N(z)} \xrightarrow{z \rightarrow z_0} 0.$$

**Notation.** In this case, we write  $f(z) \sim \sum_{n=0}^{\infty} a_n \phi_n(z)$ .

**Remark.** We do not require  $\sum a_n \phi_n(z)$  to converge. \*

**Proposition.** If  $f(z) \sim \sum_{n=0}^{\infty} a_n \phi_n(z)$  then

$$a_{N+1} = \lim_{z \rightarrow z_0} \frac{f(z) - \sum_{k=0}^N a_k \phi_k(z)}{\phi_{N+1}(z)}.$$

Consider  $f(z) \sim \sum a_n \phi_n(z)$  for  $n = 0, 1, 2, \dots$  then

$$a_0 = 0 = \lim_{z \rightarrow z_0} \frac{f(z)}{\phi_0(z)}, \dots$$

Then

$$a_n = \lim_{z \rightarrow z_0} \frac{f(z)}{\phi_n(z)} = 0.$$

**Example.** (Taylor)  $f : [a, b] \rightarrow \mathbb{R}, C^\infty$ . Then  $f(x) \sim \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$   $x_0 \in (a, b)$ .

$$\frac{f(x) - \sum_{n=0}^N \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n}{(x - x_0)^N} \xrightarrow{x \rightarrow x_0} 0.$$

We can write

$$f(x) - \sum_{n=0}^N \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n = R_N(x) = \int_{x_0}^x \frac{(x-t)^N}{N!} f^{(N+1)}(t) dt.$$

Now

$$|R_N(x)| \leq \max_{a \leq t \leq b} f^{(N+1)}(t) \int_{x_0}^x \frac{|(x-t)^N|}{N!} dt = \frac{|x - x_0|^{N+1}}{(N+1)!} \max |f^{(N+1)}(t)|.$$

So

$$\left| \frac{R_N(x)}{(x - x_0)^N} \right| \leq \frac{|x - x_0|}{(N+1)!} \max |f^{(N+1)}(t)| \xrightarrow{x \rightarrow x_0} 0.$$

**Proposition.** (i) Let  $f(z) \sim \sum_{n=0}^{\infty} a_n \phi_n(z)$  and  $g(z) \sim \sum_{n=0}^{\infty} b_n \phi_n(z)$ , then

$$\alpha f(z) + \beta g(z) \sim \sum_{n=0}^{\infty} (\alpha a_n + \beta b_n) \phi_n(z).$$

(ii) Let  $f(z) \sim \sum_{n=0}^{\infty} a_n (z - z_0)^n$  and  $g(z) \sim \sum_{n=0}^{\infty} b_n (z - z_0)^n$ , then

$$f(z)g(z) \sim \sum_{n=0}^{\infty} c_n (z - z_0)^n, c_n = \sum_{k=0}^n a_k b_{n-k}.$$

**Proposition.** The asymptotic expansion is unique, that is if  $f(z) \sim \sum_{n=0}^{\infty} a_n \phi_n(z)$  and  $f(z) \sim \sum_{n=0}^{\infty} b_n \phi_n(z)$  then  $a_n = b_n \forall n$

$$a_0 - b_0 = \frac{(a_0 - b_0)\phi_0(z)}{\phi_0(z)} = \frac{(a_0 + \phi_0(z) - f(z)) + (f(z) - b_0\phi_0(z))}{\phi_0(z)} \xrightarrow{z \rightarrow z_0} 0.$$

The rest are done by induction.

Consider

$$H(x) = \begin{cases} e^{-\frac{1}{x^2}} & x \neq 0 \\ 0 & x = 0 \end{cases}.$$

$$f^{(n)}(0) = 0$$

$$\sin x = \sum_{n=0}^{\infty} \frac{1}{(2n+1)!} x^{2n+1}.$$

But

$$\sin x + H(x) \sim \sum_{n=0}^{\infty} \frac{1}{(2n+1)!} x^{2n+1}.$$