

# Assignment 3: Data Exploration

Jack Alcorn

## OVERVIEW

This exercise accompanies the lessons in Environmental Data Analytics on Data Exploration.

## Directions

1. Change “Student Name” on line 3 (above) with your name.
2. Work through the steps, **creating code and output** that fulfill each instruction.
3. Be sure to **answer the questions** in this assignment document.
4. When you have completed the assignment, **Knit** the text and code into a single PDF file.
5. After Knitting, submit the completed exercise (PDF file) to the dropbox in Sakai. Add your last name into the file name (e.g., “Salk\_A03\_DataExploration.Rmd”) prior to submission.

The completed exercise is due on <>.

## Set up your R session

1. Check your working directory, load necessary packages (tidyverse), and upload two datasets: the ECOTOX neonicotinoid dataset (ECOTOX\_Neonicotinoids\_Insects\_raw.csv) and the Niwot Ridge NEON dataset for litter and woody debris (NEON\_NIWO\_Litter\_massdata\_2018-08\_raw.csv). Name these datasets “Neonics” and “Litter”, respectively.

```
getwd()

## [1] "Z:/ENV_872_Data_Analy/Environmental_Data_Analytics_2021/Assignments"

setwd("Z:/ENV_872_Data_Analy/Environmental_Data_Analytics_2021")
library(tidyverse)

## Warning: package 'tidyverse' was built under R version 4.0.3
## Warning: package 'ggplot2' was built under R version 4.0.3
## Warning: package 'tibble' was built under R version 4.0.3
## Warning: package 'tidyr' was built under R version 4.0.3
## Warning: package 'readr' was built under R version 4.0.3
## Warning: package 'purrr' was built under R version 4.0.3
## Warning: package 'dplyr' was built under R version 4.0.3
## Warning: package 'stringr' was built under R version 4.0.3
## Warning: package 'forcats' was built under R version 4.0.3
library(lubridate)

## Warning: package 'lubridate' was built under R version 4.0.3
```

```
Neonics <- read.csv("Z:/ENV_872_Data_Analy/Environmental_Data_Analytics_2021/Data/Raw/ECOTOX_Neonicotin
Litter <- read.csv('Z:/ENV_872_Data_Analy/Environmental_Data_Analytics_2021/Data/Raw/NEON_NIWO_Litter_ma
```

## Learn about your system

2. The neonicotinoid dataset was collected from the Environmental Protection Agency's ECOTOX Knowledgebase, a database for ecotoxicology research. Neonicotinoids are a class of insecticides used widely in agriculture. The dataset that has been pulled includes all studies published on insects. Why might we be interested in the ecotoxicology of neonicotinoids on insects? Feel free to do a brief internet search if you feel you need more background information.

Answer: Neonics have been shown to have adverse effects on honey-bee populations and to also be harmful to birds by reducing overall insect populations.

3. The Niwot Ridge litter and woody debris dataset was collected from the National Ecological Observatory Network, which collectively includes 81 aquatic and terrestrial sites across 20 ecoclimatic domains. 32 of these sites sample forest litter and woody debris, and we will focus on the Niwot Ridge long-term ecological research (LTER) station in Colorado. Why might we be interested in studying litter and woody debris that falls to the ground in forests? Feel free to do a brief internet search if you feel you need more background information.

Answer: Looking at forest litter can be useful in understanding the successional stage of a forest. It can also be used to analyze the ecosystem and animals who depend on litter and woody debris to live. Lastly, it could be used for wildfire management. Forest with an overabundance of litter and woody debris could need a prescribed burn.

4. How is litter and woody debris sampled as part of the NEON network? Read the NEON\_Litterfall\_UserGuide.pdf document to learn more. List three pieces of salient information about the sampling methods here:

Answer: Litter was collected in sites where woody vegetation is greater than 2 meters tall. 40x40m and 20x20m plots with 10x10m and 1x1m nested subplots were created to sample the litter and woody debris. \* deciduous forests were sampled during senescence and evergreen forests were sampled year round \* In sites with greater than 50% aerial cover of woody vegetation greater than 2m in height, placement of litter traps is random \* litter is defined as material that is dropped from the forest canopy and has a butt end diameter < 2cm and a length < 50 cm

## Obtain basic summaries of your data (Neonics)

5. What are the dimensions of the dataset?

```
dim(Neonics)
```

```
## [1] 4623 30
```

6. Using the `summary` function on the "Effects" column, determine the most common effects that are studied. Why might these effects specifically be of interest?

```
summary(Neonics$Effect)
```

##	Accumulation	Avoidance	Behavior	Biochemistry
##	12	102	360	11
##	Cell(s)	Development	Enzyme(s)	Feeding behavior
##	9	136	62	255
##	Genetics	Growth	Histology	Hormone(s)
##	82	38	5	1
##	Immunological	Intoxication	Morphology	Mortality
##	16	12	22	1493
##	Physiology	Population	Reproduction	

##

7

1803

197

Answer: Population and mortality are the the most common effects studied. These are important to monitor the number of insects/honey-bees in the ecosystem. Severe declines in population or severe increases in mortality of insects/honey-bees can have disastrous effects on animals and plants that depend on these insects and honey bees.

7. Using the `summary` function, determine the six most commonly studied species in the dataset (common name). What do these species have in common, and why might they be of interest over other insects? Feel free to do a brief internet search for more information if needed.

```
summary(Neonics$Species.Common.Name)
```

##	Honey Bee	Parasitic Wasp
##	667	285
##	Buff Tailed Bumblebee	Carniolan Honey Bee
##	183	152
##	Bumble Bee	Italian Honeybee
##	140	113
##	Japanese Beetle	Asian Lady Beetle
##	94	76
##	Euonymus Scale	Wireworm
##	75	69
##	European Dark Bee	Minute Pirate Bug
##	66	62
##	Asian Citrus Psyllid	Parastic Wasp
##	60	58
##	Colorado Potato Beetle	Parasitoid Wasp
##	57	51
##	Erythrina Gall Wasp	Beetle Order
##	49	47
##	Snout Beetle Family, Weevil	Sevenspotted Lady Beetle
##	47	46
##	True Bug Order	Buff-tailed Bumblebee
##	45	39
##	Aphid Family	Cabbage Looper
##	38	38
##	Sweetpotato Whitefly	Braconid Wasp
##	37	33
##	Cotton Aphid	Predatory Mite
##	33	33
##	Ladybird Beetle Family	Parasitoid
##	30	30
##	Scarab Beetle	Spring Tiphia
##	29	29
##	Thrip Order	Ground Beetle Family
##	29	27
##	Rove Beetle Family	Tobacco Aphid
##	27	27
##	Chalcid Wasp	Convergent Lady Beetle
##	25	25
##	Stingless Bee	Spider/Mite Class
##	25	24
##	Tobacco Flea Beetle	Citrus Leafminer
##	24	23
##	Ladybird Beetle	Mason Bee

##	23	22
##	Mosquito	Argentine Ant
##	22	21
##	Beetle	Flatheaded Appletree Borer
##	21	20
##	Horned Oak Gall Wasp	Leaf Beetle Family
##	20	20
##	Potato Leafhopper	Tooth-necked Fungus Beetle
##	20	20
##	Codling Moth	Black-spotted Lady Beetle
##	19	18
##	Calico Scale	Fairyfly Parasitoid
##	18	18
##	Lady Beetle	Minute Parasitic Wasps
##	18	18
##	Mirid Bug	Mulberry Pyralid
##	18	18
##	Silkworm	Vedalia Beetle
##	18	18
##	Araneoid Spider Order	Bee Order
##	17	17
##	Egg Parasitoid	Insect Class
##	17	17
##	Moth And Butterfly Order	Oystershell Scale Parasitoid
##	17	17
##	Hemlock Woolly Adelgid Lady Beetle	Hemlock Woolly Adelgid
##	16	16
##	Mite	Onion Thrip
##	16	16
##	Western Flower Thrips	Corn Earworm
##	15	14
##	Green Peach Aphid	House Fly
##	14	14
##	Ox Beetle	Red Scale Parasite
##	14	14
##	Spined Soldier Bug	Armoured Scale Family
##	14	13
##	Diamondback Moth	Eulophid Wasp
##	13	13
##	Monarch Butterfly	Predatory Bug
##	13	13
##	Yellow Fever Mosquito	Braconid Parasitoid
##	13	12
##	Common Thrip	Eastern Subterranean Termite
##	12	12
##	Jassid	Mite Order
##	12	12
##	Pea Aphid	Pond Wolf Spider
##	12	12
##	Spotless Ladybird Beetle	Glasshouse Potato Wasp
##	11	10
##	Lacewing	Southern House Mosquito
##	10	10
##	Two Spotted Lady Beetle	Ant Family

```
##               10               9
##           Apple Maggot         (Other)
##               9             670
```

Answer: The Honey Bee, Parasitic Wasp, Buff Tailed Bumblebee, Carniolan Honey Bee, Bumble Bee, and Italian Honeybee are the six most studied species in the data set. All of these insects are pollinators and would be of importance to crops. Without these pollinators, crops would not be as productive.

8. Concentrations are always a numeric value. What is the class of `Conc.1..Author.` in the dataset, and why is it not numeric?

```
class(Neonics$Conc.1..Author.)
```

```
## [1] "factor"
```


Answer: It is a character. It could be because the concentrations are in different units.

## Explore your data graphically (Neonics)

9. Using `geom_freqpoly`, generate a plot of the number of studies conducted by publication year.

```
ggplot(Neonics) + geom_freqpoly(aes(x=Publication.Year))
```


```
## `stat_bin()` using `bins = 30`. Pick better value with `binwidth`.
```



10. Reproduce the same graph but now add a color aesthetic so that different `Test.Location` are displayed as different colors.

```
ggplot(Neonics) + geom_freqpoly(aes(x=Publication.Year, color = Test.Location))
```

```
## `stat_bin()` using `bins = 30`. Pick better value with `binwidth`.
```




Interpret this graph. What are the most common test locations, and do they differ over time?

Answer: The most common test locations are the lab and the natural field. The natural field has been pretty constant with a sharp spike around 2010. Lab locations have increase from 1990 to 2015 and decreased from 2015 to 2020.

11. Create a bar graph of Endpoint counts. What are the two most common end points, and how are they defined? Consult the ECOTOX\_CodeAppendix for more information.

```
ggplot(Neonics, aes(x=Endpoint)) +geom_bar() + theme(axis.text.x = element_text(angle = 90))
```



Answer: The two most common endpoints are LOEL and NOEL. LOEL is Lowest-observable-effect-level: lowest dose (concentration) producing effects that were significantly different and NOEL is No-observable-effect-level: highest dose (concentration) producing effects not significantly different from responses of controls.

## Explore your data (Litter)

12. Determine the class of collectDate. Is it a date? If not, change to a date and confirm the new class of the variable. Using the `unique` function, determine which dates litter was sampled in August 2018.

```
class(Litter$collectDate)
```

```
## [1] "factor"
```

```
Litter$collectDate <- as.Date(Litter$collectDate, format = "%Y-%m-%d")
```

```
class(Litter$collectDate)
```

```
## [1] "Date"
```

```
?unique
```

```
## starting httpd help server ... done
```

```
unique(Litter$collectDate)
```

```
## [1] "2018-08-02" "2018-08-30"
```

13. Using the `unique` function, determine how many plots were sampled at Niwot Ridge. How is the information obtained from `unique` different from that obtained from `summary`?

```
unique(Litter$plotID)
```

```
## [1] NIWO_061 NIWO_064 NIWO_067 NIWO_040 NIWO_041 NIWO_063 NIWO_047 NIWO_051  
## [9] NIWO_058 NIWO_046 NIWO_062 NIWO_057  
## 12 Levels: NIWO_040 NIWO_041 NIWO_046 NIWO_047 NIWO_051 NIWO_057 ... NIWO_067
```


```
summary(Litter$plotID)
```

```
## NIWO_040 NIWO_041 NIWO_046 NIWO_047 NIWO_051 NIWO_057 NIWO_058 NIWO_061  
##      20      19      18      15      14       8      16      17  
## NIWO_062 NIWO_063 NIWO_064 NIWO_067  
##      14      14      16      17
```

Answer: 12 different plots were sampled at NIWOT Ridge. The unique function tells you how many unique values are in a column. It eliminates duplicate values. The summary function tells us how many samples in each plot were taken.

14. Create a bar graph of functionalGroup counts. This shows you what type of litter is collected at the Niwot Ridge sites. Notice that litter types are fairly equally distributed across the Niwot Ridge sites.


```
ggplot(Litter, aes(x = functionalGroup)) + geom_bar()
```




15. Using `geom_boxplot` and `geom_violin`, create a boxplot and a violin plot of `dryMass` by `functionalGroup`.

```
ggplot(Litter) +  
  geom_boxplot(aes(x = functionalGroup, y = dryMass))
```





```
ggplot(Litter) +  
  geom_violin(aes(x = functionalGroup, y = dryMass))
```



Why is the boxplot a more effective visualization option than the violin plot in this case?

Answer: The box plot allows us to see the mean and interquartile ranges whereas the violin plot shows us the ammount of points in those ranges. The violin plot is not very useful in this situation becuase there is not an abundant amount of dry mass values for the functional group. Thus the box plot gives us a better visual becuase it shows the range of values.

What type(s) of litter tend to have the highest biomass at these sites?

Answer: Needles and “mixed” tend to have the highest biomass at the sites.