

## Appendix 2. Cyclic groups, group generators

The bonus problem 2 (1p) in Moodle is based on this appendix.

Diffie Hellman key exchange and ECDHE key exchange are both based on cyclic groups. In this appendix, we review the concepts of group and cyclic group and use their properties to find a group generator of multiplicative group  $Z_p$ .

### Basic concepts

---

**Definition 1:** A set  $G$  with operation  $*$  defined in  $G$  is called a group, if it has the following properties.

G1)  $a*b \in G$  for all  $a, b \in G$

G2)  $(a*b)*c = a*(b*c)$  for  $a, b, c \in G$

G3)  $G$  has a neutral element  $e$  for which  $a*e = e*a = a$  for all  $a \in G$

G4) Every element  $a \in G$  has an inverse element  $a^{-1}$  for which  $a^{-1} * a = a * a^{-1} = e$

---

Example : The number set  $Z_p^*$  of integers  $\{1, 2, \dots, p-1\}$  is a group where the group operation is multiplication  $a*b \pmod{p}$ .

The neutral element of  $Z_p^*$  is number 1. All elements  $a \in Z_p^*$  have an inverse element  $a^{-1} \in Z_p^*$ . Indeed, modulus  $p$  is a prime and therefore  $\gcd(a, p) = 1$  for all  $a \in Z_p^*$ . Inverse can be calculated using Euclid's extended GDC algorithm.

---

**Definition 2:** Let  $G$  be a finite group of  $n$  elements.

If there is such an element  $g \in G$  that the set of its powers  $\{g, g^2, \dots, g^n\}$  includes all elements of  $G$ , we say that group  $G$  is **cyclic** and the element  $g$  is a **generator** of group  $G$ .

---

If  $p$  is prime, the number set  $Z_p^*$  of integers  $\{1, 2, \dots, p-1\}$  is a cyclic group. Diffie Hellman key exchange uses this group.

Example: Group  $Z_{13}^*$  is cyclic. For example number 7 is a generator of  $Z_{13}^*$ , because the set of powers  $\{7^1 \pmod{13}, 7^2 \pmod{13}, \dots, 7^{12} \pmod{13}\} = \{7, 10, 5, 9, 11, 12, 6, 3, 8, 4, 2, 1\}$  contains all the elements of  $Z_{13}^*$ .

---

**Definition 3:** If  $H$  is a group and  $H$  is a subset of group  $G$ , we say that  $H$  is a **subgroup** of group  $G$ .

---

Example: Set  $H = \{1, 3, 9\}$  is a subgroup of  $Z_{13}^*$ .

The multiplication table, where operation is  $a*b \pmod{13}$  shows that all group properties G1, ..., G4 hold.

	1	3	9
1	1	3	9
3	3	9	1
9	9	1	3

This group is cyclic, because 3 generate all its elements:  $\{3, 3^2, 3^3\} \pmod{13} = \{3, 9, 1\}$

## Properties of multiplicative groups $Z_p^*$

1. All elements  $a$  of  $Z_p^* = \{1, 2, \dots, p-1\}$  generate a cyclic subgroup of  $Z_p^*$
2. The size of the subgroup generated by element  $a$  is called **the order of element  $a$**  and denoted **Ord( $a$ )**.
3. Lagrange's theorem: Ord( $a$ ) is a divisor of  $p-1$  for all  $a \in Z_p^*$ .
4. If Ord( $g$ ) is  $p-1$ , *element  $g$  is called a generator of  $Z_p^*$  or primitive root of  $Z_p^*$*

The next property follows from properties 1 – 4.

5. Let  $d_1, d_2, \dots, d_n$  be the list of divisors of  $p-1$  in ascending order (where  $d_1=1$  and  $d_n=p-1$ )

Then, the generators are those elements  $g$  of  $Z_p^*$ , for which only the last power of  $g$  in the sequence  $g^{d_1}, g^{d_2}, \dots, g^{d_n}$  equals 1 (mod  $p$ ).

Example. Below is a table of powers ( $a^k \bmod 11$ ) of elements of  $Z_{11}^*$ .

$a$	$a^1$	$a^2$	$a^3$	$a^4$	$a^5$	$a^6$	$a^7$	$a^8$	$a^9$	$a^{10}$	Subgroup size
1	1	1	1	1	1	1	1	1	1	1	1
2	2	4	8	5	10	9	7	3	6	1	10 (generator)
3	3	9	5	4	1	3	9	5	4	1	5
4	4	5	9	3	1	4	5	9	3	1	5
5	5	3	4	9	1	5	3	4	9	1	5
6	6	3	7	9	10	5	8	4	2	1	10 (generator)
7	7	5	2	3	10	4	6	9	8	1	10 (generator)
8	8	9	6	4	10	3	2	5	7	1	10 (generator)
9	9	4	3	5	1	9	4	3	5	1	5
10	10	1	10	1	10	1	10	1	10	1	2

There are 4 generators:  $\{2, 6, 7, 8\}$

All the subgroup sizes 1, 2, 5, and 10 are divisors of 10 (in general  $p-1$ ) as Lagrange's theorem predicts.

The previous theory provides a tool for finding a group generator of  $Z_p^*$

### Case1: Generator test, when $p$ is a relatively small prime

Let  $d_1, d_2, \dots, d_n$  be the divisors of  $p-1$  in ascending order.

An integer  $g \in Z_p^*$  is a generator if and only if the last integer in the list of powers  $g^{d_1}, g^{d_2}, \dots, g^{d_n}$  equals 1.

Example: a) Test if number 5 a generator of  $Z_{29}^*$  or not?

Divisors of  $p-1 = 28$  are  $\{1, 2, 4, 7, 14, 28\}$ . Raising 5 to all the powers in the divisor list gives (using WolframAlpha)  $5^{\{1,2,4,7,14,28\} \bmod 29} = \{5, 25, 16, 28, 1, 1\}$ . Number 5 is not a generator, because the last two numbers are ones. (WolframAlpha allows to calculate all powers with only one command line)

$2^{\{1,2,4,7,14,28\} \bmod 29} = \{2, 4, 16, 12, 28, 1\}$ . Number 2 is a generator of  $Z_{29}^*$

## Case2: Finding a generator, when p is very large prime

When prime  $p$  is very large, for example 1000 bit integer, it is often impossible to factor  $p - 1$ . Some divisors are trivial: 1, 2 and  $(p-1)/2$  (2 is a divisor, because  $p - 1$  is even). However, if  $p - 1$  has very large factors, some of the divisors may be not be found.

**Example.** If  $p = 265738830135992486377941683556469254997964098756853$ , then  $p - 1 = 265738830135992486377941683556469254997964098756852$ . Attempts to factor  $p - 1$  further fail.

## Strong primes

---

**Definition:** A prime  $p$  is called a "**strong prime**", if also  $(p-1)/2$  is a prime.

In other words: If  $p$  is a strong prime, then  $p - 1$  has only four divisors:  $\{1, 2, (p-1)/2, p - 1\}$

---

It is recommended that Diffie-Hellman key exchange protocol should use strong primes as modulus  $p$ . To find these primes, one can write a Python program or utilize the help of chatGPT. Such a program generated 1033 primes, out of which 75 were strong primes within the range of 10000-20000. In practise, the system parameter  $g$  can also be any element of  $Z_p^*$  that generates an adequately large subgroup.

Generator test is easy to do for  $Z_p^*$ , if  $p$  is a strong prime.

Example. Integer  $p = 200087$  is a strong prime. Divisors of  $p - 1$  are  $\{1, 2, 100043, 200086\}$ . Test if number 5 is a generator of  $Z_{200087}^*$ .

Calculation of powers  $5^{\{1, 2, 100043, 200086\}} \bmod 200087$  gives  $\{5, 25, 200086, 1\}$ , which shows that number 5 is a generator of  $Z_{200087}^*$ .

**Rule: Number of generators of  $Z_p^* = \phi(p-1)$ , where  $\phi$  is Euler's totient function.**

(More general rule is that for any divisor  $d$  of  $p - 1$  the number of elements of order  $d$  is  $\phi(d)$ )

If  $p$  is a strong prime, then  $p - 1 = 2 \cdot r$ , where  $r = (p-1)/2$  is also a prime. Now  $\phi(2 \cdot r) = (2-1)(r-1) = (p-1)/2 - 1 \approx (p-1)/2$ .

Hence: If  $p$  is a large prime, nearly 50% of elements of  $Z_p^*$  are generators.

---