

# trapezoid: The Trapezoidal Distribution

Jeremy Thoms Hetzel

September 12, 2011

## 1 Introduction

The trapezoidal distribution is defined by minimum, lower mode, upper mode, and maximum parameters. The generalized trapezoidal distribution adds three more parameters: the growth rate, decay rate, and boundary ratio parameters. Van Dorp and Kotz[2] and van Dorp and colleagues[1] formally describe the generalized trapezoidal distribution, representing the minimum, lower mode, upper mode, maximum, growth rate, decay rate, and boundary ratio with parameters  $a, b, c, d, m, n$ , and  $\alpha$ , respectively.

The probability density function of the generalized trapezoidal distribution with parameters  $a, b, c, d, m, n$ , and  $\alpha$  is given by:

$$f_X(x | \theta) = \mathcal{C}(\Theta) \times \begin{cases} \alpha \left( \frac{x-a}{b-a} \right)^{m-1}, & \text{for } a \leq x < b \\ (1-\alpha) \left( \frac{x-b}{c-b} \right) + \alpha, & \text{for } b \leq x < c \\ \left( \frac{d-x}{d-c} \right)^{n-1}, & \text{for } c \leq x \leq d \end{cases}$$

with the normalizing constant  $\mathcal{C}(\Theta)$  defined as:

$$\mathcal{C}(\Theta) = \frac{2mn}{2\alpha(b-a)n + (\alpha+1)(c-b)mn + 2(d-c)m}$$

and where the parameter vector  $\Theta = \{a, b, c, d, m, n, \alpha\}$ ,  $a \leq b \leq c \leq d$ , and  $m, n, \alpha > 0$ .

The **trapezoid** package provides functions for the probability density function (**dtrapezoid**), cumulative distribution function (**ptrapezoid**), quantile function (**qtrapezoid**), and random generation (**rtrapezoid**). The parameters  $a, b, c, d, m, n$ , and  $\alpha$  are specified by the arguments **min**, **mode1**, **mode2**, **max**, **n1**, **n3**, and **alpha**, respectively. The argument names were chosen to avoid conflicts with names that commonly have specific meaning in R functions, such as **c** and **n**.


## 2 Examples

### 2.1 Trapezoid

The generalized trapezoidal distribution simplifies to the trapezoidal distribution when  $m = n = 2$  and  $\alpha = 1$ .

```
> require(trapezoid)
> require(ggplot2)
> x <- seq(from = 0, to = 1, by = 0.01)
> density <- dttrapezoid(x, min = 0, mode1 = 1/3, mode2 = 2/3,
  max = 1, n1 = 2, n3 = 2, alpha = 1)
> trapezoid <- ggplot(data = data.frame(x, density), aes(x = x,
  y = density)) + geom_line() + theme_bw()
> print(trapezoid)
```

Figure 1: A trapezoidal distribution.




## 2.2 Triangle

The trapezoidal distribution further simplifies to the triangular distribution when  $b = c$ .

```
> x <- seq(from = 0, to = 1, by = 0.01)
> density <- dtrapezoid(x, min = 0, mode1 = 1/2, mode2 = 1/2,
  max = 1, n1 = 2, n3 = 2, alpha = 1)
> triangle <- ggplot(data = data.frame(x, density), aes(x = x,
  y = density)) + geom_line() + theme_bw()
> print(triangle)
```

Figure 2: A triangular distribution.




## 2.3 Generalized trapezoidal distribution

Parameters  $m$ ,  $n$ , and  $\alpha$  control the growth rate, decay rate, and boundary ratio, respectively, of the distribution. In the `trapezoid` package, these

parameters are controlled by the `n1`, `n3`, and `alpha` arguments. To demonstrate the effects of these three parameters, van Dorp and Kotz[2] generated eight distributions with varying parameter values. The distributions are approximately re-generated here.

```
> x <- seq(from = 0, to = 1, by = 0.01)
> arguments <- list()
> arguments[["A"]] <- list(x = x, n1 = 2, n3 = 2, alpha = 0.8)
> arguments[["B"]] <- list(x = x, n1 = 1.5, n3 = 1.5, alpha = 1)
> arguments[["C"]] <- list(x = x, n1 = 2.5, n3 = 2.5, alpha = 1.5)
> arguments[["D"]] <- list(x = x, n1 = 1.5, n3 = 2.5, alpha = 0.5)
> arguments[["E"]] <- list(x = x, n1 = 2.5, n3 = 1.5, alpha = 1)
> arguments[["F"]] <- list(x = x, n1 = 0.5, n3 = 0.5, alpha = 1.5)
> arguments[["G"]] <- list(x = x, n1 = 1.5, n3 = 0.5, alpha = 0.5)
> arguments[["H"]] <- list(x = x, n1 = 2.5, n3 = 0.5, alpha = 1)
> arguments[["I"]] <- list(x = x, n1 = 0.5, n3 = 1.5, alpha = 1.5)
> arguments[["J"]] <- list(x = x, n1 = 0.5, n3 = 2.5, alpha = 0.5)
> plot.data <- ldply(arguments, function(z) {
  x <- z$x
  density <- dtrapezoid(x = z$x, min = 0, mode1 = 0.2,
    mode2 = 0.8, max = 1, n1 = z$n1, n3 = z$n3, alpha = z$alpha)
  args <- paste("n1 = ", z$n1, ", n3 = ", z$n3, ", alpha = ",
    z$alpha, sep = "", collapse = "")
  out <- data.frame(x, density, args)
})
> plot.data$label <- paste(plot.data$id, ": ", plot.data$args,
  sep = "")
> generalizedTrapezoids <- ggplot(data = plot.data, aes(x = x,
  y = density)) + geom_line() + theme_bw() + facet_wrap(~label,
  ncol = 2, scales = "free_y")
> print(generalizedTrapezoids)
```

Figure 3: Examples of generalized trapezoidal distributions.



## References

- [1] J. R. van Dorp, S. C. Rambaud, J. G. Perez, and R. H. Pleguezuelo. An Elicitation Procedure for the Generalized Trapezoidal Distribution with a Uniform Central Stage. *Decision Analysis*, 4(3):156–166, September 2007.
- [2] J René van Dorp and Samuel Kotz. Generalized Trapezoidal Distributions. *Metrika*, 58(1):85–97, 2003.