

Barcast Model

March 27, 2014

1 Barcast Model

1.1 Data Model

$$\mathbf{W}_t = \mathbf{H}_t \mathbf{T}_t + \mathbf{B}_t + \boldsymbol{\eta}_t \quad \boldsymbol{\eta}_t \sim \mathcal{N}(\mathbf{0}, \boldsymbol{\Sigma}_t) \quad \boldsymbol{\Sigma}_t = \begin{pmatrix} \boldsymbol{\Sigma}_{It} & \mathbf{0} \\ \mathbf{0} & \boldsymbol{\Sigma}_{Pt} \end{pmatrix}$$

where $\mathbf{W}_t = \begin{pmatrix} \mathbf{W}_{It} \\ \mathbf{W}_{Pt} \end{pmatrix}$ is a vector of instrumental observations \mathbf{W}_{It} and proxy observations \mathbf{W}_{Pt} at time t , $\mathbf{T}_t = \begin{pmatrix} \mathbf{T}_{It} \\ \mathbf{T}_{Pt} \end{pmatrix}$ is a vector of latent climate variables, $\mathbf{H}_t = \begin{pmatrix} \mathbf{H}_{It} \\ \beta_1 \mathbf{H}_{Pt} \end{pmatrix}$, $\mathbf{B}_t = \begin{pmatrix} \mathbf{H}_{It} \\ \mathbf{H}_{Pt} \end{pmatrix} \begin{pmatrix} \mathbf{0}_{N_I} \\ \beta_0 \mathbf{1}_{N_P} \end{pmatrix}$, $\boldsymbol{\Sigma}_{It} = \tau_I^2 \mathbf{I}_{N_{It}}$, $\boldsymbol{\Sigma}_{Pt} = \tau_P^2 \mathbf{I}_{N_{Pt}}$, N_I is the total number of instrumental observations, N_P is the total number of proxy observations, N_{It} is the number of instrumental observations at time t , N_{Pt} is the number of proxy observations at time t , \mathbf{H}_{It} and \mathbf{H}_{Pt} are selection matrices indicating whether the instrumental or proxy variable was measured at time t , and β_0 and β_1 are regression coefficients relating the proxy observations and latent field.

1.2 Process Model

$$\mathbf{T}_t - \mu \mathbf{1}_N = \alpha (\mathbf{T}_{t-1} - \mu \mathbf{1}_N) + \boldsymbol{\epsilon}_t \quad \boldsymbol{\epsilon}_t \sim \mathcal{N}(\mathbf{0}, \boldsymbol{\Sigma}_\epsilon) \quad \boldsymbol{\Sigma}_\epsilon = \sigma^2 \exp(-\phi \mathbf{D})$$

where \mathbf{D} is the distance matrix between observation points

1.3 Parameter Model

$$\begin{aligned}
\mathbf{T}_0 &\sim \mathcal{N}(\tilde{\boldsymbol{\mu}}_0, \tilde{\boldsymbol{\Sigma}}_0) & \tilde{\boldsymbol{\mu}}_0 &= \mathbf{0} & \tilde{\boldsymbol{\Sigma}}_0 &= \tilde{\sigma}_0^2 \mathbf{I} \\
\alpha &\sim \mathcal{U}(0, 1,) \\
\mu &\sim \mathcal{N}(\mu_0, \sigma_0^2) & \text{For PDSI } \mu_0 &= 0 & \sigma_0^2 &= 1 \\
\sigma^2 &\sim \text{IG}(\alpha_{\sigma^2}, \beta_{\sigma^2}) \\
\phi &\sim \text{IG}(\alpha_\phi, \beta_\phi) \\
\tau_I &\sim \text{IG}(\alpha_I, \beta_I) \\
\tau_P &\sim \text{IG}(\alpha_P, \beta_P) \\
\beta_1 &\sim \mathcal{N}(\mu_{\beta_1}, \sigma_{\beta_1}^2) & \mu_{\beta_1} &= \left(\frac{(1 - \tau_P^2)(1 - \alpha^2)}{\sigma^2} \right)^{-\frac{1}{2}} & \sigma_{\beta_1}^2 &= 8 \\
\beta_0 &\sim \mathcal{N}(\mu_{\beta_0}, \sigma_{\beta_0}^2) & \mu_{\beta_0} &= -\mu_{\beta_1} & \sigma_{\beta_1}^2 &= 8
\end{aligned}$$

Note that the prior values are set as the prior modes (e.g. μ_{β_1} is a function of the prior modes for τ_P^2, α , and σ^2).

2 Posterior

$$\begin{aligned}
\prod_{t=1}^T [\mathbf{T}_t, \beta_0, \beta_1, \mu, \alpha, \tau_I^2, \tau_P^2, \sigma^2, \phi | \mathbf{W}_t, T] &\propto \prod_{t=1}^T [\mathbf{W}_t | \mathbf{T}_t, \beta_0, \beta_1, \tau_I^2, \tau_P^2] \propto [\mathbf{T}_t | \mathbf{T}_t, \mu, \alpha, \sigma^2, \phi] [\mathbf{T}_0] \\
&\times [\beta_0] [\beta_1] [\tau_I^2] [\tau_P^2] [\mu] [\alpha] [\sigma^2] [\phi]
\end{aligned}$$

3 Full Conditionals

3.1 Full Conditional for \mathbf{T}_0

$$\begin{aligned}
[\mathbf{T}_0 | \cdot] &\propto [\mathbf{T}_1 | \mathbf{T}_0, \mu, \alpha, \sigma^2, \phi] [\mathbf{T}_0] \\
&\propto \exp \left\{ -\frac{1}{2} (\mathbf{T}_1 - \alpha \mathbf{T}_0 - (1 - \alpha) \mu \mathbf{1}_N)^T \boldsymbol{\Sigma}_\epsilon^{-1} (\mathbf{T}_1 - \alpha \mathbf{T}_0 - (1 - \alpha) \mu \mathbf{1}_N) \right\} \\
&\quad \times \exp \left\{ -\frac{1}{2} (\mathbf{T}_0 - \tilde{\boldsymbol{\mu}}_0)^T \tilde{\boldsymbol{\Sigma}}_0^{-1} (\mathbf{T}_0 - \tilde{\boldsymbol{\mu}}_0) \right\} \\
&\propto \exp \left\{ -\frac{1}{2} \left[\mathbf{T}_0^T \left(\alpha^2 \boldsymbol{\Sigma}_\epsilon^{-1} + \tilde{\boldsymbol{\Sigma}}_0^{-1} \right) \mathbf{T}_0 - 2 \mathbf{T}_0^T \left(\alpha \boldsymbol{\Sigma}_\epsilon^{-1} (\mathbf{T}_1 - (1 - \alpha) \mu \mathbf{1}_N) + \tilde{\boldsymbol{\Sigma}}_0^{-1} \tilde{\boldsymbol{\mu}} \right) \right] \right\}
\end{aligned}$$

which is $\text{MVN}(\mathbf{A}^{-1} \mathbf{b}, \mathbf{A}^{-1})$ with

$$\mathbf{A}^{-1} = \left(\alpha^2 \boldsymbol{\Sigma}_\epsilon^{-1} + \tilde{\boldsymbol{\Sigma}}_0^{-1} \right)^{-1}$$

$$\mathbf{b} = \alpha \boldsymbol{\Sigma}_\epsilon^{-1} (\mathbf{T}_1 - (1 - \alpha) \mu \mathbf{1}_N) + \tilde{\boldsymbol{\Sigma}}_0^{-1} \tilde{\boldsymbol{\mu}}.$$

3.2 Full Conditional for \mathbf{T}_t

For $t = 1, \dots, T - 1$,

$$\begin{aligned} [\mathbf{T}_t | \cdot] &\propto [\mathbf{W}_t | \mathbf{T}_t, \beta_0, \beta_1, \tau_I^2, \tau_P^2] [\mathbf{T}_t | \mathbf{T}_{t-1}, \mu, \alpha, \sigma^2, \phi] [\mathbf{T}_{t+1} | \mathbf{T}_t, \mu, \alpha, \sigma^2, \phi] \\ &\propto \exp \left\{ -\frac{1}{2} (\mathbf{W}_t - (\mathbf{H}_t \mathbf{T}_t + \mathbf{B}_t))^T \boldsymbol{\Sigma}_t^{-1} (\mathbf{W}_t - (\mathbf{H}_t \mathbf{T}_t + \mathbf{B}_t)) \right\} \\ &\quad \times \exp \left\{ -\frac{1}{2} (\mathbf{T}_t - \alpha \mathbf{T}_{t-1} - (1 - \alpha) \mu \mathbf{1}_N)^T \boldsymbol{\Sigma}_\epsilon^{-1} (\mathbf{T}_t - \alpha \mathbf{T}_{t-1} - (1 - \alpha) \mu \mathbf{1}_N) \right\} \\ &\quad \times \exp \left\{ -\frac{1}{2} (\mathbf{T}_{t+1} - \alpha \mathbf{T}_t - (1 - \alpha) \mu \mathbf{1}_N)^T \boldsymbol{\Sigma}_\epsilon^{-1} (\mathbf{T}_{t+1} - \alpha \mathbf{T}_t - (1 - \alpha) \mu \mathbf{1}_N) \right\} \\ &\propto \exp \left\{ -\frac{1}{2} \mathbf{T}_t^T \left(\mathbf{H}_t^T \boldsymbol{\Sigma}_t^{-1} \mathbf{H}_t + (\alpha^2 + 1) \boldsymbol{\Sigma}_\epsilon^{-1} \right) \mathbf{T}_t \right\} \\ &\quad \times \exp \left\{ -\mathbf{T}_t^T \left(\mathbf{H}_t^T \boldsymbol{\Sigma}_t^{-1} (\mathbf{W}_t - \mathbf{B}_t) + \boldsymbol{\Sigma}_\epsilon^{-1} (\alpha \mathbf{T}_{t-1} + (1 - \alpha) \mu \mathbf{1}_N) + \alpha \boldsymbol{\Sigma}_\epsilon^{-1} (\mathbf{T}_{t+1} - (1 - \alpha) \mu \mathbf{1}_N) \right) \right\} \end{aligned}$$

which is $\text{MVN}(\mathbf{A}^{-1} \mathbf{b}, \mathbf{A}^{-1})$ with

$$\mathbf{A}^{-1} = \left(\mathbf{H}_t^T \boldsymbol{\Sigma}_t^{-1} \mathbf{H}_t + (\alpha^2 + 1) \boldsymbol{\Sigma}_\epsilon^{-1} \right)^{-1}$$

$$\mathbf{b} = \mathbf{H}_t^T \boldsymbol{\Sigma}_t^{-1} (\mathbf{W}_t - \mathbf{B}_t) + \boldsymbol{\Sigma}_\epsilon^{-1} \left(\alpha (\mathbf{T}_{t+1} + \mathbf{T}_{t-1}) + (1 - \alpha)^2 \mu \mathbf{1}_N \right).$$

For $t = T$,

$$\begin{aligned} [\mathbf{T}_T | \cdot] &\propto [\mathbf{W}_T | \mathbf{T}_T, \beta_0, \beta_1, \tau_I^2, \tau_P^2] [\mathbf{T}_T | \mathbf{T}_{T-1}, \mu, \alpha, \sigma^2, \phi] \\ &\propto \exp \left\{ -\frac{1}{2} (\mathbf{W}_T - (\mathbf{H}_T \mathbf{T}_T - \mathbf{B}_T))^T \boldsymbol{\Sigma}_T^{-1} (\mathbf{W}_T - (\mathbf{H}_T \mathbf{T}_T - \mathbf{B}_T)) \right\} \\ &\quad \times \exp \left\{ -\frac{1}{2} (\mathbf{T}_T - \alpha \mathbf{T}_{T-1} - (1 - \alpha) \mu \mathbf{1}_N)^T \boldsymbol{\Sigma}_\epsilon^{-1} (\mathbf{T}_T - \alpha \mathbf{T}_{T-1} - (1 - \alpha) \mu \mathbf{1}_N) \right\} \\ &\propto \exp \left\{ -\frac{1}{2} \mathbf{T}_T^T \left(\mathbf{H}_T^T \boldsymbol{\Sigma}_T^{-1} \mathbf{H}_T + \boldsymbol{\Sigma}_\epsilon^{-1} \right) \mathbf{T}_T \right\} \\ &\quad \times \exp \left\{ -\mathbf{T}_T^T \left(\mathbf{H}_T^T \boldsymbol{\Sigma}_T^{-1} (\mathbf{W}_T - \mathbf{B}_T) + \boldsymbol{\Sigma}_\epsilon^{-1} (\alpha \mathbf{T}_{T-1} + (1 - \alpha) \mu \mathbf{1}_N) \right) \right\} \end{aligned}$$

which is $\text{MVN}(\mathbf{A}^{-1}\mathbf{b}, \mathbf{A}^{-1})$ with

$$\begin{aligned}\mathbf{A}^{-1} &= \left(\mathbf{H}_T^T \Sigma_T^{-1} \mathbf{H}_T + \Sigma_\epsilon^{-1} \right)^{-1} \\ \mathbf{b} &= \mathbf{H}_T^T \Sigma_T^{-1} (\mathbf{W}_T - \mathbf{B}_T) + \Sigma_\epsilon^{-1} (\alpha \mathbf{T}_{T-1} + (1 - \alpha) \mu \mathbf{1}_N).\end{aligned}$$

3.3 Full conditional for β_0

$$\begin{aligned}[\beta_0|\cdot] &\propto \prod_{t=1}^T [\mathbf{W}_t | \mathbf{T}_t, \beta_0, \beta_1, \tau_P^2, \tau_P^2] [\beta_0] \\ &\propto \prod_{t=1}^T \exp \left\{ -\frac{1}{2} (\mathbf{W}_t - (\mathbf{H}_t \mathbf{T}_t + \mathbf{B}_t))^T \Sigma_t^{-1} (\mathbf{W}_t - (\mathbf{H}_t \mathbf{T}_t + \mathbf{B}_t)) \right\} \\ &\quad \times \exp \left\{ -\frac{1}{2} \frac{(\beta_0 - \mu_{\beta_0})^2}{\sigma_{\beta_0}^2} \right\} \\ &\propto \exp \left\{ -\frac{1}{2} \beta_0^2 \left(\sum_{t=1}^T \mathbf{1}_{N_{Pt}}^T \Sigma_{Pt}^{-1} \mathbf{1}_{N_{Pt}} + \frac{1}{\sigma_{\beta_0}^2} \right) - \beta_0 \left(\sum_{t=1}^T \mathbf{1}_{N_{Pt}}^T \Sigma_{Pt}^{-1} (\mathbf{W}_{Pt} - \beta_1 \mathbf{H}_{Pt} \mathbf{T}_t) + \frac{\mu_{\beta_0}}{\sigma_{\beta_0}^2} \right) \right\} \\ &\propto \exp \left\{ -\frac{1}{2} \beta_0^2 \left(\frac{M_P}{\tau_P^2} + \frac{1}{\sigma_{\beta_0}^2} \right) - \beta_0 \left(\frac{1}{\tau_P^2} \sum_{t=1}^T \mathbf{1}_{N_{Pt}}^T (\mathbf{W}_{Pt} - \beta_1 \mathbf{H}_{Pt} \mathbf{T}_t) + \frac{\mu_{\beta_0}}{\sigma_{\beta_0}^2} \right) \right\}\end{aligned}$$

which is $\text{MVN}(\mathbf{A}^{-1}\mathbf{b}, \mathbf{A}^{-1})$ with

$$\begin{aligned}\mathbf{A}^{-1} &= \left(\frac{M_P}{\tau_P^2} + \frac{1}{\sigma_{\beta_0}^2} \right)^{-1} \\ \mathbf{b} &= \frac{1}{\tau_P^2} \sum_{t=1}^T \mathbf{1}_{N_{Pt}}^T (\mathbf{W}_{Pt} - \beta_1 \mathbf{H}_{Pt} \mathbf{T}_t) + \frac{\mu_{\beta_0}}{\sigma_{\beta_0}^2}.\end{aligned}$$

3.4 Full conditional for β_1

$$\begin{aligned}
[\beta_1|\cdot] &\propto \prod_{t=1}^T [\mathbf{W}_t | \mathbf{T}_t, \beta_0, \beta_1, \tau_I^2, \tau_P^2] [\beta_1] \\
&\propto \prod_{t=1}^T \exp \left\{ -\frac{1}{2} (\mathbf{W}_t - (\mathbf{H}_t \mathbf{T}_t + \mathbf{B}_t))^T \boldsymbol{\Sigma}_t^{-1} (\mathbf{W}_t - (\mathbf{H}_t \mathbf{T}_t + \mathbf{B}_t)) \right\} \\
&\quad \times \exp \left\{ -\frac{1}{2} \frac{(\beta_1 - \mu_{\beta_1})^2}{\sigma_{\beta_1}^2} \right\} \\
&\propto \exp \left\{ -\frac{1}{2} \beta_1^2 \left(\sum_{t=1}^T (\mathbf{H}_{Pt} \mathbf{T}_t)^T \boldsymbol{\Sigma}_{Pt}^{-1} (\mathbf{H}_{Pt} \mathbf{T}_t) + \frac{1}{\sigma_{\beta_1}^2} \right) \right\} \\
&\quad \times \exp \left\{ -\beta_1 \left(\sum_{t=1}^T (\mathbf{H}_{Pt} \mathbf{T}_t)^T \boldsymbol{\Sigma}_{Pt}^{-1} (\mathbf{W}_{Pt} - \beta_0 \mathbf{1}_{N_{Pt}}) + \frac{\mu_{\beta_1}}{\sigma_{\beta_1}^2} \right) \right\} \\
&\propto \exp \left\{ -\frac{1}{2} \beta_1^2 \left(\frac{1}{\tau_P^2} \sum_{t=1}^T (\mathbf{H}_{Pt} \mathbf{T}_t)^T \boldsymbol{\Sigma}_{Pt}^{-1} (\mathbf{H}_{Pt} \mathbf{T}_t) + \frac{1}{\sigma_{\beta_1}^2} \right) \right\} \\
&\quad \times \exp \left\{ -\beta_1 \left(\frac{1}{\tau_P^2} \sum_{t=1}^T (\mathbf{H}_{Pt} \mathbf{T}_t)^T (\mathbf{W}_{Pt} - \beta_0 \mathbf{1}_{N_{Pt}}) + \frac{\mu_{\beta_1}}{\sigma_{\beta_1}^2} \right) \right\}
\end{aligned}$$

which is $\text{MVN}(\mathbf{A}^{-1} \mathbf{b}, \mathbf{A}^{-1})$ with

$$\begin{aligned}
\mathbf{A}^{-1} &= \left(\frac{1}{\tau_P^2} \sum_{t=1}^T (\mathbf{H}_{Pt} \mathbf{T}_t)^T (\mathbf{H}_{Pt} \mathbf{T}_t) + \frac{1}{\sigma_{\beta_1}^2} \right)^{-1} \\
\mathbf{b} &= \frac{1}{\tau_P^2} \sum_{t=1}^T (\mathbf{H}_{Pt} \mathbf{T}_t)^T (\mathbf{W}_{Pt} - \beta_0 \mathbf{1}_{N_{Pt}}) + \frac{\mu_{\beta_1}}{\sigma_{\beta_1}^2}.
\end{aligned}$$

NOTE: My derivation includes \mathbf{H}_{Pt} in the \mathbf{A}^{-1} and \mathbf{b} terms but is not present in Martin's paper... This is worth pursuing further. It seems that Martin uses \mathbf{T}_{Pt} to denote $\mathbf{H}_{Pt} \mathbf{T}_t$??

3.5 Full conditional for μ

$$\begin{aligned}
[\mu|\cdot] &\propto \prod_{t=1}^T [\mathbf{T}_t | \mathbf{T}_{t-1}, \mu, \alpha, \sigma^2, \phi] [\mu] \\
&\propto \prod_{t=1}^T \exp \left\{ -\frac{1}{2} (\mathbf{T}_t - \alpha \mathbf{T}_{t-1} - (1-\alpha) \mu \mathbf{1}_N)^T \boldsymbol{\Sigma}_\epsilon^{-1} (\mathbf{T}_t - \alpha \mathbf{T}_{t-1} - (1-\alpha) \mu \mathbf{1}_N) \right\} \\
&\quad \times \exp \left\{ -\frac{1}{2} \frac{(\mu - \mu_0)^2}{\sigma_0^2} \right\} \\
&\propto \exp \left\{ -\frac{1}{2} \mu^2 \left(\sum_{t=1}^T (1-\alpha)^2 \mathbf{1}_N^T \boldsymbol{\Sigma}_\epsilon^{-1} \mathbf{1}_N + \frac{1}{\sigma_0^2} \right) - \mu \left((1-\alpha) \mathbf{1}_N^T \boldsymbol{\Sigma}_\epsilon^{-1} \sum_{t=1}^T (\mathbf{T}_t - \alpha \mathbf{T}_{t-1}) + \frac{\mu_0}{\sigma_0^2} \right) \right\} \\
&\propto \exp \left\{ -\frac{1}{2} \mu^2 \left(T(1-\alpha)^2 \frac{N}{\sigma^2} + \frac{1}{\sigma_0^2} \right) - \mu \left(\frac{(1-\alpha)}{\sigma^2} \mathbf{1}_N^T \sum_{t=1}^T (\mathbf{T}_t - \alpha \mathbf{T}_{t-1}) + \frac{\mu_0}{\sigma_0^2} \right) \right\}
\end{aligned}$$

which is $\text{MVN}(\mathbf{A}^{-1} \mathbf{b}, \mathbf{A}^{-1})$ with

$$\begin{aligned}
\mathbf{A}^{-1} &= \left(T(1-\alpha)^2 \frac{N}{\sigma^2} + \frac{1}{\sigma_0^2} \right)^{-1} \\
\mathbf{b} &= \frac{(1-\alpha)}{\sigma^2} \mathbf{1}_N^T \sum_{t=1}^T (\mathbf{T}_t - \alpha \mathbf{T}_{t-1}) + \frac{\mu_0}{\sigma_0^2}
\end{aligned}$$

where $N = N_I + N_P = 33$ for the Hudson Valley PDSI

3.6 Full conditional for α

$$\begin{aligned}
[\alpha|\cdot] &\propto \prod_{t=1}^T [\mathbf{T}_t | \mathbf{T}_{t-1}, \mu, \alpha, \sigma^2, \phi] [\alpha] \\
&\propto \prod_{t=1}^T \exp \left\{ -\frac{1}{2} (\mathbf{T}_t - \alpha \mathbf{T}_{t-1} - (1-\alpha) \mu \mathbf{1}_N)^T \boldsymbol{\Sigma}_\epsilon^{-1} (\mathbf{T}_t - \alpha \mathbf{T}_{t-1} - (1-\alpha) \mu \mathbf{1}_N) \right\} I\{\alpha \in (0, 1)\} \\
&\propto \exp \left\{ -\frac{1}{2} \alpha^2 \sum_{t=1}^T (\mathbf{T}_{t-1} - \mu \mathbf{1}_N)^T \boldsymbol{\Sigma}_\epsilon^{-1} (\mathbf{T}_{t-1} - \mu \mathbf{1}_N) - \alpha \sum_{t=1}^T (\mathbf{T}_{t-1} - \mu \mathbf{1}_N)^T \boldsymbol{\Sigma}_\epsilon^{-1} (\mathbf{T}_t - \mu \mathbf{1}_N) \right\} \\
&\quad \times I\{\alpha \in (0, 1)\} \\
&\propto \exp \left\{ -\frac{1}{2} \alpha^2 \frac{1}{\sigma^2} \sum_{t=1}^T (\mathbf{T}_{t-1} - \mu \mathbf{1}_N)^T (\mathbf{T}_{t-1} - \mu \mathbf{1}_N) - \alpha \frac{1}{\sigma^2} \sum_{t=1}^T (\mathbf{T}_{t-1} - \mu \mathbf{1}_N)^T (\mathbf{T}_t - \mu \mathbf{1}_N) \right\} \\
&\quad \times I\{\alpha \in (0, 1)\}
\end{aligned}$$

which is truncated $N(\mathbf{A}^{-1}\mathbf{b}, \mathbf{A}^{-1})$, , restricted to $\alpha \in (0, 1)$, with

$$\mathbf{A}^{-1} = \left(\frac{1}{\sigma^2} \sum_{t=1}^T (\mathbf{T}_{t-1} - \mu \mathbf{1}_N)^T (\mathbf{T}_{t-1} - \mu \mathbf{1}_N) \right)^{-1}$$

$$\mathbf{b} = \frac{1}{\sigma^2} \sum_{t=1}^T (\mathbf{T}_{t-1} - \mu \mathbf{1}_N)^T (\mathbf{T}_t - \mu \mathbf{1}_N).$$

3.7 Full conditional for τ_I^2

$$\begin{aligned} [\tau_I^2 | \cdot] &\propto \prod_{t=1}^T [\mathbf{W}_t | \mathbf{T}_t, \beta_0, \beta_1, \tau_I^2, \tau_P^2] [\tau_I^2] \\ &\propto \prod_{t=1}^T |\boldsymbol{\Sigma}_t|^{-\frac{1}{2}} \exp \left\{ -\frac{1}{2} (\mathbf{W}_t - (\mathbf{H}_t \mathbf{T}_t + \mathbf{B}_t))^T \boldsymbol{\Sigma}_t^{-1} (\mathbf{W}_t - (\mathbf{H}_t \mathbf{T}_t + \mathbf{B}_t)) \right\} \\ &\quad \times (\tau_I^2)^{-\alpha_I - 1} \exp \left\{ \frac{\beta_I}{\tau_I^2} \right\} \\ &\propto (\tau_I^2)^{-\frac{1}{2} \sum_{t=1}^T N_{It}} \exp \left\{ -\frac{1}{2} \sum_{t=1}^T (\mathbf{W}_{It} - (\mathbf{H}_{It} \mathbf{T}_t + \mathbf{B}_{It}))^T \boldsymbol{\Sigma}_{It}^{-1} (\mathbf{W}_{It} - (\mathbf{H}_{It} \mathbf{T}_t + \mathbf{B}_{It})) \right\} \\ &\quad \times (\tau_I^2)^{-\alpha_I - 1} \exp \left\{ \frac{\beta_I}{\tau_I^2} \right\} \\ &\propto (\tau_I^2)^{-\alpha_I - \frac{M_I}{2} - 1} \exp \left\{ -\frac{1}{\tau_I^2} \left(\beta_I + \frac{1}{2} \sum_{t=1}^T (\mathbf{W}_{It} - (\mathbf{H}_{It} \mathbf{T}_t))^T (\mathbf{W}_{It} - (\mathbf{H}_{It} \mathbf{T}_t)) \right) \right\} \end{aligned}$$

which is $\text{IG}(\alpha_I + \frac{M_I}{2}, \beta_I + \frac{1}{2} \sum_{t=1}^T (\mathbf{W}_{It} - (\mathbf{H}_{It} \mathbf{T}_t))^T (\mathbf{W}_{It} - (\mathbf{H}_{It} \mathbf{T}_t)))$

3.8 Full conditional for τ_P^2

$$\begin{aligned}
[\tau_P^2 | \cdot] &\propto \prod_{t=1}^T [\mathbf{W}_t | \mathbf{T}_t, \beta_0, \beta_1, \tau_I^2, \tau_P^2] [\tau_P^2] \\
&\propto \prod_{t=1}^T |\Sigma_t|^{-\frac{1}{2}} \exp \left\{ -\frac{1}{2} (\mathbf{W}_t - (\mathbf{H}_t \mathbf{T}_t + \mathbf{B}_t))^T \Sigma_t^{-1} (\mathbf{W}_t - (\mathbf{H}_t \mathbf{T}_t + \mathbf{B}_t)) \right\} \\
&\quad \times (\tau_P^2)^{-\alpha_P - 1} \exp \left\{ \frac{\beta_P}{\tau_P^2} \right\} \\
&\propto (\tau_P^2)^{-\frac{1}{2} \sum_{t=1}^T N_{Pt}} \\
&\quad \times \exp \left\{ -\frac{1}{2} \sum_{t=1}^T (\mathbf{W}_{Pt} - (\beta_1 \mathbf{H}_{Pt} \mathbf{T}_t + \mathbf{B}_{Pt}))^T \Sigma_{Pt}^{-1} (\mathbf{W}_{Pt} - (\beta_1 \mathbf{H}_{Pt} \mathbf{T}_t + \mathbf{B}_{Pt})) \right\} \\
&\quad \times (\tau_P^2)^{-\alpha_P - 1} \exp \left\{ \frac{\beta_P}{\tau_P^2} \right\} \\
&\propto (\tau_P^2)^{-\alpha_P - \frac{M_P}{2} - 1} \\
&\quad \times \exp \left\{ -\frac{1}{\tau_P^2} \left(\beta_P + \frac{1}{2} \sum_{t=1}^T (\mathbf{W}_{Pt} - (\beta_1 \mathbf{H}_{Pt} \mathbf{T}_t + \beta_0 \mathbf{1}_{N_{Pt}}))^T (\mathbf{W}_{Pt} - (\beta_1 \mathbf{H}_{Pt} \mathbf{T}_t + \beta_0 \mathbf{1}_{N_{Pt}})) \right) \right\}
\end{aligned}$$

which is $\text{IG} \left(\alpha_P + \frac{M_P}{2}, \beta_P + \frac{1}{2} \sum_{t=1}^T (\mathbf{W}_{Pt} - (\beta_1 \mathbf{H}_{Pt} \mathbf{T}_t + \beta_0 \mathbf{1}_{N_{Pt}}))^T (\mathbf{W}_{Pt} - (\beta_1 \mathbf{H}_{Pt} \mathbf{T}_t + \beta_0 \mathbf{1}_{N_{Pt}})) \right)$

3.9 Full conditional for σ^2

$$\begin{aligned}
[\sigma^2 | \cdot] &\propto \prod_{t=1}^T [\mathbf{T}_t | \mathbf{T}_{t-1}, \mu, \alpha, \sigma^2, \phi] [\sigma^2] \\
&\propto \prod_{t=1}^T |\Sigma_\epsilon|^{-\frac{1}{2}} \exp \left\{ -\frac{1}{2} (\mathbf{T}_t - \alpha \mathbf{T}_{t-1} - (1-\alpha) \mu \mathbf{1})^T \Sigma_\epsilon^{-1} (\mathbf{T}_t - \alpha \mathbf{T}_{t-1} - (1-\alpha) \mu \mathbf{1}) \right\} \\
&\quad \times (\sigma^2)^{-\alpha_{\sigma^2}-1} \exp \left\{ -\frac{\beta_{\sigma^2}}{\sigma^2} \right\} \\
&\propto (\sigma^2)^{-\frac{NT}{2}} \exp \left\{ -\frac{1}{\sigma^2} \left(\frac{1}{2} \sum_{t=1}^T (\mathbf{T}_t - \alpha \mathbf{T}_{t-1} - (1-\alpha) \mu \mathbf{1})^T (\mathbf{T}_t - \alpha \mathbf{T}_{t-1} - (1-\alpha) \mu \mathbf{1}) \right) \right\} \\
&\quad \times (\sigma^2)^{-\alpha_{\sigma^2}-1} \exp \left\{ -\frac{\beta_{\sigma^2}}{\sigma^2} \right\} \\
&\propto (\sigma^2)^{-\alpha_{\sigma^2} - \frac{NT}{2} - 1} \\
&\quad \times \exp \left\{ -\frac{1}{\sigma^2} \left(\beta_{\sigma^2} + \frac{1}{2} \sum_{t=1}^T (\mathbf{T}_t - \alpha \mathbf{T}_{t-1} - (1-\alpha) \mu \mathbf{1})^T (\mathbf{T}_t - \alpha \mathbf{T}_{t-1} - (1-\alpha) \mu \mathbf{1}) \right) \right\}
\end{aligned}$$

which is $\text{IG} \left(\alpha_{\sigma^2} + \frac{NT}{2}, \beta_{\sigma^2} + \frac{1}{2} \sum_{t=1}^T (\mathbf{T}_t - \alpha \mathbf{T}_{t-1} - (1-\alpha) \mu \mathbf{1})^T (\mathbf{T}_t - \alpha \mathbf{T}_{t-1} - (1-\alpha) \mu \mathbf{1}) \right)$

4 Extensions to the model

- Predictive Process for Σ_ϵ in the spatial model
- Include the number of trees used to create a chronology in the model
- Maybe we need to model a covariance between the proxy chronologies instead of a spatial covariance? e.g. $\mathbf{R}(\phi)$ is not a spatial covariance but a covariance between proxy records at a given site
- ???